



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Extreme parameter sensitivity in quasidilaton massive gravity

Stefano Anselmi, Diana López Nacir, and Glenn D. Starkman

Phys. Rev. D **92**, 084033 — Published 15 October 2015

DOI: [10.1103/PhysRevD.92.084033](https://doi.org/10.1103/PhysRevD.92.084033)

Extreme parameter sensitivity in quasidilaton massive gravity

Stefano Anselmi,^{1,*} Diana López Nacir,^{2,3,†} and Glenn D. Starkman^{1,‡}

¹*Department of Physics, Case Western Reserve University, Cleveland, OH 44106-7079 – USA*

²*The Abdus Salam International Center for Theoretical Physics, Strada Costiera 11, I-34151 Trieste – Italy*

³*Departamento de Física and IFIBA, FCEyN UBA, Facultad de Ciencias Exactas y Naturales, Ciudad Universitaria, Pabellón I, 1428 Buenos Aires – Argentina*

We reanalyze the behavior of Friedmann-Lemaître-Robertson-Walker cosmologies in the recently proposed quasidilaton massive-gravity model, and discover that the background dynamics present hitherto unreported features that require unexpected fine-tuning of the additional fundamental parameters of the theory for an observationally consistent background cosmology. We also identify new allowed regions in the parameters space and exclude some of the previously considered ones. The evolution of the mass of gravitational waves reveals non-trivial behavior, exhibiting a mass-squared that may be negative in the past, and that presently, while positive, is larger than the square of the Hubble parameter, H_0^2 . These properties of the gravity-wave mass have the potential to lead to observational tests of the theory. While quasidilaton massive gravity is known to have issues with stability at short distances, the current analysis is a first step toward the investigation of the more stable extended quasidilaton massive gravity theory, with some expectation that both the fine-tuning of parameters and the interesting behavior of the gravity-wave mass will persist.

Keywords: Cosmology, modified theories of gravity.

I. INTRODUCTION

The standard cosmological model, Λ CDM, describes the acceleration of the universe by properly adjusting the cosmological constant Λ . While this simple model is consistent with current observational data, other models provide alternative explanations of this acceleration. For example, some models attribute the acceleration to the presence of a dynamical component known as dark energy [1–3], and others to a modification of the gravitational laws on cosmological distances [4–8]. The questions will be to what extent it is possible to discriminate among the different models from observations, and whether any of the models are better at fitting the data than what is currently the most parsimonious explanation, Λ .

The next generation of experiments (such as EUCLID [9] or DESI [10]) will provide an unprecedented amount of observational data. However, there is now a wide range of candidate theories. For instance, different modifications of general relativity primarily in the infrared have been considered by many authors (see [11] for a recent review), and probably still others have yet to be proposed. Ultimately, the predictions of each candidate model must be confronted with data. This includes not just cosmological data but data on all scales where the models make calculable predictions that can be tested observationally or experimentally.

Within one interesting class of theories, the current acceleration era is associated to the presence of a mass term for the graviton (for a historical overview, motivations and an updated description of different proposed

massive gravity theories, see [12, 13]). Here we consider a particular modification of general relativity known as quasidilaton massive gravity (QDMG), which we summarize in the Section II. This theory was proposed in [14], as an extension of the dRGT theory of massive gravity [15, 16], and contains an additional scalar degree of freedom: the quasidilaton. The main motivation for such extension is the absence of isotropic and homogeneous cosmological background solutions in dRGT [17]. Indeed, it has been shown that QDMG has solutions with spatially flat Friedmann-Lemaître-Robertson-Walker (FLRW) background metrics [14]. Moreover, it has been found that (even in the absence of a cosmological constant) there are solutions for which at late times the metric approaches to a de Sitter metric, providing a plausible (self-accelerating) explanation of the accelerated expansion of the universe [14, 18, 19]. The quasidilaton theory has three parameters more than Λ CDM.

In this paper we will perform a careful analysis of the background cosmological evolution, taking into account the main goal of describing the observed expansion history of the universe. While other authors have made preliminary investigations [18, 19] of the background evolution in QDMG, a more detailed reexamination reveals important new insights. The allowed set of parameters split into two disconnected regions characterized by “low” and “high” values of a dimensionless parameter of the theory, ω (which multiplies the kinetic term of the quasidilaton). In the region with low values of ω , while viable background solutions exist for a wide range of values of the Lagrangian parameter nominally called the graviton mass m_g , with $m_g \sim \mathcal{O}(H_0)$, a careful fine-tuning of the dimensionless constants α_3 and α_4 is required. The permitted values of α_3 , α_4 and m_g thus describe a very thin 2-dim surface $m_g(\alpha_3, \alpha_4)$ in the $\{\alpha_3, \alpha_4, m_g\}$ parameter space. In the other region, the parameter m_g is constrained to be much smaller than H_0 , and the larger it

* stefano.anselmi@case.edu

† dlopez_n@ictp.it

‡ glenn.starkman@case.edu

is, the narrower the 2-dim surface of allowed α_3 and α_4 .

The paper is organized as follows. After summarizing the theory QDMG in Section II, in Section III we present the dynamical equations. In Section IV we analyze the existence of viable de Sitter fixed-point attractors. By exploring the 4-dimensional parameter space of the theory, in Section V, we assess the viability of a self-accelerating explanation of the current expansion of the universe. An important outcome of our analysis is that, in order to reproduce an expansion history consistent with data, the graviton mass parameter must also be fine-tuned to a value that depends on other parameters of the model.

In Section VI we study the evolution of the mass of gravitational waves M_{GW} for the allowed set of parameters. We find the current value of M_{GW} to be generically larger than the current Hubble constant H_0 even when we set the graviton mass parameter $m_g \ll H_0$. In the past (for example at redshifts relevant for the Cosmic Microwave Background) M_{GW} can be either real or imaginary. For a conservative choice of $0 < m_g \leq H_0$, $|M_{GW}(t)| < H(t)$ in the past, with $|M_{GW}| \lesssim 10^{-2}H$ at last scattering. While this precludes the development of a catastrophic instability when M_{GW} is imaginary, nevertheless potentially there could be observable cosmological signatures. These merit further investigation [20–22].

II. THEORY OF QUASIDILATON MASSIVE GRAVITY

We consider the action for the quasidilaton theory [14]:

$$\begin{aligned} S &= S_{EH} + S_\sigma \\ &= \frac{M_{\text{Pl}}^2}{2} \int d^4x \sqrt{-g} \left[R - \frac{\omega}{M_{\text{Pl}}^2} \partial_\mu \sigma \partial^\mu \sigma \right. \\ &\quad \left. + 2m_g^2 (\mathcal{L}_2 + \alpha_3 \mathcal{L}_3 + \alpha_4 \mathcal{L}_4) \right], \end{aligned} \quad (1)$$

where M_{Pl} is the Planck mass and, in addition to the Einstein-Hilbert action S_{EH} , a contribution S_σ characterizes the quasidilaton scalar field σ . In addition to the quasidilaton kinetic term, S_σ includes three interaction terms: Here

$$\mathcal{L}_2 \equiv \frac{1}{2} ([\mathcal{K}]^2 - [\mathcal{K}^2]), \quad (2)$$

$$\mathcal{L}_3 \equiv \frac{1}{6} ([\mathcal{K}]^3 - 3[\mathcal{K}][\mathcal{K}^2] + 2[\mathcal{K}^3]), \quad (3)$$

$$\begin{aligned} \mathcal{L}_4 &\equiv \frac{1}{24} ([\mathcal{K}]^4 - 6[\mathcal{K}]^2[\mathcal{K}^2] + 3[\mathcal{K}^2]^2 + \\ &\quad + 8[\mathcal{K}][\mathcal{K}^3] - 6[\mathcal{K}^4]), \end{aligned} \quad (4)$$

with square brackets denoting a trace, and

$$\mathcal{K}^\mu_\nu \equiv \delta^\mu_\nu - e^{\sigma/M_{\text{Pl}}} \left(\sqrt{g^{-1}f} \right)^\mu_\nu. \quad (5)$$

The non-dynamical “fiducial metric” is built from four Stückelberg fields ϕ^a ($a = 0, \dots, 3$),

$$f_{\mu\nu} \equiv \eta_{ab} \partial_\mu \phi^a \partial_\nu \phi^b. \quad (6)$$

In the space of Stückelberg fields, the theory enjoys the Poincare symmetry [14]

$$\phi^a \rightarrow \phi^a + c^a, \quad \phi^a \rightarrow \Lambda_b^a \phi^b, \quad (7)$$

and in addition, there is a global symmetry given by

$$\sigma \rightarrow \sigma + \sigma_0, \quad \phi^a \rightarrow e^{-\sigma_0/M_{\text{Pl}}} \phi^a, \quad (8)$$

with σ_0 an arbitrary constant.

The addition of S_σ to the action, introduces four new parameters: the dimensionless kinetic coupling ω , the graviton mass parameter m_g , and the coupling constants α_3 and α_4 . As shown below, the cosmological solution depends sensitively on the values of these parameters.

III. THE BACKGROUND COSMOLOGICAL EQUATIONS

We consider a spatially flat Friedmann-Lemaître-Robertson-Walker ansatz, for which

$$ds^2 = -N(t)^2 dt^2 + a(t)^2 \delta_{ij} dx^i dx^j, \quad (9)$$

$$\phi^0 = \phi^0(t), \quad (10)$$

$$\phi^i = x^i, \quad (11)$$

$$\sigma = \bar{\sigma}(t). \quad (12)$$

The fiducial metric $f_{\mu\nu}$ reduces to

$$f_{00} = -n(t)^2, \quad f_{ij} = \delta_{ij}, \quad (13)$$

where

$$n(t)^2 \equiv (\dot{\phi}^0)^2. \quad (14)$$

The minisuperspace action for the background metric and fields can now be written as

$$\begin{aligned} S/V &= M_{\text{Pl}}^2 \int dt \left\{ -3 \frac{a^3}{N} \left(\frac{\dot{a}}{a} \right)^2 + a^3 \frac{w}{M_{\text{Pl}}} \frac{\dot{\sigma}^2}{2N} \right. \\ &\quad \left. + N a^3 m_g^2 (\mathcal{L}_2 + \alpha_3 \mathcal{L}_3 + \alpha_4 \mathcal{L}_4) \right\}, \end{aligned} \quad (15)$$

where

$$\mathcal{L}_2 = 3(X - 1)(-2 + X(1 + r)), \quad (16)$$

$$\mathcal{L}_3 = -(X - 1)^2(-4 + X(1 + 3r)), \quad (17)$$

$$\mathcal{L}_4 = (X - 1)^3(-1 + rX), \quad (18)$$

and we have defined

$$X \equiv \frac{e^{\bar{\sigma}/M_{\text{Pl}}}}{a}, \quad (19)$$

$$r \equiv \frac{n}{N} a. \quad (20)$$

Varying the action with respect to $\phi^0(t)$ leads to

$$\partial_t \left[\frac{\dot{\phi}^0}{n} a^4 G_2(X) \right] = \partial_t [a^4 G_2(X)] = 0, \quad (21)$$

where $G_2(X) = X(1-X)J(X)$, with

$$J(X) \equiv 3 + 3(1-X)\alpha_3 + (1-X)^2\alpha_4. \quad (22)$$

We use time reparameterization freedom to set $N = 1$.

In summary, the independent background equations are:

- the constraint equation (21), or its integral

$$G_2(X) = \frac{C}{a^4}. \quad (23)$$

- the Friedman equation,

$$3H^2 = \frac{\omega}{2} \left(\frac{\dot{\sigma}}{M_{\text{Pl}}} \right)^2 + 3m_g^2 G_1(X) + \frac{\rho_m + \rho_r}{M_{\text{Pl}}^2}, \quad (24)$$

where

$$G_1(X) \equiv \frac{X-1}{3} [\alpha_3(X-1)^2 - 3(X-1) + J(X)] \quad (25)$$

and we have included the contributions of matter and radiation;

- the conservation of the stress-energy tensor obtained from S_σ

$$(\ddot{\sigma} + 3H\dot{\sigma})\omega\dot{\sigma} + 3M_{\text{Pl}}m_g^2(\dot{\sigma} - rHM_{\text{Pl}})XG_1'(X) = 0, \quad (26)$$

where a prime means derivative with respect to X ;

- the conservation of the stress-energy tensors of matter, $\dot{\rho}_m = -3H\rho_m$ and of radiation $\dot{\rho}_r = -4H\rho_r$.

(Note that using the constraint equation (21) one can show that the equation obtained by taking the variation of S_σ with respect to σ is not an independent equation.)

IV. DE SITTER FIXED POINT ANALYSIS

We start by investigating the future background evolution of the quasidilaton massive gravity model. The Λ CDM concordance model predicts the universe will approach a de Sitter phase in the future. Though we do not know the future of the universe, we require our model to reproduce this prediction, consistent with recent practice [18, 19].

We rewrite eq. (24) in terms of the relative energy densities

$$1 = \Omega_{DE} + \Omega_m + \Omega_r, \quad (27)$$

where

$$\Omega_{DE} = \Omega_\Lambda + \Omega_\sigma, \quad (28)$$

and

$$\Omega_m = \frac{\rho_m}{3M_{\text{Pl}}^2 H^2}, \quad (29)$$

$$\Omega_r = \frac{\rho_r}{3M_{\text{Pl}}^2 H^2}, \quad (30)$$

$$\Omega_\Lambda = \frac{m_g^2}{H^2} G_1(X), \quad (31)$$

$$\Omega_\sigma = \frac{\omega}{6H^2} \left(\frac{\dot{\sigma}}{M_{\text{Pl}}} \right)^2. \quad (32)$$

Employing eq. (21) and assuming that $X \neq 0$ we obtain

$$\dot{\sigma} = M_{\text{Pl}}H \left(1 - \frac{4G_2(X)}{XG_2'(X)} \right). \quad (33)$$

Eq. (23) implies that as $a \rightarrow \infty$, $X \rightarrow \text{constant}$. Therefore the set of variables $\{\Omega_m, \Omega_r, X, \Omega_\Lambda\}$ will approach constants in the asymptotic future. We study the dynamical stability of the system by means of the following equations:

$$\frac{d\Omega_r}{dN} = -2\Omega_r \left(2 + \frac{\dot{H}}{H^2} \right), \quad (34)$$

$$\frac{d\Omega_\Lambda}{dN} = -2\Omega_\Lambda \left(2 \frac{G_2 G_1'}{G_1 G_2'} + \frac{\dot{H}}{H^2} \right), \quad (35)$$

$$\frac{dX}{dN} = -4 \frac{G_2}{G_2'}, \quad (36)$$

where $N = \ln a$ is the number of e-foldings. $\frac{\dot{H}}{H^2}$ can be obtained by differentiating the first Friedmann equation (24) to obtain

$$\frac{\dot{H}}{H^2} = \frac{9\Omega_m + 12\Omega_r + 12\frac{G_2}{G_2'} \left[\frac{G_1'}{G_1} \Omega_\Lambda + \frac{\omega}{6} \frac{d}{dX} \left(1 - 4 \frac{G_2}{XG_2'} \right)^2 \right]}{\omega \left[1 - 4 \frac{G_2}{XG_2'} \right]^2 - 6}. \quad (37)$$

As noted above, we focus on de Sitter fixed points, and require that these critical points are attractors. The de Sitter critical points relative to the system (34), (35) and (36) are given in Table I, where

$$X_\pm \equiv 1 + \frac{3\alpha_3}{2\alpha_4} \pm \sqrt{\frac{9\alpha_3^2}{4\alpha_4^2} - \frac{3}{\alpha_4}}. \quad (38)$$

To assess the stability we compute the matrix form of the perturbation equations linearized around each of the fixed points. Then, the linear asymptotic stability of each fixed point can be studied by analyzing the signs of the eigenvalues of that matrix. If the sign of the real part of every eigenvalues is negative, then the critical point is an attractor. The results are shown in the last column of Table I.

This analysis indicates there are three possible late-time de Sitter fixed points, A , B and C . For each, constraints on the parameters α_3 and α_4 are obtained by

F. P.	Ω_r	Ω_Λ	X	Ω_m	Ω_σ	Existence	Stability	Eigenvalues
A	0	$1 - \omega/6$	X_+	0	$\omega/6$	$0 < \omega < 6, 0 \leq X_+, X_+ \in \mathcal{R}$	Attractor	-4,-4,-3
B	0	$1 - \omega/6$	X_-	0	$\omega/6$	$0 < \omega < 6, 0 \leq X_-, X_- \in \mathcal{R}$	Attractor	-4,-4,-3
C	0	$1 - 3\omega/2$	0	0	$3\omega/2$	$0 < \omega < 2/3$	Attractor	-4,-4,-3

TABLE I. De Sitter fixed points.




FIG. 1. The blue line shows the $G_2(X)$ function, while the red points are the $G_2(X_0) = C$ values for $\{m_g = 0.4H_0, \omega = 0.01, \alpha_3 = 0.75, \alpha_4 = 0.345\}$.

requiring that $H^2 > 0$ and $X \geq 0$. For the point A we obtain $\alpha_3 > 0$ and $0 < \alpha_4 < 2\alpha_3^2/3$; for B , $\alpha_3 < -3$ and $-3 - 3\alpha_3 \leq \alpha_4 \leq 2\alpha_3^2/3$; for C , $\alpha_4 < -6 - 4\alpha_3$. Noticing that one must insist that $X \geq 0$, we found different constraints than [18, 19].

Consider more closely the fixed point B . Given the α_3 and α_4 constraints for B , we obtain $0 < X_- < X_+ < 1$. The constraint equation (23) implies that in the asymptotic future $G_2(X) = 0$. Moreover (23) requires that G_2 should be unbounded either above or below in order to have a past history. G_2 is a polynomial in X , so this is impossible if $0 < X_- < 1$ [23] as it is. Thus B cannot be a well-defined fixed point. Recalling that $X \geq 0$, a similar argument can be applied to the point C .

The allowed $\{\alpha_3, \alpha_4\}$ parameter region for A entails that $1 < X_- < X_+$. Therefore A is the only well-defined de Sitter fixed point for the QDMG theory. We emphasize that our findings now differ from those of [18, 19], in that we exclude the points B, C .

V. COSMOLOGICAL EVOLUTION AND PARAMETER FIXING

The aim of this section is to study the evolution of the relevant background quantities in agreement with the results of the previous section and with the observed cosmological history. That depends on the initial conditions,

on the expansion history and on the fixed point A . By means of this analysis we constrain the four parameters of quasidilaton massive gravity: $\{m_g, \omega, \alpha_3, \alpha_4\}$.

Given that we are dealing with the background energy density evolution, we can consider neutrinos to be relativistic, since the value of Ω_r is not negligible only in the radiation era when neutrinos were indeed relativistic. The spectrum of the CMB today is precisely measured, so we accurately determine $\Omega_{\gamma,0}$. For relativistic neutrinos, $\Omega_{\nu,0}$ is proportional to $\Omega_{\gamma,0}$. Therefore we assume $\Omega_{r,0}$ is known and we fix it by $\Omega_{r,0} = \Omega_{\gamma,0} + \Omega_{\nu,0} = 0.0000851$.

To be definite we also fix $\Omega_{DE,0} = 0.72$, close to the best fit value [24]. That corresponds in Λ CDM to $z_{eq} \approx 3300$. We stress that this choice will not qualitatively affect our conclusions on the quasidilaton massive gravity background evolution.

To fix the initial conditions we require $\Omega_{DE,0} = \Omega_{\sigma,0} + \Omega_{\Lambda,0}$, where $\Omega_{\sigma,0}$ and $\Omega_{\Lambda,0}$ are given by (31), (32). In this way we obtain a 9-th order polynomial that has no analytical solutions. In Fig. 1 we plot the $G_2(X)$ function for $\{m_g = 0.4H_0, \omega = 0.01, \alpha_3 = 0.75, \alpha_4 = 0.345\}$. The red points represent the values of X_0 and $G_2(X_0)$ where the initial conditions are satisfied. From eq. (23) we obtain $C = G_2(X_0)$. Finally, notice that G_2 is unbounded as $X \rightarrow \infty$; this implies that for our model the correct past evolution of the background is allowed only if $X_0 > X_+$.

The dark energy equation-of-state parameter $w_{DE,0}$ is constrained by observations. To compute w for the QDMG model we first define the total effective equation-of-state parameter

$$w_{eff} = -1 - \frac{2}{3} \frac{\dot{H}}{H^2}, \quad (39)$$

and consequently

$$w_{DE} = \frac{w_{eff} - w_m \Omega_m - w_r \Omega_r}{\Omega_\Lambda + \Omega_\sigma}. \quad (40)$$

We must require that $-1.2 < w_{DE,0} < -0.9$ in agreement with the current limits [25].

The quasidilaton massive gravity model shows a particular feature – Ω_{DE} scales as matter at early times [18]. Indeed, from the analysis above we have $\alpha_4 > 0$ and $C > 0$. At early times $G_2 \sim \alpha_4 X^4 = C/a^4$. Therefore we find [26]

$$\Omega_{DE} H^2 \simeq \Omega_\Lambda H^2 \simeq m_g^2 \left(\frac{C}{\alpha_4} \right)^{3/4} \frac{\alpha_3 + \alpha_4}{3} a^{-3}. \quad (41)$$




FIG. 2. (REG1): $m_g^2/H_0^2 - \omega$ constraints after marginalizing over $\alpha_3 - \alpha_4$.




FIG. 4. (REG2): $m_g^2/H_0^2 - \omega$ constraints after marginalizing over $\alpha_3 - \alpha_4$.




FIG. 3. (REG1): $\alpha_3 - \alpha_4$ constraints for different m_g^2/H_0^2 values (*black bold numbers*). We marginalized over ω . The blue line corresponds to the boundary of the region $\alpha_4 < 2\alpha_3^2/3$, which is the existence condition we obtained from the fixed point analysis.

It follows that, for redshifts $z \gtrsim 10$, Ω_{DE} would contribute to the effective matter energy density! Therefore Ω_{DE} should be negligible in the radiation era in order to have a viable expansion history. We demand that $\Omega_{DE}(z_{eq}) < 0.01$.

In order to identify the allowed ranges of the four parameters of QDMG, the main computational obstacle is to find the solutions of the initial condition, namely $\Omega_{DE,0} = 0.72$. In principle $\Omega_{DE,0} = 0.72$ could have from 1 to 9 allowed solutions for each value of the QDMG parameters. However, after enforcing all the observational conditions, we find that there is never more than 1 viable solution. We identify two disconnected allowed regions

in the 4-dimensional space of parameters, one shows just low- ω values (hereafter REG1) and the other one high- ω values (hereafter REG2).

In Fig. 2 and 3 we present the constraints for the (REG1) parameter space. After marginalizing over $\alpha_3 - \alpha_4$ we find that ω is constrained to $0 < \omega \lesssim 1.2$ as we report in Fig. 2. On the other hand, marginalizing over ω , the contour plot reported in Fig. 3 shows that the $\alpha_3 - \alpha_4$ values are tightly related to the m_g^2/H_0^2 value (*black bold numbers*). Once we know two of the three $\{m_g, \alpha_3, \alpha_4\}$ parameters, the other one is determined to a good approximation. In other words the quasidilaton massive gravity theory presents a fine-tuning of the parameters.

Repeating the procedure for (REG2) we find a different behavior. In Fig. 4 we see that $m_g^2/H_0^2 \lesssim 0.008$ while the ω allowed interval depends on m_g and it becomes larger as m_g decreases. The $\alpha_3 - \alpha_4$ region is again m_g dependent, however the dependence now is different than for (REG1) as we show in Fig. 5. For (REG2) if $m_g^2/H_0^2 \sim 0.1$ then $\omega \sim 6$ and α_4 becomes effectively a function of α_3 , so we find a fine-tuning of 2 parameters. On the other hand if the graviton mass is small, i.e. $m_g^2/H_0^2 \lesssim 0.001$, the other parameters are no longer strongly constrained.

Notice that we find different results than [18, 19]. In particular they allowed ω to be negative and they obtain $\omega \lesssim 0.3$.

In our analysis we did not compute the whole expansion history for each point in the four dimensional parameter space for practical computational reasons. As an illustrative example, we choose two set of allowed parameters for (REG1) and (REG2) and we plot the evolution of the energy densities in Fig. 6. As expected, the two panels, are consistent with the observed expansion history.

The parameter fine-tuning we found practically reduces from four to three the effective parameters of the




FIG. 5. (REG2): $\alpha_3 - \alpha_4$ constraints for different m_g^2/H_0^2 values (*black bold numbers*). We marginalized over ω . The blue line corresponds to the boundary of the region $\alpha_4 < 2\alpha_3^2/3$, which is the existence condition we obtained from the fixed point analysis.

quasidilaton massive gravity theory. We expect that studying the perturbations will further constrain the theory. Some of those perturbations will be unstable.

VI. GRAVITY WAVES

In this section we focus on the evolution of the mass of the gravitational waves. We consider tensor perturbations around the background metric solutions,

$$\delta g_{ij} = a^2 h_{ij}^{TT} = a^2 \int \frac{d^3 k}{(2\pi)^{3/2}} h_{ij,\vec{k}}^{TT} \exp(i\vec{k} \cdot \vec{x}) + cc, \quad (42)$$

with $\delta^{ij} h_{ij}^{TT} = 0$, and $\partial^j h_{ij}^{TT} = 0$. After a straightforward calculation, one gets the quadratic Lagrangian for $h_{ij,\vec{k}}^{TT}$

$$\mathcal{L}_{GW} = \frac{M_{\text{Pl}}^2}{8} a^3 \left[|\dot{h}_{ij}^{TT}|^2 - \left(\frac{k^2}{a^2} + M_{GW}^2 \right) |h_{ij,\vec{k}}^{TT}|^2 \right], \quad (43)$$

where the mass of the gravitational waves M_{GW} is given by [27]

$$M_{GW}^2 = m_g^2 X (3 + 3\alpha_3 + \alpha_4 - (1 + 2\alpha_3 + \alpha_4)(1 + r)X + (\alpha_3 + \alpha_4)rX^2). \quad (44)$$

We start by computing the ratio M_{GW}^2/H^2 at redshift $z = 1100$, relevant for CMB, for the two disconnected regions (REG1) and (REG2) defined in the previous section. An exploration of the values computed reveals there is a minimum and maximum M_{GW}^2/H^2 for each of the regions. The results are presented in Table II. We see that a real mass as large as $M_{GW} \sim 10^{-2}H$ can be obtained, even for our conservative choice $m_g \leq H_0$. For both parameter regions (REG1) and (REG2), we note the mass can be imaginary. However, the maximum absolute values turn out to be much smaller than the Hubble rate, preventing the development of a full instability. It is worth noting that so far signatures in the Cosmic Microwave Background (CMB) due to a non-vanishing M_{GW} have been studied assuming this mass is always real [20–22]. Our results suggest that one should explore




FIG. 6. *Upper panel:* expansion history for $\{m_g^2/H_0^2 = 0.05, \omega = 0.9484, \alpha_3 = 7.059, \alpha_4 = 10.63\}$. *Lower panel:* expansion history for $\{m_g^2/H_0^2 = 0.0005, \omega = 5.4211, \alpha_3 = 10.797, \alpha_4 = 17.680\}$

also the possibility of having cosmological gravitational waves with a small but imaginary mass at the relevant redshifts for CMB.

In Fig. 7 we plotted the evolution of the ratio M_{GW}^2/H^2 for the parameters given in Table II. We notice that at low redshifts (and in particular at $z = 0$) the mass becomes positive, and is larger than H_0 , despite the fact that $m_g \leq H_0$. In order to assess the generality of this result we computed the ratio M_{GW}^2/H_0^2 at $z = 0$ varying the parameters in the two disconnected allowed regions of the 4-dimensional space, and we obtained its maximum and minimum value. The results are shown in Table III. We see that M_{GW} is larger than H_0 , even for values of $m_g \sim 10^{-2}H_0$, and it is up to a factor of ~ 5 larger than H_0 for $m_g \leq H_0$. The existence of a minimum value of M_{GW} that is larger than H_0 is remarkable, since this represents a motivated observational threshold. That is, if one could constrain M_{GW} to be smaller than H_0 one would be able to rule out a self-accelerating explanation of the current acceleration of the universe within the QDMG theory. It would be interesting to see whether an analogous result holds for other theories that

also aim to provide a self-accelerating explanation. Unfortunately, current experiments are still far from probing $M_{GW} \sim H_0$ [28]. Moreover, the upper limits one can obtain are in general model dependent, since they are based on assumptions involving different scales of the theory. This represents a challenge for both theory and observations, and highlights the ongoing importance of working out predictions within the framework of specific models of modified gravity.

$z = 1100$	(REG1)		(REG2)	
	Max.	Min.	Max.	Min.
M_{GW}^2/H^2	2.5×10^{-4}	-9.2×10^{-5}	1.5×10^{-5}	-8.5×10^{-6}
m_g^2/H_0^2	1	1	10^{-4}	8.45×10^{-3}
ω	1.03	0.32	5.99	5.95
α_3	6.60	6.80	5.83	3.56
α_4	19.61	19.87	20	8.44
X_0	1.82	1.85	1.73	1.94

TABLE II. Maximum and minimum values of M_{GW}^2/H^2 at $z = 1100$ and the corresponding values of the parameters $\{m_g, \omega, \alpha_3, \alpha_4\}$. For completeness, the value of X at $z = 0$ is also presented.




FIG. 7. Evolution of the mass-squared of the gravitational waves in units of the Hubble rate for the set of parameters given in Table II. Solid (Dashed) lines correspond to the values of parameters for which we found the maximum (minimum) value of M_{GW}^2/H^2 at $z = 1100$.

VII. CONCLUSIONS

The combination of General Relativity and the Standard Model of particle physics are demonstrably and remarkably successful descriptions of the world on scales up to and including the solar system. On larger scales, there is a need either to modify the theory of gravity

$z = 0$	(REG1)		(REG2)	
	Max.	Min.	Max.	Min.
M_{GW}^2/H_0^2	25	2.5	18	5
m_g^2/H_0^2	1	10^{-4}	10^{-4}	10^{-4}
ω	0.08	0.08	6	5.05
α_3	0.86	3.46	5.83	1.70
α_4	0.48	0.55	20	0.49
X_0	4.67	19.59	1.73	10.99

TABLE III. Maximum and minimum values of M_{GW}^2/H_0^2 at $z = 0$, the corresponding values of the parameters $\{m_g, \omega, \alpha_3, \alpha_4\}$, and the value of X at $z = 0$.

or to introduce new forms of dark matter and dark energy. The most parsimonious solution would be to identify candidates for the latter in the Standard Model, and such candidates may exist for dark matter (see for example [29, 30]) and evade existing constraints [31], although the phenomenological successes of MOND (see for example [32]) cannot be entirely dismissed as an indication of the need to modify gravity on galactic scales. For the observed cosmic acceleration, the situation is even less clear. A cosmological constant is the canonical explanation, but despite decades of attempts has as yet no clear explanation in the Standard Model. The need for observational probes of possible dark energy and modified gravity explanations is thus paramount.

One possibility would be to develop some general phenomenological classification of possible deviations of gravity from GR. The Parametrized post-Newtonian approach is one such program, in the context of almost-Schwarzschild backgrounds. Such generic approaches have also been attempted in the cosmological context (eg. [33, 34]). However, in the context of a highly non-linear theory such as GR, the observational consequences of small theoretical departures from GR can be quite idiosyncratic. While phenomenological parametrization of observables may be convenient, and even useful, they may not capture (or may capture poorly) the specific phenomena or behaviour that result from actual models. Careful examination of specific individual models can therefore be both instructive and essential.

In this paper, we have studied the (homogeneous) cosmological solutions of quasidilaton massive gravity. A study of the linear perturbations around the asymptotic self-accelerated cosmological solution of this theory (which corresponds to a De Sitter background metric) has been done in [35, 36]. These studies have revealed that the kinetic term of one of the scalar perturbations becomes negative for short wavelengths, indicating that the theory may have a ghost instability that shows up at short distances. This is indeed the case at linear level. Several authors [37, 38] have therefore extended the theory by allowing for a new coupling, which can be properly adjusted to make the scalar sector stable at linear level. This extended quasidilaton massive gravity theory

(EQDMG), has been considered by other authors [39–41].

Although this current reconsideration of the background cosmological solutions of QDMG was performed as a first step for a full analysis of the EQDMG, it revealed important attributes of the QDMG cosmology, which we expect to carry over qualitatively or in detail to EQDMG. The first is that observationally viable QDMG cosmologies require fine-tuning of parameters. In particular, the allowed values of the graviton mass parameter, m_g is a tightly constrained function of the coupling constants α_3 and α_4 , with only a very narrow tolerance around a central value $m_g(\alpha_3, \alpha_4)$. This fine-tuning, and the precise value of $m_g(\alpha_3, \alpha_4)$, is dictated by observational constraints on the dark energy properties [42].

The second observation is that some small (but possibly non-negligible) fraction of what manifests as Ω_{DE} (i.e. $p/\rho \simeq -1$) today, was Ω_m (i.e. $p/\rho \simeq 0$) in the past. The transition from one equation of state to the other was sudden and probably not well-captured by a linear parametrization of $w(z)$. The expected difference between Ω_m at high redshift (as measured in the CMB) and at low redshift (as measured, say in large scale structure), could be the source of recently noted tensions in different determinations of Ω_m [43, 44]. While the details of these behaviors of the background cosmology are likely to be altered in EQDMG, it is plausible that these qualitative features are robust.

We have also analyzed the phenomenology of the graviton. The governing equations for the graviton mass M_{GW} (which is not equal to the graviton mass parameter m_g) are the same in QDMG and EQDMG. We therefore expect to gain useful insights for the extended model provided that the background solutions do not depend sensitively on the new parameter α_σ of the extended theory. We find that the graviton mass-squared typically is negative at redshifts well above $z = 1$, indicating an instability. This includes redshifts $z \simeq 10^3 - 10^4$ where such physics may well imprint itself on the CMB. At any given time $|M_{GW}^2| \ll H^2$, so we do not expect the instability to lead to many e-foldings of growth. Nevertheless, if this persists in EQDMG, it may be another opportunity to see evidence of modified gravity in CMB observations.

Regarding vector perturbations, according to equations (4.16) and (4.17) of [41], the square of the speed of propagation, c_V^2 , can be recast as $c_V^2 = \kappa_V/M_{GW}^2$, where the absence of ghost instability is guaranteed provided $\kappa_V > 0$ [45]. Notice in particular that when M_{GW}^2 becomes negative, the absence of ghost instability implies c_V^2 becomes also negative. Therefore, we expect that a detail analysis of the perturbations will further reduce the region of allowed parameters.

In a future work, we will therefore extend our analysis to the EQDMG theory, taking into account the constraints from the study of the perturbations, anticipating hopefully that these observable effects will indeed persist.

ACKNOWLEDGMENTS

We thank C. de Rham for helpful comments, and G. Gabadadze, M. Fasiello, D. Müller and A. Tolley for discussions. DLN is thankful the Case Western Reserve

University for hospitality when this work was initiated. We acknowledge the use of the xAct - xPand package for Mathematica [46, 47]. SA and GDS are supported by a Department of Energy grant DE-SC0009946 to the particle astrophysics theory group at CWRU.

-
- [1] P. Creminelli, G. D’Amico, J. Norena, and F. Vernizzi, *JCAP* **0902**, 018 (2009), arXiv:0811.0827 [astro-ph].
- [2] I. Zlatev, L. Wang, and P. J. Steinhardt, *Physical Review Letters* **82**, 896 (1999), astro-ph/9807002.
- [3] T. Chiba, T. Okabe, and M. Yamaguchi, *Phys. Rev. D* **62**, 023511 (2000), astro-ph/9912463.
- [4] N. A. Hamed, H. S. Cheng, M. A. Luty, and S. Mukohyama, *Journal of High Energy Physics* **5**, 074 (2004), hep-th/0312099.
- [5] S. Nojiri and S. D. Odintsov, *eConf* **C0602061**, 06 (2006), arXiv:hep-th/0601213 [hep-th].
- [6] T. P. Sotiriou and V. Faraoni, *Reviews of Modern Physics* **82**, 451 (2010), arXiv:0805.1726 [gr-qc].
- [7] G. Dvali, G. Gabadadze, and M. Porrati, *Physics Letters B* **485**, 208 (2000), hep-th/0005016.
- [8] R. Maartens, *Living Reviews in Relativity* **7**, 7 (2004), gr-qc/0312059.
- [9] <http://sci.esa.int/euclid/>.
- [10] <http://desi.lbl.gov/>.
- [11] T. Clifton, P. G. Ferreira, A. Padilla, and C. Skordis, *Phys.Rept.* **513**, 1 (2012), arXiv:1106.2476 [astro-ph.CO].
- [12] C. de Rham, *Living Reviews in Relativity* **17**, 7 (2014), arXiv:1401.4173 [hep-th].
- [13] A. J. Tolley, *Lect.Notes Phys.* **892**, 203 (2015).
- [14] G. D’Amico, G. Gabadadze, L. Hui, and D. Pirtskhalava, *Phys. Rev. D* **87**, 064037 (2013), arXiv:1206.4253 [hep-th].
- [15] C. de Rham and G. Gabadadze, *Phys. Rev. D* **82**, 044020 (2010), arXiv:1007.0443 [hep-th].
- [16] C. de Rham, G. Gabadadze, and A. J. Tolley, *Physical Review Letters* **106**, 231101 (2011), arXiv:1011.1232 [hep-th].
- [17] G. D’Amico, C. de Rham, S. Dubovsky, G. Gabadadze, D. Pirtskhalava, and A. J. Tolley, *Phys. Rev. D* **84**, 124046 (2011).
- [18] R. Gannouji, M. W. Hossain, M. Sami, and E. N. Saridakis, *Phys. Rev. D* **87**, 123536 (2013), arXiv:1304.5095 [gr-qc].
- [19] K. Bamba, M. W. Hossain, R. Myrzakulov, S. Nojiri, and M. Sami, *Phys. Rev. D* **89**, 083518 (2014), arXiv:1309.6413 [hep-th].
- [20] S. Dubovsky, R. Flauger, A. Starobinsky, and I. Tkachev, *Phys. Rev. D* **81**, 023523 (2010).
- [21] A. Emir Gümürükçüoğlu, S. Kuroyanagi, C. Lin, S. Mukohyama, and N. Tanahashi, *Classical and Quantum Gravity* **29**, 235026 (2012), arXiv:1208.5975 [hep-th].
- [22] D. Bessada and O. D. Miranda, *JCAP* **8**, 033 (2009), arXiv:0908.1360 [astro-ph.CO].
- [23] Notice that $X = 0$ and $X = 1$ cannot be crossed in the past history.
- [24] Planck Collaboration, P. A. R. Ade, N. Aghanim, M. Arnaud, M. Ashdown, J. Aumont, C. Baccigalupi, A. J. Banday, R. B. Barreiro, J. G. Bartlett, and et al., ArXiv e-prints (2015), arXiv:1502.01589.
- [25] Planck Collaboration, P. A. R. Ade, N. Aghanim, M. Arnaud, M. Ashdown, J. Aumont, C. Baccigalupi, A. J. Banday, R. B. Barreiro, N. Bartolo, and et al., ArXiv e-prints (2015), arXiv:1502.01590.
- [26] To be consistent with observations, $H^2 \sim a^{-3}$ or $H^2 \sim a^{-4}$ in the past, so $\Omega_\sigma H^2 \sim a^{-1}$ or $\Omega_\sigma H^2 \sim a^{-2}$.
- [27] Note that this expression coincides with the one computed for the extended quasidilaton in [41].
- [28] K. Olive *et al.* (Particle Data Group), *Chin.Phys.* **C38**, 090001 (2014).
- [29] E. Witten, *Phys. Rev. D* **30**, 272 (1984).
- [30] B. W. Lynn, A. E. Nelson, and N. Tetradis, *Nuclear Physics B* **345**, 186 (1990).
- [31] D. M. Jacobs, G. D. Starkman, and B. W. Lynn, *MNRAS* **450**, 3418 (2015).
- [32] S. S. McGaugh, *Canadian Journal of Physics* **93**, 250 (2015), arXiv:1404.7525.
- [33] A. Lue, R. Scoccimarro, and G. Starkman, *Phys. Rev. D* **69**, 044005 (2004), astro-ph/0307034.
- [34] W. Hu and I. Sawicki, *Phys. Rev. D* **76**, 104043 (2007), arXiv:0708.1190.
- [35] G. D’Amico, G. Gabadadze, L. Hui, and D. Pirtskhalava, *Classical and Quantum Gravity* **30**, 184005 (2013), arXiv:1304.0723 [hep-th].
- [36] A. Emir Gumrukcuoglu, K. Hinterbichler, C. Lin, S. Mukohyama, and M. Trodden, ArXiv e-prints (2013), arXiv:1304.0449 [hep-th].
- [37] A. De Felice and S. Mukohyama, *Physics Letters B* **728**, 622 (2014), arXiv:1306.5502 [hep-th].
- [38] A. De Felice, A. E. Gümürükçüoğlu, and S. Mukohyama, *Phys. Rev. D* **88**, 124006 (2013), arXiv:1309.3162 [hep-th].
- [39] H. Motohashi and W. Hu, *Phys. Rev. D* **90**, 104008 (2014), arXiv:1408.4813 [hep-th].
- [40] T. Kahniashvili, A. Kar, G. Lavrelashvili, N. Agarwal, L. Heisenberg, and A. Kosowsky, *Phys. Rev. D* **91**, 041301 (2015), arXiv:1412.4300.
- [41] L. Heisenberg, *JCAP* **4**, 010 (2015), arXiv:1501.07796 [hep-th].
- [42] Notice that if the Galileon type terms considered in [38] are added, the constraint equation (23) and consequently eq. (41) would not change. Therefore the new parameters would enter in the computation only through the information encoded by the cosmological constraints at redshift zero. We thus expect that, after the marginalization over the extra parameters, the allowed region would be broader.
- [43] Planck Collaboration, P. A. R. Ade, N. Aghanim, M. Arnaud, M. Ashdown, J. Aumont, C. Baccigalupi, A. J. Banday, R. B. Barreiro, J. G. Bartlett, and et al., ArXiv e-prints (2015), arXiv:1502.01597.
- [44] T. Delubac, J. E. Bautista, N. G. Busca, J. Rich, D. Kirkby, S. Bailey, A. Font-Ribera, A. Slosar, K.-

- G. Lee, M. M. Pieri, J.-C. Hamilton, É. Aubourg, M. Blomqvist, J. Bovy, J. Brinkmann, W. Carithers, K. S. Dawson, D. J. Eisenstein, S. G. A. Gontcho, J.-P. Kneib, J.-M. Le Goff, D. Margala, J. Miralda-Escudé, A. D. Myers, R. C. Nichol, P. Noterdaeme, R. O'Connell, M. D. Olmstead, N. Palanque-Delabrouille, I. Pâris, P. Petitjean, N. P. Ross, G. Rossi, D. J. Schlegel, D. P. Schneider, D. H. Weinberg, C. Yèche, and D. G. York, *Astron. Astrophys.* **574**, A59 (2015), arXiv:1404.1801.
- [45] We note this inequality is satisfied in the QDMG case for parameters in the allowed regions.
- [46] C. Pitrou, X. Roy, and O. Umeh, *Classical and Quantum Gravity* **30**, 165002 (2013), arXiv:1302.6174 [astro-ph.CO].
- [47] J. M. Martín-García, <http://www.xact.es>.