encounter a disagreement with experiment in weak interactions. Then, one must instead overcome difficulties associated with the massless boson. I am indebted to Dr. Th. A. J. Maris for pointing this out.

π^-p Interactions at 683 MeV/c, R. A. BURNSTEIN, G. R. CHARLTON, T. P. DAY, G. QUARENI, A. QUARENI-VIGNUDELLI, G. B. YODH, AND I. NADELHAFT [Phys. Rev. **137**, B1044 (1965)]. There were some omissions in Ref. 7 of this paper, which should read: 'M. Olsson and G. B. Yodh, Phys. Rev. Letters **10**, 353 (1963); Bull. Am. Phys. Soc. **9**, 27 (1964); University of Maryland Technical Report No. 358, 1964 (unpublished) and M. Olsson (thesis), University of Maryland Technical Report No. 379, 1964 (unpublished). The model herein is based upon the isobar model of S. J. Lindenbaum and R. M. Sternheimer, Phys. Rev. **105**, 1874 (1957); **106**, 1107 (1957); R. M. Sternheimer and S. J. Lindenbaum, Phys. Rev. **109**, 1723 (1958); **123**, 333 (1961).

Frequency Dependence of the Two-Magnon Ferromagnetic Resonance Linewidth, P. E. SEIDEN AND M. SPARKS [Phys. Rev. **137**, A1278 (1965)]. Reference 1 should read: E. Schlömann, in Proceedings of the Conference on Magnetism and Magnetic Materials, 1956, Boston, Massachusetts (unpublished). In the sentence after Eq. (14) on p. A1280 the factor $\cos\theta_k$ should be $\cos\theta_u$.

Electromagnetic Structure of the Giant Resonance in Oxygen-16, F. H. LEWIS, JR. [Phys. Rev. **134**,

FIG. 1.

FIG. 2.

FIG. 4.

B331 (1964)]. Because of a numerical error in the original calculations, the graphs shown below should replace Figs. 1, 2, and 4 of the paper, as indicated. The discussion of the results should remain unchanged except for the remarks in the last paragraph starting on p. B335 concerning the state at 17.3 MeV. The theoretical cross section for this