ERRATA

Erratum: Quantum theory of the stability region of an ion in a Paul trap [Phys. Rev. A 52, 1419 (1995)]

Wang Kelin, Feng Mang, and Wu Juhao

PACS number(s): 32.80.Pj, 03.65.Ge, 33.80.Ps, 42.50.Vk, 99.10.+g

We would like to correct a typographical error in our publication. Equation (19) should be revised as

$$B_0 = (mc/u)^{1/4}, \quad (19)$$

$$B_k = \frac{1}{mk^2\omega^2 B_0^3 - 4uB_0^3} [-m\omega^2 F_1^k + m\omega^2 F_2^{k+2} + uF_4^k + vF_3^{k-1}].$$

1050-2947/96/53(2)/1200(1)/\$06.00

© 1996 The American Physical Society

Erratum: Distribution of electrons in double photoionization of helium and heavier atoms in the asymptotic region [Phys. Rev. A 52, 3910 (1995)]

E. G. Drukarev

Petersburg Nuclear Physics Institute, Gatchina, St. Petersburg 188350, Russia

To avoid misunderstanding, the following insertions should be made in the text:

1. This paragraph should follow the one containing Eq. (6): To obtain the energy distribution it is crucial to take into account that the electrons can exchange by hard photons carrying large momenta of the order $p_1 \gg \eta$ [1]. While $J/\varepsilon_i \ll 1$ the interaction can be included in the lowest order of perturbative theory. This is illustrated by Figs. 1(b) and 1(c), where the final-state and intermediate-state electrons are described by free functions. The initial state is described by the function $\Psi(r_1, r_2)$.

2. After Eq. (23) it should read: Here, as well as in Eq. (4), the hard photon exchange is included in the function $\psi(r_1, r_2)$. This leads to $i(e\nabla_1)\Psi(r_1, r_2) \sim 4\pi\alpha m[(e\rho)/\rho]\Psi(r_1, r_1)$.

[1] M. Ya. Amusla, E. G. Drukarev, V. G. Gorshkov, and M. P. Kazachkov, J. Phys. B **8**, 1248 (1975).