Erratum: Electron-ion collision spectroscopy: Lithium-like xenon ions
[Phys. Rev. A 91, 012710 (2015)]

D. Bernhardt, C. Brandau, Z. Harman, C. Kozhuharov, S. Böhm, F. Bosch, S. Fritzsche, J. Jacobi, S. Kieslich, H. Knopp,
 F. Nolden, W. Shi, Z. Stachura, M. Steck, Th. Stöhlker, S. Schippers, and A. Müller

(Received 18 July 2016; published 8 August 2016)

DOI: [10.1103/PhysRevA.94.029903](https://doi.org/10.1103/PhysRevA.94.029903)

In our publication, in Tables IV and V as well as in Fig. 10, the experimental resonance strengths S and the theoretical resonance strengths S_T were scaled according to $S' = Sp^2n^3$ and $S'_T = S_T p_T^2 n^3$, respectively, to account for the dominant expected proportionalities. In these equations, p is the momentum in the electron-ion center-of-mass frame, and n the principal quantum number of the shell into which the incident electron is captured. In the calculation of $p^2 = (E_{\text{c.m.}}/c)^2 + 2m_0 E_{\text{c.m.}}$ by mistake the mass of the ion, that is, $m_{0,i} = 136 \times 931.494\,061 \times 10^6 \text{ eV}/c^2$ was used instead of the mass of the electron $m_{0,e} = 510.998\,928 \times 10^3 \text{ eV}/c^2$. Thus, within the three digits given, all scaled resonance strengths S' and S'_T in Tables IV and V should be divided by a constant value of 2.48×10^5 . Since Fig. 10 is a plot of the scaled resonance strengths, this correction also applies to Fig. 10.

Please note, that this correction only applies to the scaled resonance strengths S' and S'_T and not to dielectronic recombination rate coefficients or unscaled resonance strengths used elsewhere in the paper. In addition, none of the conclusions of our paper are influenced by this correction.