**Erratum: Atomic cluster expansion for accurate and transferable interatomic potentials
[Phys. Rev. B **99**, 014104 (2019)]**

Ralf Drautz

(Received 11 June 2019; published 23 December 2019)

DOI: [10.1103/PhysRevB.100.249901](https://doi.org/10.1103/PhysRevB.100.249901)

In Eq. (C6), the relative weight of the different J couplings was omitted. The symbol $\begin{bmatrix} j_1 & j_2 & j_3 & j_4 \\ m_1 & m_2 & m_3 & m_4 \end{bmatrix}$ depends on the relative contribution of the different couplings J . The correct form of Eq. (C6) reads

$$\begin{bmatrix} j_1 & j_2 & j_3 & j_4 \\ m_1 & m_2 & m_3 & m_4 \end{bmatrix} = \sum_J c_J \sum_{M=-J}^J (-1)^M C_{J j_1 j_2}^{M m_1 m_2} C_{J j_3 j_4}^{-M m_3 m_4}, \quad (\text{C6})$$

with constants c_J that determine the relative contributions of the different couplings J and $\sum_J c_J^2 = 1$. The coefficients c_J , in general, differ for different basis functions indexed by $n_1 n_2 n_3 n_4 j_1 j_2 j_3 j_4$.

In analogy, the correct expression for Eq. (C8) is given by

$$\begin{bmatrix} j_1 & j_2 & j_3 & j_4 & j_5 \\ m_1 & m_2 & m_3 & m_4 & m_5 \end{bmatrix} = \sum_{J_1 J_2} c_{J_1 J_2} \sum_{M_1 M_2} (-1)^{M_2} C_{J_1 j_1 j_2}^{M_1 m_1 m_2} C_{J_2 j_3 j_4}^{M_2 m_3 m_4} C_{J_2 j_4 j_5}^{-M_2 m_4 m_5}, \quad (\text{C8})$$

with $\sum_{J_1 J_2} c_{J_1 J_2}^2 = 1$.

None of the conclusions or results of the paper are affected by these typing errors.

I am indebted to Dr. A. Shapeev from Skoltech, Russia for pointing out the mistake in Eq. (C6).