Strain-induced pseudomagnetic field and quantum oscillations in kagome crystals

Tianyu Liu

Max-Planck-Institut für Physik komplexer Systeme, 01187 Dresden, Germany

(Received 21 February 2020; revised 12 July 2020; accepted 15 July 2020; published 30 July 2020)

A kagome lattice is composed of corner-sharing triangles arranged on a honeycomb lattice such that each honeycomb bond hosts a kagome site while each kagome triangle encloses a honeycomb site. Such close relation implies that the two lattices share common features. We predict here that a kagome crystal, similar to the honeycomb lattice graphene, reacts to elastic strain in a unique way that the bulk electronic states in the vicinity of Dirac points are reorganized by the strain-induced pseudomagnetic field into pseudo-Landau levels, while the degenerate edge states in the undeformed crystal become separated in the energy dimension. When the strain is tuned continuously, the resulting scanning pseudomagnetic field gives rise to quantum oscillations in both the density of states and the electrical conductivity.

DOI: [10.1103/PhysRevB.102.045151](https://doi.org/10.1103/PhysRevB.102.045151)

I. INTRODUCTION

A kagome lattice is a hexagonal Bravais lattice with a three-site basis. The unique lattice site arrangement renders the kagome lattice an ideal platform to study the geometric frustration [1] and the resulting exotic quantum states of matter known as quantum spin liquids [2–10]. Also for structural reasons, the wave function associated with a hexagonal ring in the kagome lattice becomes completely localized due to the destructive interference of the wave functions of the corner sites [11,12], resulting in highly degenerate dispersionless bands [13,14] stable against disorder [15]. The existence of these flat bands has recently been verified using the scanning tunneling microscope (STM) in layered silicene [16] and $\text{Co}_3\text{Sn}_2\text{S}_2$ [17]. In the presence of the spin-orbit coupling, the band degeneracy is lifted and the flat bands acquire a nonzero Chern number, giving rise to the fractional quantum Hall effect (FQHE) when partially filled [18]. The spin-orbit coupling can also gap out the two dispersive bands above the flat band, producing the quantum anomalous Hall effect (QAHE) [19,20] or the quantum spin Hall effect (QSHE) [21,22].

In the absence of the spin-orbit coupling and next-nearest-neighbor hoppings, the two dispersive bands linearly touch at the corners of the Brillouin zone. Such band crossings have been theoretically acknowledged for a long time but experimentally observed only recently in FeSn [23]. Materials with linear energy band crossings react to elastic strain in a unique way by generating in the vicinity of the crossings a chiral gauge field, which is first proposed in graphene [24–26] and later generalized to Dirac and Weyl semimetals [27–33], Weyl

and d -wave superconductors [34–38], and various bosonic Dirac materials [39–41]. However, whether a similar strain-induced gauge field occurs in kagome crystals has not yet been systematically investigated before, presumably because the additional quadratic band touching of the dispersive bands and the flat band [42,43] produces an extra degree of freedom obscuring the Dirac physics derived from the dispersive bands.

In this paper, by projecting to a subspace associated with the linear band crossings, we are able to get rid of the extra degree of freedom and extract the Dirac physics. Through a combination of analytical calculation and numerical simulation, we elaborate that a properly engineered strain can induce a uniform pseudomagnetic field, which engenders pseudo-Landau levels and quantum oscillations. To support these findings, we organize this paper as follows. In Sec. II, we study the band structure of a kagome crystal with only nearest-neighbor hopping terms and analytically extract the Dirac physics from the sublattice space Hamiltonian. In Sec. III, we investigate the response of a kagome crystal to elastic strain with our focus on the bulk Dirac cones as well as the edge states residing on a pair of sawtooth boundaries. In Sec. IV, we demonstrate the quantum oscillations of the density of states (DOS) and the longitudinal electrical conductivity in the kagome crystal. Section V concludes the paper and discusses the experimental implementation of the strain-induced quantum oscillations.

II. MODEL

We consider a toy model of a periodic kagome crystal with only nearest-neighbor hopping terms. The unit cell of the lattice contains three sites as illustrated by the shaded triangle in Fig. 1(a). The tight-binding Hamiltonian then reads

$$H = \sum_r (t_1 b_r^\dagger a_r + t_1 a_{r+\beta_1}^\dagger b_r + t_2 c_r^\dagger b_r + t_2 b_{r+\beta_2}^\dagger c_r + t_3 a_r^\dagger c_r + t_3 c_{r+\beta_3}^\dagger a_r) + \text{H.c.}, \quad (1)$$

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by the Max Planck Society.

FIG. 1. (a) Schematic plot of a periodic kagome lattice which is a hexagonal Bravais lattice with a three-site basis illustrated by the shaded triangle. The associated primitive vectors are $\beta_1 = 2a\hat{x}$, $\beta_2 = -a\hat{x} + \sqrt{3}a\hat{y}$, and $\beta_3 = -a\hat{x} - \sqrt{3}a\hat{y}$, with a being the lattice constant. (b) The first Brillouin zone of the kagome lattice with high-symmetry points labeled. (c) Band structure of a kagome crystal plotted along the path in (b). There are two linear band crossings at the corners (K/K') of the Brillouin zone and one quadratic band crossing in the center (Γ) of the Brillouin zone. The energy bands (red dashed curves) of the linearized Hamiltonian [Eq. (5)] are overlaid onto the band crossings at K and K' .

where the summation of $\mathbf{r}$ runs over all unit cells and $t_{i=1,2,3}$ is the hopping parameter between the nearest neighbors along the direction of the primitive vector $\beta_{i=1,2,3}$. Apply Fourier transform

$$\begin{pmatrix} a_r \\ b_r \\ c_r \end{pmatrix} = \frac{1}{\sqrt{N_{uc}}} \sum_k e^{ik \cdot r} \begin{pmatrix} a_k \\ b_k \\ c_k \end{pmatrix}, \quad (2)$$

where N_{uc} is the number of unit cells. We can then rewrite the Hamiltonian [Eq. (1)] in the sublattice basis $\psi_k = (a_k, b_k, c_k)^T$ as $H = \sum_k \psi_k^\dagger \mathcal{H}_k \psi_k$, where the first quantized Bloch Hamiltonian matrix reads

$$\mathcal{H}_k = \begin{pmatrix} 0 & t_1 + t_1 e^{-ik \cdot \beta_1} & t_3 + t_3 e^{ik \cdot \beta_3} \\ & 0 & t_2 + t_2 e^{-ik \cdot \beta_2} \\ & & 0 \end{pmatrix}. \quad (3)$$

In the absence of anisotropy, we have $t_{i=1,2,3} = t$, in which case, $\mathcal{H}_k$ produces three bands

$$\begin{aligned} \epsilon_{k,0} &= -2t, \\ \epsilon_{k,1} &= t - t\sqrt{3 + 2\lambda_k}, \\ \epsilon_{k,2} &= t + t\sqrt{3 + 2\lambda_k}, \end{aligned} \quad (4)$$

where $\lambda_k = \sum_i \cos(\mathbf{k} \cdot \beta_i)$. The flat band $\epsilon_{k,0}$ results from the destructive interference in the hexagonal rings [11,12], while the two dispersive bands $\epsilon_{k,1}$ and $\epsilon_{k,2}$ are similar to the bands of graphene [44,45], whose Bravais lattice is also hexagonal but with a two-site basis. Band $\epsilon_{k,1}$ and band $\epsilon_{k,2}$ cross at the corners of the Brillouin zone $\mathbf{K}_\eta = (\eta \frac{2\pi}{3a}, 0)$ at energy $\epsilon_{\mathbf{K}_\eta, i=1,2} = t$. To scrutinize the band crossings, we study the Hamiltonian in the vicinity of the Brillouin zone corners by

projecting $\mathcal{H}_{\mathbf{K}_\eta + \mathbf{q}}$ onto the space spanned by the eigenvectors $|\phi_{\mathbf{K}_\eta, i=1,2}\rangle$ of $\mathcal{H}_{\mathbf{K}_\eta}$ associated with eigenenergies $\epsilon_{\mathbf{K}_\eta, i=1,2} = t$. Explicitly, we have

$$\begin{aligned} \langle \mathcal{H}_{\mathbf{K}_\eta + \mathbf{q}} \rangle_\phi &= \begin{pmatrix} \langle \phi_{\mathbf{K}_\eta, 1} | \mathcal{H}_{\mathbf{K}_\eta + \mathbf{q}} | \phi_{\mathbf{K}_\eta, 1} \rangle & \langle \phi_{\mathbf{K}_\eta, 1} | \mathcal{H}_{\mathbf{K}_\eta + \mathbf{q}} | \phi_{\mathbf{K}_\eta, 2} \rangle \\ \langle \phi_{\mathbf{K}_\eta, 2} | \mathcal{H}_{\mathbf{K}_\eta + \mathbf{q}} | \phi_{\mathbf{K}_\eta, 1} \rangle & \langle \phi_{\mathbf{K}_\eta, 2} | \mathcal{H}_{\mathbf{K}_\eta + \mathbf{q}} | \phi_{\mathbf{K}_\eta, 2} \rangle \end{pmatrix} \\ &\approx t\sigma^0 + \sqrt{3}\eta at q_x \sigma^x + \sqrt{3}at q_y \sigma^y, \end{aligned} \quad (5)$$

where we have used $|\phi_{\mathbf{K}_\eta, 1}\rangle = \frac{1}{\sqrt{3}}(1, 1, 1)^T$ and $|\phi_{\mathbf{K}_\eta, 2}\rangle = \frac{1}{\sqrt{3}}(\frac{1}{2} + \frac{\sqrt{3}}{2}i\eta, \frac{1}{2} - \frac{\sqrt{3}}{2}i\eta, -1)^T$. $\sigma^{x,y}$ and σ^0 are the Pauli matrices and the identity matrix in the space $\{|\phi_{\mathbf{K}_\eta, 1}\rangle, |\phi_{\mathbf{K}_\eta, 2}\rangle\}$, in which $\langle \mathcal{H}_{\mathbf{K}_\eta + \mathbf{q}} \rangle_\phi$ is a standard Dirac Hamiltonian encoding at the corners of the Brillouin zone two Dirac cones [red dashed curves, Fig. 1(c)] characterized by velocity parameters $(v_x^\eta, v_y^\eta) = (\sqrt{3}\eta at/\hbar, \sqrt{3}at/\hbar)$. Such Dirac cones can well capture the spectral features of the first quantized Bloch Hamiltonian [Eq. (3)] in the vicinity of K and K' as illustrated in Fig. 1(c).

We have extracted the Dirac physics from the bulk of the Hamiltonian [Eq. (1)]. We now briefly discuss kagome crystals with open boundary conditions. For a nanoribbon with a pair of sawtooth boundaries [Fig. 2(a)], we observe that a doubly degenerate arc state connects the two Dirac points while the other doubly degenerate arc state emerges from and terminates at the quadratic band touching arising from the flat band and the lower dispersive band [Fig. 2(b)]. The edge origination of these arc states can be confirmed by calculating the spectral function on each sawtooth edge

$$A_s(E, k_x) = -\frac{1}{\pi} \sum_{y \in \text{edge}} \lim_{\delta \rightarrow 0} \Im[E + i\delta - \mathcal{H}_{yy'}(k_x)]_{y=y'}^{-1}. \quad (6)$$

The spectral function associated with the upper edge is plotted in Fig. 2(c), while the spectral function associated with the lower edge is identical as illustrated in Fig. 2(d). For a nanoribbon with a pair of zigzag edges [Fig. 2(e)], we also find two pairs of doubly degenerate edge states connecting Dirac points and the quadratic band touchings [Fig. 2(f)], respectively. The edge origination is confirmed by calculating the spectral function on each zigzag edge

$$A_z(E, k_y) = -\frac{1}{\pi} \sum_{x \in \text{edge}} \lim_{\delta \rightarrow 0} \Im[E + i\delta - \mathcal{H}_{xx'}(k_y)]_{x=x'}^{-1}, \quad (7)$$

which is plotted in Fig. 2(g) [Fig. 2(h)] for the left (right) edge.

III. STRAIN-INDUCED PSEUDOMAGNETIC FIELDS

In Sec. II, we have studied the band structure of kagome crystals. The linear energy band crossings at Brillouin zone corners indicate that kagome crystals belong in the family of Dirac matter. One of the most important features of Dirac matter is that elastic strain is equivalent to a gauge field inducing Landau quantization [24–41]. In this section, we will study the reaction of kagome crystals to a properly engineered strain.

The most important effect of elastic strain is that it alters the position of each lattice site, resulting in spatial modulation of the overlap of electron clouds from neighboring sites [46]. Explicitly, the hopping parameter $t_{\mathbf{R}', \mathbf{R}}$ between site $\mathbf{R}'$ and $\mathbf{R}$ will be varied by an amount of $\delta t_{\mathbf{R}', \mathbf{R}} = t_{\mathbf{R}' + \mathbf{u}(\mathbf{R}'), \mathbf{R} + \mathbf{u}(\mathbf{R})} - t_{\mathbf{R}', \mathbf{R}}$,

FIG. 2. Band structure of kagome nanoribbons. (a) A kagome nanoribbon with a pair of sawtooth edges along the x direction. Each unit cell (between the two dashed zigzag lines) contains $2N = 50$ shaded triangles and an extra c site c_0 on the lower edge. (b) The band structure of the sawtooth kagome nanoribbon in (a). Each sawtooth edge hosts two edge states. One of them is located between the two dispersive bands, connecting the Dirac points, while the other is located between the flat band and the lower dispersive band. (c) The spectral function [Eq. (6)] of the upper sawtooth edge. (d) The spectral function [Eq. (6)] of the lower sawtooth edge. (e) A kagome nanoribbon with a pair of zigzag edges along the y direction. Each unit cell (between the two dashed lines) contains $2N - 1 = 49$ shaded triangles. On the left edge, there are an extra b site b_0 and an extra c site c_0 . On the right edge, there are an extra a site a_{2N} and an extra c site c_{2N} . (f) The band structure of the zigzag kagome nanoribbon in (e). Each zigzag edge hosts two edge states. One of them is located between the two dispersive bands, while the other is located between the flat band and the lower dispersive band. (g) The spectral function [Eq. (7)] of the left zigzag edge. (h) The spectral function [Eq. (7)] of the right zigzag edge. It is worth noting that the flat bands in all spectral function plots are bulk states rather than edge states. The appearance of the flat bands can be attributed to the fact that both the sawtooth edges and the zigzag edges contain sites from the hexagonal rings where the flat states are trapped.

where $\mathbf{u}(\mathbf{R})$ is the displacement of the lattice site located at $\mathbf{R}$. To the linear order, it reads

$$\begin{aligned} \delta t_{\mathbf{R}',\mathbf{R}} &\approx (\mathbf{R}' - \mathbf{R}) \cdot \nabla_{\mathbf{r}} \mathbf{u}(\mathbf{r})|_{\mathbf{R}} \cdot \nabla_{\mathbf{r}'-\mathbf{r}} t_{\mathbf{r}',\mathbf{r}}|_{\mathbf{R}'-\mathbf{R}} \\ &= -\frac{g t_{\mathbf{R}',\mathbf{R}}}{|\mathbf{R}' - \mathbf{R}|^2} (\mathbf{R}' - \mathbf{R}) \cdot \nabla_{\mathbf{r}} \mathbf{u}(\mathbf{r})|_{\mathbf{R}} \cdot (\mathbf{R}' - \mathbf{R}), \end{aligned} \quad (8)$$

where we have adopted exponentially decaying overlap integrals $t_{\mathbf{r}',\mathbf{r}} = t_{\mathbf{R}',\mathbf{R}} \exp[-g(|\mathbf{r}' - \mathbf{r}| - |\mathbf{R}' - \mathbf{R}|)]/|\mathbf{R}' - \mathbf{R}|$, in which g is the Grüneisen parameter [25]. Without loss of generality, we use $g = 1$ in the following. Consequently, in the presence of strain, the overlap integrals in Eq. (1) are modulated as $t_i \rightarrow t_i + \delta t_i$ with

$$\begin{aligned} \delta t_1 &= -t u_{xx}, \\ \delta t_2 &= -t \left(\frac{1}{4} u_{xx} + \frac{3}{4} u_{yy} - \frac{\sqrt{3}}{2} u_{xy} \right), \\ \delta t_3 &= -t \left(\frac{1}{4} u_{xx} + \frac{3}{4} u_{yy} + \frac{\sqrt{3}}{2} u_{xy} \right), \end{aligned} \quad (9)$$

where $u_{ij} = \frac{1}{2}(\partial_i u_j + \partial_j u_i)$ is the symmetrized strain tensor whose value should be taken at lattice site $\mathbf{R}$, from which an electron hops to the neighboring sites $\mathbf{R}' = \mathbf{R} + \boldsymbol{\beta}_i/2$. Under strain, the first quantized Bloch Hamiltonian [Eq. (3)] acquires

an extra term

$$\delta \mathcal{H}_{\mathbf{k}} = \begin{pmatrix} 0 & \delta t_1 + \delta t_1 e^{-ik \cdot \boldsymbol{\beta}_1} & \delta t_3 + \delta t_3 e^{ik \cdot \boldsymbol{\beta}_3} \\ & 0 & \delta t_2 + \delta t_2 e^{-ik \cdot \boldsymbol{\beta}_2} \\ & & 0 \end{pmatrix}. \quad (10)$$

To figure out how this strain-induced Hamiltonian can affect the Dirac cones, we project $\mathcal{H}_{\mathbf{K}_\eta + \mathbf{q}} + \delta \mathcal{H}_{\mathbf{K}_\eta + \mathbf{q}}$ onto the space spanned by $|\phi_{\mathbf{K}_\eta, i=1,2}\rangle$ and obtain

$$\begin{aligned} \langle \mathcal{H}_{\mathbf{K}_\eta + \mathbf{q}} + \delta \mathcal{H}_{\mathbf{K}_\eta + \mathbf{q}} \rangle_\phi &\approx \sqrt{3} \eta a t \left(q_x + \frac{e}{\hbar} \mathcal{A}_x \right) \sigma^x \\ &\quad + \sqrt{3} a t \left(q_y + \frac{e}{\hbar} \mathcal{A}_y \right) \sigma^y + (t + \mathcal{U}) \sigma^0. \end{aligned} \quad (11)$$

Consequently, the strain alters the strain-free Dirac Hamiltonian [Eq. (5)] in two ways. First, the onsite energy acquires an extra term $\mathcal{U} = -\frac{1}{2}t(u_{xx} + u_{yy})$, which may be interpreted as a strain-induced elastic scalar potential whose gradient corresponds to a strain-induced pseudoelectric field $\boldsymbol{\mathcal{E}} = -\frac{1}{e} \nabla \mathcal{U} = \frac{t}{2e} \nabla (u_{xx} + u_{yy})$. Second, compared to the strain-free Dirac Hamiltonian [Eq. (5)], the Dirac points of this projected Dirac Hamiltonian are shifted in the momentum

FIG. 3. Schematic plot of deformed kagome nanoribbons (solid) in contrast to undeformed nanoribbons (dashed). (a) A nanoribbon deformed by the properly engineered displacement field $\mathbf{u} = (\frac{C}{a}xy, -\frac{C}{2a}x^2 - \frac{C}{2a}y^2)$. (b) A nanoribbon deformed by the circular bend $\mathbf{u}' = (\frac{2C}{a}xy, -\frac{C}{a}x^2)$. The two displacement fields induce identical pseudomagnetic field [Eq. (13)], which is parallel (antiparallel) to the z axis at the K (K') point.

space through a standard Peierls substitution $\mathbf{q} \rightarrow \mathbf{q} + \frac{e}{\hbar} \mathcal{A}$. The strain-induced elastic vector potential can be read off as

$$\mathcal{A} = \eta \frac{\hbar}{2\sqrt{3}ea} (u_{yy} - u_{xx}, 2u_{xy}). \quad (12)$$

In contrast to the ordinary magnetic vector potential $\mathbf{A}$ that affects all the momenta $\mathbf{k}$ in the Brillouin zone through the Peierls substitution $\mathbf{k} \rightarrow \mathbf{k} + \frac{e}{\hbar} \mathbf{A}$, the elastic vector potential $\mathcal{A}$ only applies to the momenta $\mathbf{q}$ in the vicinity of Dirac points. Moreover, $\mathcal{A}$ takes opposite signs at different Dirac points due to the valley index η in Eq. (12). But similar to the ordinary magnetic vector potential, $\mathcal{A}$ can Landau-quantize electron bands if $\nabla \times \mathcal{A} \neq 0$. This requires the strain tensors u_{ij} to be space dependent which seems to contradict with the aforementioned derivation assuming constant strain. However, we argue that even when u_{ij} is spatially varying, the strain effect can still be treated as an elastic gauge field, as long as it varies slowly on the lattice scale.

To support our argument, we consider a properly engineered lattice deformation characterized by the displacement field $\mathbf{u} = (\frac{C}{a}xy, -\frac{C}{2a}x^2 - \frac{C}{2a}y^2)$, under which a rectangular nanoribbon is reshaped as is depicted in Fig. 3(a). According to Ref. [24], such reshaping may be accomplished by applying along the boundary either normal forces whose magnitudes are spatially modulated or isometric forces whose orientations are spatially modulated. The detailed derivation of such spatial modulations is beyond the scope of the present paper. In the following, we will only study the induced elastic gauge field resulting from the displacement. Specifically, the properly engineered displacement field $\mathbf{u}$ cancels the strain-induced scalar potential $\mathcal{U}$ but produces a strain-induced vector potential $\mathcal{A} = -\frac{\eta\hbar C}{\sqrt{3}ea^2} (y, 0)$, which leads to a homogeneous strain-induced pseudomagnetic field

$$\mathcal{B} = \nabla \times \mathcal{A} = \eta \frac{\hbar}{\sqrt{3}ea^2} C \hat{z}. \quad (13)$$

Alternatively, the same pseudomagnetic field can be induced by a circular bend [Fig. 3(b)] resulting from the displacement field $\mathbf{u}' = (\frac{2C}{a}xy, -\frac{C}{a}x^2)$ [47], whose strain-induced vector potential $\mathcal{A}' = -\frac{\eta\hbar C}{\sqrt{3}ea^2} (y, 0)$ is the same as that induced by

the properly engineered displacement $\mathbf{u}$. It is worth noting that such a displacement field also induces a nonvanishing elastic scalar potential $\mathcal{U}' = -\frac{C}{a}y$ giving rise to a pseudoelectric field, which will tilt the pseudo-Landau levels and thus obscure the associated transport [48]. For simplicity, we will focus on the properly engineered displacement field $\mathbf{u}$ in the following.

Due to the strain-induced pseudomagnetic field arising from the properly engineered displacement field, the Dirac cones are reorganized into pseudo-Landau levels

$$\varepsilon_n = \text{sgn}(n) \sqrt{2n \frac{e\mathcal{B}}{\hbar} \hbar v_x^\eta \hbar v_y^\eta}, \quad n = 0, \pm 1, \pm 2, \dots \quad (14)$$

As the common feature of Dirac matter, the presence of such strain-induced pseudo-Landau levels only relies on the existence of Dirac points, which are protected by crystalline symmetries in kagome crystals [23,49,50]. Therefore, these pseudo-Landau levels can survive the symmetry-preserving interactions (e.g., the onsite Hubbard interaction in Ref. [50]). On the other hand, if the crystalline symmetries protecting the linear band crossings are broken by weak interactions and/or the spin-orbit coupling, the Dirac cones are generally gapped out. Consequently, the pseudo-Landau levels will acquire a mass term m and become $\varepsilon_n \rightarrow \pm \sqrt{\varepsilon_n^2 + m^2}$. Since the interactions in kagome crystals are unlikely to qualitatively alter the Dirac cones and thus the strain-induced pseudo-Landau levels, we will ignore them for simplicity in the following. Detailed investigations regarding the impact of interactions on pseudo-Landau levels are provided in Refs. [51–53].

We have numerically verified these pseudo-Landau levels by applying the hopping substitution $t_i \rightarrow t_i + \delta t_i$ with δt_i listed in Eq. (9) to the tight-binding Hamiltonian of a nanoribbon with a sawtooth edge and a flat edge. Such a nanoribbon can be obtained by removing the c_0 sites in the nanoribbon illustrated in Fig. 2(a) and thus contains $2N$ shaded triangles. We emphasize that the profile of the boundaries does not affect the pseudo-Landau levels, which result from the strain-induced pseudomagnetic field coupled to the Dirac cones deep in the bulk. Indeed, we observe that the pseudo-Landau levels [Eq. (14)] capture the feature of the spectrum and the DOS in the vicinity of Dirac points ($\epsilon_{K_{\eta}, i=1,2} = t$) as illustrated in Fig. 4(a). For comparison, we also plot the spectrum and the DOS of an unstrained nanoribbon in the presence of an ordinary magnetic field which coincides with the pseudomagnetic field at the K point. Similar results are shown in Fig. 4(b).

Though the pseudomagnetic field [Eq. (13)] and the pseudo-Landau levels [Eq. (14)] are purely bulk effects, the elastic strain from which they are derived can affect the edge states as well because the properly engineered displacement field we have employed deforms the two edges differently [Fig. 3(a)]. Therefore, the energy band degeneracy arising from edge states will generally be lifted. To confirm this, we numerically study a sawtooth kagome nanoribbon, which hosts doubly degenerate edge states connecting the band crossings [Figs. 2(c) and 2(d)]. In the presence of strain, we notice that the two edge states residing on the upper (lower) edge of the nanoribbon become more (less) separated as illustrated in Fig. 5(b) [Fig. 5(c)]. Specifically, for the edge states between the two dispersive bands, we find that the one

FIG. 4. Numerically calculated Landau levels of a kagome nanoribbon with a sawtooth edge and a flat edge with $2N = 600$ shaded triangles. (a) Pseudo-Landau levels (blue curves) due to a strain-induced pseudomagnetic field [Eq. (13)] with $\sqrt{3}eBa^2/4\hbar = 8.66 \times 10^{-5}$ flux quanta per shaded triangle. The red dotted lines are the theoretically predicted pseudo-Landau levels [Eq. (14)], while the red curve is the numerically calculated DOS. For comparison, the numerically calculated Landau levels and DOS in the presence of an ordinary magnetic field $\mathbf{B} = B\hat{z}$ with $\sqrt{3}eBa^2/4\hbar = 8.66 \times 10^{-5}$ are plotted in (b).

on the upper edge always has higher energy than that on the lower edge. This is similar to the edge state separation in bent graphene nanoribbons [54].

To obtain more insights, we take a closer look in the vicinity of Dirac points to see how the edge states emerge from the bulk. Without deformation, the edge states connect Dirac points K and K' , similar to the Fermi arcs in Weyl semimetals. When the nanoribbon is gradually deformed, due

to the induced pseudomagnetic field, the two Dirac points at K and K' extend into the zeroth pseudo-Landau levels, whose outer (inner) ends are connected by the edge state on the upper (lower) sawtooth edge, resulting in edge state separation in the energy dimension as illustrated in Fig. 5(d). In the presence of an ordinary magnetic field, the edge state on the upper (lower) edge connects the left (right) ends of the two zeroth Landau levels as illustrated in Fig. 5(e), leading to edge state separation in the momentum dimension.

IV. STRAIN-INDUCED QUANTUM OSCILLATIONS

In Sec. III, we have demonstrated that a properly engineered elastic strain can induce a uniform pseudomagnetic field in kagome crystals. The electronic states near the Dirac points are reorganized by the strain into pseudo-Landau levels, implying that the magnetic transport associated with Landau levels can be reproduced by the applied elastic strain. In the present section, we will numerically study the quantum oscillations in kagome crystals in the presence of the strain-induced pseudomagnetic field.

Though predicted to be flat as in Eq. (14), the actual strain-induced pseudo-Landau levels in a kagome nanoribbon are dispersive as illustrated in Fig. 4(a). This is because the Peierls substitution $\mathbf{q} \rightarrow \mathbf{q} + \frac{e}{\hbar}\mathbf{A}$ for the Dirac Hamiltonian under strain [Eq. (11)] is only rigorously valid at the Dirac points, away from which the higher order correction terms $O(u_{ij}q_k)$ and $O(q_k q_l)$ in the strain-induced Hamiltonian [Eq. (10)] begin to deform the pseudo-Landau levels.

To appropriately incorporate the effect from the dispersive pseudo-Landau levels, we employ the tetrahedron method [55] to calculate the quantum oscillations of the DOS and the electrical conductivity. In particular, we first divide the 1D Brillouin zone into a set of discretized momenta k_x^i with $i = 1, 2, \dots, i_{\max}$ such that each interval $[k_x^i, k_x^{i+1}]$ is sufficiently narrow. Therefore, the pseudo-Landau levels in such an interval can be approximated to disperse linearly

$$\varepsilon_n(k_x) = \frac{\varepsilon_n^{i+1} - \varepsilon_n^i}{k_x^{i+1} - k_x^i} k_x + \frac{k_x^{i+1} \varepsilon_n^i - k_x^i \varepsilon_n^{i+1}}{k_x^{i+1} - k_x^i}, \quad (15)$$

FIG. 5. Edge state separation in a sawtooth kagome nanoribbon under strain. (a) Band structure of a kagome nanoribbon with a pair of sawtooth edges in the presence of strain. The degenerate edge states in the undeformed nanoribbon are now separated. (b) The spectral function of the upper sawtooth edge. The two edge states are further separated by the strain. (c) The spectral function of the lower sawtooth edge. The spacing between the two edge states is reduced by the strain. (d) Band structure in the vicinity of Dirac points under strain. The Dirac points are broadened by the strain-induced pseudomagnetic field into flat zeroth pseudo-Landau levels, whose outer (inner) ends are connected by the edge state residing on the upper (lower) sawtooth edge. The edge states are separated in the energy dimension. (e) Band structure in the vicinity of Dirac points under an ordinary magnetic field of the same magnitude. The Dirac points extend into flat zeroth Landau levels whose left (right) ends are connected by the edge state hosted by the upper (lower) sawtooth edge. The edge states are thus separated in the momentum dimension.

where we have denoted $\varepsilon_n^i = \varepsilon_n(k_x^i)$ for transparency. Consequently, the DOS is

$$\begin{aligned} g(\mu) &= \frac{L_x}{2\pi} \sum_n \sum_{i=1}^{i_{\max}-1} \int_{k_x^i}^{k_x^{i+1}} dk_x \delta[\mu - \varepsilon_n(k_x)] \\ &= \frac{L_x}{2\pi} \sum_n \sum_{i=1}^{i_{\max}-1} \frac{k_x^{i+1} - k_x^i}{\varepsilon_n^{i+1} - \varepsilon_n^i} [\Theta(\mu - \varepsilon_n^i) - \Theta(\mu - \varepsilon_n^{i+1})], \end{aligned} \quad (16)$$

where L_x is the length of the nanoribbon and $\Theta(\cdot)$ is the Heaviside step function. The electrical conductivity can be calculated using the Boltzmann formalism [56] as

$$\begin{aligned} \sigma_{xx}(\mu) &= \frac{L_x}{2\pi} \sum_n \sum_i^{i_{\max}-1} \int_{k_x^i}^{k_x^{i+1}} dk_x e^2 \tau_n(\varepsilon_n) (v_n^x)^2 \left(-\frac{\partial f}{\partial \varepsilon} \right)_{\varepsilon_n(k_x)} \\ &= \frac{e^2 L_x}{2\pi \hbar^2} \tau(\mu) \sum_n \sum_i^{i_{\max}-1} \frac{\varepsilon_n^{i+1} - \varepsilon_n^i}{k_x^{i+1} - k_x^i} \\ &\quad \times [\Theta(\mu - \varepsilon_n^i) - \Theta(\mu - \varepsilon_n^{i+1})], \end{aligned} \quad (17)$$

where $v_n^x = \frac{1}{\hbar} \frac{\partial \varepsilon_n}{\partial k_x}$ is the band group velocity and $f(\varepsilon) = [e^{(\varepsilon - \mu)/k_B T} + 1]^{-1}$ is the Fermi distribution function. We have assumed temperature $T \rightarrow 0$ and identical relaxation time $\tau_n(\mu) = \tau(\mu)$ for all the pseudo-Landau levels. Explicitly, the scattering rate can be approximated by the lowest order Born approximation [57]

$$\frac{1}{\tau(\mu)} = 2\pi g(\mu) n_{\text{imp}} C, \quad (18)$$

where n_{imp} is the impurity concentration and C depends on the strength of the scattering.

We have numerically calculated the DOS and the electrical conductivity at different pseudomagnetic fields and find they both exhibit oscillations periodic in $1/B$ at the Fermi level $\mu = 0.9565t$ as illustrated in Fig. 6. This is because the pseudo-Landau levels successively pass through the Fermi level when their spacing is continuously tuned by the varying pseudomagnetic field, resulting in periodic population of electrons on the Fermi level. We have also verified other Fermi energies and observed similar oscillations. For comparison, the quantum oscillations of the DOS and the electrical conductivity due to an ordinary magnetic field of the same magnitude are overlaid in Fig. 6. The match of the periodicity further confirms the similarity of the strain-induced pseudomagnetic field to the ordinary magnetic field.

V. CONCLUSIONS

In this paper, by a proper projection of the sublattice basis, we have studied the Dirac physics of kagome crystals with our focus on its response to the applied elastic strain and the associated transport in the form of quantum oscillations. We first analyze the band structure of a kagome crystal with only nearest-neighbor hopping terms. By projecting the sublattice basis Hamiltonian onto the space spanned by the eigenvectors associated with the Dirac points, we are able to drop off the degree of freedom related to the flat band and analytically

FIG. 6. Quantum oscillations of kagome crystals. The upper panel shows the quantum oscillations of the DOS at the Fermi level $\mu = 0.9565t$ resulting from the strain-induced pseudomagnetic field (red solid curve) and the ordinary magnetic field (green dashed curve). The blue crosses mark the value of $1/B$ and $1/B$ where Landau levels hit the Fermi level. The lower panel shows the quantum oscillations of the electrical conductivity, i.e., Shubnikov-de Haas (SdH) oscillations, caused by the strain-induced pseudomagnetic field (red solid curve) and the ordinary magnetic field (green dashed curve). Both conductivity curves are broadened by convolving a Lorentzian of width $\delta = 0.001t$ to consider the impurity effect in the relaxation time [Eq. (18)]. The DOS curves are also broadened but by a much smaller Lorentzian width $\delta' = 0.08\delta$ in order to better characterize the magnetic fields at which Landau levels coincide with the Fermi level. The magnetic field is measured in the unit of $10^{-4}\hbar/ea^2$.

extract the Dirac physics, which is also numerically simulated for nanoribbons with sawtooth edges and zigzag edges. For both cases, we find two pairs of doubly degenerate edge states emerging from and terminating at band crossings, i.e., the Dirac points between the two dispersive bands and the quadratic band touching between the flat band and the lower dispersive band.

Then, we incorporate elastic strain by hopping integral substitutions and elucidate that the most important effect of the strain to the bulk states is to shift the Dirac points oppositely. Therefore, a properly engineered strain can be interpreted as a uniform chiral pseudomagnetic field resulting in Landau quantization. By numerically studying a sawtooth kagome nanoribbon, we find the strain modulates the edges differently, thus lifting the edge state degeneracy. In particular, the edge states connecting the Dirac points become completely separated in the energy dimension due to the unique way they connect to the zeroth pseudo-Landau levels broadened from the Dirac points. Similar separation, however, is absent in the presence of an ordinary magnetic field due to the different connection of the edge states to the zeroth Landau levels, reflecting the fundamental difference between the strain-induced chiral pseudomagnetic field and the ordinary magnetic field.

Lastly, we study the DOS and the longitudinal electrical conductivity of a kagome nanoribbon under different values of the strain-induced pseudomagnetic field. The tetrahedron method is adopted in order to appropriately treat the Landau level dispersion resulting from the higher order corrections in

the strain-induced Hamiltonian. Though departing from the ideal flat Landau levels, the strain-induced dispersive pseudo-Landau levels can produce quantum oscillations in both the DOS and the longitudinal electrical conductivity.

To experimentally implement the strain-induced quantum oscillations, we first require a kagome crystal such as $\text{Co}_3\text{Sn}_2\text{S}_2$ [17,58,59], FeSn [23], and Fe_2Sn_3 [49,60,61] in which ordinary magnetic field Shubnikov-de Haas oscillations and de Haas-van Alphen oscillations have been observed [23,58,60]. Second, the candidate materials must be sufficiently flexible to sustain the strain designed for

pseudo-Landau levels. However, the mechanical properties regarding the flexibility of the aforementioned materials are not complete and further experimental work is needed to verify whether one or more of these materials are suitable for the strain-induced quantum oscillation experiment.

ACKNOWLEDGMENTS

The author is indebted to S. Fujimoto and Z. Shi for insightful discussions. I particularly thank R. Moessner and P. A. McClarty for the valuable suggestions for this work.

-
- [1] C. Lacroix, P. Mendels, and F. Mila, *Introduction to Frustrated Magnetism: Materials, Experiments, Theory* (Springer-Verlag, Berlin, Heidelberg, 2011).
 - [2] S.-H. Lee, H. Kikuchi, Y. Qiu, B. Lake, Q. Huang, K. Habicht, and K. Kiefer, *Nat. Mater.* **6**, 853 (2007).
 - [3] L. Balents, *Nature (London)* **464**, 199 (2010).
 - [4] S. Yan, D. A. Huse, and S. R. White, *Science* **332**, 1173 (2011).
 - [5] H.-C. Jiang, Z. Wang, and L. Balents, *Nat. Phys.* **8**, 902 (2012).
 - [6] T.-H. Han, J. S. Helton, S. Chu, D. G. Nocera, J. A. Rodriguez-Rivera, C. Broholm, and Y. S. Lee, *Nature (London)* **492**, 406 (2012).
 - [7] M. R. Norman, *Rev. Mod. Phys.* **88**, 041002 (2016).
 - [8] H.-J. Liao, Z.-Y. Xie, J. Chen, Z.-Y. Liu, H.-D. Xie, R.-Z. Huang, B. Normand, and T. Xiang, *Phys. Rev. Lett.* **118**, 137202 (2017).
 - [9] Y. Zhou, K. Kanoda, and T.-K. Ng, *Rev. Mod. Phys.* **89**, 025003 (2017).
 - [10] A. M. Läuchli, J. Sudan, and R. Moessner, *Phys. Rev. B* **100**, 155142 (2019).
 - [11] D. L. Bergman, C. Wu, and L. Balents, *Phys. Rev. B* **78**, 125104 (2008).
 - [12] B.-J. Yang and N. Nagaosa, *Phys. Rev. Lett.* **112**, 246402 (2014).
 - [13] A. Mielke, *J. Phys. A: Math. Gen.* **24**, 3311 (1991).
 - [14] A. Mielke, *J. Phys. A: Math. Gen.* **25**, 4335 (1992).
 - [15] T. Bilitewski and R. Moessner, *Phys. Rev. B* **98**, 235109 (2018).
 - [16] Z. Li, J. Zhuang, L. Wang, H. Feng, Q. Gao, X. Xu, W. Hao, X. Wang, C. Zhang, K. Wu *et al.*, *Sci. Adv.* **4**, eaau4511 (2018).
 - [17] J.-X. Yin, S. S. Zhang, G. Chang, Q. Wang, S. S. Tsirkin, Z. Guguchia, B. Lian, H. Zhou, K. Jiang, I. Belopolski *et al.*, *Nat. Phys.* **15**, 443 (2019).
 - [18] E. Tang, J.-W. Mei, and X.-G. Wen, *Phys. Rev. Lett.* **106**, 236802 (2011).
 - [19] Z.-Y. Zhang, *J. Phys.: Condens. Matter* **23**, 365801 (2011).
 - [20] G. Xu, B. Lian, and S.-C. Zhang, *Phys. Rev. Lett.* **115**, 186802 (2015).
 - [21] H.-M. Guo and M. Franz, *Phys. Rev. B* **80**, 113102 (2009).
 - [22] Z. Wang and P. Zhang, *New J. Phys.* **12**, 043055 (2010).
 - [23] M. Kang, L. Ye, S. Fang, J.-S. You, A. Levitan, M. Han, J. I. Facio, C. Jozwiak, A. Bostwick, E. Rotenberg *et al.*, *Nat. Mater.* **19**, 163 (2020).
 - [24] F. Guinea, M. Katsnelson, and A. Geim, *Nat. Phys.* **6**, 30 (2010).
 - [25] M. A. Vozmediano, M. Katsnelson, and F. Guinea, *Phys. Rep.* **496**, 109 (2010).
 - [26] N. Levy, S. Burke, K. Meaker, M. Panlasigui, A. Zettl, F. Guinea, A. C. Neto, and M. Crommie, *Science* **329**, 544 (2010).
 - [27] A. Cortijo, Y. Ferreirós, K. Landsteiner, and M. A. H. Vozmediano, *Phys. Rev. Lett.* **115**, 177202 (2015).
 - [28] H. Sumiyoshi and S. Fujimoto, *Phys. Rev. Lett.* **116**, 166601 (2016).
 - [29] D. I. Pikulin, A. Chen, and M. Franz, *Phys. Rev. X* **6**, 041021 (2016).
 - [30] A. G. Grushin, J. W. F. Venderbos, A. Vishwanath, and R. Ilan, *Phys. Rev. X* **6**, 041046 (2016).
 - [31] A. Cortijo, D. Kharzeev, K. Landsteiner, and M. A. H. Vozmediano, *Phys. Rev. B* **94**, 241405(R) (2016).
 - [32] V. Arjona, E. V. Castro, and M. A. Vozmediano, *Phys. Rev. B* **96**, 081110(R) (2017).
 - [33] T. Liu, D. Pikulin, and M. Franz, *Phys. Rev. B* **95**, 041201 (2017).
 - [34] T. Liu, M. Franz, and S. Fujimoto, *Phys. Rev. B* **96**, 224518 (2017).
 - [35] T. Matsushita, T. Liu, T. Mizushima, and S. Fujimoto, *Phys. Rev. B* **97**, 134519 (2018).
 - [36] T. Kobayashi, T. Matsushita, T. Mizushima, A. Tsuruta, and S. Fujimoto, *Phys. Rev. Lett.* **121**, 207002 (2018).
 - [37] G. Massarelli, G. Wachtel, J. Y. Wei, and A. Paramekanti, *Phys. Rev. B* **96**, 224516 (2017).
 - [38] E. M. Nica and M. Franz, *Phys. Rev. B* **97**, 024520 (2018).
 - [39] M. C. Rechtsman, J. M. Zeuner, A. Tünnermann, S. Nolte, M. Segev, and A. Szameit, *Nat. Photon.* **7**, 153 (2013).
 - [40] Y. Ferreiros and M. A. Vozmediano, *Phys. Rev. B* **97**, 054404 (2018).
 - [41] T. Liu and Z. Shi, *Phys. Rev. B* **99**, 214413 (2019).
 - [42] L. Du, X. Zhou, and G. A. Fiete, *Phys. Rev. B* **95**, 035136 (2017).
 - [43] Y. Ren, T.-S. Zeng, W. Zhu, and D. Sheng, *Phys. Rev. B* **98**, 205146 (2018).
 - [44] A. C. Neto, F. Guinea, N. M. Peres, K. S. Novoselov, and A. K. Geim, *Rev. Mod. Phys.* **81**, 109 (2009).
 - [45] S. D. Sarma, S. Adam, E. Hwang, and E. Rossi, *Rev. Mod. Phys.* **83**, 407 (2011).
 - [46] H. Shapourian, T. L. Hughes, and S. Ryu, *Phys. Rev. B* **92**, 165131 (2015).
 - [47] F. Guinea, A. Geim, M. Katsnelson, and K. Novoselov, *Phys. Rev. B* **81**, 035408 (2010).
 - [48] É. Lantagne-Hurtubise, X.-X. Zhang, and M. Franz, *Phys. Rev. B* **101**, 085423 (2020).

- [49] L. Ye, M. Kang, J. Liu, F. Von Cube, C. R. Wicker, T. Suzuki, C. Jozwiak, A. Bostwick, E. Rotenberg, D. C. Bell *et al.*, [Nature \(London\)](#) **555**, 638 (2018).
- [50] I. I. Mazin, H. O. Jeschke, F. Lechermann, H. Lee, M. Fink, R. Thomale, and R. Valentí, [Nat. Commun.](#) **5**, 4261 (2014).
- [51] B. Roy and I. F. Herbut, [Phys. Rev. B](#) **88**, 045425 (2013).
- [52] B. Roy and J. D. Sau, [Phys. Rev. B](#) **90**, 075427 (2014).
- [53] B. Roy, F. F. Assaad, and I. F. Herbut, [Phys. Rev. X](#) **4**, 021042 (2014).
- [54] S. Stuij, P. Jacobse, V. Juričić, and C. M. Smith, [Phys. Rev. B](#) **92**, 075424 (2015).
- [55] P. E. Blöchl, O. Jepsen, and O. K. Andersen, [Phys. Rev. B](#) **49**, 16223 (1994).
- [56] N. W. Ashcroft and N. D. Mermin, *Solid State Physics* (Saunders College, Philadelphia, 1976).
- [57] S. Doniach and E. Sondheimer, *Green's Functions for Solid State Physicists* (Imperial College Press, London, 1998).
- [58] E. Liu, Y. Sun, N. Kumar, L. Muechler, A. Sun, L. Jiao, S.-Y. Yang, D. Liu, A. Liang, Q. Xu *et al.*, [Nat. Phys.](#) **14**, 1125 (2018).
- [59] Q. Wang, Y. Xu, R. Lou, Z. Liu, M. Li, Y. Huang, D. Shen, H. Weng, S. Wang, and H. Lei, [Nat. Commun.](#) **9**, 3681 (2018).
- [60] L. Ye, M. K. Chan, R. D. McDonald, D. Graf, M. Kang, J. Liu, T. Suzuki, R. Comin, L. Fu, and J. G. Checkelsky, [Nat. Commun.](#) **10**, 4870 (2019).
- [61] J.-X. Yin, S. S. Zhang, H. Li, K. Jiang, G. Chang, B. Zhang, B. Lian, C. Xiang, I. Belopolski, H. Zheng *et al.*, [Nature \(London\)](#) **562**, 91 (2018).