Erratum: Quasi-two-dimensional magnetic correlations in $\text{Ni}_2\text{P}_2\text{S}_6$ probed by ^{31}P NMR [Phys. Rev. B **102**, 064429 (2020)]

A. P. Dioguardi , S. Selter, U. Peeck, S. Aswartham, M.-I. Sturza, R. Murugesan, M. S. Eldeeb, L. Hozoi, B. Büchner, and H.-J. Grafe

(Received 23 August 2021; published 3 September 2021)

DOI: [10.1103/PhysRevB.104.099901](https://doi.org/10.1103/PhysRevB.104.099901)

In the original version of our published paper, we erroneously calculated the high-temperature limits of the spin-lattice relaxation rates $\lim_{T \rightarrow \infty}(T_1^{-1})$ based on the raw high-temperature hyperfine coupling parameters, i.e., the values extracted by fitting the raw NMR shift K_{raw} vs χ without first correcting for bulk magnetic effects. The corrected hyperfine coupling values perpendicular to the applied field were stated in the paper— $|A_{\perp}(H_0 \parallel c^*)| = 0.5 \pm 0.2 \text{ T}/\mu_B$ and $|A_{\perp}(H_0 \perp c^*)| = 0.4 \pm 0.1 \text{ T}/\mu_B$ —but the raw values were used in the calculation. Using the corrected values, we find $\lim_{T \rightarrow \infty}(T_1^{-1})_{\parallel} = 200 \pm 100 \text{ s}^{-1}$ for $H_0 \parallel c^*$ and $\lim_{T \rightarrow \infty}(T_1^{-1})_{\perp} = 120 \pm 60 \text{ s}^{-1}$ for $H_0 \perp c^*$. We show the spin-lattice relaxation rates T_1^{-1} (markers) and the updated high-temperature limits (horizontal dashed lines) in the corrected Fig. 12. This correction does not change any conclusions drawn from the data or calculations.

FIG. 12. Markers show the spin-lattice relaxation rate T_1^{-1} vs temperature compared to the calculated high-temperature limiting values (dashed lines; shaded bands indicate uncertainty) for a local moment magnetic system as discussed in the text.