Erratum: Chiral gauge field and axial anomaly in a Weyl semimetal
[Phys. Rev. B 87, 235306 (2013)]

Chao-Xing Liu, Peng Ye, and Xiao-Liang Qi
 (Received 28 August 2015; published 18 September 2015)

DOI: [10.1103/PhysRevB.92.119904](https://doi.org/10.1103/PhysRevB.92.119904) PACS number(s): 71.90.+q, 03.65.Vf, 73.43.-f, 75.50.Pp, 99.10.Cd

A small revision for the anomaly equation $\partial_\mu j^{\mu L(R)} = (-)\frac{1}{32\pi^2}\epsilon^{\lambda\rho\mu\nu}f_{\lambda\rho}^{L(R)}f_{\mu\nu}^{L(R)}$, as well as the definition of the field strength $f_{\mu\nu}^{L(R)}$, in the paragraph just above Eqs. (2) and (3) is required. The anomaly equation should be $\partial_\mu j^{\mu L(R)} = -(+)\frac{1}{32\pi^2}\epsilon^{\lambda\rho\mu\nu}f_{\lambda\rho}^{L(R)}f_{\mu\nu}^{L(R)}$ for left-handed (right-handed) fermions and the corresponding field strength for left-handed and right-handed gauge field should be defined as $f_{\mu\nu}^{L(R)} = \partial_\mu(A_\nu + a_{\nu L(R)}) - \partial_\nu(A_\mu + a_{\mu L(R)})$. Our main results, i.e., Eqs. (2) and (3) as well as other parts of the paper, are valid.