Erratum: Probing multistrange dibaryons with proton-omega correlations in high-energy heavy ion collisions [Phys. Rev. C **94**, 031901(R) (2016)]

Kenji Morita , Akira Ohnishi, Faisal Etminan, and Tetsuo Hatsuda

(Received 25 November 2019; published 20 December 2019)

DOI: [10.1103/PhysRevC.100.069902](https://doi.org/10.1103/PhysRevC.100.069902)

The original paper is found to contain a typographical error in the binding energy of $p\Omega$ with the $V_{II} + \text{Coulomb}$ potential. The binding energy value of 6.3 MeV shown in Table I of the original paper should be corrected to be 0.63 MeV. The conclusions and other results are not affected.

TABLE I. Revised version of Table I in the original paper. The binding energy (E_B), the scattering length (a_0), and the effective range (r_{eff}) with and without the Coulomb attraction in the $p\Omega$ system. Physical masses of the proton and Ω are used.

Spin-2 $N\Omega$ Potentials		V_I	V_{II}	V_{III}
Without Coulomb	E_B (MeV)		0.05	24.8
	a_0 (fm)	−1.0	23.1	1.60
	r_{eff} (fm)	1.15	0.95	0.65
With Coulomb	E_B (MeV)		0.63	26.9
	a_0 (fm)	−1.12	5.79	1.29
	r_{eff} (fm)	1.16	0.96	0.65