Erratum: Dissecting reaction calculations using halo effective field theory and *ab initio* input
[Phys. Rev. C 98, 034610 (2018)]P. Capel , D. R. Phillips , and H.-W. Hammer

(Received 1 December 2021; published 25 January 2022)

DOI: [10.1103/PhysRevC.105.019901](https://doi.org/10.1103/PhysRevC.105.019901)

In the original paper, we have presented a new analysis of the breakup of one-neutron halo nuclei by coupling a model of the projectile motivated by halo effective field theory with a fully dynamical treatment of the reaction using the dynamical eikonal approximation. The agreement with data at intermediate energy on the archetypical halo nucleus ^{11}Be is excellent. It is independent of the cutoff in the effective ^{10}Be - n interaction used to describe the halo structure. The extension of this paper to other reactions [1,2] has shown there is a typographical error in Table I of the original paper: for the cutoff $\sigma = 2$ fm, the depth of the surface term of the next-to-leading order ^{10}Be - n interaction should read $V_{s1/2}^{(2)} = +3 \text{ MeV fm}^{-2}$. The units for that term in Tables I, II, IV, V, and VI should read MeV fm^{-2} . None of these changes affect the results presented in the paper.

We thank C. Hebborn and I. Alnamlah for bringing these issues to our attention and checking our results.

-
- [1] L. Moschini and P. Capel, *Phys. Lett. B* **790**, 367 (2019).
[2] C. Hebborn and P. Capel, *Phys. Rev. C* **104**, 024616 (2021).