Spinodal instabilities in asymmetric nuclear matter based on realistic NN interactions

E. N. E. van Dalen and H. Mütter

Institut für Theoretische Physik, Universität Tübingen, Auf der Morgenstelle 14, D-72076 Tübingen, Germany

(Received 15 August 2012; published 21 February 2013)

A density dependent relativistic mean-field model is determined to reproduce the components of the nucleon self-energy at low densities. This model is used to investigate spinodal instabilities in isospin asymmetric nuclear matter at finite temperatures. The inhomogeneous density distributions in the spinodal region are investigated through calculations in a cubic box with periodic boundary conditions. Compared to results obtained in phenomenological calculations the spinodal region is large, i.e., the spinodal region at zero temperature can reach densities above 0.12 fm^{-3} . The predicted spinodal region is strongly concentrated around isospin symmetric nuclear matter and the critical temperature is considerably lower than in the previous microscopic based investigation within a non-relativistic Brueckner-Hartree-Fock approach.

DOI: [10.1103/PhysRevC.87.024317](https://doi.org/10.1103/PhysRevC.87.024317)

PACS number(s): 21.30.Fe, 21.65.Cd, 24.10.Jv, 25.70.Pq

I. INTRODUCTION

The properties of isospin asymmetric nuclear matter systems like their instabilities and phase transitions are presently a topic of great interest [1–7]. At either high energy or high baryon density one explores the phase transition to quark matter, i.e., the hadrons dissolve into a phase of deconfined quarks. Below a critical temperature or density, the homogeneous fluid consisting of nucleons may condense into nuclei. This transition can be regarded to be of the liquid-gas type. Heavy ion collisions are well suited to investigate these transitions, since the multifragmentation phenomenon occurs in these experiments. In the multifragmentation phenomenon, highly excited composed nuclei are formed in a gas of evaporated particles. Therefore, this multifragmentation phenomenon can be interpreted as the coexistence of a liquid and a gas phase. This liquid-gas phase transition is important in the description of the nucleation processes. Furthermore, it plays an important role in the collapse of supernovae into neutron stars.

The hadrons involved in this transition can be either neutrons or protons. Therefore, one should consider nuclear matter as composed of two different fluids. These phase transitions in nuclear matter are related to the thermodynamic instability regions, which are limited by so-called spinodals. This spinodal instability is dominated by mechanical or isoscalar density fluctuations which lead to a liquid-gas phase separation with restoration of the isospin symmetry in the dense phase. This phenomenon is known as the isospin distillation or fragmentation effect [8,9], i.e., the formation of large droplets of high density symmetric nuclear matter within a neutron gas with only a very small fraction of protons.

In the past systematic analysis of the stability conditions of asymmetric nuclear matter against the liquid-gas phase transition have been performed using different methods. These approaches can roughly be divided into phenomenological

methods and microscopic based methods. Phenomenological studies of spinodal instabilities involve mean-field calculations with effective forces of the Skyrme or Gogny type [1,10–12] and relativistic mean-field calculations using constant and density dependent coupling parameters [3,4,7,13,14]. Some of these models predict that the total density at which spinodal instability appears will decrease if the asymmetry increases, whereas for other models it will increase up to large asymmetry and will finally decrease. However, a common feature of these models is that the spinodal region is in general situated at densities below $\rho_B < 0.1 \text{ fm}^{-3}$. The microscopic based studies are a calculation with low-density functionals [5] based on the Dirac-Brueckner-Hartree-Fock (DBHF) approach [15] with a realistic Bonn-A potential [16] and a calculation within a microscopic Brueckner-Hartree-Fock (BHF) approach [6] using the realistic Argonne V18 potential [17] plus a three-body force of Urbana type. Both calculations predict larger spinodal regions compared to those obtained in the phenomenological calculations, i.e., the spinodal region can reach densities above 0.12 fm^{-3} close to symmetric nuclear matter. The critical temperature in the BHF calculation is $T_c = 17.5 \text{ MeV}$, whereas finite temperature effects are not considered in the work of Ref. [5].

In the present work, we construct a density dependent relativistic mean-field (DDRMF) model, i.e., a density dependent relativistic Hartree (DDRH) model, which is based on microscopic DBHF calculations [18] using the realistic Bonn-A potential [16]. Apart from the application of the results of this DBHF calculation to spinodal instabilities [5], it has also already been used for other applications such as investigations of the nonhomogeneous structures in the neutron star crust [19], investigations of hybrid stars [20], and studies of nucleon-nucleus scattering [21]. In the previous DBHF based investigation of the spinodal instability [5], a functional is constructed on a density expansion of the relativistic mean-field theory with fixed coupling constants. However, in the present work the approximation of an expansion is not made and the effective coupling functions, which are obtained from the DBHF self-energy components, are density dependent. Special attention is devoted to reproduce the DBHF self-energy also at very low densities, i.e., $\rho_B < 0.10 \text{ fm}^{-3}$.

Published by the American Physical Society under the terms of the Creative Commons Attribution 3.0 License. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

In the framework of this DDRH theory based on a DBHF approach, the equation of state at zero and finite temperature can then be determined from an iteration procedure until convergence is reached. Within this model, the temperature effects and the nature of spinodal instabilities are investigated. In addition to these investigations in infinite nuclear matter calculations, the density distribution in the spinodal region is studied in cubic box calculations.

The paper is organized as follows. The density dependent relativistic Hartree (DDRH) theory is treated in Sec. II. The parametrization for the DDRH theory based on the DBHF results is discussed in Sec. III. Section IV is devoted to the theory of the spinodal instabilities. The results for the spinodal instabilities are presented and discussed in Sec. V. Finally, we end with a summary and the conclusion in Sec. VI.

II. DENSITY DEPENDENT RELATIVISTIC HARTREE THEORY

A Lagrangian density of an interacting many-particle system consisting of nucleons and mesons is the starting point of a DDRH theory. The Lagrangian density of the DDRH theory used for our investigation includes as well the isoscalar mesons σ and ω as the isovector mesons δ and ρ . Therefore, the Lagrangian density has the following parts: the free baryon Lagrangian density $\mathcal{L}_B$, the free meson Lagrangian density $\mathcal{L}_M$, and the interaction Lagrangian density $\mathcal{L}_{\text{int}}$:

$$\mathcal{L} = \mathcal{L}_B + \mathcal{L}_M + \mathcal{L}_{\text{int}}, \quad (1)$$

which can be written as

$$\begin{aligned} \mathcal{L}_B &= \bar{\Psi}(i\gamma_\mu \partial^\mu - M)\Psi, \\ \mathcal{L}_M &= \frac{1}{2} \sum_{i=\sigma,\delta} (\partial_\mu \Phi_i \partial^\mu \Phi_i - m_i^2 \Phi_i^2) \\ &\quad - \frac{1}{2} \sum_{\kappa=\omega,\rho} \left(\frac{1}{2} F_{(\kappa)\mu\nu} F_{(\kappa)}^{\mu\nu} - m_\kappa^2 A_{(\kappa)\mu} A_{(\kappa)}^\mu \right), \\ \mathcal{L}_{\text{int}} &= -g_\sigma \bar{\Psi} \Phi_\sigma \Psi - g_\delta \bar{\Psi} \boldsymbol{\tau} \boldsymbol{\Phi}_\delta \Psi - g_\omega \bar{\Psi} \gamma_\mu A_{(\omega)}^\mu \Psi \\ &\quad - g_\rho \bar{\Psi} \boldsymbol{\tau} \gamma_\mu \mathbf{A}_{(\rho)}^\mu \Psi. \end{aligned} \quad (2)$$

In the above Lagrangian density the field strength tensor for the vector mesons is defined as $F_{(\kappa)\mu\nu} = \partial_\mu A_{(\kappa)\nu} - \partial_\nu A_{(\kappa)\mu}$. The nucleon field is Ψ and the nucleon rest mass is M . The meson fields are denoted by $\Phi_\sigma, \boldsymbol{\Phi}_\delta, A_{(\omega)}$, and $\mathbf{A}_{(\rho)}$. Moreover, the bold symbols denote vectors in the isospin space. The mesons couple to the nucleons with the strength of the coupling constants $g_\sigma, g_\delta, g_\omega$, and g_ρ . The rest masses of these mesons are $m_\sigma, m_\omega, m_\delta$, and m_ρ .

By minimizing the action for variations of the fields $\bar{\Psi}$ included in the Lagrangian density of Eq. (1), one obtains the field equations for the nucleons,

$$\delta \int_{t_0}^{t_1} dt \int d^3x \mathcal{L}(\bar{\Psi}(x), \partial_\mu \bar{\Psi}(x), t) = 0. \quad (5)$$

Finally the following field equation can be derived:

$$\frac{\partial}{\partial x^\mu} \frac{\partial \mathcal{L}}{\partial (\partial_\mu \bar{\Psi})} - \frac{\partial \mathcal{L}}{\partial \bar{\Psi}} - \frac{\partial \mathcal{L}}{\partial \rho} \frac{\delta \rho}{\delta \bar{\Psi}} = 0. \quad (6)$$

Without density dependent meson-baryon vertices the third term vanishes in Eq. (6) and the normal Euler-Lagrange field equation is obtained. From the evaluation of Eq. (6), we obtain the Dirac equation

$$(i\gamma_\mu \partial^\mu - M - (\Sigma + \Sigma^{(r)}\gamma_0)) \Psi = 0, \quad (7)$$

with Σ the nucleon self-energy and $\Sigma^{(r)}$ the so-called rearrangement contribution to the nucleon self-energy.

The self-energy Σ for the Lagrangian density in Eqs. (1)–(4) can be written as

$$\Sigma = g_\sigma \Phi_\sigma + g_\delta \boldsymbol{\tau} \boldsymbol{\Phi}_\delta + g_\omega \gamma_\mu A_{(\omega)}^\mu + g_\rho \boldsymbol{\tau} \gamma_\mu \mathbf{A}_{(\rho)}^\mu. \quad (8)$$

This self-energy can be split into different parts with well-defined behavior under Lorentz transformations,

$$\Sigma(k) = \Sigma_s(k) - \gamma_0 \Sigma_o(k) + \boldsymbol{\gamma} \cdot \mathbf{k} \Sigma_v(k). \quad (9)$$

The Σ_s, Σ_o , and Σ_v components are Lorentz scalar functions. Hence, one can define the following effective quantities:

$$k^* = k(1 + \text{Re}\Sigma_v(k)), \quad (10)$$

$$m^*(k) = M + \text{Re}\Sigma_s(k), \quad (11)$$

and

$$E^*(k) = E(k) + \text{Re}\Sigma_o(k). \quad (12)$$

In the on-shell case, the effective energy can also be given by

$$E^*(k)^2 = k^{*2} + m^*(k)^2. \quad (13)$$

The contributions to the self-energy of the Lagrangian density presented in Eqs. (1)–(4) can be written as

$$\Sigma_{s,i}(k) = -\left(\frac{g_\sigma}{m_\sigma}\right)^2 (\rho_{s,n} + \rho_{s,p}) - \left(\frac{g_\delta}{m_\delta}\right)^2 \sum_{j=n,p} (\rho_{s,i} - \rho_{s,j})$$

and

$$\Sigma_{o,i}(k) = -\left(\frac{g_\omega}{m_\omega}\right)^2 (\rho_n + \rho_p) - \left(\frac{g_\rho}{m_\rho}\right)^2 \sum_{j=n,p} (\rho_i - \rho_j),$$

with

$$\rho_{s,i} = \frac{1}{\pi^2} \int f_i(k) k^2 dk \frac{m_i^*}{E_i(k)^*} \quad (14)$$

and

$$\rho_i = \frac{1}{\pi^2} \int f_i(k) k^2 dk. \quad (15)$$

The thermal occupation factors in Eqs. (14) and (15) are given by

$$f_i(k) = \frac{1}{1 + \exp[(E_i^* - \varepsilon_{F,i})/T]}, \quad i = p, n \quad (16)$$

in the “no-sea” approximation.

The $\Sigma^{(r)}$ term in Eq. (7) is generated by the third term in Eq. (6). It will only give a nonzero contribution, if meson-baryon vertices are density dependent. This rearrangement contribution $\Sigma^{(r)}$ can be written as

$$\begin{aligned} \Sigma^{(r)} &= \left(\frac{\partial g_\sigma}{\partial \rho} \bar{\Psi} \Phi_\sigma \Psi + \frac{\partial g_\delta}{\partial \rho} \bar{\Psi} \boldsymbol{\tau} \boldsymbol{\Phi}_\delta \Psi \right. \\ &\quad \left. + \frac{\partial g_\omega}{\partial \rho} \bar{\Psi} \gamma_\mu A_{(\omega)}^\mu \Psi + \frac{\partial g_\rho}{\partial \rho} \bar{\Psi} \boldsymbol{\tau} \gamma_\mu \mathbf{A}_{(\rho)}^\mu \Psi \right). \end{aligned} \quad (17)$$

TABLE I. Parameter set for the DDRH model from the DBHF approach in Ref. [18].

Meson i	m [MeV]	a_i	b_i	c_i	d_i	e_i
σ	550	14.0089064	9.91395378	-2.1599853	0.280160069	-13.3718958
ω	782.6	13.5288515	7.86182976	-1.8871752	0.244318038	-9.42372704
ρ	769	4.90660906	-14.9182243	3.93277073	-0.498567462	13.1945915
δ	983	-8.32733536	-43.6584053	10.4067049	-1.26465869	50.3025665

Rearrangement contributions are essential to provide a symmetry conserving approach [22,23], i.e., an approach satisfying energy-momentum conservation and thermodynamic consistency like the Hugenholtz-van Hove theorem.

III. PARAMETRIZATION

We have constructed a density dependent relativistic mean-field (DDRMF) model, i.e., a density dependent relativistic Hartree (DDRH) model. The effective density dependent coupling functions of this DDRH model are determined by the requirement that the results of a DBHF calculation at zero temperature using Bonn-A [18] for the components of the nucleon self-energy at the Fermi momentum are reproduced. The rearrangement terms of DDRH are not included in the fit, since the DBHF approach has no rearrangement contributions.

The density dependence of the coupling functions are parametrized by

$$g_i(\rho_B) = a_i + b_i x + c_i x^2 + d_i x^3 + e_i \sqrt{x}, \quad (18)$$

with $x = \rho_B/\rho_0$ and $\rho_0 = 0.16 \text{ fm}^{-3}$. The DBHF data at densities of $\rho_B = 0.017, 0.100, 0.197, 0.313$, and 0.467 fm^{-3} at a proton fraction of $Y_p = 0.4$ are used to obtain these parameters of our DDRH model. The meson masses are chosen to be identical to those of the Bonn-A potential. All parameters are given in Table I.

The density dependence of these coupling functions is displayed in Fig. 1. The density dependence is rather weak except at very low densities. The strength for the isovector mesons (ρ and δ) is significantly smaller than for the isoscalar

FIG. 1. (Color online) Density dependence of the coupling functions of the σ meson (solid), the ω meson (dotted), the ρ meson (dashed), and the δ meson (dashed-dotted).

mesons (σ and ω). All coupling functions tend to decrease with increasing density. This may reflect the quenching of the DBHF G matrix underlying this fit due to Pauli and dispersive effects in the Bethe-Goldstone equation.

In Fig. 2 the binding energy of the DDRH model is shown for neutron matter and symmetric nuclear matter. The results are in reasonable agreement with the DBHF results. Note, that the DDRH parametrization has solely been based on the DBHF self-energies at the corresponding Fermi momenta for a proton fraction $Y_p = 0.4$. No attempt has been made to account for the momentum and further asymmetry dependence of the DBHF self-energies, which enter the calculated energies for symmetric nuclear matter and neutron matter. Both approaches show a nonlinear convergence to zero when the density decreases which is in qualitative agreement with a study of low-density nuclear matter [24] based on a virial expansion that includes protons, neutrons, and α -particle degrees of freedom. In particular at densities below $\rho < 0.1 \text{ fm}^{-3}$, the DDRH model presented here is in better agreement with the DBHF result than those models constructed in Ref. [25].

From Table II, one can see that the saturation density ρ_{sat} is shifted to lower densities and the binding energy E/A of the saturation point is weaker in the DDRH model compared to the original DBHF results. The compression modulus K is somewhat smaller and symmetry energy a_s is in a fairly good agreement with the DBHF results.

IV. SPINODAL INSTABILITIES

The stability conditions for isospin asymmetric nuclear matter at finite temperatures are obtained from the free energy density $\mathcal{F}$. This free energy density is defined as

$$\mathcal{F} = \mathcal{E} - TS, \quad (19)$$

where $\mathcal{E}$ is the energy density and S denotes the entropy density. The energy density $\mathcal{E}$ is obtained from the

TABLE II. Saturation properties of nuclear matter for the DDRH model compared to the DBHF results of Refs. [15,18]. The quantities listed include the saturation density ρ_{sat} , the binding energy E/A , the compressibility modulus K , and the symmetry energy a_s . The symmetry energy a_s has been obtained assuming a quadratic dependence of the EOS on the asymmetry parameter $\beta = (\rho_n - \rho_p)/(\rho_n + \rho_p)$.

		DDRH	DBHF
ρ_{sat}	[fm^{-3}]	0.1759	0.181
E/A	[MeV]	-14.39	-16.15
K	[MeV]	193	230
a_s	[MeV]	35.03	34.36

FIG. 2. (Color online) Binding energy is plotted as a function of the density for the DBHF approach and the DDRH model. (a) Isospin symmetric nuclear matter. (b) Pure neutron matter.

energy-momentum tensor

$$\mathcal{E} = \langle T^{00} \rangle = \frac{1}{\pi^2} \sum_{i=p,n} \int f_i(k) k^2 dk \sqrt{k^2 + m_i^{*2}} - \frac{1}{2} \sum_{i=p,n} (\Sigma_{s,i} \rho_{s,i} + \Sigma_{o,i} \rho_i) - M\rho, \quad (20)$$

where $f_i(k)$ is the thermal occupation probability. The rearrangement term $\Sigma^{(r)}$ does not appear in Eq. (20), since rearrangement does not contribute to the energy density. However, note that the rearrangement does modify other quantities such as the pressure. Furthermore, the entropy density $\mathcal{S}$ is given by

$$\mathcal{S} = -\frac{1}{\pi^2} \sum_{i=p,n} \int f_i(k) k^2 dk [f_i(k) \ln(f_i(k)) + (1 - f_i(k)) \ln(1 - f_i(k))]. \quad (21)$$

The necessary ingredients to analyze the stability of isospin asymmetric nuclear matter against separation into two phases are now introduced. The nuclear system will be stable if the free energy of a single phase is lower than the free energy in any two-phase configurations. This stability condition implies that the free energy density is a convex function of the neutron and proton densities ρ_n and ρ_p . This means that the curvature matrix,

$$[\mathcal{F}_{ij}] = \left[\frac{\partial^2 \mathcal{F}}{\partial \rho_i \partial \rho_j} \right]_T, \quad (22)$$

is positive-definite and has two positive eigenvalues $\lambda^\pm$, which are associated to eigenvectors $\delta\rho^\pm$. In the spinodal region, the λ^- can become negative and the eigenvector associated with this negative eigenvalue indicates the direction of the instability. Below we characterize this direction by the ratio

$$\frac{\delta\rho_p^-}{\delta\rho_n^-} = \frac{\mathcal{F}_{np}}{\lambda^- - \mathcal{F}_{pp}} = \frac{\lambda^- - \mathcal{F}_{nn}}{\mathcal{F}_{np}}. \quad (23)$$

V. RESULTS AND DISCUSSION

In the present work, we have investigated the spinodal region in the case of isospin asymmetric nuclear matter at finite temperature. The projection of the spinodal contour in the neutron-proton density plane for several temperatures ranging from $T = 0$ to $T = 10$ MeV is depicted in Fig. 3. Keep in mind that the isospin symmetry of the nucleon-nucleon interaction guarantees the symmetry of the spinodal contours with respect to the interchange of the neutron density ρ_n and the proton density ρ_p .

Let us first consider the results for zero temperature ($T = 0$ MeV). One finds that the spinodal region extends up to a maximum density of 0.127 fm^{-3} . This is considerably larger than the maximum density obtained in phenomenological

FIG. 3. (Color online) The projection of the spinodal contour in the neutron-proton density plane at several temperatures. The arrows at the boundary of the spinodal region indicate the direction of instability for $T = 6$ MeV.

models [1–4,7,10–14], which indicate that the spinodal region is situated at densities below $\rho_B < 0.1 \text{ fm}^{-3}$. The large spinodal region, however, is in agreement with microscopic calculations [5,6]. This difference between the phenomenological and the microscopic based models is of course related to the larger saturation densities, which are obtained in the microscopic calculations. Note, however, that the present study yields a larger maximal density than the nonrelativistic BHF calculations of Vidana and Polls [6], although the saturation density for the present approach ($\rho_0 = 0.176 \text{ fm}^{-3}$) is smaller than the one obtained in the BHF calculation ($\rho_0 = 0.182 \text{ fm}^{-3}$). Therefore, our value of the ratio between the maximum density ρ_{max} of spinodal zone and the saturation density ρ_0 is higher ($\rho_{\text{max}}/\rho_0 = 0.72$) than that in the nonrelativistic BHF calculation of Vidana and Polls [6] with $\rho_{\text{max}}/\rho_0 \approx 0.66$. The ratio ρ_{max}/ρ_0 obtained in the present study is also larger than the corresponding ratios obtained in the phenomenological models presented in Ref. [3] which exhibit ratios between 0.62 and 0.68.

The shape of the spinodal region for $T = 0$ projected in the neutron-proton density plane, is similar to the shapes obtained in calculations before. In highly asymmetric nuclear matter at low temperatures the spinodal contour starts to run parallel to the axis, which happens in our calculation near $\rho_{n/p} = 0.06 \text{ fm}^{-3}$. In detail, however, there is a difference: The region of instability is more confined to isospin symmetric nuclear matter. Furthermore, due to this fact we have a more abrupt change compared to most other calculations, when the contour starts to run parallel to the axis.

Also the dependence of the region of instability on temperature in Fig. 3 is very similar to the one observed in previous calculations [1–4,6,7,10–14]. The spinodal region shrinks with increasing temperature, until the critical temperature T_c is reached. In the present approach no spinodal region can be obtained above $T = 11 \text{ MeV}$. Therefore, the critical temperature is considerably lower than the value $T_c = 17.5 \text{ MeV}$ obtained in the other microscopic based calculation [6] or the values for T_c reported in the phenomenological studies [4]. It is worth mentioning that the spinodal region obtained here at temperatures close to T_c occurs at slightly higher densities than in the other studies.

It has been shown that the spinodal instabilities in isospin asymmetric nuclear matter appear as a mixture of isoscalar and isovector instabilities [1,2]. Isoscalar or mechanical instabilities are related to density fluctuations. Pure isoscalar instabilities conserve the proton fraction and fulfill the relation $\delta\rho_p^-/\delta\rho_n^- = 1$. On the other hand, pure isovector or chemical instabilities are related to fluctuations in the proton fraction at fixed density and fulfil the relation $\delta\rho_p^-/\delta\rho_n^- = -1$. Therefore, the instabilities will be predominantly of the isoscalar nature, if the neutrons and protons move in phase $\delta\rho_p^-/\delta\rho_n^- > 0$, or of the isovector nature, if they move out of phase $\delta\rho_p^-/\delta\rho_n^- < 0$. In Fig. 4 the results obtained from our model for this ratio $\delta\rho_p^-/\delta\rho_n^-$ is plotted as a function of the asymmetry parameter $\beta = (\rho_n - \rho_p)/(\rho_n + \rho_p)$ for several temperatures ranging from $T = 0$ up to $T = 8 \text{ MeV}$ and for two different densities. It can be observed from Fig. 4 that the ratio $\delta\rho_p^-/\delta\rho_n^-$ is positive in all considered cases. This result implies that the spinodal instability is dominated by mechanical or

isoscalar density fluctuations, which is in agreement with previous investigations [1,2,5,6]. Since $\delta\rho_p^-/\delta\rho_n^-$ is larger than the ratio ρ_p/ρ_n presented by the black dotted line in Fig. 4, the dense liquid phase of the system is driven towards a more isospin symmetric composition and the gas phase will be more isospin asymmetric. These effects are less strong for higher temperatures. The liquid phase is in general also less strongly driven towards a isospin symmetric composition, when the density ρ_B or the asymmetry parameter β increase.

Up to this point we have determined the spinodal zone from the evaluation of the curvature matrix $[\mathcal{F}_{ij}]$ in Eq. (22) for the free energy of the homogeneous system of nuclear matter. Having identified the regions of instability we will now try to verify these instabilities and determine the density profile in this area.

For that purpose we consider the very same DDRH and perform variational calculations in a cubic box with periodic boundary conditions. The size and shape of the box should be determined in a variational manner, since the results will depend to some extent on the box-size chosen. In nuclear matter studies, ignoring the Coulomb repulsion of the protons, one cannot expect a periodic structure, as suggested by the periodic boundary condition, as the system will prefer to generate for each box size one part in the liquid phase, embedded in a gas phase. However, in a kind of explorative study we have chosen the length of the cubic box to be 20 fm in each direction with the origin of the coordinate system in the center of the box. Also we have restricted the study to isospin symmetric systems. In order to be consistent with our infinite nuclear matter calculations, the Coulomb energy is not included in the box calculations. Therefore the proton density distributions are identical to the corresponding neutron ones displayed in Fig. 5.

The transition from isolated nuclei to a phase of homogeneous baryon matter can be observed in the various panels

FIG. 4. (Color online) Ratio $\delta\rho_p^-/\delta\rho_n^-$ as a function of asymmetry parameter $\beta = (\rho_n - \rho_p)/(\rho_n + \rho_p)$ for several temperatures. (a) The results for a fixed density of $\rho_B = 0.03 \text{ fm}^{-3}$. (b) The results for a fixed density of $\rho_B = 0.06 \text{ fm}^{-3}$. The ratio ρ_p/ρ_n is given by the black dotted line.

FIG. 5. (Color online) Neutron density distributions at $T = 0$ MeV resulting from box calculations for isospin symmetric nuclear matter as a function of the Cartesian coordinates x, y , and z . The panels refer to densities of $\rho = 0.005 \text{ fm}^{-3}$, 0.05 fm^{-3} , 0.10 fm^{-3} , and 0.125 fm^{-3} .

of Fig. 5 for zero temperature. Figure 5 a shows the density distribution at a density of $\rho = 0.005 \text{ fm}^{-3}$, i.e., only 20 protons and 20 neutrons (^{40}Ca) are present in the cubic box. At these low densities, isolated nuclei will be present. The density profiles are identical in all three Cartesian directions, which corresponds to crystal of spherical nuclei located at the centers of the cubic box.

However, at higher densities this is not the case anymore. At a density of $\rho = 0.05 \text{ fm}^{-3}$, the density profile in z direction deviates a little from that in the x and y direction. Furthermore, one has a high-density region at the center of the cubic box and a low-density region along the connection lines to the nearest neighbor at the boundary of the box.

However, the density distribution is not as simple as suggested by Fig. 5(b). This can be seen from the complete density distribution in the $z = 0$ plane as displayed in Fig. 6(b). Apart from the low density regions at the boundary of the box along the connection lines between the centers of the box, one has also regions in which the density is almost zero. Therefore, the geometrical structure at this density can be described by large quasinuclei located in the centers of the cubic box, which are connected to each other by low-density bridges.

At even higher densities, i.e., around $\rho = 0.10 \text{ fm}^{-3}$, the opposite density distribution can be observed, i.e., isolated low-density regions completely surrounded by high-density matter. At a density of $\rho = 0.125 \text{ fm}^{-3}$, so just a little bit below the critical density ($\rho_c = 0.127 \text{ fm}^{-3}$) predicted from the study of the curvature matrix in the infinite nuclear matter calculation, one obtains a grid structure with small fluctuations, which resembles homogeneous nuclear matter.

Therefore, within the numerical accuracy one can regard this result to be in agreement with the result of the infinite nuclear matter calculation in Fig. 3.

Performing DDRH calculations in the same kind of cubic box at the finite temperature $T = 8$ MeV, we obtain the neutron

FIG. 6. (Color online) Neutron density distribution in fm^{-3} at $T = 0$ MeV in the $z = 0$ plane resulting from box calculations for isospin symmetric nuclear matter. The panels refer to densities of $\rho = 0.005 \text{ fm}^{-3}$, 0.05 fm^{-3} , 0.10 fm^{-3} , and 0.125 fm^{-3} .

FIG. 7. (Color online) Neutron density distribution in fm^{-3} at $T = 8$ MeV in the $z = 0$ plane resulting from box calculations for isospin symmetric nuclear matter. The panels refer to densities of $\rho = 0.05 \text{ fm}^{-3}$, 0.10 fm^{-3} , and 0.125 fm^{-3} .

density distributions displayed in Fig. 7. It is clear that finite temperature effects tend to dissolve the geometrical structures observed for $T = 0$ MeV. The structures at $T = 8$ MeV are less pronounced than those for $T = 0$ MeV in Fig. 6 and the transition density is lower at $T = 8$ MeV. These results are in agreement with the results presented in Fig. 3 of the infinite nuclear matter calculations.

VI. SUMMARY AND CONCLUSION

In the present work, we construct a density dependent relativistic mean field model (DDRH) to parametrize the results of microscopic Dirac Brueckner Hartree Fock (DBHF) calculations [18] using the realistic Bonn-A potential [16]. Special attention is devoted to reproduce the DBHF self-energy in particular at densities below the saturation density. The equation of state at zero and finite temperature can then be determined in the framework of this DDRH theory for homogeneous matter as well as inhomogeneous structures. The spinodal instabilities of homogeneous matter including their temperature effects are investigated within this model.

Our investigations confirm the main results of earlier studies based on phenomenological [1,2] as well as realistic models of the NN interaction [6], that the spinodal instability is dominated by mechanical or isoscalar density fluctuations which lead to a liquid-gas phase separation with restoration of the isospin symmetry in the liquid phase. However, this restoration is less strong for higher temperatures.

Furthermore, the DDRH model predicts some details, which are different from earlier studies: A large spinodal region, which can reach densities above 0.12 fm^{-3} at zero temperature. This is quite different from the predictions of the phenomenological calculations [1–4,7,10–14], but in line with the microscopic studies of Vidana and Polls [6]. It can to some extent be attributed to the differences in the predicted saturation point of nuclear matter. Differences are also observed with respect to finite temperature and details of the spinodal region. The present DDRH approach yields a smaller critical temperature (T_c less than 11 MeV) and higher densities for the spinodal region at temperatures just below T_c . The predicted spinodal region obtained from our calculation is more strongly concentrated around isospin symmetric nuclear matter.

The studies of the spinodal regions which are based on the curvature of the free energy of homogeneous matter are supplemented by variational calculations allowing for density fluctuations explicitly. Calculations employing the same DDRH model are performed in a cubic box with periodic boundary conditions by minimizing the total energy to obtain density distributions in isospin symmetric nuclear matter. The results of these calculations in a finite box must be considered with some caution, as details depend on the box size considered. Nevertheless for a box size of 20 fm in each direction a geometrical structure resembling homogeneous nuclear matter can be found at a density of 0.125 fm^{-3} and larger densities. At lower densities, nonhomogeneous nuclear

matter structures occur. Finite temperature effects tend to dissolve the geometrical structures observed, i.e., the structures at finite temperature are less pronounced and the transition density to homogeneous nuclear matter is lower.

ACKNOWLEDGMENTS

This work has been supported by the Deutsche Forschungsgemeinschaft (DFG) under Contract no. Mu 705/7-1.

-
- [1] J. Margueron and P. Chomaz, *Phys. Rev. C* **67**, 041602(R) (2003).
 - [2] P. Chomaz, M. Colonna, and J. Randrup, *Phys. Rep.* **389**, 263 (2004).
 - [3] S. S. Avancini, L. Brito, D. P. Menezes, and C. Providência, *Phys. Rev. C* **70**, 015203 (2004).
 - [4] S. S. Avancini, L. Brito, Ph. Chomaz, D. P. Menezes, and C. Providência, *Phys. Rev. C* **74**, 024317 (2006).
 - [5] J. Margueron, E. van Dalen, and C. Fuchs, *Phys. Rev. C* **76**, 034309 (2007).
 - [6] I. Vidana and A. Polls, *Phys. Lett. B* **666**, 232 (2008).
 - [7] H. Pais, A. Santos, and C. Providência, *Phys. Rev. C* **80**, 045808 (2009).
 - [8] J. Randrup and S. E. Koonin, *Nucl. Phys. A* **356**, 223 (1981).
 - [9] H. S. Xu *et al.*, *Phys. Rev. Lett.* **85**, 716 (2000).
 - [10] C. Ducoin, Ph. Chomaz, and F. Gulminelli, *Nucl. Phys. A* **771**, 68 (2006).
 - [11] C. Ducoin, Ph. Chomaz, and F. Gulminelli, *Nucl. Phys. A* **781**, 407 (2007).
 - [12] C. Ducoin, Ph. Chomaz, and F. Gulminelli, *Nucl. Phys. A* **789**, 403 (2007).
 - [13] B. Liu, V. Greco, V. Baran, M. Colonna, and M. Di Toro, *Phys. Rev. C* **65**, 045201 (2002).
 - [14] C. Providência, L. Brito, S. S. Avancini, D. P. Menezes, and Ph. Chomaz, *Phys. Rev. C* **73**, 025805 (2006).
 - [15] E. N. E. van Dalen, C. Fuchs, and A. Faessler, *Nucl. Phys. A* **744**, 227 (2004).
 - [16] R. Machleidt, K. Holinde, and Ch. Elster, *Phys. Rep.* **149**, 1 (1987).
 - [17] R. B. Wiringa, V. G. J. Stoks, and R. Schiavilla, *Phys. Rev. C* **51**, 38 (1995).
 - [18] E. N. E. van Dalen, C. Fuchs, and A. Faessler, *Eur. Phys. J. A* **31**, 29 (2007).
 - [19] P. Gögelein, E. N. E. van Dalen, C. Fuchs, and H. Mütter, *Phys. Rev. C* **77**, 025802 (2008).
 - [20] T. Klahn *et al.*, *Phys. Lett. B* **654**, 170 (2007).
 - [21] R. Xu, Z. Ma, E. N. E. van Dalen, and H. Mütter, *Phys. Rev. C* **85**, 034613 (2012).
 - [22] C. Fuchs, H. Lenske, and H. H. Wolter, *Phys. Rev. C* **52**, 3043 (1995).
 - [23] E. N. E. van Dalen and H. Mütter, *Int. J. Mod. Phys. E* **19**, 2077 (2010).
 - [24] C. J. Horowitz and A. Schwenk, *Nucl. Phys. A* **776**, 55 (2006).
 - [25] E. N. E. van Dalen and H. Mütter, *Phys. Rev. C* **84**, 024320 (2011).