**Publisher's Note: Properties of γ -decaying isomers and isomeric ratios in the ^{100}Sn region
[Phys. Rev. C **96**, 044311 (2017)]**

J. Park, R. Krücken, D. Lubos, R. Gernhäuser, M. Lewitowicz, S. Nishimura, D. S. Ahn, H. Baba, B. Blank, A. Blazhev, P. Boutachkov, F. Browne, I. Čeliković, G. de France, P. Doornenbal, T. Faestermann, Y. Fang, N. Fukuda, J. Giovinazzo, N. Goel, M. Górska, H. Grawe, S. Ilieva, N. Inabe, T. Isobe, A. Jungclaus, D. Kameda, G. D. Kim, Y.-K. Kim, I. Kojouharov, T. Kubo, N. Kurz, G. Lorusso, K. Moschner, D. Murai, I. Nishizuka, Z. Patel, M. M. Rajabali, S. Rice, H. Sakurai, H. Schaffner, Y. Shimizu, L. Sinclair, P.-A. Söderström, K. Steiger, T. Sumikama, H. Suzuki, H. Takeda, Z. Wang, H. Watanabe, J. Wu, and Z. Y. Xu

(Received 13 October 2017; published 19 October 2017)

DOI: [10.1103/PhysRevC.96.049901](https://doi.org/10.1103/PhysRevC.96.049901)

This paper was published online on 9 October 2017 with a typographical error on page 6. On page 6, the last sentence of the right-hand column should read as “The revised level scheme was presented in previous works [51,52] supported by SM calculations employing the “SR88MHJM” interaction [6,32]; this model includes final monopole corrections to the “SR88MHJ” interaction [64], and was used to calculate the $B(E2)$ value of the 107-keV transition.” The paper has been corrected as of 16 October 2017. The text is correct in the printed version of the journal.