Erratum: Neutron-unbound excited states of ^{23}N [Phys. Rev. C **95, 044323 (2017)]**

M. D. Jones, T. Baumann, J. Brett, J. Bullaro, P. A. DeYoung, J. E. Finck, N. Frank, K. Hammerton, J. Hinnefeld, Z. Kohley, A. N. Kuchera, J. Pereira, A. Rabeh, J. K. Smith, A. Spyrou, S. L. Stephenson, K. Stiefel, M. Tuttle-Timm, R. G. T. Zegers, and M. Thoennessen

(Received 9 October 2017; published 15 November 2017)

DOI: [10.1103/PhysRevC.96.059902](https://doi.org/10.1103/PhysRevC.96.059902)

We have found that the target was modeled incorrectly with beryllium rather than liquid deuterium, which resulted in an incorrect energy loss and resolution. The corrected simulations and resolution are shown in Fig. 1. The correct resolution is 60% higher than previously reported at 1 MeV with a FWHM of 1 MeV. The best-fit parameters remain unchanged, and the conclusions are unaffected.

FIG. 1. Two-body decay energy for $^{22}\text{N} + 1n$. The best fit includes two-channel Breit-Wigner distributions resulting from two states at 1.1 MeV (dashed-red curve) and 2.4 MeV (dot-dashed-blue curve). Background contributions are in shaded gray. The efficiency and resolution are shown in the inset as the blue histogram (left scale) and red-dashed line (right scale), respectively.