Three-dimensional superconformal quantum mechanics with $sl(2|1)$ dynamical symmetry

Ivan E. Cunha^{*} and Francesco Toppan[†]

CBPF, Rua Dr. Xavier Sigaud 150, Urca, cep 22290-180 Rio de Janeiro (RJ), Brazil

(Received 2 September 2019; published 2 December 2019)

We construct a three-dimensional superconformal quantum mechanics (and its associated de Alfaro-Fubini-Furlan deformed oscillator) possessing an $sl(2|1)$ dynamical symmetry. At a coupling parameter $\beta \neq 0$, the Hamiltonian contains a $\frac{1}{r^2}$ potential and a spin-orbit (hence, a first-order differential operator) interacting term. At $\beta = 0$, four copies of undeformed three-dimensional oscillators are recovered. The Hamiltonian gets diagonalized in each sector of total j and orbital l angular momentum (the spin of the system is $\frac{1}{2}$). The Hilbert space of the deformed oscillator is given by a direct sum of $sl(2|1)$ lowest-weight representations. The selection of the admissible Hilbert spaces at given values of the coupling constant β is discussed. The spectrum of the model is computed. The vacuum energy (as a function of β) consists of a recursive zigzag pattern. The degeneracy of the energy eigenvalues grows linearly up to $E \sim \beta$ (in proper units) and quadratically for $E > \beta$. The orthonormal energy eigenstates are expressed in terms of the associated Laguerre polynomials and the spin spherical harmonics. The dimensional reduction of the model to $d = 2$ produces two copies (for β and $-\beta$, respectively) of the two-dimensional $sl(2|1)$ deformed oscillator. The dimensional reduction to $d = 1$ produces the one-dimensional $D(2, 1; \alpha)$ deformed oscillator, with α determined by β .

DOI: [10.1103/PhysRevD.100.125002](https://doi.org/10.1103/PhysRevD.100.125002)

I. INTRODUCTION

In this paper, we construct a three-dimensional superconformal quantum mechanics having, as an input, an $sl(2|1)$ dynamical symmetry. The associated (denoted as “DFF”; see [1]) de Alfaro-Fubini-Furlan type of deformed oscillator, with $sl(2|1)$ as spectrum-generating superalgebra, is presented in detail. These quantum models (both the superconformal and the DFF one) are defined in terms of a dimensionless deformation parameter β which, without the loss of generality, can be assumed to belong to the $\beta \geq 0$ interval. The $\beta = 0$ undeformed DFF Hamiltonian corresponds to four copies of the three-dimensional isotropic oscillator; when $\beta > 0$, a spin-orbit term enters the Hamiltonians. It is a first-order differential operator which can be diagonalized in each sector of given total and orbital angular momenta.

The list of the main results is the following:

- (i) Once derived from the $sl(2|1)$ lowest-weight representations, alternative admissible Hilbert spaces

that can be associated with the quantum models at a given β are constructed. Consistency conditions require the wave functions to be normalized and the Hamiltonian to be self-adjoint. The procedure is an extension of the approach and results discussed in Refs. [2,3] for conformal quantum mechanics. The spectrum of the deformed oscillators is computed. When varying the deformation parameter β (which, in physical applications, can play the role of an external control parameter), certain lowest-weight representations of $sl(2|1)$ can be “switched on” as admissible in the given Hilbert space, while certain other lowest-weight representations can be “switched off.” One of the consequences is the production, for the vacuum energies, of the recursive zigzag patterns observed in Figs. 1 and 2.

- (ii) An unexpected feature concerning the degeneracy of the energy eigenvalues of the deformed oscillator is derived. For β integer or half-integer, the energy spectrum is a shifted version of the spectrum of the undeformed oscillator. The β -deformed oscillator realizes an interpolation between two different regimes. Up to energy $E \sim \beta$ [measured in natural units, by setting $\Delta(E) = 1$, where $\Delta(E) = E_1 - E_{\text{vac}}$ is the energy difference between the first excited state and the vacuum], the degeneracy grows linearly, mimicking the behavior of a two-dimensional oscillator; starting from $E \sim \beta$, the degeneracy grows

^{*}ivanec@cbpf.br

[†]toppan@cbpf.br

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

FIG. 1. The vacuum energy $E_{\text{vac}}(\beta)$ of the model is portrayed in the y axis, with β up to $\beta \leq 5$ depicted in the x axis. This diagram refers to the Hilbert space admitting singular but normalized wave functions at the origin. Starting from $\beta > \frac{1}{2}$, the graph is composed by a triangle wave of half-open line segments plus isolated points at $\beta = \frac{1}{2} + \mathbb{N}$. From $\beta \geq \frac{1}{2}$, the models with β separated by an integer number and, therefore, with identical $\mu = \{\beta - \frac{1}{2}\}$ possess the same spectrum but different degeneracies of their eigenvalues. For $\mu \neq 0, \frac{1}{2}$, the same spectrum is also recovered (with different degeneracies) under the $\mu \leftrightarrow 1 - \mu$ duality. The vacuum energy $E_{\text{vac}}(\beta)$ is always positive [$E_{\text{vac}}(\beta) > 0$].

quadratically [we recall that the degeneracy $d(n)$ of the n th excited energy eigenvalue of an ordinary D -dimensional oscillator grows as $d(n) \propto n^{D-1}$]. This behavior has been computed in Sec. VI and visually presented in Fig. 3.

- (iii) The orthonormal eigenstates are expressed in terms of two classical functions: the associated Laguerre polynomials dependent on the (square of) the radial

FIG. 2. The vacuum energy $E_{\text{vac}}(\beta)$ of the model is portrayed in the y axis, with β up to $\beta \leq 5$ depicted in the x axis. This diagram refers to the Hilbert space satisfying the condition that its wave functions are regular at the origin. For $\beta > 0$, the vacuum energy is always comprised in the interval $\frac{3}{2} < E_{\text{vac}}(\beta) \leq \frac{5}{2}$. The isolated points are now encountered at integer values of β for $\beta = 2, 3, 4, \dots$. The features of this diagram are discussed in the main text.

FIG. 3. The energy degeneracy (y axis) for the $\mathcal{H}_{\text{norm}}$ Hilbert space at the integer values $\beta = 0, 2, 6, 16$. In the x axis are reported the 40 lowest energy eigenvalues. The “•” bullet denotes the $\beta = 0$ undeformed oscillator, while “-,” “*,” and “+” stand, respectively, for the $\beta = 2, 6$, and 16 cases. One can note the “bending” of the $\beta = 16$ curve around energy $E = 16$.

coordinate and the spin spherical harmonics (see [4]) dependent on the angular coordinates. This result has been derived in Appendix B; see formulas (B22) and (B24).

Besides the above points, further discussed features are the implementation of superselection rules, the recovery of lower-dimensional deformed oscillators via dimensional reduction, and so on.

Conformal quantum mechanics was first introduced in Ref. [5]. The superconformal extension was presented in Refs. [6,7]. Superconformal quantum mechanics and its associated DFF deformed oscillators are a very active field of investigation with a growing body of literature. There are two main motivations for that: on one side, the development of sophisticated mathematical tools (e.g., for large $\mathcal{N}$ -extended supersymmetry, the role of superconformal Lie algebras) to construct and explicitly solve models, both at the classical and quantum level, and, on the physical side, for its important applications. We mention, in particular, the motion of test particles in the proximity of the horizon of certain black holes; see [8] and the $\text{AdS}_2/\text{CFT}_1$ correspondence investigated in Refs. [9,10]. The AdS_2 holography has recently gained new attention in relation with the Sachdev-Ye-Kitaev models (see [11,12], and references therein); in this context, the connection between conformal and Schwarzian mechanics is discussed, e.g., in Refs. [13,14].

In the literature, there are three main approaches to construct superconformal quantum mechanics. The most popular one consists in quantizing classical world-line superconformal sigma models defined on supermultiplets of an $\mathcal{N}$ -extended one-dimensional supersymmetry [for $\mathcal{N} = 1, 2, 4, 8$, the relevant supermultiplets (see [15,16]) are of the type $(D, \mathcal{N}, \mathcal{N} - D)$, corresponding to D propagating bosonic fields, $\mathcal{N}$ fermionic fields, and $\mathcal{N} - D$ auxiliary bosonic fields]. In the sigma-model

interpretation, D is the dimensionality of the target manifold. These classical superconformal world-line sigma models are constructed either via superspace (see the review [17] and the references therein; more recent works are Refs. [18–21]) or via D -module representations of one-dimensional superconformal algebras, as in Refs. [22,23].

The so-called “triangular representations” (in contraposition to the ordinary “parabolic representations”) of superconformal algebras have been discussed in Ref. [24]. They induce the de Alfaro-Fubini-Furlan deformed oscillators counterparts of the superconformal mechanics. Some recent works on superconformal quantum mechanics, either “parabolic” or “trigonometric” cases, are Refs. [25–30]. Particularly relevant for our purposes here is the paper in Ref. [26]. There, it is shown that the quantization of world-line superconformal sigma models with $D \geq 3$ target dimensions cannot be performed straightforwardly but require solving nontrivial nonlinear equations. It is due to this obstruction that we apply here a more direct method (the second approach, pioneered in Refs. [28,30] for one-dimensional models) to construct a three-dimensional superconformal quantum mechanics. It is rewarding that the dimensional reductions of the three-dimensional superconformal quantum mechanics allow one to recover (see Sec. VII) the models obtained in Ref. [26] by quantizing the world-line superconformal sigma models based on the $\mathcal{N} = 4$ (1,4,3) and the $\mathcal{N} = 2$ (2,2,0) supermultiplets (for target dimensions $D = 1, 2$, respectively). One sign of the obstruction for $D = 3$ is the appearance in the Hamiltonian of the nondiagonal spin-orbit term.

The third approach to superconformal quantum mechanics is based on symmetries of the time-dependent Schrödinger equation, regarded as a partial differential equation (it will be briefly discussed in the conclusions).

We mention that several different, both classical and quantum, supersymmetric models possessing (a real form of) $sl(2|1)$ as dynamical symmetry have been investigated in the literature; see [25,26,29,31–34]. These models do not correspond to the three-dimensional Hamiltonians here presented. The first one of these papers, Ref. [31], presents the $sl(2|1)$ dynamical symmetry of a Dirac magnetic monopole with a $\frac{1}{r^2}$ potential.

Hamiltonians with a spin-orbit coupling, as the one here discussed, are not a novelty. It has to be mentioned, in particular, the paper in Ref. [35]. In that work, a particular 2×2 matrix Hamiltonian with spin-orbit coupling has been solved by showing that the system possesses an $osp(1|2)$ dynamical symmetry. The restriction to the upper-left block of the 4×4 matrix Hamiltonian (9) derived below produces, at the specific $\beta = \frac{1}{2}$ value, the Hamiltonian in formula (2.6) of Ref. [35] [a further constant λ , entering the (2.6) Hamiltonian, is set to zero]. Since the β coupling constant entering [35] is kept fixed, from that work no features can be derived concerning the zigzag pattern of vacuum energy or the interpolating regimes obtained at varying β .

The scheme of the paper is the following. In Sec. II, we introduce the three-dimensional superconformal quantum mechanics. In Sec. III, we apply the de Alfaro-Fubini-Furlan “trick” to construct the associated deformed oscillator with $sl(2|1)$ spectrum-generating superalgebra. The selection of its admissible Hilbert spaces is discussed in Sec. IV. The derivation of its spectrum is given in Sec. V. The computation of the energy degeneracy is presented in Sec. VI. The recovering of previous models from dimensional reduction is explained in Sec. VII. In the conclusions, we mention open problems and lines of future research. The paper is complemented by three Appendixes. Our notations of conventions are introduced in Appendix A. In Appendix B, the orthonormal eigenstates are computed. We illustrate in Appendix C the open problem of the reducibility of the Hilbert space with respect to the $sl(2|1)$ lowest-weight representations versus its possible irreducibility with respect to a representation of a larger algebraic structure.

II. THE THREE-DIMENSIONAL SUPERCONFORMAL QUANTUM MECHANICS

We realize, following the approaches in Refs. [28,30], a superconformal dynamical symmetry in terms of first-order matrix differential operators. Several requirements have to be satisfied. The operators have to be Hermitian. The fermionic ones need to be block antidiagonal in order to be accommodated into the odd sector of the superalgebra. A supersymmetric quantum mechanics has to be constructed at first. For our purposes, the supersymmetry generators have to be the square roots of a Hamiltonian which corresponds to a deformation of the three-dimensional Laplacian of the free theory. It is easily seen that a three-dimensional Laplacian is nicely expressed in terms of quaternions (which require at least 4×4 real matrices). On the other hand, as recalled in Ref. [30], the closure with Hermitian operators of a superconformal dynamical symmetry (which contains, in particular, the conformal partner of the Hamiltonian) requires the introduction of complex matrices. All in all, 4×4 complex matrices is the minimal setup to achieve the goal (it follows from the properties of Clifford algebras discussed, e.g., in Refs. [36,37]). It allows one to produce an $\mathcal{N} = 2$ extended supersymmetric quantum mechanics, due to the presence of two block-antidiagonal matrices γ_a ($a = 1, 2$) which commute with the three imaginary quaternions h_i ($i = 1, 2, 3$). An explicit realization of the matrices γ_a and h_i and of the fermion parity operator $N_F = \gamma_3$ is given in Eq. (A2). The two supersymmetry operators Q_a are further assumed to have scaling dimension $[Q_a] = \frac{1}{2}$ if $[x_i] = -\frac{1}{2}$ is the assigned scaling dimension of the three space coordinates x_i (our notations and conventions are given in Appendix A).

A natural ansatz to produce two supersymmetry operators Q_a with the required properties consists in setting

$$Q_a = \frac{1}{\sqrt{2}} \gamma_a \left(\not{\partial} - \frac{\beta}{r^2} N_F \not{r} \right). \quad (1)$$

In the above formula, β is a real parameter and $r = \sqrt{x_1^2 + x_2^2 + x_3^2}$ is the radial coordinate, while $\not{\partial} = \partial_i h_i$ and $\not{r} = x_i h_i$ are introduced in Eq. (A7).

The superalgebra of the $\mathcal{N} = 2$ supersymmetric quantum mechanics is

$$\{Q_a, Q_b\} = 2\delta_{ab}H, \quad [H, Q_a] = 0. \quad (2)$$

The 4×4 matrix supersymmetric Hamiltonian H is given by

$$H = \begin{pmatrix} \left(-\frac{1}{2}\nabla^2 + \frac{2\beta}{r^2} \vec{S} \cdot \vec{L} + \frac{\beta(\beta+1)}{2r^2} \right) \mathbb{I}_2 & 0 \\ 0 & \left(-\frac{1}{2}\nabla^2 - \frac{2\beta}{r^2} \vec{S} \cdot \vec{L} + \frac{\beta(\beta-1)}{2r^2} \right) \mathbb{I}_2 \end{pmatrix}, \quad (3)$$

where $\nabla^2 = \partial_{x_1}^2 + \partial_{x_2}^2 + \partial_{x_3}^2$ is the three-dimensional Laplacian and $\vec{S}$ is the spin $\frac{1}{2}$ introduced in Eq. (A9). At $\beta = 0$, the Hamiltonian gets reduced to $H = -\frac{1}{2}\nabla^2 \cdot \mathbb{I}_4$. At non-vanishing β , two extra terms appear: a $\frac{1}{r^2}$ diagonal potential term proportional to $\beta(\beta \pm 1)$ in the upper and lower diagonal blocks and a nondiagonal spin-orbit interaction term proportional to $\vec{S} \cdot \vec{L}$. The latter one is a first-order differential operator.

The Hamiltonian H has scaling dimension $[H] = 1$. Based on the de Alfaro-Fubini-Furlan construction [1], we can introduce its conformal partner as the rotationally invariant operator K of scaling dimension $[K] = -1$. We can therefore set

$$K = \frac{1}{2} r^2 \mathbb{I}_4 \quad (4)$$

and verify whether the repeated (anti)commutators of the operators Q_a and K close the $\mathcal{N} = 2$ one-dimensional superconformal algebra $sl(2|1)$ (see [30,38] for a discussion of one-dimensional, $\mathcal{N}$ -extended superconformal algebras). This is indeed the case. Four extra operators ($\bar{Q}_a$, D , and R) have to be added. D is the (bosonic) dilatation operator which, together with H and K , closes the $sl(2)$ subalgebra. The two fermionic operators $\bar{Q}_a$, of scaling dimension $[\bar{Q}_a] = -\frac{1}{2}$, are introduced from the commutators $[Q_a, K]$. Finally, R is the $u(1)$ R -symmetry bosonic operator of $sl(2|1)$. It is introduced from the anticommutators $\{Q_a, \bar{Q}_b\}$ with $a \neq b$. The (anti)commutators among the eight operators H , D , K , R , Q_a , and $\bar{Q}_a$ close the $sl(2|1)$ superalgebra. We present them for completeness. The nonvanishing ones are

$$\begin{aligned} [D, H] &= -2iH, & [D, K] &= 2iK, & [H, K] &= iD, \\ [D, Q_a] &= -iQ_a, & [D, \bar{Q}_a] &= i\bar{Q}_a, \\ [H, \bar{Q}_a] &= iQ_a & [K, Q_a] &= -i\bar{Q}_a, \\ \{Q_a, Q_b\} &= 2\delta_{ab}H, & \{\bar{Q}_a, \bar{Q}_b\} &= 2\delta_{ab}K, & \{Q_a, \bar{Q}_b\} &= \delta_{ab}D + \epsilon_{ab}R, \\ [R, Q_a] &= -3i\epsilon_{ab}Q_b, & [R, \bar{Q}_a] &= -3i\epsilon_{ab}\bar{Q}_b, \end{aligned} \quad (5)$$

with the antisymmetric tensor ϵ_{ab} normalized so that $\epsilon_{12} = 1$.

Besides Q_a , H , and K , respectively, given in Eqs. (1), (3), and (4), the remaining operators are

$$\begin{aligned} \bar{Q}_a &= -\frac{i}{\sqrt{2}} \gamma_a \not{r}, \\ D &= i \left(x_j \partial_j + \frac{3}{2} \right) \cdot \mathbb{I}_4 = i \left(r \partial_r + \frac{3}{2} \right) \cdot \mathbb{I}_4, \\ R &= -\left(\frac{3}{2} N_F + \beta \cdot \mathbb{I}_4 \right). \end{aligned} \quad (6)$$

The Hamiltonian H is, by construction, Hermitian. Since the spin is $\frac{1}{2}$, the total angular momentum $\vec{J} = \vec{L} + \vec{S}$ of the quantum-mechanical system is half-integer. The Hamiltonian is nondiagonal; on the other hand, due to the relation

$$\vec{L} \cdot \vec{S} = \frac{1}{2} (\vec{J}^2 - \vec{L}^2 - \vec{S}^2) = \frac{1}{2} \left(j(j+1) - l(l+1) - \frac{3}{4} \right), \quad (7)$$

it gets diagonalized in each sector of given total j and orbital l angular momentum. In each such sector, it

corresponds to a constant kinetic term plus a diagonal potential term proportional to $\frac{1}{r^2}$.

III. THE THREE-DIMENSIONAL DEFORMED OSCILLATOR

By setting, following Ref. [1],

$$H_{osc} = H + K, \quad (8)$$

with H and K , respectively, given in Eq. (3) and (4), we produce the 4×4 matrix deformed oscillator Hamiltonian H_{osc} whose spectrum is discrete and bounded from below. By construction, the $sl(2|1)$ dynamical symmetry of the H Hamiltonian acts as a spectrum-generating superalgebra for the H_{osc} Hamiltonian.

The explicit expression of H_{osc} is

$$H_{osc} = -\frac{1}{2}\nabla^2 \cdot \mathbb{I}_4 + \frac{1}{2r^2}(\beta^2 \cdot \mathbb{I}_4 + \beta N_F(1 + 4 \cdot \mathbb{I}_2 \otimes \vec{S} \cdot \vec{L})) + \frac{1}{2}r^2 \cdot \mathbb{I}_4. \quad (9)$$

The spin of the H_{osc} quantum mechanical system is $\frac{1}{2}$. Therefore, its total angular momentum j is half-integer ($j \in \frac{1}{2} + \mathbb{N}_0$), and the relation with the orbital angular momentum $l \in \mathbb{N}_0$ is given by

$$j = l + \delta \frac{1}{2}, \quad \text{for } \delta = \pm 1. \quad (10)$$

In the given j, l sector, the operator $\vec{L} \cdot \vec{S}$ from Eq. (7) is constant. We get

$$\vec{L} \cdot \vec{S} = \frac{1}{2}\alpha, \quad \text{with } \alpha = \delta \left(j + \frac{1}{2} \right) - 1. \quad (11)$$

Each given bosonic (or fermionic) energy eigenvalue in the j, l sector is $(2j + 1)$ times degenerated, the degenerate eigenstates being labeled by the $J_z \equiv J_3$ quantum numbers $-j, j - 1, \dots, j$.

The energy eigenstates of the system are described with the help of the two-component $\mathcal{Y}_{j,l,m}(\theta, \phi)$ spin spherical harmonics (see [4]), given by

$$\begin{aligned} \mathcal{Y}_{j,j-\frac{1}{2}\delta,m}(\theta, \phi) \\ = \frac{1}{\sqrt{2j-\delta+1}} \begin{pmatrix} \delta \sqrt{j + \frac{1}{2}(1-\delta) + \delta m} Y_{j-\frac{1}{2}\delta}^{m-\frac{1}{2}}(\theta, \phi) \\ \sqrt{j + \frac{1}{2}(1-\delta) - \delta m} Y_{j-\frac{1}{2}\delta}^{m+\frac{1}{2}}(\theta, \phi) \end{pmatrix}, \end{aligned} \quad (12)$$

where $Y_l^n(\theta, \phi)$ (for $n = -l, -l + 1, \dots, l$) are the ordinary spherical harmonics.

The spin spherical harmonics $\mathcal{Y}_{j,j-\frac{1}{2}\delta,m}(\theta, \phi)$ are the eigenstates for the compatible observable operators $\vec{J} \cdot \vec{J}$, $\vec{L} \cdot \vec{L}$, and J_z , with eigenvalues $j(j+1)$, $(j-\frac{1}{2}\delta)(j-\frac{1}{2}\delta+1)$, and m , respectively.

The spectrum of the model is derived from the creation (annihilation) operators $a_b^\dagger$ (a_b), with $b = 1, 2$, which are introduced through the positions

$$a_b = Q_b + i\bar{Q}_b, \quad a_b^\dagger = Q_b - i\bar{Q}_b. \quad (13)$$

Indeed, we obtain

$$H_{osc} = \frac{1}{2}\{a_1, a_1^\dagger\} = \frac{1}{2}\{a_2, a_2^\dagger\}, \quad (14)$$

together with

$$[H_{osc}, a_b] = -a_b, \quad [H_{osc}, a_b^\dagger] = a_b^\dagger. \quad (15)$$

For completeness, we also present the commutators

$$[a_1, a_1^\dagger] = [a_2, a_2^\dagger] = 3 \cdot \mathbb{I}_4 + 4 \cdot \mathbb{I}_2 \otimes \vec{S} \cdot \vec{L} - 2\beta N_F. \quad (16)$$

They should be compared with the analogous commutators for one-dimensional deformed oscillators (see [30]), producing deformed Heisenberg algebras with diagonal operators in the right-hand side; the right-hand side of Eq. (16) is diagonalized in the j, l sector.

Let us set, for convenience, $a_b^- \equiv a_b$ and $a_b^+ \equiv a_b^\dagger$. The explicit expression of the creation and annihilation operators is

$$a_b^\pm = \frac{\not{r}}{r\sqrt{2}} \gamma_b \left(\mathbb{I}_4 \cdot (\partial_r \mp r) - \frac{2}{r} \mathbb{I}_2 \otimes \vec{S} \cdot \vec{L} - \frac{\beta}{r} N_F \right). \quad (17)$$

They can be factorized as

$$\begin{aligned} a_b^\pm &= \frac{\not{r}}{r\sqrt{2}} \gamma_b a^\pm, \\ \text{with } a^\pm &= \left(\mathbb{I}_4 \cdot (\partial_r \mp r) - \frac{2}{r} \mathbb{I}_2 \otimes \vec{S} \cdot \vec{L} - \frac{\beta}{r} N_F \right). \end{aligned} \quad (18)$$

The creation and annihilation operators $a_b^\pm$ anticommute with the fermion parity operator N_F ,

$$\{a_b^\pm, N_F\} = 0, \quad (19)$$

producing towers of alternating bosonic and fermionic energy eigenstates.

A lowest-weight state Ψ_{lws} is defined to satisfy

$$a_b^- \Psi_{\text{lws}} = 0. \quad (20)$$

Because of the Eq. (18) factorization, in both $b = 1, 2$ cases, this is tantamount to satisfying $a^- \Psi_{\text{lws}} = 0$.

A lowest-weight representation is spanned by the action of the a_b^+ creation operators on Ψ_{lws} .

If Ψ_{lws} is either bosonic or fermionic, the vectors $a_1^+ v$ and $a_2^+ v$, with v belonging to the lowest-weight representation, differ by a phase. Therefore, the action of a_1^+ and a_2^+ produces the same ray vector characterizing a physical state of the Hilbert space. This phenomenon was already observed [30] in the one-dimensional context. Without the loss of generality, we can just pick, say, a_1^+ to create or annihilate the ray vectors.

We search for solutions $\Psi_{j,\delta,m}^\epsilon(r, \theta, \phi)$ of the lowest-weight condition (20) of the form

$$\Psi_{j,\delta,m}^\epsilon(r, \theta, \phi) = f_{j,\delta}^\epsilon(r) \cdot e_\epsilon \otimes \mathcal{Y}_{j,j-\frac{1}{2}\delta,m}(\theta, \phi), \quad (21)$$

with $\epsilon = \pm 1$.

The sign of ϵ (no summation over this repeated index) refers to the bosonic (fermionic) states with respective eigenvalues $\epsilon = +1$ ($\epsilon = -1$) of the fermion parity operator N_F ; we have $e_{+1} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $e_{-1} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$.

Solutions of Eq. (20) are obtained for the radial-coordinate functions $f_{j,\delta}^\epsilon(r)$ given by

$$f_{j,\delta}^\epsilon(r) = r^{\gamma_{(j,\delta,\epsilon)}} e^{-(1/2)r^2}, \quad (22)$$

where

$$\gamma_{(j,\delta,\epsilon)}(\beta) = \alpha + \beta\epsilon = \delta \left(j + \frac{1}{2} \right) + \beta\epsilon - 1. \quad (23)$$

The corresponding lowest-weight state energy eigenvalue $E_{j,\delta,\epsilon}(\beta)$ from

$$H_{osc}(\beta) \Psi_{j,\delta,m}^\epsilon(r, \theta, \phi) = E_{j,\delta,\epsilon}(\beta) \Psi_{j,\delta,m}^\epsilon(r, \theta, \phi) \quad (24)$$

is

$$E_{j,\delta,\epsilon}(\beta) = \delta \left(j + \frac{1}{2} \right) + \beta\epsilon + \frac{1}{2}. \quad (25)$$

Since $E_{j,\delta,\epsilon}(\beta)$ does not depend on the quantum number m , this energy eigenvalue is $(2j + 1)$ times degenerate.

We discuss in the next section under which condition the vectors of a given lowest-weight representation belong to a normed Hilbert space. For the time being, we point out that the application of the creation operator a_1^+ on an energy eigenstate with energy eigenvalue E produces a ray vector with energy eigenvalue $E + 1$. As we will see, an admissible Hilbert space is defined by a direct sum of (an infinite number of) $sl(2|1)$ lowest-weight representations. The degeneracy of each energy level is finite and can be computed with a recursive formula. Let $n(E)$ be the total number of distinct, admissible, lowest-weight vectors in the Hilbert space, and let $d(E)$ be the number of degenerate

eigenstates at energy level E . At energy level $E + 1$, we obtain

$$d(E + 1) = d(E) + n(E + 1). \quad (26)$$

The $d(E)$ term in the right-hand side gives the number of descendant states obtained by applying a_1^+ to the degenerate states at energy E , while the $n(E + 1)$ term corresponds to the number of new primary states at $E + 1$. From Eq. (26), the computation of the degeneracy of the spectrum is reduced to a combinatorial problem based on the determination of the lowest-weight vectors.

IV. ALTERNATIVE HILBERT SPACES FOR THE 3D DEFORMED OSCILLATOR

Without the loss of generality, we can restrict the real parameter β to belong to the half line $\beta \geq 0$, since the mapping $\beta \leftrightarrow -\beta$ is recovered by a similarity transformation (induced by an operator S) which exchanges bosons into fermions. We have, indeed,

$$SH_{osc}(\beta)S^{-1} = H_{osc}(-\beta) \quad \text{with} \quad S = \sigma_1 \otimes \mathbb{I}_2. \quad (27)$$

At $\beta = 0$, we recover four copies of the undeformed, isotropic, three-dimensional oscillator. Therefore, $\beta > 0$ parametrizes the deformed oscillator.

To the following j , δ , ϵ , and m quantum numbers,

$$\begin{aligned} j &\in \frac{1}{2} + \mathbb{N}_0, & \delta &= \pm 1, \\ \epsilon &= \pm 1, & m &= -j, -j + 1, \dots, j, \end{aligned} \quad (28)$$

is associated an $sl(2|1)$ lowest-weight vector and its induced representation.

At a given β , the Hilbert space has to be selected by requiring, in particular, the normalization of the wave functions entering a lowest-weight representation. This puts a restriction on the admissible lowest-weight representations, the Hilbert space being constructed as the direct sum of the admissible lowest-weight representations.

The analysis of Refs. [2,3] (and also of Ref. [30]) can be extended to the present case. Two choices to select the Hilbert space naturally appear:

- (i) *Case (i).*—The wave functions can be singular at the origin, but they need to be normalized.
- (ii) *Case (ii).*—The wave functions are assumed to be regular at the origin.

We will see in a moment that case (i) corresponds in restricting the admissible lowest-weight representations to those satisfying the necessary and sufficient condition

$$2\gamma_{(j,\delta,\epsilon)}(\beta) + 3 > 0, \quad (29)$$

where $\gamma_{(j,\delta,\epsilon)}(\beta)$ was introduced in Eq. (22) and given in Eq. (23). The normalizability condition (29) is equivalent to the requirement

$$E_{j,\delta,\epsilon}(\beta) > 0 \quad (30)$$

for the lowest-weight energy $E_{j,\delta,\epsilon}(\beta)$ given in Eq. (25).

Case (ii) (regularity at the origin) corresponds in restricting the admissible lowest-weight representations to those satisfying the condition

$$\gamma_{(j,\delta,\epsilon)}(\beta) \geq 0 \quad \text{for } \beta \geq 0. \quad (31)$$

The single-valuedness of the wave functions at the origin implies that the equality $\gamma_{(j,\delta,\epsilon)}(\beta) = 0$ can be realized only with vanishing ($l = 0$) orbital angular momentum (therefore, for $\delta = 1$ and $j = \frac{1}{2}$). At $\beta = 0$, one recovers the vacuum state of the undeformed oscillator. For the deformed $\beta > 0$ oscillator, the strict inequality

$$\gamma_{(j,\delta,\epsilon)}(\beta) > 0 \quad \text{for } \beta > 0 \quad (32)$$

is required as a necessary and sufficient condition.

For illustrative purposes, it is useful to present a table (up to $j = \frac{5}{2}$) of the β range of admissible lowest-weight representations under the *norm* [case (i)] and *reg* [case (ii)] conditions. The j , δ , and ϵ quantum numbers are, respectively, given in columns 1, 2, and 3, while $\gamma_{(j,\delta,\epsilon)}(\beta)$ and $E_{j,\delta,\epsilon}(\beta)$ are presented in columns 4 and 5. We have

j	δ	ϵ	γ	E	norm	reg
$\frac{1}{2}$	+	+	β	$\frac{3}{2} + \beta$	$\beta \geq 0$	$\beta \geq 0$
$\frac{1}{2}$	+	-	$-\beta$	$\frac{3}{2} - \beta$	$0 \leq \beta < \frac{3}{2}$	$\beta = 0$
$\frac{1}{2}$	-	+	$\beta - 2$	$-\frac{1}{2} + \beta$	$\beta > \frac{1}{2}$	$\beta > 2$
$\frac{1}{2}$	-	-	$-\beta - 2$	$-\frac{1}{2} - \beta$	$\times$	$\times$
$\frac{3}{2}$	+	+	$\beta + 1$	$\frac{5}{2} + \beta$	$\beta \geq 0$	$\beta \geq 0$
$\frac{3}{2}$	+	-	$-\beta + 1$	$\frac{5}{2} - \beta$	$0 \leq \beta < \frac{5}{2}$	$0 \leq \beta < 1$
$\frac{3}{2}$	-	+	$\beta - 3$	$-\frac{3}{2} + \beta$	$\beta > \frac{3}{2}$	$\beta > 3$
$\frac{3}{2}$	-	-	$-\beta - 3$	$-\frac{3}{2} - \beta$	$\times$	$\times$
$\frac{5}{2}$	+	+	$\beta + 2$	$\frac{7}{2} + \beta$	$\beta \geq 0$	$\beta \geq 0$
$\frac{5}{2}$	+	-	$-\beta + 2$	$\frac{7}{2} - \beta$	$0 \leq \beta < \frac{7}{2}$	$0 \leq \beta < 2$
$\frac{5}{2}$	-	+	$\beta - 4$	$-\frac{5}{2} + \beta$	$\beta > \frac{5}{2}$	$\beta > 4$
$\frac{5}{2}$	-	-	$-\beta - 4$	$-\frac{5}{2} - \beta$	$\times$	$\times$

(33)

For the undeformed $\beta = 0$ oscillator, the restrictions from either case (i) or case (ii) select the same Hilbert space. It is the direct sum of all lowest-weight representations with j , ϵ , and m satisfying Eq. (28), while δ is restricted to be

$$\delta = +1. \quad (34)$$

For the $\beta > 0$ deformed oscillators, the Hilbert spaces $\mathcal{H}_{\text{norm}}$ and $\mathcal{H}_{\text{reg}}$ are direct sums of the lowest-weight representations with $j \in \frac{1}{2} + \mathbb{N}_0$ satisfying (depending on δ, ϵ)

		$\mathcal{H}_{\text{norm}} :$	$\mathcal{H}_{\text{reg}} :$
$\delta = +1$	$\epsilon = +1$	Any j	Any j
$\delta = +1$	$\epsilon = -1$	$j > \beta - 1$	$j > \beta + \frac{1}{2}$
$\delta = -1$	$\epsilon = +1$	$j < \beta$	$j < \beta - \frac{3}{2}$
$\delta = -1$	$\epsilon = -1$	No j	No j

(35)

The vacuum energy of the system is obtained by comparing the smallest lowest-weight energies [obtained from Eq. (25)] in each one of the three contributing sectors ($\delta = 1$ with $\epsilon = 1$, $\delta = 1$ with $\epsilon = -1$, and $\delta = -1$ with $\epsilon = 1$) for the admissible values of j . One should take into account that for $\delta = +1$, at a given $\epsilon = \pm 1$, the smallest value of energy is encountered at the smallest admissible value of j . Conversely, for $\delta = -1$ and $\epsilon = +1$, the smallest lowest-weight energy is encountered at the largest admissible value of j .

The normalizability condition (29) comes from the requirement to have a finite norm for the lowest-weight vector (21), such that

$$\int d^3x \text{Tr}(\Psi_{j,\delta,m}^\epsilon(r, \theta, \phi)^\dagger \Psi_{j,\delta,m}^\epsilon(r, \theta, \phi)) < \infty. \quad (36)$$

Passing to spherical coordinates, the only potentially troublesome integral is

$$\int_0^{+\infty} dr r^2 r^{2\gamma_{(j,\delta,\epsilon)}} e^{-r^2}, \quad (37)$$

which is finite at the origin, provided that $2\gamma_{(j,\delta,\epsilon)} + 2 > -1$, namely, the Eq. (29) condition. A simple inspection shows that, acting with the a_1^+ creation operators, no further condition is necessary to ensure the normalizability of the excited states of the lowest-weight representation.

The normalizability condition discussed in the papers in Refs. [2,3,30] (which deal with one-dimensional systems) implies that the normalized wave functions are defined on the real line, with the possible “smooth”

singularity at the origin. The regular wave functions (for $\beta > 0$) are, on the other hand, defined on the half line and satisfy the Dirichlet boundary condition at the origin. We stress that the mathematical framework in the present work coincides with the one introduced in Refs. [2,3,30]. For the singular Hilbert space $\mathcal{H}_{\text{norm}}$, this means that one can extend the wave functions (defined by the spherical coordinates r, θ, ϕ) in two regions, namely, I_+ (parametrized by $r \geq 0$) and I_- (parametrized by $r \leq 0$), so that their intersection $I_+ \cap I_- = \{O\}$ is the origin. Mathematically, this construction works fine. On a physical ground, one has to reconcile the interpretation of the wave functions defined on $I = I_+ \cup I_-$ with the wave functions defined on the ordinary three-dimensional space. This can be achieved by taking into account that the $\mathbf{Z}_2$ transformation

$$Z: r \mapsto -r \quad (\text{with } \theta, \phi \text{ unchanged and } Z^2 = 1) \quad (38)$$

is a symmetry of the H_{osc} Hamiltonian. The induced operator $\hat{Z}$, acting on the wave functions defined on $I_+ \cup I_-$, satisfies the

$$\hat{Z}^2 = \mathbb{I} \quad (39)$$

condition, is Hermitian, and possesses ± 1 eigenvalues, provided that

$$(-1)^{\gamma(j,\delta,\epsilon)} = \pm 1, \quad (40)$$

implying that $\gamma_{(j,\delta,\epsilon)}(\beta)$ (and, consequently, β) is an integer. If Eq. (40) is satisfied for the lowest-weight vector, $\hat{Z}^\dagger = \hat{Z}$ and Eq. (39) are satisfied for all states belonging to the associated lowest-weight representation. For such a well-defined $\hat{Z}$ operator, one can impose the $\{I_+ \cup I_-\}/\mathbf{Z}_2$ coset construction, by identifying

$$(r, \theta, \phi) \equiv (-r, \theta, \phi). \quad (41)$$

Under this coset construction, one can set

$$\{I_+ \cup I_-\}/\mathbf{Z}_2 \approx \mathbb{R}^3. \quad (42)$$

The $\mathbf{Z}_2$ coset construction entails, from Eqs. (40) and (23), the quantization of β which, under this condition, is restricted to be integer:

$$\beta \in \mathbb{Z}. \quad (43)$$

We stress the fact that the Hilbert space can be consistently defined for any real value of β . It is the $\mathbf{Z}_2$ coset construction with the associated interpretation which requires β to be an integer.

V. THE SPECTRUM OF THE β -DEFORMED OSCILLATOR

In this section, we present, for H_{osc} , the computation of the spectrum for the alternative choices of $\mathcal{H}_{\text{norm}}$ and $\mathcal{H}_{\text{reg}}$ Hilbert spaces. Without the loss of generality, β is taken to

be $\beta \geq 0$. We start with the $\mathcal{H}_{\text{norm}}$ case (normalized wave functions).

A. The spectrum for the $\mathcal{H}_{\text{norm}}$ Hilbert space of normalized wave functions

For $\beta \geq \frac{1}{2}$, it is convenient to introduce, via the floor function, the parameter μ , defined as

$$\mu = \left\{ \beta - \frac{1}{2} \right\} = \left(\beta - \frac{1}{2} \right) - \left\lfloor \beta - \frac{1}{2} \right\rfloor, \quad p = \left\lfloor \beta - \frac{1}{2} \right\rfloor,$$

$$\text{so that } \mu \in [0, 1], \quad p \in \mathbb{N}_0, \quad \text{and} \quad \beta = \frac{1}{2} + \mu + p. \quad (44)$$

The results for the spectrum split into six different cases which have to be separately analyzed:

- (i) case I.— $\beta = 0$ (the undeformed oscillator),
- (ii) case II.— $\beta = 1 + p$, with $p \in \mathbb{N}_0$ ($p = 0, 1, 2, \dots$),
- (iii) case III.— $\beta = \frac{1}{2} + p$, with $p \in \mathbb{N}_0$,
- (iv) case IV.— $0 < \beta < \frac{1}{2}$,
- (v) case V.— $0 < \mu < \frac{1}{2}$, and, therefore, $\beta = \frac{1}{2} + \mu + p$, with $p \in \mathbb{N}_0$,
- (vi) case VI.— $\frac{1}{2} < \mu < 1$, therefore $\beta = \frac{1}{2} + \mu + p$, with $p \in \mathbb{N}_0$.

The energy eigenvalues corresponding to the above cases are as follows.

- (i) Case I.— $E_n = \frac{3}{2} + n$, where $n \in \mathbb{N}_0$ is a non-negative integer. The vacuum energy is $E_{\text{vac}} = \frac{3}{2}$; the ground state is four times degenerated, with two bosonic and two fermionic eigenstates (hence, “ $2_B + 2_F$ ”). The vacuum lowest-weight vectors are specified by the quantum numbers $j = \frac{1}{2}$, $\delta = +1$, $\epsilon = \pm 1$, and (here and in the following) all compatible values $m = -j, \dots, j$.
- (ii) Case II.— $E_n = \frac{1}{2} + n$, with $n \in \mathbb{N}_0$. The vacuum energy is $E_{\text{vac}} = \frac{1}{2}$; the degeneration of the ground state is $2(p+1)$, with $p+1$ bosonic and $p+1$ fermionic eigenstates, and is therefore denoted as “ $(p+1)_B + (p+1)_F$ ”. The vacuum lowest-weight vectors are specified by $j = \frac{1}{2} + p$, with either $\delta = +1$, $\epsilon = -1$ or $\delta = -1$, $\epsilon = +1$.
- (iii) Case III.— $E_n = 1 + n$, with $n \in \mathbb{N}_0$. The vacuum energy is $E_{\text{vac}} = \frac{1}{2}$; the degeneration of the ground state is $4p+2$, with $2p$ bosonic and $2(p+1)$ fermionic eigenstates, and is therefore denoted as “ $(2p)_B + (2p+2)_F$ ”. For $p = 0$, the two vacuum lowest vectors are specified by $j = \frac{1}{2}$, $\delta = +1$, and $\epsilon = -1$. For $p > 0$, the vacuum lowest vectors are specified either by $j = \frac{1}{2} + p$, $\delta = +1$, and $\epsilon = -1$ or by $j = p - \frac{1}{2}$, $\delta = -1$, and $\epsilon = +1$.
- (iv) Case IV.—Two series of energy eigenvalues $E_n^\pm = \frac{3}{2} \pm \beta + n$, with $n \in \mathbb{N}_0$, are encountered. The vacuum energy is $E_{\text{vac}} = \frac{3}{2} - \beta$; the ground state is fermionic and doubly degenerated (“ 2_F ”). The two

vacuum lowest-weight vectors are specified by $j = \frac{1}{2}$, $\delta = +1$, and $\epsilon = -1$.

- (v) Case V.—Two series of energy eigenvalues $E_n^- = \mu + n$ and $E_n^+ = 1 - \mu + n$, with $n \in \mathbb{N}_0$, are encountered. The vacuum energy is $E_{\text{vac}} = \mu$; the ground state is bosonic and $(2p + 2)$ times degenerated [hence, “ $(2p + 2)_B$ ”]. The vacuum lowest-weight vectors are specified by $j = \frac{1}{2} + p$, $\delta = -1$, and $\epsilon = +1$.
- (vi) Case VI.—Two series of energy eigenvalues $E_n^- = 1 - \mu + n$ and $E_n^+ = \mu + n$, with $n \in \mathbb{N}_0$, are encountered. The vacuum energy is $E_{\text{vac}} = 1 - \mu$; the ground state is fermionic and $(2p + 2)$ times degenerated [hence, “ $(2p + 2)_F$ ”]. The vacuum lowest-weight vectors are specified by $j = \frac{1}{2} + p$, $\delta = +1$, and $\epsilon = -1$.

There is an important remark. The energy spectrum of cases V and VI coincides under a

$$\mu \leftrightarrow 1 - \mu, \quad \text{with } \mu \neq 0, \frac{1}{2}, \quad (45)$$

duality transformation. Under this duality transformation, the parity (bosonic and fermionic) of the ground state is exchanged. On the other hand, the degeneracies of the energy eigenvalues above the ground state are not respected by the duality transformation. It is sufficient to discuss a specific example to show it. Let us take $\mu = \frac{1}{4}$ with $p = 0$ and consider the dually related $\beta = \frac{3}{4}$ and $\beta = \frac{5}{4}$ cases.

The lowest-weight vectors appearing in the first five energy levels (from the vacuum energy $E_{\text{vac}} = \frac{1}{4}$ to $E = \frac{9}{4}$) are presented in the following table:

E	$\beta = \frac{3}{4}$	$\beta = \frac{5}{4}$
$\frac{9}{4}$	$\frac{1}{2} + B$	$\frac{5}{2} + F$
$\frac{7}{4}$	$\frac{3}{2} + F$	$\times$
$\frac{5}{4}$	$\times$	$\frac{3}{2} + F$
$\frac{3}{4}$	$\frac{1}{2} + F$	$\frac{1}{2} - B$
$\frac{1}{4}$	$\frac{1}{2} - B$	$\frac{1}{2} + F$

(46)

The lowest-weight vectors are identified by their quantum numbers j , the $\pm$ sign referring to δ , while B (standing for bosons) and F (standing for fermions) correspond to $\epsilon = +1$ and $\epsilon = -1$, respectively.

The degeneracy $d_\beta(E)$ of a given energy level is computed with the recursive formula (26) involving primary (lowest-weight vectors) and descendant states. The results are presented in the table

E	$d_{\beta=\frac{3}{4}}(E)$	$d_{\beta=\frac{5}{4}}(E)$
$\frac{9}{4}$	4	12
$\frac{7}{4}$	6	2
$\frac{5}{4}$	2	6
$\frac{3}{4}$	2	2
$\frac{1}{4}$	2	2

(47)

One can see that $\frac{5}{4}$ is the first energy level where an inequality of the degeneracies is produced:

$$d_{\beta=3/4}\left(\frac{5}{4}\right) \neq d_{\beta=5/4}\left(\frac{5}{4}\right). \quad (48)$$

The β dependence of the vacuum energy is graphically presented in Fig. 1.

B. The spectrum for the $\mathcal{H}_{\text{reg}}$ Hilbert space of regular wave functions

The $\beta = 0$ case was already discussed since, for this value, the $\mathcal{H}_{\text{norm}}$ and $\mathcal{H}_{\text{reg}}$ Hilbert spaces coincide. For $\beta > 0$, the following cases have to be separately treated:

- (i) case (i).— $0 < \beta < \frac{1}{2}$,
- (ii) case (ii).— $\beta = \frac{1}{2}$,
- (iii) case (iii).— $\frac{1}{2} < \beta < 1$,
- (iv) case (iv).— $\beta = 1$,
- (v) case (v).— $1 < \beta < 2$, with $\beta \neq \frac{3}{2}$,
- (vi) case (vi).— $\beta = \frac{3}{2}$,
- (vii) case (vii).— $\beta = 2$,
- (viii) case (viii).— $\beta = 2 + \mu + p$, with $0 < \mu < \frac{1}{2}$ and $p \in \mathbb{N}_0$,
- (ix) case (ix).— $\beta = \frac{5}{2} + p$, with $p \in \mathbb{N}_0$,
- (x) case (x).— $\beta = 2 + \mu + p$, with $\frac{1}{2} < \mu < 1$ and $p \in \mathbb{N}_0$, and
- (xi) case (xi).— $\beta = 3 + p$, with $p \in \mathbb{N}_0$.

The energy eigenvalues corresponding to the above cases are as follows.

- (i) Case (i).—Two series of energy eigenvalues, $E_n^+ = \frac{3}{2} + \beta + n$ and $E_n^- = \frac{5}{2} - \beta + n$, with $n \in \mathbb{N}_0$, are encountered. The vacuum energy is $E_{\text{vac}} = \frac{3}{2} + \beta$; the ground state is bosonic and doubly degenerate (hence, denoted as “ 2_B ”). The vacuum lowest-weight vectors are specified by the quantum numbers $j = \frac{1}{2}$ and $\delta = \epsilon = +1$.
- (ii) Case (ii).—The energy eigenvalues are $E_n = 2 + n$, with $n \in \mathbb{N}_0$. The vacuum energy is $E_{\text{vac}} = 2$; the degeneration of the ground state is 6, with two

bosonic and four fermionic eigenstates (“ $2_B + 4_F$ ”). The vacuum lowest-weight vectors are specified by $j = \frac{1}{2}$, with $\delta = \epsilon = +1$ and by $j = \frac{3}{2}$ with $\delta = +1$ and $\epsilon = -1$.

- (iii) Case (iii).—Two series of energy eigenvalues, $E_n^+ = \frac{3}{2} + \beta + n$ and $E_n^- = \frac{5}{2} - \beta + n$, with $n \in \mathbb{N}_0$, are encountered. The vacuum energy is $E_{\text{vac}} = \frac{5}{2} - \beta$; the ground state is fermionic and 4 times degenerate (“ 4_F ”). The vacuum lowest-weight vectors are specified by $j = \frac{3}{2}$, $\delta = +1$, and $\epsilon = -1$.
- (iv) Case (iv).—The energy eigenvalues are $E_n = \frac{5}{2} + n$, with $n \in \mathbb{N}_0$. The vacuum energy is $E_{\text{vac}} = \frac{5}{2}$; the degeneration of the ground state is 8, with two bosonic and six fermionic eigenstates (“ $2_B + 6_F$ ”). The vacuum lowest-weight vectors are specified by $j = \frac{1}{2}$ with $\delta = \epsilon = +1$ and by $j = \frac{5}{2}$ with $\delta = +1$ and $\epsilon = -1$.
- (v) Case (v).—Two series of energy eigenvalues, $E_n^+ = \frac{3}{2} + \beta + n$ and $E_n^- = \frac{7}{2} - \beta + n$, with $n \in \mathbb{N}_0$, are encountered. The vacuum energy is $E_{\text{vac}} = \frac{7}{2} - \beta$; the ground state is fermionic and 6 times degenerate (“ 6_F ”). The vacuum lowest-weight vectors are specified by $j = \frac{5}{2}$, $\delta = +1$, and $\epsilon = -1$.
- (vi) Case (vi).—the energy eigenvalues are $E_n = 2 + n$, with $n \in \mathbb{N}_0$. The vacuum energy is $E_{\text{vac}} = 2$; the degeneration of the ground state is fermionic and 6 times degenerate (“ 6_F ”). The vacuum lowest-weight vectors are specified by $j = \frac{5}{2}$ with $\delta = +1$ and $\epsilon = -1$.
- (vii) Case (vii).—The energy eigenvalues are $E_n = \frac{5}{2} + n$, with $n \in \mathbb{N}_0$. The vacuum energy is $E_{\text{vac}} = \frac{5}{2}$; the ground state is fermionic and 8 times degenerate (“ 8_F ”). The vacuum lowest-weight vectors are specified by $j = \frac{7}{2}$ with $\delta = +1$ and $\epsilon = -1$.
- (viii) Case (viii).—Two series of energy eigenvalues, $E_n^- = \frac{3}{2} + \mu + n$ and $E_n^+ = \frac{5}{2} - \mu + n$, with $n \in \mathbb{N}_0$, are encountered. The vacuum energy is $\frac{3}{2} + \mu$; the ground state is bosonic and $2(p+1)$ degenerate [“(2p+2) $_B$ ”]. The vacuum lowest vectors are specified by $j = \frac{1}{2} + p$, with $\delta = -1$ and $\epsilon = +1$.
- (ix) Case (ix).—The energy eigenvalues are $E_n = 2 + n$, with $n \in \mathbb{N}_0$. The vacuum energy is $E_{\text{vac}} = 2$; the degeneration of the ground state is $10 + 4p$ with $2(p+1)$ bosonic and $8 + 2p$ fermionic eigenstates [“(2p+2) $_B$ + (8+2p) $_F$ ”]. The vacuum lowest-weight vectors are specified by $j = \frac{1}{2} + p$ with $\delta = -1$ and $\epsilon = +1$ and by $j = \frac{7}{2} + p$ with $\delta = +1$ and $\epsilon = -1$.
- (x) Case (x).—Two series of energy eigenvalues, $E_n^+ = \frac{3}{2} + \mu + n$ and $E_n^- = \frac{5}{2} - \mu + n$, with $n \in \mathbb{N}_0$, are encountered. The vacuum energy is $\frac{5}{2} - \mu$; the ground state is fermionic and $2(p+4)$ degenerate [“(2p+8) $_B$ ”]. The vacuum lowest vectors are specified by $j = \frac{7}{2} + p$ with $\delta = +1$ and $\epsilon = -1$.
- (xi) Case (xi).—The energy eigenvalues are $E_n = \frac{5}{2} + n$, with $n \in \mathbb{N}_0$. The vacuum energy is $E_{\text{vac}} = \frac{5}{2}$; the degeneration of the ground state is $12 + 4p$ with $2(p+1)$ bosonic and $10 + 2p$ fermionic eigenstates [“(2p+2) $_B$ + (2p+10) $_F$ ”]. The vacuum lowest-weight vectors are specified by $j = \frac{1}{2} + p$ with $\delta = -1$ and $\epsilon = +1$ and by $j = \frac{9}{2} + p$ with $\delta = +1$ and $\epsilon = -1$. The β dependence of the vacuum energy is graphically presented in Fig. 2.

C. About superselecting the wave functions

It is beyond the scope of this work to present an investigation of the admissible superselection rules which can be implemented on the $\mathcal{H}_{\text{norm}}$ and $\mathcal{H}_{\text{reg}}$ Hilbert spaces. We limit ourselves to mention here a natural type of superselection for its consequence on the spectrum. As we have seen, when the deformation parameter β is integer or half-integer, the spectrum coincides with a shifted version of the undeformed oscillator (more on that in Sec. VI). A natural superselection consists in imposing the ground state and the even excited states to be bosonic, with fermionic odd excited states (or vice versa, exchanging the role of bosons and fermions). For $\beta \in \mathbb{Z}$ or $\beta \in \frac{1}{2} + \mathbb{Z}$, this superselection is implemented by the projectors

$$P_{\pm} = \frac{1}{2} (\mathbb{I}_4 \pm N_F e^{\pi i (H_{\text{osc}} - E_{gr})}) \quad (P_{\pm}^2 = P_{\pm}), \quad (49)$$

where E_{gr} denotes the given ground energy. The “+” (“−”) sign is associated with the fermionic (bosonic) ground state. The superselected wave functions Ψ are constrained to satisfy

$$\text{either } P_+ \Psi = 0 \quad \text{or} \quad P_- \Psi = 0. \quad (50)$$

VI. COMBINATORICS OF THE ENERGY EIGENSTATE DEGENERACY

For β integer or half-integer, the spectrum of the deformed oscillator is a shifted version of the $\beta = 0$ undeformed oscillator. A second dissimilarity consists in the different degeneracies of the energy eigenvalues for the corresponding ground states and n th excited states. When $\beta \notin \{\mathbb{Z}\} \cup \{\frac{1}{2} + \mathbb{Z}\}$, the spectrum is a combination of two differently shifted spectra of the undeformed oscillator [see cases (IV), (V), and (VI) and cases (i), (iii), (v), (viii), and (x) above].

In order to facilitate the comparison with the undeformed oscillator and highlight the differences, for β integer and half-integer we solve the combinatorics of the degenerate eigenstates provided by the (26) recursive relation with the (35) input of primary (i.e., lowest-weight vectors) states. The same technique can be applied for any real value of β (the results for this latter case will not be reported to avoid excessively burdening the paper).

Let $E_{\beta,\text{norm}}^n$ and $E_{\beta,\text{reg}}^n$ denote the energy eigenvalues for β integer or half-integer in correspondence with, respectively, the $\mathcal{H}_{\text{norm}}$ and $\mathcal{H}_{\text{reg}}$ choices of the Hilbert space ($E_{\beta,\text{norm}}^0$ and $E_{\beta,\text{reg}}^0$ are the corresponding vacuum energies). The energy shifts $\Delta_{\beta,\text{norm}}$ and $\Delta_{\beta,\text{reg}}$ with respect to the undeformed oscillator can be read from Figs. 1 and 2. We have

$$\begin{aligned} E_{\beta,\text{norm}}^n &= E_{\beta=0}^n + \Delta_{\beta,\text{norm}} = \frac{3}{2} + n + \Delta_{\beta,\text{norm}}, \\ E_{\beta,\text{reg}}^n &= E_{\beta=0}^n + \Delta_{\beta,\text{reg}} = \frac{3}{2} + n + \Delta_{\beta,\text{reg}}, \end{aligned} \quad (51)$$

with, for $p \in \mathbb{N}_0$,

$$\begin{aligned} \Delta_{\frac{1}{2}+p,\text{norm}} &= -\frac{1}{2}, & \Delta_{p,\text{norm}} &= -1, \\ \Delta_{\frac{1}{2}+p,\text{reg}} &= \frac{1}{2}, & \Delta_{p,\text{reg}} &= 1. \end{aligned} \quad (52)$$

We start our analysis of the eigenvalue degeneracies by recalling, at first, the features of the $\beta = 0$ undeformed oscillator.

A. Degeneracies of the $\beta = 0$ undeformed oscillator

Since, at $\beta = 0$, the Eq. (9) Hamiltonian H_{osc} corresponds to four copies of the ordinary isotropic three-dimensional oscillator, its degeneracy $d_{\beta=0}(n)$ is

$$d_{\beta=0}(n) = 4 \cdot d(n), \quad \text{with} \quad d(n) = \frac{1}{2}(n^2 + 3n + 2). \quad (53)$$

Here $d(n)$ denotes the eigenvalues degeneracy of the ordinary three-dimensional oscillator. It produces the 1, 3, 6, 10, 15, ... tower of states, the vacuum being nondegenerate.

The rotational invariance of the ordinary three-dimensional oscillator implies that at each energy level the degenerate states can be accommodated into multiplets (denoted as “[l]”) of non-negative integer orbital angular momentum l , each [l] multiplet containing $2l + 1$ states. We get, at the lowest orders,

$$\begin{aligned} n = 0: & \quad [0], & \text{so that } d(0) &= 1, \\ n = 1: & \quad [1], & \text{so that } d(1) &= 3, \\ n = 2: & \quad [0] \oplus [2], & \text{so that } d(2) &= 1 + 5 = 6, \\ n = 3: & \quad [1] \oplus [3], & \text{so that } d(3) &= 3 + 7 = 10, \\ n = 4: & \quad [0] \oplus [2] \oplus [4], & \text{so that } d(4) &= 1 + 5 + 9 = 15, \\ \dots: & \quad \dots, & & \dots. \end{aligned} \quad (54)$$

The spectrum of the ordinary oscillator can be recovered by imposing, to the H_{osc} Hamiltonian at $\beta = 0$, two independent superselection rules defined by the fermion parity operator N_F and by the spin operator $\mathbb{I}_2 \otimes S_3$ (since the

spin-orbit coupling is absent at $\beta = 0$, for this particular value of β the Hermitian operator $\mathbb{I}_2 \otimes S_3$ commutes with the H_{osc} Hamiltonian). The corresponding projector operators $P_{\pm}^N$ and $P_{\pm}^S$ which induce the superselection rules are different from the projectors introduced in Eq. (49); they are given by

$$P_{\pm}^N = \frac{1}{2}(\mathbb{I}_4 \pm N_F), \quad P_{\pm}^S = \frac{1}{2}(\mathbb{I}_4 \pm 2\mathbb{I}_2 \otimes S_3) \quad (55)$$

and satisfy

$$\begin{aligned} (P_{\pm}^N)^2 &= P_{\pm}^N, & (P_{\pm}^S)^2 &= P_{\pm}^S, \\ [P_{\pm}^N, P_{\pm}^S] &= 0, & [P_{\pm}^N, P_{\mp}^S] &= 0. \end{aligned} \quad (56)$$

The two compatible superselection rules at $\beta = 0$ can be imposed by restricting the H_{osc} wave functions Ψ to satisfy

$$P_{+}^N \Psi = P_{+}^S \Psi = \Psi. \quad (57)$$

One should note that the $\beta = 0$ rotational invariance is, from the $sl(2|1)$ spectrum-generating superalgebra viewpoint, an emergent symmetry, being not immediately derivable from the $sl(2|1)$ data.

B. Degeneracies for $\beta = \frac{1}{2} + \mathbb{N}_0$ and $\beta = 1 + \mathbb{N}_0$ with $\mathcal{H}_{\text{norm}}$ Hilbert space

We present here the results of the degeneracies for the Hilbert space $\mathcal{H}_{\text{norm}}$ of normalized wave functions. We have the following cases.

Case (a).— $\beta = \frac{1}{2} + p$ (energy levels $E_n = n + 1$) with $p, n \in \mathbb{N}_0$.

For $\beta = \frac{1}{2}$, the degeneracy $d_{\beta=\frac{1}{2}}(E_n)$ of the n th level is simply

$$d_{\beta=1/2}(E_n) = 2 \cdot (n + 1)^2. \quad (58)$$

The above formula can be recovered from the following most general case at given integer p . The degeneracy $d_{\beta=(1/2)+p}(E_n)$ grows linearly (mimicking a two-dimensional oscillator) up to $n = p$; it then grows quadratically starting from $n = p + 1$:

$$\begin{aligned} d_{\beta=(1/2)+p}(E_n) &= 2(n + 1)(2p + 1) \quad \text{for } n = 0, 1, 2, \dots, p, \\ d_{\beta=(1/2)+p}(E_n) &= 2 \cdot (q^2 + 2(p + 1)q + (p + 1)(2p + 1)) \\ &\quad \text{for } n = p + q \quad \text{with } q = 0, 1, 2, \dots \end{aligned} \quad (59)$$

Case (b).— $\beta = 1 + p$ (energy levels $E_n = n + \frac{1}{2}$) with $p, n \in \mathbb{N}_0$.

As in the previous case, the degeneracy $d_{\beta=1+p}(E_n)$ grows linearly (mimicking a two-dimensional oscillator) up to $n = p$; it then grows quadratically starting from $n = p + 1$:

$$\begin{aligned}
d_{\beta=1+p}(E_n) &= 4(n+1)(p+1) \quad \text{for } n=0,1,2,\dots,p, \\
d_{\beta=1+p}(E_n) &= 2 \cdot (q^2 + (2p+1)q + 2(p+1)^2) \\
&\quad \text{for } n=p+q \quad \text{with } q=0,1,2,\dots \quad (60)
\end{aligned}$$

We provide in Fig. 3 a graphical description of the degeneracy.

C. Degeneracies for $\beta = \frac{1}{2} + \mathbb{N}_0$ and $\beta = 1 + \mathbb{N}_0$ with $\mathcal{H}_{\text{reg}}$ Hilbert space

We present here the results of the degeneracies for the Hilbert space $\mathcal{H}_{\text{reg}}$ of regular wave functions. We have the following cases.

Case (a).— β half-integer with energy levels $E_n = n + 2$.

The $\beta = \frac{1}{2}$ and $\beta = \frac{3}{2}$ cases present the same degeneracy, the functions $d_{\beta=1/2}(E_n)$, $d_{\beta=3/2}(E_n)$ being given by

$$d_{\beta=1/2}(E_n) = d_{\beta=3/2}(E_n) = 2 \cdot (n^2 + 4n + 3). \quad (61)$$

The $\beta = \frac{3}{2}$ case can also be recovered from the most general formula for $\beta = \frac{3}{2} + p$, with $p \in \mathbb{N}_0$. We have

$$\begin{aligned}
d_{\beta=(3/2)+p}(E_n) &= 2 \cdot (n+1)(2p+3) \quad \text{for } n=0,1,2,\dots,p, \\
d_{\beta=(3/2)+p}(E_n) &= 2 \cdot (q^2 + (2p+4)q + (2p+3)(p+1)) \\
&\quad \text{for } n=p+q \quad \text{with } q=0,1,2,\dots \quad (62)
\end{aligned}$$

We recover the same feature observed for normalized wave functions belonging to $\mathcal{H}_{\text{norm}}$, namely, a linear growth followed by a quadratic growth.

Case (b).— β integer with energy levels $E_n = n + \frac{5}{2}$.

The $\beta = 1$ and $\beta = 2$ cases present the same degeneracy, the functions $d_{\beta=1}(E_n)$ and $d_{\beta=2}(E_n)$ being given by

$$d_{\beta=1}(E_n) = d_{\beta=2}(E_n) = 2 \cdot (n^2 + 5n + 4). \quad (63)$$

The $\beta = 2$ case can also be recovered from the most general formula for $\beta = 2 + p$, with $p \in \mathbb{N}_0$. We have

$$\begin{aligned}
d_{\beta=2+p}(E_n) &= 4 \cdot (p+2)(n+1) \quad \text{for } n=0,1,2,\dots,p, \\
d_{\beta=2+p}(E_n) &= 2 \cdot (q^2 + (2p+5)q + 2(p+2)(p+1)) \\
&\quad \text{for } n=p+q \quad \text{with } q=0,1,2,\dots \quad (64)
\end{aligned}$$

Here again we observe a linear growth up to $E \approx \beta$, followed by a quadratic growth.

VII. DIMENSIONAL REDUCTIONS OF THE 3D DEFORMED OSCILLATOR

In this section, we discuss how to recover two-dimensional and one-dimensional models of superconformal quantum mechanics (originally introduced in Ref. [26] via the quantization of world-line superconformal sigma models, with two and one target coordinates, respectively)

by dimensional reduction of the three-dimensional superconformal quantum mechanics. We explicitly treat the case of the de Alfaro-Fubini-Furlan deformed oscillator (the construction can be repeated step by step, with analogous results, in the absence of the oscillatorial term).

A. The 3D $\rightarrow$ 2D dimensional reduction

To obtain this dimensional reduction, we have to freeze the dependence on the third coordinate x_3 in the wave functions. This can be achieved by setting ∂ , $\not{r}$, and r entering the operators (1), (3), (4), and (6) to be restricted to

$$\partial = h_1 \partial_1 + h_2 \partial_2, \quad \not{r} = x_1 h_1 + x_2 h_2, \quad r = \sqrt{x_1^2 + x_2^2} \quad (65)$$

[the last equality holds for D in formula (6)]. With the above positions, the $sl(2|1)$ (anti)commutators (5) are maintained. The $\vec{S} \cdot \vec{L}$ operator entering the Hamiltonians (3) and (9) is now given by $S_3 L_3$. In the two-dimensional case, being proportional to σ_3 , this operator is diagonal.

The resulting Hamiltonian $H_{2D,osc}$ of the deformed oscillator corresponds to two copies of the two-dimensional 2×2 matrix Hamiltonians derived in Ref. [26] from the quantization of the $sl(2|1)$ world-line sigma model with two propagating bosonic and two propagating fermionic fields. We have

$$\begin{aligned}
H_{2D,osc} &= -\frac{1}{2}(\partial_{x_1}^2 + \partial_{x_2}^2) \cdot \mathbb{I}_4 \\
&\quad + \frac{1}{2r^2}(\beta^2 \mathbb{I}_4 + \beta N_F(1 + 2 \cdot \mathbb{I}_2 \otimes \sigma_3 L_3)) + \frac{1}{2}r^2 \mathbb{I}_4. \quad (66)
\end{aligned}$$

The two nonvanishing upper-left and bottom-right blocks of the given 4×4 Hamiltonian present the deformation parameters $+\beta$ and $-\beta$, respectively.

There is an important remark. For the deformed oscillator, the $sl(2|1)$ spectrum-generating superalgebra induced by the (1), (3), (4), and (6) operators with the (65) restriction does not coincide with the $sl(2|1)$ spectrum-generating superalgebra introduced in Ref. [26]. The reason is that the ladder operators constructed with Q_a and $\bar{Q}_a$ connect the upper-left and bottom-right blocks, while the $sl(2|1)$ ladder operators of Ref. [26] act inside each 2×2 component blocks. Therefore, the $sl(2|1)$ spectrum-generating superalgebra of Ref. [26], as well as its charge-conjugated $sl(2|1)$ superalgebra (also in Ref. [26]) are, from the point of view of the dimensionally reduced theory, emergent spectrum-generating superalgebras which are not directly connected with the original $sl(2|1)$ superalgebra.

B. The 3D $\rightarrow$ 1D dimensional reduction

To operate this dimensional reduction, it is convenient to freeze the dependence on the x_1 and x_2 coordinates by setting

$$\partial = h_3 \partial_3, \quad \not{x} = x_3 h_3, \quad r = \sqrt{x_3^2}. \quad (67)$$

The (1), (3), (4), and (6) operators continue, with the above positions, to satisfy the $sl(2|1)$ (anti)commutators (5). The resulting $H_{1D,osc}$ deformed oscillator, given by (we set, for simplicity, $x = x_3$)

$$H_{1D,osc} = -\frac{1}{2} \partial_x^2 \cdot \mathbb{I}_4 + \frac{1}{2x^2} (\beta^2 \cdot \mathbb{I}_4 + \beta N_F) + \frac{1}{2} x^2 \cdot \mathbb{I}_4, \quad (68)$$

coincides with the model derived in Ref. [26] from the quantization of the world-line sigma model induced by the (1,4,3) supermultiplet (namely, with one propagating boson, four propagating fermions, and three auxiliary fields). The $H_{1D,osc}$ Hamiltonian possesses the larger $D(2, 1; \alpha)$ spectrum-generating superalgebra, with $\alpha = \beta - \frac{1}{2}$. On the other hand, as recalled in Ref. [30], the $sl(2|1) \subset D(2, 1; \alpha)$ generators are sufficient to determine the ray vectors of the Hilbert space (the vectors produced by the remaining generators differ by an inessential phase). From the dimensional reduction viewpoint, the extra generators entering $D(2, 1; \alpha)$ are associated with an emergent symmetry, not manifest from the $sl(2|1)$ construction.

VIII. CONCLUSIONS

We provided a direct construction of a three-dimensional superconformal quantum mechanics and of its associated de Alfaro-Fubini-Furlan deformed oscillator. Our approach allowed us to overcome the difficulties (pinpointed in Ref. [26]) in quantizing world-line superconformal sigma models with $D > 2$ target dimensions. It is rewarding that a dimensional reduction of the model allows us to recover the $D = 1, 2$ quantized world-line superconformal sigma models introduced in Ref. [26]. For the 3D deformed oscillator, $sl(2|1)$ acts as a spectrum-generating superalgebra; the complete spectrum is obtained from a (infinite) direct sum of $sl(2|1)$ lowest-weight representations. Depending on the coupling constant β , different Hilbert spaces can be consistently selected by suitably restricting the allowed $sl(2|1)$ lowest-weight representations.

Even if $sl(2|1)$ is sufficient to completely determine the spectrum, extra symmetries can play a role (possibly as enhanced symmetries at given values of the coupling constant β). This is worth investigating, since the recalled arbitrariness in the selection of the Hilbert space can be eliminated if an enlarged algebra is responsible for extra conditions to be imposed.

In this paper, we were able to determine the three-dimensional quantum Hamiltonians [given in Eqs. (3) and (9) for the superconformal and, respectively, deformed oscillator cases]. Once the Hamiltonian has been individuated, there is a standard method to be applied (see [39]) to derive the symmetry operators of the associated partial differential equation (for the case at hand, the partial

differential equation is the time-dependent Schrödinger equation for the given Hamiltonian). By applying this method, the most general dynamical symmetry of the model can be derived through a very lengthy but straightforward procedure. We plan to address this investigation in a future work. It allows one to answer the open question about the possible existence of extra symmetries and of the role they could play. It is worth mentioning that a similar analysis, conducted for a one-dimensional superconformal quantum mechanics, led to the surprising result [40] that the AdS_2 higher-spin superalgebra [41] is a dynamical symmetry of the model. A tantalizing possibility is that the extra symmetries, as those mentioned in Appendix C, of the superconformal quantum mechanics could be linked to a Bondi-Metzner-Sachs (BMS)/CFT correspondence (see [42]), the BMS transformations being associated with the “large” diffeomorphisms acting on asymptotically flat space-times [43].

ACKNOWLEDGMENTS

The authors are grateful to Naruhiko Aizawa and Zhanna Kuznetsova for helpful discussions. This work is supported by CNPq (PQ Grant No. 308095/2017-0).

APPENDIX A: NOTATIONS AND CONVENTIONS

We collect here for convenience the relevant notations and conventions used throughout the paper.

The $n \times n$ identity matrix is denoted as “ $\mathbb{I}_n$.” The three Pauli matrices σ_i are

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (A1)$$

The following 4×4 complex matrices γ_j and h_j ($j = 1, 2, 3$) are introduced:

$$\gamma_j = \sigma_j \otimes \mathbb{I}_2, \quad h_j = i\mathbb{I}_2 \otimes \sigma_j. \quad (A2)$$

The matrices γ_1, γ_2 are block antidiagonal, while $\gamma_3 = -i\gamma_1\gamma_2$ is the fermion parity operator N_F which defines bosons and fermions as its eigenvectors with eigenvalues $+1$ and -1 , respectively. When we need to stress this property, we use the notation

$$N_F = \gamma_3. \quad (A3)$$

The imaginary unit “ i ” is introduced in the definition of the matrices h_j , so that they furnish a representation of the three imaginary quaternions. Indeed, the composition law

$$h_i \cdot h_j = -\delta_{ij} \mathbb{I}_4 - \epsilon_{ijk} h_k \quad (A4)$$

is satisfied. The totally antisymmetric tensor ϵ_{ijk} is normalized so that $\epsilon_{123} = 1$. Throughout the paper, the

Einstein convention of summation over the repeated indices is understood unless otherwise specified.

By construction, the matrices γ_i commute with the quaternionic matrices h_j :

$$[\gamma_i, h_j] = 0 \quad \forall i, j = 1, 2, 3. \quad (\text{A5})$$

In Cartesian coordinates, the position vector $\vec{r}$ has components x_i . The orbital angular momentum $\vec{L}$ has components L_i , where

$$L_i = -i\epsilon_{ijk}x_j\partial_k. \quad (\text{A6})$$

For convenience, we introduced the slashed notation for the Euclidean version of the Dirac operator and a matrix-valued space coordinates operator, by setting

$$\not{\partial} := \vec{\nabla} \cdot \vec{h} = \partial_i h_i, \quad \not{r} := \vec{r} \cdot \vec{h} = x_i h_i. \quad (\text{A7})$$

It follows, in particular, that $\not{\partial}^2 = -\nabla^2$ and $\not{r}^2 = -r^2$, where $r = \sqrt{x_1^2 + x_2^2 + x_3^2}$ is the radial coordinate. In our conventions, the spherical coordinates r, θ, ϕ (restricted to $r \geq 0$, $0 \leq \theta \leq \pi$, and $0 \leq \phi < 2\pi$) are introduced through the positions

$$x_1 = r \sin \theta \cos \phi, \quad x_2 = r \sin \theta \sin \phi, \quad x_3 = r \cos \theta. \quad (\text{A8})$$

The spin $\vec{S}$ has components $\frac{1}{2}\sigma_i$, so that

$$\vec{S} = \frac{1}{2}\vec{\sigma}; \quad \text{then } \vec{S}^2 = \frac{3}{4}\mathbb{I}_2 = s(s+1)\mathbb{I}_2 \quad \text{with } s = \frac{1}{2}. \quad (\text{A9})$$

APPENDIX B: THE ORTHONORMAL EIGENSTATES

We present for completeness the orthonormal eigenstates. The derivation of the orthonormal conditions is based on some key observations and a lengthy but straightforward inductive proof concerning the excited states. We outline the main points and give the final results.

At first, one has to mention that the spin spherical harmonics $\mathcal{Y}_{j,l,m}(\theta, \phi)$ introduced in Eq. (12) are orthonormal, satisfying

$$\int d\Omega \mathcal{Y}_{j,l,m}^\dagger(\theta, \phi) \mathcal{Y}_{j',l',m'}(\theta, \phi) = \delta_{jj'} \delta_{ll'} \delta_{mm'}, \quad (\text{B1})$$

with $d\Omega$ an infinitesimal solid angle.

By setting $y = r^2$, the integral (37) produces a Gamma function, so that

$$\begin{aligned} \int_0^{+\infty} dr r^2 r^{2\gamma_{(j,\delta,\epsilon)}} e^{-r^2} &= \frac{1}{2} \int_0^{+\infty} dy y^{\gamma_{(j,\delta,\epsilon)} + (1/2)} e^{-y} \\ &= \frac{1}{2} \Gamma\left(\gamma_{(j,\delta,\epsilon)} + \frac{3}{2}\right). \end{aligned} \quad (\text{B2})$$

The lowest-weight vectors $\Psi_{j,\delta,m}^\epsilon(r, \theta, \phi)$ from Eq. (21) which satisfy, at given β , the (20) condition are orthogonal. The orthonormal wave functions $\Psi_{N,j,\delta,m}^\epsilon(r, \theta, \phi)$ are expressed in terms of the normalizing factor $M_{j,\delta}^{\epsilon,\beta}$. We have

$$\begin{aligned} M_{j,\delta}^{\epsilon,\beta} &= \sqrt{\frac{2}{\Gamma(\epsilon\beta + \delta j + \frac{1}{2}(1 + \delta))}}, \\ \Psi_{N,j,\delta,m}^\epsilon(r, \theta, \phi) &= M_{j,\delta}^{\epsilon,\beta} \Psi_{j,\delta,m}^\epsilon(r, \theta, \phi), \end{aligned} \quad (\text{B3})$$

so that

$$\langle \Psi_{N,j,\delta,m}^\epsilon(r, \theta, \phi) | \Psi_{N,j',\delta',m'}^{\epsilon'}(r, \theta, \phi) \rangle = \delta_{jj'} \delta_{mm'} \delta_{\epsilon\epsilon'} \delta_{\delta\delta'}. \quad (\text{B4})$$

The excited eigenstates $(a_1^+)^k \Psi_{j,\delta,m}^\epsilon(r, \theta, \phi)$, obtained by applying k times the a_1^+ creation operator (17), are orthogonal, as can be easily verified. The computation of their normalization factors which make the wave functions orthonormal is more cumbersome. It involves the computation of Rodrigues-type formulas (see [44]) for recursive polynomials in the radial coordinate r . These recursive polynomials can be recovered from the associated Laguerre polynomials. In order to proceed, we recall that

$$a_1^+ = \frac{1}{\sqrt{2}} \gamma_1 \not{r} \left(\mathbb{I}_4 \cdot (\partial_r - r) - \frac{2}{r} \mathbb{I}_2 \otimes \vec{S} \cdot \vec{L} - \frac{\beta}{r} N_F \right) \quad (\text{B5})$$

and that the (unnormalized) lowest-weight vectors $\Psi_{j,\delta,m}^\epsilon(r, \theta, \phi)$ from Eq. (21) are

$$\begin{aligned} \Psi_{j,\delta,m}^\epsilon(r, \theta, \phi) &= e_\epsilon \otimes \mathcal{Y}_{j,j-(1/2)\delta,m}(\theta, \phi) \cdot r^{\epsilon\beta + \delta j + (1/2)\delta - 1} e^{-(1/2)r^2}. \end{aligned} \quad (\text{B6})$$

The action of $\vec{S} \cdot \vec{L}$ can be read from Eq. (7), while the action of $\not{r}$ can be read from the

$$\frac{\vec{r} \cdot \vec{\sigma}}{r} \mathcal{Y}_{j,j-(1/2)\delta,m}(\theta, \phi) = -\mathcal{Y}_{j,j+(1/2)\delta,m}(\theta, \phi) \quad (\text{B7})$$

identity. The even and odd excited states are therefore, respectively, given by

$$\begin{aligned} (a_1^+)^{2k} \Psi_{j,\delta,m}^\epsilon(r, \theta, \phi) &= e_\epsilon \otimes \mathcal{Y}_{j,j-(1/2)\delta,m}(\theta, \phi) \cdot (-2)^k p_{2k,j}^{\epsilon,\delta,\beta}(r) r^{\epsilon\beta + \delta j + (1/2)\delta - 1} e^{-(1/2)r^2}, \\ (a_1^+)^{2k+1} \Psi_{j,\delta,m}^\epsilon(r, \theta, \phi) &= i\sqrt{2} e_{-\epsilon} \otimes \mathcal{Y}_{j,j+(1/2)\delta,m}(\theta, \phi) \cdot (-2)^k p_{2k+1,j}^{\epsilon,\delta,\beta}(r) r^{\epsilon\beta + \delta j + (1/2)\delta - 1} e^{-(1/2)r^2}, \end{aligned} \quad (\text{B8})$$

where $p_{2k,j}^{\epsilon,\delta,\beta}(r)$ and $p_{2k+1,j}^{\epsilon,\delta,\beta}(r)$ are r -dependent polynomials recursively determined by the Rodrigues-type formulas

$$\begin{aligned} p_{2k,j}^{\epsilon,\delta,\beta}(r) &= \frac{1}{2^{2k}} (r^{-\bar{\gamma}} e^{r^2/2} \quad 0) \begin{pmatrix} 0 & \partial_r - r + \frac{\bar{\gamma}+2}{r} \\ \partial_r - r - \frac{\bar{\gamma}}{r} & 0 \end{pmatrix}^{2k} \begin{pmatrix} r^{\bar{\gamma}} e^{-r^2/2} \\ 0 \end{pmatrix}, \\ p_{2k+1,j}^{\epsilon,\delta,\beta}(r) &= \frac{1}{2^{2k+1}} (r^{-\bar{\gamma}} e^{r^2/2} \quad 0) \begin{pmatrix} 0 & \partial_r - r + \frac{\bar{\gamma}+2}{r} \\ \partial_r - r - \frac{\bar{\gamma}}{r} & 0 \end{pmatrix}^{2k+1} \begin{pmatrix} 0 \\ r^{\bar{\gamma}} e^{-r^2/2} \end{pmatrix}, \end{aligned} \quad (\text{B9})$$

where [see (23)]

$$\bar{\gamma} \equiv \gamma_{(j,\delta,\epsilon)}(\beta) = \epsilon\beta + \delta j + \frac{1}{2}\delta - 1. \quad (\text{B10})$$

The connection with the associated Laguerre polynomials is as follows [we discuss it explicitly for the even polynomials $p_{2k,j}^{\epsilon,\delta,\beta}(r)$, and the derivation for the odd polynomials $p_{2k+1,j}^{\epsilon,\delta,\beta}(r)$ is made along the same lines]. We obtain, from Eq. (B9), the relation

$$\begin{aligned} p_{2(k+1),j}^{\epsilon,\delta,\beta}(r) &= \frac{1}{4} r^{-\bar{\gamma}} e^{r^2/2} \left(\partial_r - r + \frac{\bar{\gamma}+2}{r} \right) \left(\partial_r - r - \frac{\bar{\gamma}}{r} \right) \\ &\quad \times p_{2k,j}^{\epsilon,\delta,\beta}(r) r^{\bar{\gamma}} e^{-r^2/2}. \end{aligned} \quad (\text{B11})$$

It follows, in particular, from $p_{0,j}^{\epsilon,\delta,\beta}(r) = 1$, that

$$p_{2,j}^{\epsilon,\delta,\beta}(r) = r^2 - \bar{\gamma} - \frac{3}{2}. \quad (\text{B12})$$

The associated Laguerre polynomials $L_k^{(\gamma)}(x)$ are introduced through the position

$$L_k^{(\gamma)}(x) = \frac{x^{-\gamma} e^x}{k!} \left(\frac{d}{dx} \right)^k x^{\gamma+k} e^{-x}. \quad (\text{B13})$$

They satisfy the identities

$$\begin{aligned} L_k^{(\gamma)}(x) &= L_k^{(\gamma+1)}(x) - L_{k-1}^{(\gamma+1)}(x), \\ x L_{k-1}^{(\gamma+1)}(x) &= (\gamma + k) L_{k-1}^{(\gamma)}(x) - k L_k^{(\gamma)}(x). \end{aligned} \quad (\text{B14})$$

Since

$$L_1^{(\gamma)}(x) = -x + \gamma - 1, \quad (\text{B15})$$

by setting

$$x = r^2, \quad \gamma = \bar{\gamma} + \frac{1}{2}, \quad (\text{B16})$$

we can identify

$$p_{2,j}^{\epsilon,\delta,\beta}(r) = -L_1^{\bar{\gamma}+(1/2)}(r^2). \quad (\text{B17})$$

By assuming the ansatz

$$p_{2k,j}^{\epsilon,\delta,\beta}(r) = C_k L_k^{\bar{\gamma}+(1/2)}(r^2), \quad (\text{B18})$$

after lengthy computations and the use of the (B14) identities, one arrives at the inductive proof that Eq. (B18) is satisfied, provided that

$$C_k = (-1)^k k! \quad (\text{B19})$$

The $p_{2k,j}^{\epsilon,\delta,\beta}(r)$ even and $p_{2k+1,j}^{\epsilon,\delta,\beta}(r)$ odd polynomials are expressed, in terms of the associated Laguerre polynomials, as

$$\begin{aligned} p_{2k,j}^{\epsilon,\delta,\beta}(r) &= (-1)^k k! L_k^{\bar{\gamma}+(1/2)}(r^2), \\ p_{2k+1,j}^{\epsilon,\delta,\beta}(r) &= (-1)^{k+1} k! r L_k^{\bar{\gamma}+(3/2)}(r^2). \end{aligned} \quad (\text{B20})$$

By plugging these expressions into the (B8) formulas, we are able to determine the normalizing factors that need to be used to construct orthonormal excited eigenstates. The normalizing factors are recovered from the orthogonal relations for the associated Laguerre polynomials, given by

$$\int_0^{+\infty} dx x^\gamma e^{-x} L_n^{(\gamma)}(x) L_m^{(\gamma)}(x) = \frac{\Gamma(n+\gamma+1)}{n!} \delta_{nm}. \quad (\text{B21})$$

The final expressions for the orthonormal wave functions $\Psi_{N,k,j,\delta,m}^\epsilon(r, \theta, \phi)$ are

$$\begin{aligned} \Psi_{N,2k,j,\delta,m}^\epsilon(r, \theta, \phi) \\ = e_\epsilon \otimes \mathcal{Y}_{j,j-(1/2)\delta,m}(\theta, \phi) \cdot M_{2k}^{\bar{\gamma}} L_k^{\bar{\gamma}+(1/2)}(r^2) \cdot r^{\bar{\gamma}} e^{-r^2/2} \end{aligned} \quad (\text{B22})$$

with

$$M_{2k}^{\bar{\gamma}} = \sqrt{\frac{(k!) \cdot 2}{\Gamma(k + \bar{\gamma} + \frac{3}{2})}} \quad (\text{B23})$$

and

$$\begin{aligned} \Psi_{N,2k+1,j,\delta,m}^\epsilon(r, \theta, \phi) \\ = e_{-\epsilon} \otimes \mathcal{Y}_{j,j+(1/2)\delta,m}(\theta, \phi) \cdot M_{2k+1}^{\bar{\gamma}} L_k^{\bar{\gamma}+(3/2)}(r^2) \cdot r^{\bar{\gamma}+1} e^{-r^2/2} \end{aligned} \quad (\text{B24})$$

with

$$M_{2k+1}^{\tilde{\gamma}} = \sqrt{\frac{(k!) \cdot 2}{\Gamma(k + \tilde{\gamma} + \frac{5}{2})}}. \quad (\text{B25})$$

The parameter $\tilde{\gamma} = \gamma_{(j,\delta,e)}(\beta)$ was introduced in Eq. (B10). One should note that $M_0^{\tilde{\gamma}}$ reproduces, as it should, the normalizing factor given in formula (B3).

APPENDIX C: A GRAPHICAL ILLUSTRATION OF AN OPEN PROBLEM

The three-dimensional deformed oscillator can be completely solved and its energy spectrum and associated degeneracy computed. On the other hand, as discussed in Secs. V and VI, alternative admissible choices of Hilbert space lead to different results corresponding to different quantum models. This means that the spectrum-generating superalgebra provides a (relevant) piece of information, but it does not allow one to uniquely deduce the spectrum and the degeneracy of the β -deformed model. Some extra information should be supplemented. It is quite possible that (yet to be understood) larger algebraic structures could be responsible for that, in such a way that they allow one to uniquely pinpoint a given quantum model. The source of this ambiguity is the fact that the Hilbert space is not expressed by a single irreducible representation but by a direct sum of several (infinite) lowest-weight representations of the $sl(2|1)$ spectrum-generating superalgebra. Consistent selections of subsets of the set of lowest-weight representations therefore lead to different Hilbert spaces.

The open problem of searching for enlarged algebraic structures is not specific to the three-dimensional deformed oscillator; it applies to the whole class of related (deformed) oscillators. It already appears in the case of the two-dimensional oscillator which corresponds [see formula (66)] to a dimensional reduction of the 3D oscillator. It is quite appealing to focus on this simpler case, because it offers a nice graphical visualization of the question at hand.

For our illustrative purpose, it is sufficient to take a superselected version of the undeformed two-dimensional theory with a unique bosonic vacuum and integerly spaced energy excited states (the energy degeneracy growing linearly) arranged in a triangular shape as shown in Figs. 4 and 5 below. As discussed in Sec. III, the ray vectors are determined by the $osp(1|2) \subset sl(2|1)$ spectrum-generating subalgebra [the action of the remaining $sl(2|1)$ generators produce inessential phases]. In Ref. [26], it was shown that a new set of operators, obtained by conjugating the original operators of the $sl(2|1)$ spectrum-generating superalgebra, produce another $sl(2|1)$ superalgebra, denoted as $sl(2|1)_C$. The details of the Ref. [26] construction are not relevant here. What is relevant is that the conjugated operators admit a nice

FIG. 4. Superselected 2D oscillator. The bosonic (fermionic) eigenstates are represented by black (white) dots. The y axis labels the integerly spaced energy eigenvalues, while the x axis labels the $so(2)$ spin components. The black dot at the bottom corresponds to the bosonic vacuum. The solid edges represent the action of the creation operator from the $osp(1|2) \subset sl(2|1)$ subalgebra. Infinite $osp(1|2)$ lowest-weight representations (a new lowest-weight vector at any given energy eigenvalue) are required to produce the spectrum of the theory.

FIG. 5. A mirror dual of the previous figure. The dashed edges represent the action of the creation operator from the $osp(1|2)_C \subset sl(2|1)_C$ subalgebra, where $sl(2|1)_C$ is an $sl(2|1)$ superalgebra produced by a new set of operators obtained by “mirroring” the operators entering the original $sl(2|1)$ spectrum-generating superalgebra. As before, infinite $osp(1|2)_C$ lowest-weight representations are required to produce the spectrum. On the other hand, any energy eigenstate can be obtained from the bosonic vacuum through a path combining both solid and dashed edges.

“mirror” interpretation, visualized in Fig. 5. The huge enveloping algebra induced by the operators entering both $sl(2|1)$ and $sl(2|1)_C$ uniquely determines the spectrum of the theory, as explained in the comment about Fig. 5.

An enlarged algebraic structure, producing the three-dimensional counterpart of this simpler two-dimensional setting, has to be properly investigated.

- [1] V. de Alfaro, S. Fubini, and G. Furlan, *Nuovo Cimento A* **34**, 569 (1976).
- [2] H. Miyazaki and I. Tsutsui, *Ann. Phys. (Amsterdam)* **299**, 78 (2002).
- [3] L. Fehér, I. Tsutsui, and T. Fülöp, *Nucl. Phys. B* **715**, 713 (2005).
- [4] L. C. Biedenharn and J. D. Louck, Angular momentum in quantum physics: Theory and application, in *Encyclopedia of Mathematics* (Addison-Wesley, Reading, MA, 1981), p. 283, Vol. 8, ISBN:0-201-13507-8.
- [5] F. Calogero, *J. Math. Phys. (N.Y.)* **10**, 2191 (1969).
- [6] S. Fubini and E. Rabinovici, *Nucl. Phys. B* **245**, 17 (1984).
- [7] V. P. Akulov and A. I. Pashnev, *Teor. Mat. Fiz.* **56**, 344 (1983) [*Theor. Math. Phys.* **56**, 862 (1983)].
- [8] R. Britto-Pacumio, J. Michelson, A. Strominger, and A. Volovich, Lectures on superconformal quantum mechanics and multi-black hole moduli spaces, in *Progress in String Theory and M-Theory*, NATO Science Series Vol. 564 (Kluwer Academic, Dordrecht, 2001), p. 235.
- [9] A. Sen, *J. High Energy Phys.* **11** (2008) 075.
- [10] C. Chamon, R. Jackiw, S.-Y. Pi, and L. Santos, *Phys. Lett. B* **701**, 503 (2011).
- [11] J. Maldacena and D. Stanford, *Phys. Rev. D* **94**, 106002 (2016).
- [12] A. Kitaev, A simple model of quantum holography, in *Proceedings of the KITP*, <http://online.kitp.ucsb.edu/online/entangled15/kitaev/> (2015) and <http://online.kitp.ucsb.edu/online/entangled15/kitaev2/> (2015).
- [13] J. Maldacena, D. Stanford, and Z. Yang, *Prog. Theor. Exp. Phys.* **2016**, 12C104 (2016).
- [14] M. Cárdenas, O. Fuentealba, H. A. González, D. Grumiller, C. Valcárcel, and D. Vassilevich, *J. High Energy Phys.* **11** (2018) 077.
- [15] A. Pashnev and F. Toppan, *J. Math. Phys. (N.Y.)* **42**, 5257 (2001).
- [16] Z. Kuznetsova, M. Rojas, and F. Toppan, *J. High Energy Phys.* **03** (2006) 098.
- [17] S. Fedoruk, E. Ivanov, and O. Lechtenfeld, *J. Phys. A* **45**, 173001 (2012).
- [18] E. Ivanov and S. Sidorov, *Classical Quantum Gravity* **31**, 075013 (2014).
- [19] S. Fedoruk and E. Ivanov, *J. High Energy Phys.* **10** (2015) 087.
- [20] E. Ivanov, O. Lechtenfeld, and S. Sidorov, *J. High Energy Phys.* **11** (2016) 031.
- [21] E. Ivanov, O. Lechtenfeld, and S. Sidorov, *J. High Energy Phys.* **08** (2018) 193.
- [22] Z. Kuznetsova and F. Toppan, *J. Math. Phys. (N.Y.)* **53**, 043513 (2012).
- [23] S. Khodae and F. Toppan, *J. Math. Phys. (N.Y.)* **53**, 103518 (2012).
- [24] N. L. Holanda and F. Toppan, *J. Math. Phys. (N.Y.)* **55**, 061703 (2014).
- [25] E. Ivanov and S. Sidorov, *J. Phys. A* **47**, 292002 (2014).
- [26] I. E. Cunha, N. L. Holanda, and F. Toppan, *Phys. Rev. D* **96**, 065014 (2017).
- [27] S. Fedoruk, E. Ivanov, and S. Sidorov, *J. High Energy Phys.* **01** (2018) 132.
- [28] N. Aizawa, Z. Kuznetsova, and F. Toppan, *J. Math. Phys. (N.Y.)* **59**, 022101 (2018).
- [29] S. Fedoruk, E. Ivanov, O. Lechtenfeld, and S. Sidorov, *J. High Energy Phys.* **04** (2018) 043.
- [30] N. Aizawa, I. Cunha, Z. Kuznetsova, and F. Toppan, *J. Math. Phys. (N.Y.)* **60**, 042102 (2019).
- [31] E. D'Hoker and L. Vinet, *Commun. Math. Phys.* **97**, 391 (1985).
- [32] E. Ivanov, S. Sidorov, and F. Toppan, *Phys. Rev. D* **91**, 085032 (2015).
- [33] E. Ivanov and S. Sidorov, *Classical Quantum Gravity* **33**, 055001 (2016).
- [34] N. Kozyrev, S. Krivonos, O. Lechtenfeld, and A. Sutulin, *J. High Energy Phys.* **05** (2018) 175.
- [35] A. B. Balantekin, *Ann. Phys. (N.Y.)* **164**, 277 (1985).
- [36] S. Okubo, *J. Math. Phys. (N.Y.)* **32**, 1657 (1991).
- [37] H. L. Carrion, M. Rojas, and F. Toppan, *J. High Energy Phys.* **04** (2003) 040.
- [38] F. Toppan, in *Symmetries and Groups in Contemporary Physics, Proceedings of the XXIX International Colloquium on Group-Theoretical Methods in Physics, Tianjin, China, 2012*, Nankai Series in Pure, Applied Mathematics and Theoretical Physics Vol. 11 (World Scientific, Singapore, 2013), pp. 417–422.
- [39] P. J. Olver, *Applications of Lie Groups to Differential Equations*, 2nd ed. (Springer, New York, 1991).
- [40] F. Toppan and M. Valenzuela, *Adv. Math. Phys.* **2018**, 6263150 (2018).
- [41] M. A. Vasiliev, *Int. J. Mod. Phys. A* **06**, 1115 (1991).
- [42] G. Barnich and C. Troessaert, *J. High Energy Phys.* **05** (2010) 062.
- [43] A. Strominger, *J. High Energy Phys.* **07** (2014) 152.
- [44] G. Szegő, *Orthogonal Polynomials*, 4th ed. (American Mathematical Society, Providence, 1975).