Correlation measures and distillable entanglement in AdS/CFT

Joshua Levin^{*,†}, Oliver DeWolfe[†], and Graeme Smith[‡]

*Department of Physics and Center for Theory of Quantum Matter, University of Colorado,
Boulder, Colorado 80309, USA*

(Received 18 December 2019; accepted 24 January 2020; published 13 February 2020)

Recent developments have exposed close connections between quantum information and holography. In this paper, we explore the geometrical interpretations of the recently introduced Q -correlation and R -correlation, E_Q and E_R . We find that E_Q admits a natural geometric interpretation via the surface-state correspondence: it is a minimal mutual information between a surface region A and a cross section of A 's entanglement wedge with B . We note a strict trade-off between this minimal mutual information and the symmetric side-channel-assisted distillable entanglement from the environment E to A , $I^{ss}(E:A)$. We also show that the R -correlation, E_R , coincides holographically with the entanglement wedge cross section. This further elucidates the intricate relationship between entanglement, correlations, and geometry in holographic field theories.

DOI: [10.1103/PhysRevD.101.046015](https://doi.org/10.1103/PhysRevD.101.046015)

I. INTRODUCTION

Information measures quantify the various types of information and disorder contained in the state of a quantum system. Familiar information measures such as the mutual information are built as linear combinations of entropies; others, such as the squashed entanglement [1], involve the optimization over additional auxiliary degrees of freedom (d.o.f.). A particularly useful class of information measures, which we refer to as *correlation measures*, are those that are monotonically decreasing under local quantum operations. Such a monotonic measure captures properties of the state that individual parties cannot create on their own, and thus quantifies correlations between parties.

The entanglement of purification is an especially interesting information measure [2]. It is defined as a minimization over purifications of the state ρ_{AB} to $|\psi\rangle_{AaBb}$ of the entropy $S(Aa)$,

$$E_P(A:B) = \inf_{|\psi\rangle_{AaBb}} S(Aa), \quad (1)$$

and can be understood operationally as the entanglement cost of creating the state ρ_{AB} with asymptotically vanishing communication between parties. E_P is monotonic under

quantum operations on either party [2,3], and is thus a correlation measure. In Ref. [3], two similar bipartite correlation measures were identified: the Q -correlation $E_Q(A:B)$ and the R -correlation $E_R(A:B)$ (defined below). In addition to monotonicity all three of these correlation measures satisfy the inequalities

$$\begin{aligned} \frac{1}{2} I(A:B) &\leq E_\alpha(A:B) \leq \min(S(A), S(B)), \\ E_\alpha(A:BC) &\geq \frac{1}{2} (I(A:B) + I(A:C)), \end{aligned} \quad (2)$$

for $\alpha = P, Q, R$, where $I(A:B) = S(A) + S(B) - S(AB)$ is the mutual information. It is of considerable interest to better characterize the new correlation measures, their operational significance, and their relationship to E_P .

In recent years, powerful new tools for exploring quantum information have arisen using holography. The AdS/CFT correspondence relates d -dimensional conformal field theories (CFTs) to theories of gravity living in asymptotically anti-de Sitter (AdS) spacetimes in $d+1$ dimensions, where the CFT is thought of as living at the boundary of AdS. In AdS/CFT, the entropy of a spatial region A in the CFT can be calculated in terms of the gravity dual using the Ryu-Takayanagi (RT) formula [4], which states that the entropy is proportional to the area of the minimal bulk surface Γ_A living in a Cauchy slice Σ of AdS homologous to A , whose boundary is ∂A^1 :

$$S(A) = \frac{\text{Area}(\Gamma_A)}{4G_N}, \quad (3)$$

¹Here we restrict to time-independent states where this definition is sufficient.

^{*}joshua.t.levin@colorado.edu
[†]oliver.dewolfe@colorado.edu
[‡]graeme.smith@colorado.edu

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

where G_N is the gravitational constant. Inequalities satisfied by information measures such as weak and strong subadditivity can be shown to follow geometrically from the RT formula. The region in Σ bounded by A and Γ_A is called the *entanglement wedge* of A , $EW(A)$.

It is natural to ask whether correlation measures involving the optimization over ancillary d.o.f. such as E_P , E_Q and E_R have a holographic presentation. A proposal for the entanglement of purification was made in Refs. [5,6], as follows. Consider two disjoint, connected spatial CFT regions A and B , which for the sake of concreteness we can take to live on the boundary of a spatial slice in an asymptotically AdS_3 geometry as depicted in Fig. 1, where “surfaces” and “curves” are interchangeable. For sufficiently small A and B one has that $S(AB) = S(A) + S(B)$ (with RT surfaces depicted by Γ_A and Γ_B in Fig. 1), and hence the entanglement wedge $EW(AB)$ is the sum of the disconnected components $EW(A)$ and $EW(B)$. On the other hand, for sufficiently large A and B the minimal surface giving $S(AB)$ becomes the dashed curves in Fig. 1, leading to a single, connected entanglement wedge spanning the space from A to B .

If $EW(AB)$ is connected then we can define a quantity called the *entanglement wedge cross section*, denoted $E_W(A:B)$, as the minimal area of a surface partitioning $EW(AB)$ into two regions: one containing A in its boundary and the other containing B . We denote this minimal surface by $EWCS(AB)$. The surfaces bounding the entanglement wedge are necessarily geodesics, and it is a fact of two-dimensional hyperbolic geometry that any two complete parallel geodesics (which do not meet at infinity) have a unique common perpendicular, which minimizes the distance between the two geodesics; this common perpendicular, labeled M in Fig. 1, is therefore $EWCS(AB)$. The proposal is then that the entanglement of purification can be calculated holographically as proportional to $E_W(A:B)$ [5,6]:

$$E_P(A:B) = \frac{E_W(A:B)}{4G_N}. \quad (4)$$

Interestingly, the primary evidence for this E_P - E_W correspondence was monotonicity and the inequalities (2). Given that E_Q and E_R are now known to satisfy the same properties, it is useful to have another way of thinking about the issue, to suggest whether E_P is indeed the correct dual for E_W , and to identify the proper gravity dual for the other correlation measures. This involves understanding how an information measure involving an optimization over additional d.o.f. should be calculated holographically.

Comparing the definition of E_P to the geometry in Fig. 1 provides a hint. In calculating E_P we are instructed to purify the system ρ_{AB} , and then find the partition of the new d.o.f. into a and b such that $S(Aa)$ is minimized. Consider for concreteness the state ρ_{AB} obtained from the vacuum of the CFT by tracing out other d.o.f. Clearly one purification of this is the vacuum state itself, which geometrically corresponds to connecting A and B using the remainder of the boundary circle, denoted E in Fig. 1; however there is no reason to believe this is the optimal purification. Looking at the geometry, if we allow ourselves to think of the dashed RT surfaces as spatial regions corresponding to purifications of ρ_{AB} , then we can associate a with the left half and b with the right half as indicated in the figure, and then E_W does indeed extremize the area and gives exactly the entropy $S(Aa)$, leading back to the E_P - E_W conjecture (4). However, for this to make sense we must have a way of associating states not just with boundary regions, but with surfaces in the bulk of the gravity theory.

The surface-state correspondence (SSC) [7] is a generalization of AdS/CFT in which a Hilbert space $\mathcal{H}_{\text{tot}}$ is associated not only to the AdS boundary, but to surfaces in the interior of the geometry as well. A pure state on $\mathcal{H}_{\text{tot}}$ is dual to a space-like closed convex surface of codimension 2 (an embedding of S^{n-2} into a Cauchy slice Σ of AdS_n whose image is convex in Σ), while a mixed state is dual to a strict subset of a closed convex surface, i.e., a pure state with a subsystem traced out (erased). The entropy of a mixed state is given by the area of the smallest surface which closes the dual surface; note that for states whose dual surfaces lie in the AdS boundary, this reduces exactly to the RT formula. An important implication of this is that a state which is dual to a geodesic in AdS must have no correlation between any of its subsystems; dividing the geodesic into disjoint pieces a and b and applying the entropy rule, one sees that $I(a:b) = 0$. A heuristic way to view the SSC is that given any space-like closed convex surface Ω of codimension 2, we have a nested instance of ordinary AdS/CFT , where the “boundary” is now Ω and the bulk is its interior.

Using SSC, we can understand better the previous arguments for the conjectured duality of E_W and E_P . Given disjoint regions A and B of the boundary of a Cauchy slice Σ

FIG. 1. The geometry of the entanglement wedge of ρ_{AB} . Γ_A and Γ_B are the RT surfaces of A and B , respectively, M is the entanglement wedge cross section, and $a_1 \cup a_2 \cup b_1 \cup b_2$ is the RT surface of AB .

of AdS, any curves connecting the boundaries of A and B into a closed convex surface correspond to a purification. To minimize $S(Aa)$ we must choose the purification and partition which minimizes correlations between a and b . Since geodesics have no correlation between their subsystems, the correct geometric purification corresponds to $\partial EW(AB)$, for which $I(a:b) = 0$ for any partition. SSC then tells us that $S(Aa)$ is given by the length of the curve M , whose minimization over the choice of $\{a, b\}$ is E_W by definition.

Tensor network models of AdS/CFT can offer further insight into this process (see e.g., Refs. [8,9]). The process of deforming a boundary region A into its bulk RT surface Γ_A can be thought of as a process of entanglement distillation, where the d.o.f. on the boundary are sorted into those entangled with other d.o.f. (tensors adjacent to the RT surface) and those that are not entangled, and are left behind. The RT surface for AB is also the RT surface Γ_E for the complement of A and B on the boundary, labeled E in Fig. 1. Thus we can view selecting $\partial EW(AB)$ as the optimal purification as a process of distilling the minimum set of d.o.f. needed to provide the entanglement with the d.o.f. in the entanglement wedge (and hence still represent a purification).

Several proposals besides E_P have been conjectured for the boundary dual of E_W [10–12]. In Ref. [10], it was argued that E_W should be dual to half of the *reflected entropy*, an upper bound for E_P that corresponds to a fixed configuration of the optimization in Eq. (1). In Ref. [11] the log negativity [13] was conjectured, while in Ref. [12] a quantity called the odd entanglement entropy was proposed to correspond to E_W . We do not explore these conjectures further in this paper, but restrict our attention to studying the implications of SSC (and the associated tensor-network models) for more general entropic formulas than E_P . We hope that our findings here can be generalized to give further geometric intuitions from perspectives such as those pursued in Refs. [10–12], but leave such exploration for future work.

In what follows, we will assume the validity of SSC, and apply these methods to calculating holographic duals for E_Q and E_R in the vacuum state of $1+1$ -dimensional holographic CFT. We shall see that in this case E_R is also given in terms of E_W using the same formula as E_P , Eq. (4), while E_Q has a different (but related) holographic realization.

II. Q - AND R -CORRELATION

A correlation measure is necessarily monotonically decreasing under local operations. Optimized correlation measures were studied axiomatically in Ref. [3], and two especially interesting ones found there were the Q -correlation and R -correlation, which can be defined as minimizations over purifications of the state ρ_{AB} to $|\psi\rangle_{AaBb}$:

$$\begin{aligned} E_Q(A:B) &= \inf_{|\psi\rangle_{AaBb}} f^Q(A, B, a), \\ E_R(A:B) &= \inf_{|\psi\rangle_{AaBb}} f^R(A, B, a) \end{aligned} \quad (5)$$

where the functions being minimized are

$$\begin{aligned} f^Q(A, B, a) &= \frac{1}{2}[S(A) + S(B) + S(Aa) - S(Ba)], \\ f^R(A, B, a) &= \frac{1}{2}[S(AB) + 2S(Aa) - S(ABa) - S(a)]. \end{aligned} \quad (6)$$

While we will work with these formulas, it is worth noting we may recast the definitions in terms of conditional entropies,

$$\begin{aligned} f^Q &= \frac{1}{2}[2S(A) + S(a|A) - S(a|B)], \\ f^R &= \frac{1}{2}[S(AB) + S(A|a) - S(B|aA)] \end{aligned} \quad (7)$$

or in terms of mutual informations,

$$\begin{aligned} f^Q &= \frac{1}{2}[2S(A) + I(B:a) - I(A:a)], \\ f^R &= \frac{1}{2}[2S(A) + I(B:a|A) - I(A:a)]. \end{aligned} \quad (8)$$

It also follows from this that the difference between the two is simply half the tripartite information,

$$f^Q - f^R = \frac{1}{2}I_3(A:B:a). \quad (9)$$

While the tripartite information in general can have either sign, in holographic states one can show that $I_3(A:B:C) \leq 0$, called the monogamy of mutual information (MMI) [14]. This implies that

$$E_Q(A:B) \leq E_R(A:B) \quad (10)$$

for any holographic state ρ_{AB} .

We now use SSC to identify holographic duals of E_Q and E_R in the vacuum state of a two-dimensional CFT, calculating entropies using the generalization of the RT formula to surfaces in the bulk.

A. Conformal invariance of E_Q and E_R

We can argue that in the vacuum state, conformal invariance means it is sufficient to consider a particular subclass of A and B regions without loss of generality. In general, the entanglement entropy of a region in quantum field theory is infinite, and the d.o.f. must be regulated to obtain a finite result. In the gravity dual, the divergences correspond to the infinite area of surfaces stretching all the way out to the boundary, and so these must be cut off

geometrically. In general $S(A)$ then depends on the cutoff, and conformal invariance is not directly realized.

For certain quantities however, this cutoff dependence cancels. An elementary example is the mutual information of two regions, $I(A:B) = S(A) + S(B) - S(AB)$. Geometrically, the diverging area of $S(A)$ and $S(B)$ is canceled by that of $S(AB)$, since each surface headed to the boundary with a plus sign in the RT expression for $I(A:B)$, is *balanced* by a corresponding surface headed to the same boundary point with a minus sign. As a result, $I(A:B)$ is cutoff independent and hence conformally invariant. Since it is associated to the four points bounding the A and B regions, $I(A:B)$ depends only on the single conformally invariant cross ratio of these four points.

For the quantities we are interested in, the story is the same. Calculations either involve surfaces that do not go to the boundary, and hence are cutoff independent, or go to the boundary at a given point in pairs with opposite signs, so that the balancing condition is satisfied and cutoff dependence cancels. As a result, once the partition M is chosen, and is allowed to transform under a conformal map via the associated bulk isometry, f^Q and f^R are also conformally invariant and depend only on the cross ratio of the four defining points. Therefore E_Q and E_R are also conformally invariant.

For ordered points on the real line, we define the cross ratio as $(x_2 - x_1)(x_4 - x_3)/(x_3 - x_1)(x_4 - x_2)$. Mapping the real line (thought of as the boundary of the upper half plane) to the boundary of the unit disk using the Cayley transform $f(z) = (z - i)/(z + i)$, the cross ratio in terms of angles on the circle is

$$\frac{\sin(\theta_{12}/2) \sin(\theta_{34}/2)}{\sin(\theta_{13}/2) \sin(\theta_{24}/2)}, \quad (11)$$

where $\theta_{ij} \equiv \theta_i - \theta_j$. This cross ratio can take any value between 0 and 1.

We can obtain a representative configuration for every possible cross ratio by considering only the “symmetric” cases where A and B are of equal size with diametrically opposite centers (as in Fig. 1), in which case the cross ratio reduces to just $\sin^2(\theta/2)$, where θ is the angular size of each region. Any other configuration of A and B shares its cross ratio value with one of these cases, and hence E_Q and E_R can always be calculated by mapping onto a symmetric case, using a bulk isometry that induces a boundary conformal map. (This would no longer be true for a configuration with less symmetry, like one containing a black hole.) Hence, in what follows, we will restrict to symmetric cases when necessary. Since the results are cutoff independent, we will always use a spatially uniform cutoff at constant radial coordinate.

B. Geometric duals of E_Q and E_R

Following the arguments in the previous subsection, we will state clearly when we are restricting to symmetric cases. In addition we are interested in geometries with

connected entanglement wedges ($\theta > \pi/2$ for symmetric cases). Following the arguments given in the Introduction, we assume the optimal purification of ρ_{AB} into $|\psi\rangle_{AaBb}$ corresponds to completing the A and B regions into a closed convex surface by connecting them along their RT surface Γ_{AB} . We then leave to be determined how to partition Γ_{AB} into surfaces a_1, a_2 associated to A and surfaces b_1, b_2 associated to B , corresponding to a partition of the purifying system into subsystems $\{a, b\}$. The following proofs are somewhat technical, and the reader interested in the results may find a discussion in the next subsection.

Consider first E_R . For each term in f^R we must use SSC to find the sum of bulk surfaces dual to the term. Using the labels in Fig. 1, $S(AB)$ is always given by $|a_1| + |b_1| + |a_2| + |b_2|$, and $S(Aa)$ is always given by $|M|$. For $S(ABa)$, there are two options according to SSC: $|b_1| + |b_2|$ or $|\Gamma_B| + |M|$. Similarly, there are two options for $S(a)$: $|a_1| + |a_2|$ or $|\Gamma_A| + |M|$. The following fact fixes these options for every partition.

Proposition 1: Given a connected $EW(AB)$ and any partition $\{a, b\}$ of the RT surface Γ_{AB} (as in Fig. 1), we have $|a_1| + |a_2| < |\Gamma_A| + |M|$ and $|b_1| + |b_2| < |\Gamma_B| + |M|$.

Proof.—We argue that $|a_1| + |a_2| < |\Gamma_A| + |M|$, and the proof for the other statement is identical. We will show that

$$|\Gamma_A| + |M| - |a_1| - |a_2| > I(A:B), \quad (12)$$

which will prove the claim since $I(A:B) \geq 0$. Writing $I(A:B)$ as a linear combination of lengths using the RT formula, Eq. (12) becomes

$$\begin{aligned} & |\Gamma_A| + |M| - |a_1| - |a_2| \\ & > |\Gamma_A| + |\Gamma_B| - |a_1| - |b_1| - |a_2| - |b_2| \end{aligned} \quad (13)$$

which is equivalent to

$$|M| > |\Gamma_B| - |b_1| - |b_2|, \quad (14)$$

or

$$|M| + |b_1| + |b_2| > |\Gamma_B|, \quad (15)$$

which is true since Γ_B is defined to be the minimal area surface connecting the two boundary points of B . ■

Corollary 1: Let $\{a, b\}$ be a partition of the RT surfaces bounding $EW(AB)$ (as in Fig. 1). Then $I(a:b) = 0$.

Proposition 1 tells us that in all cases, $S(ABa)$ is given by $|b_1| + |b_2|$ and $S(a)$ is given by $|a_1| + |a_2|$. Combining all four terms of f^R , we are left with only surface M , and therefore we conclude that for disjoint and connected boundary regions A and B ,

$$E_R(A:B) = \frac{E_W(A:B)}{4G_N}. \quad (16)$$

Thus for holographic vacuum states, E_R and E_P have the same realization as the entanglement wedge cross section. The additivity of E_R [3], as compared to the conjectured

nonadditivity of E_P [15], and its equivalence to E_W in a holographic scenario strengthens the hope of Ref. [5] that for holographic states E_P is in fact additive. It may also suggest that E_R should be considered the more fundamental dual of the entanglement wedge cross section. It would be interesting to find holographic states for which these two measures differ.

Now we address E_Q . For any partition $\{a, b\}$ of Γ_{AB} , $S(A)$ and $S(B)$ are always given by their RT surfaces ($|\Gamma_A|$ and $|\Gamma_B|$), and $S(Aa)$ is given by $|M|$. There are five possible bulk surface configurations for $S(Ba)$. For the three systems $\{a_1, a_2, B\}$, either all mutual informations are zero, each has positive mutual information with the other two, or one pair has positive mutual information while the pair has zero mutual information with the third. The latter case consists of three subcases: $I(a_1 : a_2) > 0$, $I(a_1 : B) > 0$, and $I(a_2 : B) > 0$. We will label these five cases by listing all systems in $\{a_1, a_2, B\}$ which have zero mutual information with the other two: $\{a_1, a_2, B\}$, $\emptyset$, $\{B\}$, $\{a_2\}$, and $\{a_1\}$, in the order mentioned in this paragraph. Proposition 1 implies that $\{B\}$ is never the case. In the four remaining cases, $S(Ba)$ is given by the bulk surfaces shown in Fig. 2. The linear combinations of bulk surfaces for f^Q in

each of the four cases are shown in Fig. 3. In cases $\{a_1, a_2, B\}$, $\emptyset$, $\{a_2\}$, and $\{a_1\}$, we denote by ΔA , ΔB , Δa_2 , and Δa_1 (respectively) the linear combinations of bulk surfaces which give f^Q . Then, since the only term in f^Q which has multiple options for its bulk surfaces is subtracted (and the correct choice is the minimum option), we have

$$f^Q = \frac{1}{2} \max\{\Delta A, \Delta B, \Delta a_2, \Delta a_1\} \quad (17)$$

for any partition. Therefore

$$E_Q(A : B) = \frac{1}{2} \inf_{|\psi\rangle_{AaBb}} \max\{\Delta A, \Delta B, \Delta a_2, \Delta a_1\}. \quad (18)$$

Now, if we can show that for any $\{a_2\}$ or $\{a_1\}$ partition, there is always a $\emptyset$ or $\{a_1, a_2, B\}$ partition which achieves a smaller value of $\max\{\Delta A, \Delta B\}$, we will have proven the following.

Proposition 2: For disjoint and connected boundary regions A and B ,

$$E_Q(A : B) = \frac{1}{2} \inf_{|\psi\rangle_{AaBb}} [\max\{\Delta A, \Delta B\}]. \quad (19)$$

FIG. 2. Bulk surface configurations for $S(Ba)$ in each of the four possible cases.

FIG. 3. Bulk surface configurations for f^Q in each of the four possible cases. Green curves are added, and red curves are subtracted. In the order shown, we refer to these quantities as ΔA , ΔB , Δa_2 , Δa_1 .

Proof.—Denote the points partitioning the two components of Γ_{AB} into $\{a, b\}$ by p and q . We need to show that for any $\{a_1\}$ or $\{a_2\}$ partition $\{p, q\}$ we can find a $\{a_1, a_2, B\}$ or $\emptyset$ partition $\{p^*, q^*\}$ for which

$$\begin{aligned} f^Q(\{p^*, q^*\}) &\equiv \frac{1}{2} \max\{\Delta A(\{p^*, q^*\}), \Delta B(\{p^*, q^*\})\} \\ &\leq \frac{1}{2} \max\{\Delta A(\{p, q\}), \Delta B(\{p, q\})\}. \end{aligned} \quad (20)$$

Here we will restrict to the case of symmetric A and B . As always, we use a uniform cutoff, since all relevant quantities are cutoff independent.²

Without loss of generality, let $\{p, q\}$ be a $\{a_2\}$ partition for symmetric boundary regions A and B . Let p_{EW} and q_{EW} be the points on the upper and lower surfaces where $EWCS(AB)$ is anchored (see Fig. 4). Invoking the

² f^Q satisfies the balancing condition in all cases, and is therefore cutoff independent. The difference of any two quantities depicted in Fig. 2 satisfies the balancing condition and is cutoff independent, and therefore the minimizing partition type ($\{a_1, a_2, B\}$, $\emptyset$, $\{a_2\}$, or $\{a_1\}$) is also cutoff independent.

symmetry of A and B and the cutoff, it must be the case that p is on the B side of p_{EW} (right, in the figures) and q is on the A side (left). If we had p on the left and q on the right, then it is clear that $\Delta a_1 > \Delta a_2$, so $\{p, q\}$ cannot be a $\{a_2\}$ partition. If both p and q are on the right, then looking at Fig. 2, it is clear that the $\emptyset$ configuration gives a smaller value for $S(Ba)$ than the $\{a_2\}$ configuration (again by symmetry, and also the fact that a diameter is the longest possible geodesic given a uniform cutoff), so again $\{p, q\}$ cannot be a $\{a_2\}$ partition. Similarly, with both on the left, the $\{a_1, a_2, B\}$ configuration gives a smaller value of $S(Ba)$ than the $\{a_2\}$ configuration.

Without loss of generality, assume that $d(p, p_{EW}) \geq d(q, q_{EW})$. Then it is clear from symmetry that $\Delta B \geq \Delta A$, since $|b_2| \leq |a_1|$ and $|b_1| \leq |a_2|$. Now, consider what happens to ΔA and ΔB as we move p towards A . Regardless of the initial partition, this will always increase ΔA and decrease ΔB , as Fig. 5 shows.

If we continue moving point p towards A we will reach a point p_q where $\Delta A = \Delta B$ (this is move 1 in Fig. 4). By symmetry, $d(p_q, p_{EW}) = d(q, q_{EW})$. Since move 1 has decreased ΔB , $\max\{\Delta A, \Delta B\}$ has also decreased. If the

the $EWCS(AB)$ partition clearly has the property that $\Delta A = \Delta B$, and has the minimal $|M|$ by definition, we conclude that this partition achieves E_Q . ■

Thus the minimizations leading to E_W , E_Q and E_R all involve the same purification and the same partition into $\{a, b\}$. This now allows us to geometrically verify the inequality (10). To see that $E_Q(A:B) \leq E_R(A:B)$, note that

$$E_Q(A:B) - E_R(A:B) = \frac{1}{2}(|\Gamma_A| - |a_1| - |a_2| - |M|), \quad (24)$$

and the right side is a negative quantity, again due to the minimality of $|\Gamma_A|$.

We can think of the optimal partition as ensuring that a is as uncorrelated with B as possible. In particular, we can prove that for the optimal partition, $I(a:B) = I(b:A) = 0$.

Corollary 3: For disjoint and connected boundary regions A and B in the optimal partition, $I(a:B) = I(b:A) = 0$.

Proof.—Since the partition is not $\{a_1\}$ or $\{a_2\}$, we know that both $S(Ba)$ and $S(Ab)$ are given by either $|\Gamma_A| + |b_1| + |b_2|$ or $|\Gamma_B| + |a_1| + |a_2|$. But since $\Delta A = \Delta B$, we know that $|\Gamma_A| + |b_1| + |b_2| = |\Gamma_B| + |a_1| + |a_2|$. Observing that $S(B) + S(a)$ is given by $|\Gamma_B| + |a_1| + |a_2|$ and $S(A) + S(b)$ is given by $|\Gamma_A| + |b_1| + |b_2|$, the claim follows. ■

Furthermore, this leads to another way to understand the value of E_Q : it is proportional to the mutual information $I(Aa:B)$ between the Aa system and B , after all correlation between both a and B , and b and A , has been eliminated.

Corollary 4: For disjoint and connected boundary regions A and B , in the optimal partition, $I(Aa:B) = I(Bb:A) = 2E_Q(A:B)$.

Proof.—We simply observe that $I(Aa:B) = \Delta B$ and $I(Bb:A) = \Delta A$ for any partition, since $|M|$ gives both $S(Aa)$ and $S(Bb)$, $S(B)$ and $S(A)$ are given by their RT surfaces Γ_B and Γ_A , and $S(ABa)$ and $S(ABb)$ are given (using Proposition 1) by $|b_1| + |b_2|$ and $|a_1| + |a_2|$. ■

Using this, another interesting presentation of E_Q follows from SSC. Another important feature of SSC is the following condition for unitary equivalence. If two surfaces representing mixed states share the same boundary, with no extremal surface lying between them (e.g., A and Γ_A in Fig. 1) then their dual states differ by a unitary, serving to concentrate the entanglement as we move closer to a geodesic sharing their boundary. This is realized by the entropy formula in SSC, since all such surfaces are closed by the same extremal curve and therefore have the same entropy. Now, interpreting M as a system which purifies Aa or Bb , we have $\Delta A = I(A:M)$ and $\Delta B = I(B:M)$, so

$$E_Q(A:B) = \frac{1}{2} \inf_M [\max\{I(A:M), I(B:M)\}]. \quad (25)$$

Since ρ_M differs from both ρ_{Aa} and ρ_{Bb} by a unitary, we have $I(A:M) = I(A:Bb)$ and $I(B:M) = I(B:Aa)$, which again gives the claim of Corollary 4. Thus we may think

about E_Q as a minimal mutual information between a surface region A and a cross section of A 's entanglement wedge with B .

III. RELATIONSHIP BETWEEN Q -CORRELATION AND SYMMETRIC SIDE-CHANNEL-ASSISTED DISTILLABLE ENTANGLEMENT

The symmetric side-channel-assisted distillable entanglement of a state [17], denoted $I^{ss}(A)B$, is a measure of bipartite quantum correlation defined by

$$I^{ss}(A)B = \frac{1}{2} \sup_{U:A \rightarrow ab} [-S(a|Bb) + S(a|Eb)] \quad (26)$$

where E purifies AB . Here $U:A \rightarrow ab$ represents an isometry from $\mathcal{H}_A$ to a Hilbert space $\mathcal{H}_a \otimes \mathcal{H}_b$. $I^{ss}(A)B$ can be interpreted operationally as the amount of the entanglement Alice and Bob can distill from their shared state if they have a symmetric channel from Alice to Bob. Symmetric side channels are stronger than classical communications, and thus $I^{ss}(A)B$ is an upper bound on the one-way distillable entanglement, $D^{\rightarrow}(\rho_{AB})$. $I^{ss}(A)B$ is much better behaved than one-way distillable entanglement: it is convex and additive, while one-way distillable entanglement is neither. It has also proven a useful tool for finding upper bounds on $D^{\rightarrow}(\rho_{AB})$ [18].

A. Entanglement with Eve vs Q -correlation with Bob

We find the following relationship between the correlation measures E_Q and I^{ss} , independent of holography.

Proposition 3: Given a state on A and B , and an environment E which purifies the state,

$$S(A) = E_Q(A:B) + I^{ss}(E)A. \quad (27)$$

Proof.—

$$\begin{aligned} I^{ss}(E)A &= \frac{1}{2} \sup_{E \rightarrow ea} [-S(e|Aa) + S(e|Ba)] \\ &= \frac{1}{2} \sup_{E \rightarrow ea} [-S(eAa) + S(Aa) + S(eBa) - S(Ba)] \\ &= \frac{1}{2} \sup_{E \rightarrow ea} [-S(B) + S(Aa) + S(A) - S(Ba)] \\ &= S(A) + \frac{1}{2} \sup_{E \rightarrow ea} [-S(B) + S(Aa) - S(A) - S(Ba)] \\ &= S(A) + -\frac{1}{2} \inf_{E \rightarrow ea} [S(B) - S(Aa) + S(A) + S(Ba)] \\ &= S(A) - E_Q(A:B). \end{aligned}$$

Notice that since $S(A)$ is fixed, Eq. (27) implies that the partition which achieves $E_Q(A:B)$ also achieves $I^{ss}(E)A$. ■

This equation tells us that the entropy of Alice's system can be decomposed into two contributions: Q -correlation with Bob, plus entanglement with Eve that can be distilled via a symmetric channel from Eve to Alice. A similar relationship relating E_P and the dense coding advantage was noted in Ref. [19]. Equation (27) is also reminiscent of the decomposition of Ref. [20],

$$S(A) = E_f(A:B) + I^-(A:E), \quad (28)$$

which embodies the monogamy of entanglement. In this equation $E_f(A:B)$ is the entanglement of formation of ρ_{AB} and $I^-(A:E)$ is the Holevo information of A with respect to E , maximized over positive-operator-valued measurements on E . Equation (27) further clarifies the relationship between A 's distillable entanglement with E and A 's correlation with B .

B. Entanglement, Q -correlation, and the entanglement wedge

As a corollary of Proposition 3, we have the following expression for $I^{ss}(A:B)$ for a pure holographic state on ABE .

Corollary 5: If ρ_{ABE} is a pure holographic state with A and B disjoint and connected, and ρ_M is a state dual to $EWCS(AB)$ via SSC (where ab is the purification of AB determined by M , as in Fig. 1), then

$$I^{ss}(E)A = \frac{1}{2}I(A:a). \quad (29)$$

Proof.—

$$\begin{aligned} I^{ss}(E)A &= S(A) - \frac{1}{2}I(A:M) \\ &= \frac{1}{2}[S(A) + S(A|M)] \\ &= \frac{1}{2}[S(A) - S(A|a)] \\ &= \frac{1}{2}I(A:a). \end{aligned}$$

■

This can also be seen by simply constructing $I(A:a)$ out of bulk surfaces, and noting that $2S(A) - \Delta A = I(A:a)$. If $I^{ss}(E)A = \frac{1}{2}I(A:a)$ for some $U:E \rightarrow ab$, then $\rho_{aB} = \rho_a \otimes \rho_B$. To see this, note that for an optimal⁴ state $|\psi\rangle_{AaBb}$, we have

$$I^{ss}(E)A = \frac{1}{2}[-S(a|Ab) + S(a|Bb)] = \frac{1}{2}I(A:a),$$

which is equivalent [because $S(a|Bb) = -S(a|A)$] to

$$-S(a|Ab) - S(a|A) = S(a) - S(a|A).$$

⁴By optimal we mean a partition which achieves both $I^{ss}(E)A$ and $E_Q(A:B)$, which exists by Proposition 3.

In turn, this is equivalent to

$$0 = S(a) + S(a|Ab) = S(a) - S(a|B) = I(a:B).$$

Zero mutual information is exactly the condition for $\rho_{aB} = \rho_a \otimes \rho_B$. This is nicely consistent with Corollary 3. It is also interesting to note that Eq. (29) gives us a geometric dual for I^{ss} . Since $E_Q(A:B)$ and $I^{ss}(E)A$ are achieved by the same purification $|\psi\rangle_{AaBb}$, we can simply read off the RT surfaces for $I^{ss}(E)A$ from the right side of Eq. (29). We find that $I^{ss}(E)A$ is given in bulk surfaces by $\frac{1}{2}(|\Gamma_A| + |a_1| + |a_2| - |M|)$. Note that this is a cutoff-dependent quantity since it is not balanced at ∂A . The cutoff dependence of $I^{ss}(E)A$ is also evident from Eq. (27), since the left side is cutoff dependent and E_Q is cutoff independent.

IV. DISCUSSION

In this paper we have used SSC to identify the holographic duals of the optimized correlation measures E_Q and E_R introduced in Ref. [3]. We then used these geometric duals to show several properties of E_Q and E_R , including inequalities which are known to hold for these correlation measures independent of geometry. We have also shown a tradeoff relationship between $E_Q(A:B)$ and the symmetric side-channel-assisted distillable entanglement $I^{ss}(E)A$, where E purifies AB , which holds for general quantum states. This relationship allowed us to construct a holographic dual for I^{ss} as well. We have also argued that for a holographic state, the same purification and partition achieves $E_W(A:B)$, $E_Q(A:B)$, $E_R(A:B)$, and $I^{ss}(E)A$. For this we used the cutoff independence of E_Q , and the existence of a boundary conformal map (bulk isometry) taking any disjoint and connected boundary regions A and B to a pair of symmetric boundary regions.

We believe that the tradeoff between E_Q and I^{ss} embodied in Eq. (27) will find a use beyond the current setting. It tells us that the symmetric side-channel distillable entanglement from E to A consists of exactly the correlations that are not captured as Q -correlations with B . This may lead us to a deeper understanding of distillability in general and I^{ss} in particular. It may also point towards an operational understanding of E_Q . We also note that I^{ss} can be interpreted as the utility of a state when used as a quantum one-time pad [21,22], pointing towards the interpretation of $E_Q(A:B)$ as information that is “public” from the perspective of E .

While the Ryu-Takayanagi formula for holographic entanglement entropy is by now well established, the identification of surfaces in the bulk with states having a particular entropy is much less well tested. The coherence of our results, including the fact that the inequalities satisfied by E_Q and E_R independent of geometry can also be shown to be satisfied geometrically by their duals obtained via the SSC prescription, provides further

evidence for this correspondence. Furthermore, it suggests a broader context for the holographic realization of information measures involving the optimization over a purification. It would be interesting to explore this further.

In this work we have focused on the simplest holographic spacetime of pure AdS space, dual to the field theory vacuum; this was necessary for example to reduce all boundary configurations to the simple class of symmetric configurations. It would be interesting to consider the holographic realization of these correlation measures in more general backgrounds, such as black hole geometries. It would also be of interest to generalize these findings to higher-dimensional cases, since while the arguments presented here generalize to topologically simple examples, in general higher-dimensional boundary regions can have more varied topologies. Another interesting avenue to explore is a reformulation of our findings in the language of bit threads, introduced in Ref. [23] and discussed in the context of E_W in Refs. [24–27]. The bit threads formalism may provide an intuition which connects the “flow” of information from one boundary region to another to our

findings about one-way distillable entanglement in holography. In addition, the holographic duals of multipartite generalizations of E_P were considered in Ref. [28], and it would be of interest to similarly broaden our line of inquiry to multipartite systems.

ACKNOWLEDGMENTS

We are grateful to Daniel Hackett, Ethan Neil, Michael Perlin, Markus Pflaum, and Michael Walter for interesting discussions and helpful advice. We also thank Koji Umemoto for making us aware of his work prior to publication. J.L. and G.S. were partially supported by NSF CAREER Grant No. CCF 1652560. O.D. is supported by the Department of Energy under Grant No. DE-SC0010005. This work was supported by the Department of Energy under Grant No. DE-SC0020386.

Note added.—In the later stages of this work, we learned of an independent effort by Koji Umemoto which reaches partially overlapping conclusions [16].

-
- [1] M. Christandl and A. Winter, Squashed entanglement: An additive entanglement measure, *J. Math. Phys. (N.Y.)* **45**, 829 (2004).
 - [2] B. M. Terhal, M. Horodecki, D. W. Leung, and D. P. DiVincenzo, The entanglement of purification, *J. Math. Phys. (N.Y.)* **43**, 4286 (2002).
 - [3] J. Levin and G. Smith, Optimized measures of bipartite quantum correlation, [arXiv:1906.10150](#).
 - [4] S. Ryu and T. Takayanagi, Holographic Derivation of Entanglement Entropy from the Anti-de Sitter Space/Conformal Field Theory Correspondence, *Phys. Rev. Lett.* **96**, 181602 (2006).
 - [5] K. Umemoto and T. Takayanagi, Entanglement of purification through holographic duality, *Nat. Phys.* **14**, 573 (2018).
 - [6] P. Nguyen, T. Devakul, M. G. Halbasch, M. P. Zaletel, and B. Swingle, Entanglement of purification: From spin chains to holography, *J. High Energy Phys.* **01** (2018) 098.
 - [7] M. Miyaji and T. Takayanagi, Surface/state correspondence as a generalized holography, *Prog. Theor. Exp. Phys.* **2015**, 073B03 (2015).
 - [8] B. Swingle, Entanglement renormalization and holography, *Phys. Rev. D* **86**, 065007 (2012).
 - [9] F. Pastawski, B. Yoshida, D. Harlow, and J. Preskill, Holographic quantum error-correcting codes: toy models for the bulk/boundary correspondence, *J. High Energy Phys.* **06** (2015) 149.
 - [10] S. Dutta and T. Faulkner, A canonical purification for the entanglement wedge cross-section, [arXiv:1905.00577](#).
 - [11] J. Kudler-Flam and S. Ryu, Entanglement negativity and minimal entanglement wedge cross sections in holographic theories, *Phys. Rev. D* **99**, 106014 (2019).
 - [12] K. Tamaoka, Entanglement Wedge Cross Section from the Dual Density Matrix, *Phys. Rev. Lett.* **122**, 141601 (2019).
 - [13] G. Vidal and R. F. Werner, Computable measure of entanglement, *Phys. Rev. A* **65**, 032314 (2002).
 - [14] P. Hayden, M. Headrick, and A. Maloney, Holographic mutual information is monogamous, *Phys. Rev. D* **87**, 046003 (2013).
 - [15] J. Chen and A. Winter, Non-additivity of the entanglement of purification (beyond reasonable doubt), [arXiv:1206.1307](#).
 - [16] K. Umemoto, Quantum and classical correlations inside the entanglement wedge, *Phys. Rev. D* **100**, 126021 (2019).
 - [17] G. Smith, J. Smolin, and A. Winter, The quantum capacity with symmetric side channels, *IEEE Trans. Inf. Theory* **54**, 4208 (2008).
 - [18] G. Smith and J. A. Smolin, Additive extensions of a quantum channel, in *2008 IEEE Information Theory Workshop* (IEEE, New York, 2008), pp. 368–372.
 - [19] M. Horodecki and M. Piani, On quantum advantage in dense coding, *J. Phys. A* **45**, 105306 (2012).
 - [20] M. Koashi and A. Winter, Monogamy of quantum entanglement and other correlations, *Phys. Rev. A* **69**, 022309 (2004).
 - [21] F. G. Brandao and J. Oppenheim, Quantum One-Time Pad in the Presence of an Eavesdropper, *Phys. Rev. Lett.* **108**, 040504 (2012).
 - [22] F. G. Brandao and J. Oppenheim, Public quantum communication and superactivation, *IEEE Trans. Inf. Theory* **59**, 2517 (2012).

- [23] M. Freedman and M. Headrick, Bit threads and holographic entanglement, [Commun. Math. Phys.](#) **352**, 407 (2017).
- [24] N. Bao, A. Chatwin-Davies, J. Pollack, and G. N. Remmen, Towards a bit threads derivation of holographic entanglement of purification, [J. High Energy Phys.](#) **07** (2019) 152.
- [25] D.-H. Du, C.-B. Chen, and F.-W. Shu, Bit threads and holographic entanglement of purification, [J. High Energy Phys.](#) **08** (2019) 140.
- [26] C. A. Agón, J. de Boer, and J. F. Pedraza, Geometric aspects of holographic bit threads, [J. High Energy Phys.](#) **05** (2019) 075.
- [27] J. Harper and M. Headrick, Bit threads and holographic entanglement of purification, [J. High Energy Phys.](#) **08** (2019) 101.
- [28] K. Umemoto and Y. Zhou, Entanglement of purification for multipartite states and its holographic dual, [J. High Energy Phys.](#) **10** (2018) 152.