**Erratum: Sensitivity limits on heavy-light mixing $|U_{\mu N}|^2$ from
lepton number violating B meson decays
[Phys. Rev. D **96**, 035025 (2017)]**

Gorazd Cvetič and C. S. Kim

(Received 10 July 2020; published 22 July 2020)

DOI: [10.1103/PhysRevD.102.019903](https://doi.org/10.1103/PhysRevD.102.019903)

There are typos in Eqs. (10) and (11) which refer to the decay $B \rightarrow D\ell_1 N$. In both equations, the right-hand sides should be multiplied by factor 4, i.e., the equations with the correct factors are

$$|\mathcal{T}|^2 = 4|U_{\ell_1 N}|^2|V_{cb}|^2 G_F^2 |\tilde{\mathcal{T}}|^2, \quad (10)$$

$$\frac{d\Gamma(B^- \rightarrow D^0 \ell_1^- N)}{dq^2 d\Omega_{\hat{q}'} \Omega_{\hat{p}_1}} = \frac{|U_{\ell_1 N}|^2 |V_{cb}|^2 G_F^2}{M_B (4\pi)^5} |\tilde{\mathcal{T}}|^2 \lambda^{1/2} \left(1, \frac{M_D^2}{M_B^2}, \frac{q^2}{M_B^2}\right) \lambda^{1/2} \left(1, \frac{M_1^2}{q^2}, \frac{M_N^2}{q^2}\right). \quad (11)$$

In our calculations, these correct formulas were used. The explicit expression for $|\tilde{\mathcal{T}}|^2$ is given in Eq. (12) there. Integration of the above differential decay width over $\Omega_{\hat{p}_1}$ and $\Omega_{\hat{q}'}$ reproduces the explicit expression for $d\Gamma(B^- \rightarrow D^0 \ell_1^- N)/dq^2$ as given in Eq. (B6) of our earlier paper [Phys. Rev. D **94**, 053001 (2016)].