Vacuum stability conditions for Higgs potentials with $SU(2)_L$ triplets

Gilbert Moultağa^{1,*} and Michel C. Peyranère²

¹*Laboratoire Charles Coulomb (L2C), University of Montpellier,
CNRS, Montpellier, Place Eugène Bataillon CC069, France*

²*LUPM, University of Montpellier, CNRS, F-34095 Montpellier Cedex 5, France*

(Received 29 December 2020; accepted 2 March 2021; published 10 June 2021)

Tree-level dynamical stability of scalar field potentials in renormalizable theories can in principle be expressed in terms of positivity conditions on quartic polynomial structures. However, these conditions cannot always be cast in a fully analytical resolved form, involving only the couplings and being valid for all field directions. In this paper we consider such forms in three physically motivated models involving $SU(2)$ triplet scalar fields: the Type-II seesaw model, the Georgi-Machacek model, and a generalized two-triplet model. A detailed analysis of the latter model allows one to establish the full set of necessary and sufficient boundedness-from-below conditions. These can serve as a guide, together with unitarity and vacuum structure constraints, for consistent phenomenological (tree-level) studies. They also provide a seed for improved loop-level conditions and encompass in particular the leading ones for the more specific Georgi-Machacek case. Incidentally, we present complete proofs of various properties and also derive general positivity conditions on quartic polynomials that are equivalent to but much simpler than the ones used in the literature.

DOI: [10.1103/PhysRevD.103.115006](https://doi.org/10.1103/PhysRevD.103.115006)

I. INTRODUCTION

Since the experimental discovery of a Standard Model (SM)-like Higgs particle at the LHC [1,2] and the lack so far of any direct evidence for physics beyond the standard model (BSM),¹ one might ask whether the properties of the discovered 125 GeV scalar particle so close to the SM predictions (see, e.g., [4]) leave any room for BSM physics to reside below the tera-electron-volt or at the nearby few tera-electron-volts scale. If new physics is present in the electroweak symmetry breaking sector, it should be either very heavy (almost decoupled) or light but have very weak mixing with the SM Higgs. For the latter case, extensions of the scalar sector of the SM by complex or real $SU(2)_L$ triplets, or further extensions comprising left-right symmetric gauge groups, or possibly higher representation multiplets, are appealing possibilities. A typical example is the Type-II seesaw model for neutrino masses [5–10], for which an essentially SM-like physical Higgs state is unavoidable, a consequence of the very small mixing

between the doublet and triplet neutral components being set off by the tiny (Majorana) neutrino mass scale as compared to the electroweak scale. Another example is the Georgi-Machacek model [11,12] with one complex and one real triplet such that a tree-level custodial symmetry is preserved in the scalar sector through a global $SU(2)_R$.

These scenarios have triggered various activities both on the phenomenological level (including left-right symmetric or not, supersymmetric or not, scenarios; see, e.g., among the recent works [13–25] and references therein) and in experimental searches at the LHC for neutral, charged, and in particular doubly charged scalar states that are specific to such a class of models decaying either to same-sign leptons or to W boson pairs [26–29]. As for any extension of the SM, and in the absence of a unifying ultraviolet completion, these models have an increased number of free parameters and thus a large freedom in particular for the physical spectrum of the scalar sector. Theoretical conditions such as the stability of the potential, a consistent electroweak vacuum, and unitarity bounds are thus welcome as a guide together with the experimental exclusion limits to narrow future search strategies.

The present paper focuses on the potential stability issue for three models: the Type-II seesaw model, the Georgi-Machacek model, and a generalized two-triplet model. The aim is to address as thoroughly as possible the theoretical determination of necessary and sufficient (NAS) conditions on the scalar couplings that ensure a physically sound bounded from below (BFB) potential. The NAS BFB

*Corresponding author.

gilbert.moultaka@umontpellier.fr

¹Possible indirect “evidence” notwithstanding [3].

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article’s title, journal citation, and DOI. Funded by SCOAP³.

conditions have already been considered in the corresponding literature. Inspired by the approach of [30] initially proposed for the general two-Higgs doublet potential, the strategy consists in a change of parametrization of the field space reducing it to a minimal set of variables corresponding to positive-valued ratios of field magnitudes and to field orientations varying in compact domains. It is then found that in contrast with the general two-Higgs doublet case, the general doublet-triplet potential leads to a simplification that allows a fully analytical solution. A complete answer was given first in [31,32] for the Type-II seesaw model. Following the same approach the NAS BFB conditions were provided for the Georgi-Machacek model in [33]. We will nevertheless reexamine the issue for these two models, supplementing with complete proofs, for reasons that will become clear in the course of the study. Encouraged by the success of the approach, we extend it in the present paper to a generalized two-triplet model, which we will dub precustodial, for which we provide novel results by deriving the full NAS BFB conditions. Some stability constraints have already been given for this model in [34,35] corresponding, however, to specific directions in the field space, and thus to a subclass of necessary conditions. This precustodial model can be of phenomenological interest by itself, but can also serve as a guideline for the effective potential beyond tree level in the Georgi-Machacek model.

The main issue of the analysis will be to cast the conditions in a form as close as possible to a fully *resolved* one. By “fully resolved” we mean *an analytical expression that depends solely on the couplings with no reference to orientations or magnitudes in field space*. A fully resolved form, when possible, is an ideal result both technically, since no scan over the field configurations is needed, and physically, as consistency constraints are expressed directly in terms of the (physical) couplings. This was the case for the conditions derived in [31,32] while in [33] the conditions were resolved with respect to only one parameter, thus remaining in a partially unresolved form albeit with a residual field dependence reduced to a compact domain. As we will see, similar configurations arise in the precustodial model where the resolving occurs at different stages with respect to different parameters. A hindrance in the way of reaching fully resolved conditions emerges whenever dealing with a quartic polynomial that cannot be reduced to a biquadratic one. This fact motivated us to investigate further a rather mathematical question, the positivity of general quartic polynomials, for which we determine NAS conditions that are simpler than the ones found in the literature.

A word of caution is in order here: The NAS BFB conditions we are considering are obtained by requiring the tree-level potential not be unbounded from below in any direction in the field space. It is only in that sense that they are necessary and sufficient. Obviously they might be necessary only in a wider physical sense when taking into account the structure of the vacua. Moreover, going beyond

tree level would modify these conditions. As alluded to above and will be briefly discussed toward the end of the paper, the tree-level conditions can, however, encapsulate in some cases the leading loop corrections.

Several methods to treat the stability of the potential have been conceived in the literature, e.g., specifically for multi-Higgs-doublets models [36–38] including elegant geometric approaches [39], or more general methods relying on copositivity [40–42] or on other powerful mathematical techniques [43] (and references therein). As attractive as it may seem, the ability of the latter systematic methods to treat in principle any model through ready-to-use packages [44] can yet in practice run into technical difficulties when dealing with extended scalar sectors as noted in [43]. Also to the best of our knowledge a model with one triplet has been treated using copositivity [41] but for which only specific directions in field space were considered in agreement with [31], while the authors of [45] obtained with this method the all-directions conditions for the Type-II seesaw model in a form different from that of [31]. It should, however, be stressed that the copositivity method cannot always be applied to potentials with extended Higgs multiplets when all four-dimensional operators allowed by the gauge symmetries and renormalizability are considered. This was the case for the doublet extensions studied in [36–38] and will be the case here for the extensions with two Higgs triplets.² Thus, the more pedestrian and somewhat mathematically lowbrow approach we adopt in this paper remains in our opinion an efficient way of tackling the stability problem specifically for the models under consideration with two triplets.

The paper is organized as follows. In Sec. II we revisit the derivation of the NAS BFB conditions for the Type-II seesaw model finding equivalence with the conditions of [32] that corrected [31], but we stress that the conditions of [31] do remain valid, *necessary, and sufficient* when one of the couplings is negative. Adding one real $SU(2)$ triplet, the approach is extended to the general precustodial model in Sec. III, including the Georgi-Machacek model as a special case. This section contains the bulk of the new results. We recall some useful ingredients of the two model potentials in Secs. III A and III B. In Sec. III C we first identify six field dependent variables that provide a reduced parametrization of the field space suitable for the BFB study, four of which, dubbed α parameters, vary in compact domains. We then investigate the NAS BFB conditions following a procedure where the resolving with respect to these six field-dependent variables is performed step-by-step. Section III D deals with the analytical determination of the domains of variation of the α parameters as

²More precisely, the potential cannot always be cast in a bilinear form involving positive-definite independent vector components *and* an optimal space dimension to make the method advantageous; more on this at the end of Sec. V. Models with increased symmetries can be more tractable; see, e.g., [46,47].

well as all two-, three-, and four-dimensional analytical correlations between them. In Sec. III E we derive the main results identifying the fully and partially resolved branches of the NAS BFB conditions. The special case of the Georgi-Machacek model is reconsidered in Sec. III F where we relate the reduced parameters to those of the precustodial model and provide a proof of their domain of variation that was only conjectured in the literature. Section IV illustrates an unexpected feedback of the Georgi-Machacek model on the precustodial one. A wrap-up with further illustrations, comments, and a user's guide, is given in Sec. V, and we conclude in Sec. VI. Further material and detailed proofs, missing either in the literature for known properties or for the new results found in this paper, are given in Appendixes A–F. Special attention is paid, in Appendixes G and H, to the mathematical issue of deriving simple forms for the NAS positivity conditions of quartic polynomials.

II. THE TYPE-II SEESAW DOUBLET-TRIPLET HIGGS POTENTIAL

We first sketch the main ingredients, relying on the detailed analysis and notations of [31] to which the reader may refer for more details.

The potential reads

$$\begin{aligned} V(H, \Delta) = & -m_H^2 H^\dagger H + \frac{\lambda}{4} (H^\dagger H)^2 + M_\Delta^2 \text{Tr}(\Delta \Delta^\dagger) \\ & + [\mu (H^T i \sigma^2 \Delta^\dagger H) + \text{H.c.}] + \lambda_1 (H^\dagger H) \text{Tr}(\Delta \Delta^\dagger) \\ & + \lambda_2 (\text{Tr} \Delta \Delta^\dagger)^2 + \lambda_3 \text{Tr}(\Delta \Delta^\dagger)^2 + \lambda_4 H^\dagger \Delta \Delta^\dagger H. \end{aligned} \quad (2.1)$$

H denotes the standard scalar field $SU(2)_L$ doublet and Δ a colorless $SU(2)_L$ complex triplet scalar field, with charge assignments $H \sim (1, 2, 1)$ and $\Delta \sim (1, 3, 2)$ under $SU(3)_c \times SU(2)_L \times U(1)_Y$,

$$H = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix}, \Delta = \begin{pmatrix} \delta^{++} & \delta^{++} \\ \delta^0 & -\delta^+/\sqrt{2} \end{pmatrix}. \quad (2.2)$$

We have used the 2×2 traceless matrix representation for the triplet, wrote the two multiplets in terms of their complex valued scalar components, and indicated a choice of electric charges with the conventional electric charge assignment for the doublet and following $Q = I_3 + \frac{Y_\Delta}{2}$ with $I_3 = -1, 0, 1$ and $Y_\Delta = 2$ for the triplet. σ^2 denotes the second Pauli matrix. The potential $V(H, \Delta)$ is invariant under $SU(2)_L \times U(1)_Y$ field transformations $H \rightarrow e^{i\alpha} \mathcal{U}_L H$ and $\Delta \rightarrow e^{i2\alpha} \mathcal{U}_L \Delta \mathcal{U}_L^\dagger$ where $\mathcal{U}_L$ denotes an arbitrary element of $SU(2)_L$ in the fundamental representation. Since we are only interested in the issue of boundedness from below of the potential, we need not go further here into the details of the dynamics of spontaneous electroweak

symmetry breaking, the structure of the physical Higgs states and the generation of Majorana neutrino masses.

A. The BFB conditions

In order to cope generically with the shape of $V(H, \Delta)$ along all possible directions of the ten-dimensional field space, we adopt a reduced parametrization for the fields that will turn out to be particularly convenient to entirely solve the problem analytically. Following [31] we define

$$r \equiv \sqrt{H^\dagger H + \text{Tr} \Delta \Delta^\dagger}, \quad (2.3)$$

$$H^\dagger H \equiv r^2 \cos^2 \gamma, \quad (2.4)$$

$$\text{Tr} \Delta \Delta^\dagger \equiv r^2 \sin^2 \gamma, \quad (2.5)$$

$$\text{Tr}(\Delta \Delta^\dagger)^2 / (\text{Tr} \Delta \Delta^\dagger)^2 \equiv \zeta, \quad (2.6)$$

$$(H^\dagger \Delta \Delta^\dagger H) / (H^\dagger H \text{Tr} \Delta \Delta^\dagger) \equiv \xi. \quad (2.7)$$

Obviously, when H and Δ scan all the field space, the radius r scans the domain $[0, +\infty)$ and the angle $\gamma \in [0, \frac{\pi}{2}]$. With this parametrization it is straightforward to cast the quartic part of the potential, denoted hereafter by $V^{(4)}$ and given by the second line of Eq. (2.1), in the following simple form:

$$\begin{aligned} V^{(4)}(r, \tan \gamma, \xi, \zeta) = & \frac{r^4}{4(1 + \tan^2 \gamma)^2} (\lambda + 4(\lambda_1 + \xi \lambda_4) \tan^2 \gamma \\ & + 4(\lambda_2 + \zeta \lambda_3) \tan^4 \gamma). \end{aligned} \quad (2.8)$$

We stress here that the crux of the matter is the existence of a parametrization, Eqs. (2.3)–(2.7), which allows one to scan *all* the field space and in the same time recast the relevant part of the potential into a *biquadratic form* in $\tan \gamma$. It is the concomitance of these two facts that allows a tractable and complete analytical solution for the *necessary and sufficient* boundedness-from-below conditions. Indeed, the absence of linear and/or cubic powers of $\tan \gamma$ in Eq. (2.8) is anything but generic. (For instance, in a similar parametrization initially proposed in [30] to study two-Higgs-doublet models such terms do remain, hindering an easy fully analytical treatment.)

One can thus consider only the range $0 \leq \tan \gamma < +\infty$ in accordance with the above stated range for γ . Boundedness from below is then equivalent to requiring $V^{(4)} > 0$ for *all* $\tan \gamma \in [0, +\infty)$ and *all* ξ, ζ in their allowed domain. The γ -free necessary and sufficient conditions on the λ_i 's have already been given in [31]³:

³We use the conventional notations $\wedge$, $\vee$, and $\neg$ for the Boolean operators AND, OR, and NOT, respectively.

$$\lambda > 0 \wedge \lambda_2 + \zeta\lambda_3 \geq 0 \wedge \lambda_1 + \xi\lambda_4 + \sqrt{\lambda(\lambda_2 + \zeta\lambda_3)} > 0. \quad (2.9)$$

Note that the second inequality above is nonstrict. This accounts rigorously for the only possible equality among the NAS conditions that is compatible with requiring $V^{(4)}$ to be strictly positive.⁴ These inequalities are a subset of the general NAS positivity conditions for a quartic polynomial (see Appendix G). We stress here that Eq. (2.9) answers fully the question of (tree-level) boundedness from below in the totality of the ten-dimensional field space. There remains, however, the dependence on ξ and ζ that parametrize the relative magnitudes of the dimension four gauge invariant operators in Eq. (2.1) that are not controlled solely by r and γ .

One can, however, show that

$$0 \leq \xi \leq 1 \quad \text{and} \quad \frac{1}{2} \leq \zeta \leq 1. \quad (2.10)$$

(See Appendixes A 1 and A 2 for a proof.)

In [31] the authors relied on this allowed range and on the monotonic dependence on (ξ, ζ) in Eq. (2.9) to obtain Eqs. (4.21), (4.22), and (4.23) of [31] reproduced in Appendix B 2 for later discussions. The authors of [32] rightly observed that [31] had actually overlooked the fact that (ξ, ζ) being correlated cannot reach an arbitrary point in the rectangle defined by Eq. (2.10). Starting from Eq. (2.9) and using the constraint

$$2\xi^2 - 2\xi + 1 \leq \zeta \leq 1, \quad (2.11)$$

they showed that the set of conditions Eqs. (B14)–(B16) established in [31], although sufficient in all field space directions, are, in fact, not necessary, even though deviation from absolute necessity is typically at the few percent level. Although we totally agree with their general observation, we will see that despite the correlation between ξ and ζ the conditions Eqs. (B14)–(B16) *do remain sufficient and necessary whenever $\lambda_3 < 0$* ; the modification will come only for $\lambda_3 > 0$. We will come back to this point in more detail later on in Appendix B.

For now, we just add that, as shown in Appendix A 3, it is possible to cast the ξ and ζ parameters as follows:

$$\xi = \frac{1}{2}(1 + c_{2H}c_{2\Delta}), \quad (2.12)$$

$$\zeta = \frac{1}{2}(1 + c_{2\Delta}^2), \quad (2.13)$$

⁴In Sec. III C we will elaborate further on the meaning of the condition $V^{(4)} > 0$, as well as on the fact that the parameter $\tan \gamma$ varies independently of ζ and ξ .

with c_{2H} and $c_{2\Delta}$ two *independent* cosines taking any value in their allowed domain $[-1, 1]$; note also that Eq. (2.11) comes as a direct consequence of these equations.

Although the authors of [32] wrote a correct form of the necessary and sufficient BFB conditions, they only sketched a proof of their result. In Appendix B, we provide a detailed proof through a careful study of Eq. (2.9) leading to an alternative form of the fully resolved NAS BFB conditions. The latter reduce to

$$\mathcal{B}_0 \wedge \{\mathcal{B}_1 \vee \mathcal{B}_2\}, \quad (2.14)$$

where

$$\mathcal{B}_0 \Leftrightarrow \left\{ \lambda > 0 \wedge \lambda_2 + \lambda_3 \geq 0 \wedge \lambda_2 + \frac{\lambda_3}{2} \geq 0 \right\}, \quad (2.15)$$

$$\begin{aligned} \mathcal{B}_1 \Leftrightarrow & \left\{ \lambda_1 + \sqrt{\lambda(\lambda_2 + \lambda_3)} > 0 \wedge \lambda_1 + \lambda_4 \right. \\ & \left. + \sqrt{\lambda(\lambda_2 + \lambda_3)} > 0 \wedge \sqrt{\lambda\lambda_3} \leq \sqrt{(\lambda_2 + \lambda_3)\lambda_4^2} \right\}, \end{aligned} \quad (2.16)$$

and

$$\begin{aligned} \mathcal{B}_2 \Leftrightarrow & \left\{ \sqrt{\lambda\lambda_3} \geq \sqrt{(\lambda_2 + \lambda_3)\lambda_4^2} \wedge \lambda_1 + \frac{\lambda_4}{2} \right. \\ & \left. + \sqrt{\lambda\left(\lambda_2 + \frac{\lambda_3}{2}\right)\left(1 - \frac{\lambda_4^2}{2\lambda\lambda_3}\right)} > 0 \right\}. \end{aligned} \quad (2.17)$$

Note also that Eq. (2.17) implies $\lambda_3 > 0$ and $2\lambda\lambda_3 - \lambda_4^2 > 0$ so that the $\mathcal{B}_2$ part is relevant only when these conditions are satisfied simultaneously.

The above constraints are, in fact, totally equivalent to [32] although they have a slightly different form. Indeed, the equivalence is not straightforward as the two involved Boolean forms are in general not equivalent to each other. However, they become equivalent due to the implication given by Eq. (B13). The above constraints are as follows:

- (i) constitute an independent check of the results of [32].
- (ii) are written explicitly as a union of domains one of which, $\mathcal{B}_1$, is a necessary consequence of constraints Eqs. (B15) and (B16).
- (iii) allow one to understand why in some regimes the previous constraints Eqs. (B15) and (B16) would exclude only a very small part of the allowed parameter space. This is the case in particular in the regimes where $\lambda_4 \ll 1$ or $\lambda_4^2 \ll 2\lambda\lambda_3$.
- (iv) allow one to see analytically that our previous constraints Eqs. (B15) and (B16) were sufficient but not necessary. Indeed, Eq. (B14) is the same as Eq. (2.15) while one can easily check that Eqs. (B15) and (B16) always imply Eq. (2.16).

III. GENERALIZATION ADDING ONE EXTRA REAL TRIPLET

Such a generalization can be of phenomenological interest by itself, but is also motivated by the structure of the Georgi-Machacek model beyond the tree level [48].

A. The precustodial potential

Defining

$$A = \begin{pmatrix} a^+/\sqrt{2} & -a^{++} \\ a^0 & -a^+/\sqrt{2} \end{pmatrix}, \quad B = \begin{pmatrix} b^0/\sqrt{2} & -b^+ \\ -b^{+*} & -b^0/\sqrt{2} \end{pmatrix}, \quad (3.1)$$

with A a different notation for the complex triplet Δ , and H as defined in Eq. (2.2), $B = B^\dagger$ a real triplet (b^0 real valued), we write the most general renormalizable precustodial potential involving H , A , and B as follows:

$$V_{\text{p-c}} = V_{\text{p-c}}^{(2,3)} + V_{\text{p-c}}^{(4)}, \quad (3.2)$$

where the dimension-two and dimension-three operators are collected in

$$\begin{aligned} V_{\text{p-c}}^{(2,3)} = & -m_H^2 H^\dagger H + M_A^2 \text{Tr}(AA^\dagger) + M_B^2 \text{Tr}(B^2) \\ & + [\mu_A (H^T i\sigma^2 A^\dagger H) + \text{H.c.}] + \mu_B H^\dagger B H \\ & + \mu_{AB} \text{Tr}(AA^\dagger B), \end{aligned} \quad (3.3)$$

and the dimension-four operators in

$$\begin{aligned} V_{\text{p-c}}^{(4)} = & \frac{\lambda_H}{4} (H^\dagger H)^2 + \frac{\lambda_A^{(1)}}{4} (\text{Tr} AA^\dagger)^2 + \frac{\lambda_A^{(2)}}{4} \text{Tr}(AA^\dagger)^2 \\ & + \frac{\lambda_B}{4!} (\text{Tr} B^2)^2 + \lambda_{AH}^{(1)} H^\dagger H \text{Tr} AA^\dagger \\ & + \lambda_{AH}^{(2)} H^\dagger AA^\dagger H + \frac{\lambda_{BH}}{2} H^\dagger H \text{Tr} B^2 \\ & + \frac{\lambda_{AB}^{(1)}}{2} \text{Tr} AA^\dagger \text{Tr} B^2 + \frac{\lambda_{AB}^{(2)}}{2} \text{Tr} AB \text{Tr} A^\dagger B \\ & + \frac{i}{2} \lambda_{ABH} (H^\dagger \sigma^2 A^\dagger B H - H^\dagger B A \sigma^2 H^*). \end{aligned} \quad (3.4)$$

$V_{\text{p-c}}$ is invariant under $SU(2)_L \times U(1)_Y$ field transformations

$$\begin{aligned} H & \rightarrow e^{i\alpha} \mathcal{U}_L H, \\ A & \rightarrow e^{i2\alpha} \mathcal{U}_L A \mathcal{U}_L^\dagger, \\ B & \rightarrow \mathcal{U}_L B \mathcal{U}_L^\dagger, \end{aligned} \quad (3.5)$$

where $\mathcal{U}_L$ denotes an arbitrary element of $SU(2)_L$ in the fundamental representation. This potential was written

in [48] and later on in [34] with which we agree up to different normalizations and notations.⁵ All other dimension-three and dimension-four gauge invariant operators either are vanishing or can be expressed in terms of the ones listed above. (For completeness we give a proof of this in Appendix C.)

B. The Georgi-Machacek potential

This model [11,12], a special setup of the model presented in the previous subsection, allows one to extend the validity of the SM tree-level (approximate) custodial symmetry in the presence of $SU(2)_L$ triplet scalar fields. In particular, the potential reads

$$V_{\text{G-M}} = V_{\text{G-M}}^{(2,3)} + V_{\text{G-M}}^{(4)}, \quad (3.6)$$

$$\begin{aligned} V_{\text{G-M}}^{(2,3)} = & \frac{\mu_2^2}{2} \text{Tr} \Phi^\dagger \Phi + \frac{\mu_3^2}{2} \text{Tr} X^\dagger X \\ & - (M_1 \text{Tr} \Phi^\dagger \tau^a \Phi \tau^a + M_2 \text{Tr} X^\dagger t^a X t^a) (UXU^\dagger)_{ab}, \end{aligned} \quad (3.7)$$

$$\begin{aligned} V_{\text{G-M}}^{(4)} = & \hat{\lambda}_1 (\text{Tr} \Phi^\dagger \Phi)^2 + \hat{\lambda}_2 \text{Tr}(\Phi^\dagger \Phi) \text{Tr}(X^\dagger X) \\ & + \hat{\lambda}_3 \text{Tr}(X^\dagger X X^\dagger X) + \hat{\lambda}_4 (\text{Tr} X^\dagger X)^2 \\ & - \hat{\lambda}_5 \text{Tr}(\Phi^\dagger \tau^a \Phi \tau^a) \text{Tr}(X^\dagger t^a X t^a), \end{aligned} \quad (3.8)$$

where we followed the notations of [33].⁶ We hat the λ 's to distinguish them from those of Sec. II, and we define the scalar bidoublet and bitriple as

$$\begin{aligned} \Phi & \equiv \begin{pmatrix} \phi^{0*} & \phi^+ \\ -\phi^{+*} & \phi^0 \end{pmatrix} = (i\sigma^2 H^*, \quad H), \quad (3.9) \\ X & \equiv \begin{pmatrix} \chi^{0*} & \xi^+ & \chi^{++} \\ -\chi^{+*} & \xi^0 & \chi^+ \\ \chi^{++*} & -\xi^{+*} & \chi^0 \end{pmatrix} = \sqrt{2} \begin{pmatrix} a^{0*} & b^+ & a^{++} \\ -a^{+*} & b^0 & a^+ \\ a^{++*} & -b^{+*} & a^0 \end{pmatrix}, \end{aligned} \quad (3.10)$$

so that the normalization of the vacuum expectation values are the same as in [33]. Note also the sign difference in a^{++}

⁵With the field correspondence as given by Eq. (3.10) and couplings correspondence: $\lambda_H = 4\lambda$, $\lambda_A^{(i=1,2)} = 16\rho_i$, $\lambda_B = 4! \times 4\rho_3$, $\lambda_{AB}^{(i=1,2)} = 8\rho_{i+3}$, $\lambda_{AH}^{(i=1,2)} = 2\sigma_i$, $\lambda_{BH} = 4\sigma_3$, and $\lambda_{ABH} = 4\sigma_4$. Note that our normalization factors for the various couplings are chosen such that they cancel out for at least one vertex originating from each operator when symmetry factors are taken into account in the Feynman rules.

⁶In Eqs. (3.7) and (3.8) $\tau^a = \sigma^a/2$ with σ^a the Pauli matrices are the usual $SU(2)$ generators in the fundamental representation, t^a the generators in the triplet (adjoint) representation, with $a = 1, 2, 3$, and U some rotation matrix about which we skip here the details (see [33,49]) as Eq. (3.7) will not be relevant to our study.

and b^+ between Eq. (3.1) and Eq. (3.10). The potential V_{p-c} is then mapped onto V_{G-M} through the following correspondence among the couplings:

$$\begin{aligned}\hat{\lambda}_1 &= \frac{1}{16}\lambda_H, & \hat{\lambda}_2 &= \frac{1}{8}\lambda_{BH}, & \hat{\lambda}_3 &= -\frac{1}{64}\lambda_A^{(2)}, \\ \hat{\lambda}_4 &= \frac{1}{32}\lambda_{AB}^{(1)}, & \hat{\lambda}_5 &= -\frac{1}{4\sqrt{2}}\lambda_{ABH},\end{aligned}\quad (3.11)$$

provided, however, the following correlations hold for the precustodial potential couplings:

$$\begin{aligned}\lambda_A^{(1)} &= 2\lambda_{AB}^{(1)} + 3\lambda_{AB}^{(2)}, & \lambda_A^{(2)} &= -2\lambda_{AB}^{(2)}, & \lambda_{ABH} &= \sqrt{2}\lambda_{AH}^{(2)}, \\ \lambda_B &= 3(\lambda_{AB}^{(1)} + \lambda_{AB}^{(2)}), & \lambda_{BH} &= \lambda_{AH}^{(1)} + \frac{1}{2}\lambda_{AH}^{(2)}.\end{aligned}\quad (3.12)$$

The potential V_{G-M} enjoys an increased symmetry as compared to that of V_{p-c} , Eq. (3.5), with an invariance under an extra global $SU(2)$,

$$\Phi \rightarrow \mathcal{U}_L^{(2)} \Phi \mathcal{U}_R^{(2)}, \quad (3.13)$$

$$X \rightarrow \mathcal{U}_L^{(3)} X \mathcal{U}_R^{(3)}, \quad (3.14)$$

where $\mathcal{U}_{L,R}^{(n)}$ denotes n -dimensional representation of $SU(2)_{L,R}$. The correlations given by Eq. (3.12) can thus be viewed as encoding the tree-level constraints imposed by the $SU(2)_R$ global symmetry on the potential. We come back to this point in Sec. V when discussing briefly quantum effects. References [34,48] considered such correlations.⁷

C. The precustodial BFB conditions

Being a polynomial in the fields, the tree-level potential has no singularities at finite values of the fields; it follows that boundedness from below means that the potential does not become infinitely negative at infinitely large field values. This is equivalent to requiring *strict* positivity of the quartic part of the potential, Eq. (3.4), for *all field values in all field directions*. The latter requirement is *sufficient* as it implies that at infinitely large field values, where $|V_{p-c}^{(2,3)}| \ll |V_{p-c}^{(4)}|$ in Eq. (3.2), the potential does not become infinitely negative. That it is also *necessary* might not seem obvious since the last term in Eq. (3.4) is linear in A and in B , so that $V_{p-c}^{(4)}$ might be negative for some finite values of the fields without being unbounded from below. That this does not happen, and the above requirement is indeed necessary, can easily be seen as follows: If there

existed a point in field space where $V_{p-c}^{(4)} \equiv V_{p-c}^{(4)*} \leq 0$, then scaling all the fields at that point by the same real-valued amount s would have led to $V_{p-c}^{(4)} \equiv s^4 V_{p-c}^{(4)*} \leq 0$, implying unboundedness from below since s can be chosen infinitely large. Note finally that *strict* positivity is important here because a vanishing $V_{p-c}^{(4)}$ at very large field values would generically lead to the dominance of $V_{p-c}^{(3)}$ which, barring accidental cancellations in some field directions, always possesses unbounded from below directions.

The BFB conditions are thus the necessary and sufficient conditions on the ten couplings λ of Eq. (3.4) that ensure

$$V_{p-c}^{(4)} > 0, \quad \forall A, B, H. \quad (3.15)$$

Of course, loop corrections will modify the conditions on the couplings resulting from Eq. (3.15), although the effects can be partly encoded in the runnings of the couplings through a renormalization group improvement of the potential. (We will come back briefly to this point in Sec. V.) Note also that the above definition of boundedness from below does not take into account the actual pattern of spontaneous symmetry breaking that would typically lead to more stringent constraints.

The condition in Eq. (3.15) should be verified in the full 13-dimensional space of the real-valued field components of the A , B , and H multiplets. However, symmetries of the model (and possibly accidental symmetries akin to $V_{p-c}^{(4)}$) will help reduce the number of relevant degrees of freedom. Starting from Eq. (3.4) we generalize the parametrization of Eqs. (2.3)–(2.7) using spherical-like coordinates as follows:

$$H^\dagger H \equiv r^2 \cos^2 a, \quad (3.16)$$

$$\text{Tr} A A^\dagger \equiv r^2 \sin^2 a \cos^2 b, \quad (3.17)$$

$$\text{Tr}(B^2) \equiv r^2 \sin^2 a \cos^2 b, \quad (3.18)$$

$$r^2 = H^\dagger H + \text{Tr} A A^\dagger + \text{Tr}(B^2), \quad (3.19)$$

where r is a non-negative number and $a \in [-\pi/2, +\pi/2]$ and $b \in [-\pi, +\pi]$ two angles. It will also prove useful to define the following real-valued quantities,

$$T \equiv \sqrt{\frac{\text{Tr} A A^\dagger}{H^\dagger H}} = |\tan a \cos b|, \quad t \equiv \sqrt{\frac{\text{Tr}(B^2)}{\text{Tr} A A^\dagger}} = |\tan b|, \quad (3.20)$$

$$\begin{aligned}\alpha_A &\equiv \frac{\text{Tr} A A^\dagger A A^\dagger}{(\text{Tr} A A^\dagger)^2}, & \alpha_{AH} &\equiv \frac{H^\dagger A A^\dagger H}{H^\dagger H \text{Tr} A A^\dagger}, \\ \alpha_{AB} &\equiv \frac{\text{Tr} A B \text{Tr} A^\dagger B}{\text{Tr} A A^\dagger \text{Tr}(B^2)},\end{aligned}\quad (3.21)$$

⁷We agree with [34] except for a factor of 2 difference on the right-hand side of the first equation of the second line of Eq. (3.12) as compared to the first equation of the second line of Eq. (10) of [34].

$$\alpha_{ABH} \equiv i \frac{H^\top \sigma^2 A^\dagger B H - H^\dagger B A \sigma^2 H^*}{H^\dagger H \sqrt{\text{Tr} A A^\dagger} \text{Tr}(B^2)}. \quad (3.22)$$

Hereafter we will refer to the latter four parameters as the α parameters. In terms of T , t , and the α parameters, the quartic part of the potential now reads

$$V_{\text{p-c}}^{(4)} = r^4 \cos^4 a \times (\mathbf{a}_0 + \mathbf{b}_0 T^2 + \mathbf{c}_0 T^4), \quad (3.23)$$

where

$$\begin{aligned} \mathbf{a}_0 &= \frac{\lambda_H}{4}, \mathbf{b}_0 = \lambda_{AH}^{(1)} + \alpha_{AH} \lambda_{AH}^{(2)} + \frac{1}{2} (\alpha_{ABH} \lambda_{ABH} t + \lambda_{BH} t^2), \\ \mathbf{c}_0 &= \frac{1}{4} (\lambda_A^{(1)} + \alpha_A \lambda_A^{(2)}) + \frac{1}{2} (\lambda_{AB}^{(1)} + \alpha_{AB} \lambda_{AB}^{(2)}) t^2 + \frac{1}{24} \lambda_B t^4. \end{aligned} \quad (3.24)$$

It becomes evident from Eqs. (3.23) and (3.24) that the positivity of $V_{\text{p-c}}^{(4)}$ does not depend explicitly on all ten terms of the right-hand side of Eq. (3.4), but just on the reduced set of the six combinations of gauge invariant operators defined in Eqs. (3.20)–(3.22). The sought-after NAS BFB conditions on the λ 's are thus those that ensure

$$\mathbf{a}_0 + \mathbf{b}_0 T^2 + \mathbf{c}_0 T^4 > 0, \quad \forall T, t, \alpha_A, \alpha_{AH}, \alpha_{AB}, \alpha_{ABH}. \quad (3.25)$$

It is important to note that scanning independently over all values of the 13 real-valued components of the fields A , B , and H amounts to varying T , t , and the α parameters. The latter, however, do not all vary independently. For one thing, the α parameters vary in bounded domains: α_A and α_{AH} are nothing but, respectively, ζ and ξ defined in Eqs. (2.6) and (2.7). Hence

$$\alpha_A \in \left[\frac{1}{2}, 1 \right], \quad (3.26)$$

$$\alpha_{AH} \in [0, 1], \quad (3.27)$$

as shown in Appendix A. Furthermore, one can show that

$$\alpha_{AB} \in [0, 1], \quad (3.28)$$

$$\alpha_{ABH} \in [-\sqrt{2}, +\sqrt{2}]; \quad (3.29)$$

see Appendix D for details. For another, the α parameters are uncorrelated only locally. But similar to what was pointed out in [32] and discussed at length in Sec. II A for the Type-II seesaw model potential, they are correlated globally in that they cannot reach the boundaries of their respective domains independently of each other. The actual domain in the four-dimensional α parameters space is

certainly not the simple hypercube defined by Eqs. (3.26)–(3.29). One can approach the true domain by considering the projected domains on the subspaces of these parameters taken two by two. This is not trivial to establish and will be carried out in full details in Sec. III D. The more difficult task of determining fully the true domain will be discussed in Sec. III D 7.

In contrast, the variables T and t vary in $\in [0, +\infty)$ independently of each other and of the α parameters. In essence, the α parameters being ratios of different gauge invariant combinations of the fields can be seen as functions of cosines and sines of angles defined separately in the A , B , and H field spaces, where $\sqrt{\text{Tr} A A^\dagger}$, $\sqrt{\text{Tr} B^2}$, and $\sqrt{H^\dagger H}$ represent lengths. This hints at the obstruction to span the full hypercube as noted above. Whereas T and t , being two ratios of these three lengths, are clearly independent of each other and of the α parameters. It follows that T can be varied independently from $\mathbf{a}_0$, $\mathbf{b}_0$, and $\mathbf{c}_0$ in Eq. (3.25). Consequently, the NAS conditions for the strict positivity of $V_{\text{p-c}}^{(4)}$, $\forall T$, are those of a biquadratic polynomial in T , namely conditions on the λ 's satisfying

$$\begin{aligned} \mathbf{a}_0 > 0 \wedge \mathbf{c}_0 \geq 0 \wedge \mathbf{b}_0 + 2\sqrt{\mathbf{a}_0 \mathbf{c}_0} > 0, \\ \forall t, \alpha_A, \alpha_{AB}, \alpha_{AH}, \alpha_{ABH}. \end{aligned} \quad (3.30)$$

As noted previously, only the highest degree monomial coefficient can vanish. However, for the sake of simplicity we will consider in the sequel only the strict inequality $\mathbf{c}_0 > 0$. It is convenient to recast the above inequalities in the following equivalent form that disposes of the (less tractable) square root:

$$\begin{aligned} \mathbf{a}_0 > 0 \wedge \mathbf{c}_0 > 0 \\ \wedge \{ \mathbf{b}_0 > 0 \vee \{ \mathbf{b}_0 < 0 \wedge 4\mathbf{a}_0 \mathbf{c}_0 - \mathbf{b}_0^2 > 0 \} \}, \\ \forall t, \alpha_A, \alpha_{AB}, \alpha_{AH}, \alpha_{ABH}, \end{aligned} \quad (3.31)$$

which simplifies further to

$$\mathbf{a}_0 > 0 \wedge \mathbf{c}_0 > 0 \quad (3.32)$$

$\wedge$

$$\{ \mathbf{b}_0 > 0 \vee 4\mathbf{a}_0 \mathbf{c}_0 - \mathbf{b}_0^2 > 0 \}, \quad \forall t, \alpha_A, \alpha_{AB}, \alpha_{AH}, \alpha_{ABH}. \quad (3.33)$$

I. $\mathbf{a}_0 > 0 \wedge \mathbf{c}_0 > 0$

We consider first the conditions in Eq. (3.32) as they are common to the union of the two conditions of Eq. (3.33). The coefficient $\mathbf{c}_0$ being itself biquadratic in t and the latter independent of the α parameters [see Eq. (3.24)], the corresponding NAS positivity condition is in turn of the

same form as Eqs. (3.32) and (3.33). The two inequalities in Eq. (3.32) are thus equivalent to

$$\lambda_H > 0 \wedge \lambda_B > 0 \wedge \lambda_A^{(1)} + \alpha_A \lambda_A^{(2)} > 0 \quad (3.34)$$

\(\wedge\)

$$\begin{aligned} & \{\lambda_{AB}^{(1)} + \alpha_{AB} \lambda_{AB}^{(2)} > 0 \text{ (I)} \vee (\lambda_A^{(1)} + \alpha_A \lambda_A^{(2)}) \lambda_B \\ & - 6(\lambda_{AB}^{(1)} + \alpha_{AB} \lambda_{AB}^{(2)})^2 > 0 \text{ (II)}\}, \\ & \forall \alpha_A, \alpha_{AB}. \end{aligned} \quad (3.35)$$

Note that the second inequality in Eq. (3.34) and the first inequality in Eq. (3.35) depend solely on α_A or on α_{AB} . They can easily be resolved since the dependence on these parameters is monotonic; if required to be valid $\forall \alpha_A, \alpha_{AB}$ in the domains given by Eqs. (3.26) and (3.28), they become equivalent to requiring them simultaneously at the two edges of these domains, namely,

$$\lambda_A^{(1)} + \frac{\lambda_A^{(2)}}{2} > 0 \wedge \lambda_A^{(1)} + \lambda_A^{(2)} > 0, \quad (3.36)$$

for the first, and

$$\lambda_{AB}^{(1)} > 0 \wedge \lambda_{AB}^{(1)} + \lambda_{AB}^{(2)} > 0, \quad (3.37)$$

for the second. Equation (3.35)-(II) needs more care due to the nontrivial global correlation between α_A and α_{AB} (see next section and Fig. 2), and will be kept in its present form for the time being. One will also have to tackle a further complication involving the two inequalities of Eq. (3.35). Indeed, due to the “or” structure of Eq. (3.35), none of the two corresponding inequalities need to be necessarily valid for all α_A, α_{AB} in their domains; it suffices that one of the two inequalities be satisfied in a given subset of α_A, α_{AB} , and the other inequality satisfied in the complementary subset. In particular, Eq. (3.37) is only sufficient. To reach the NAS conditions one will have to consider all possible coverings of the domain by two subsets for which such a configuration holds. This issue will be solved explicitly in Sec. III E 1.

2. $\mathfrak{b}_0 > 0 \vee 4\mathfrak{a}_0\mathfrak{c}_0 - \mathfrak{b}_0^2 > 0$

We turn now to the two inequalities of Eq. (3.33). The first is quadratic in t [see Eq. (3.24)], but could in principle be treated as a biquadratic polynomial in $\sqrt{t}$, since $t \in [0, +\infty)$. The second, $4\mathfrak{a}_0\mathfrak{c}_0 - \mathfrak{b}_0^2 > 0$, is a general quartic polynomial in this same variable. This is the first place where we encounter the issue of positivity conditions for a general quartic polynomial. Relying on a classic theorem about single variable polynomials that are positive on $(-\infty, +\infty)$, we derive in Appendix G a relatively tractable form of the corresponding NAS conditions for

a quartic polynomial. However, these conditions are not directly applicable to the case at hand since the relevant variable here, t , is in $[0, +\infty)$. In this case the NAS conditions would obviously be less restrictive; see, for instance, [50,51] for recent reviews.⁸ Relying on these theorems we extend the results of Appendix G to the domain $[0, +\infty)$ in Appendix H.

However, this is not the full story. Similar to what we stated above in Sec. III C 1 regarding Eq. (3.35), the structure of Eq. (3.33) implies that it is sufficient for the two inequalities $\mathfrak{b}_0 > 0$ and $4\mathfrak{a}_0\mathfrak{c}_0 - \mathfrak{b}_0^2 > 0$ to be separately valid in two complementary subsets of the allowed t and α parameters domains. The NAS conditions will then be obtained by investigating all possible coverings of these domains for which this happens. The upshot is that the possibility of varying freely t with respect to the α parameters is not sufficient anymore. Indeed, a given subset of the α parameters where, for instance, $\mathfrak{b}_0 > 0$ (or $4\mathfrak{a}_0\mathfrak{c}_0 - \mathfrak{b}_0^2 > 0$) will necessarily be correlated with t . A strategy for an explicit resolution will be given in Sec. III E 2.

Although it will prove unavoidable to deal with positivity conditions of quartic polynomials on subdomains of $(-\infty, +\infty)$, it will still be useful for the subsequent discussions to replace from the onset $t \in [0, +\infty)$ by a variable on $(-\infty, +\infty)$ if possible. This is indeed the case if one considers the variable Z defined as

$$Z = \alpha_{ABH} \times t \quad (3.38)$$

since α_{ABH} can take either sign [cf. Eq. (3.29)]. However, in order to apply safely the NAS positivity conditions on a polynomial in Z , one should make sure that Z is not correlated with the other parameters, α_A, α_{AH} , and α_{AB} appearing in the inequalities, even though these parameters are globally correlated with α_{ABH} .

It is obviously the case for $|Z|$ since t is uncorrelated with the other parameters and allows one to scan independently of the value of $|\alpha_{ABH}|$ the full $[0, +\infty)$ range. However, the sign of Z is controlled by α_{ABH} which is globally correlated with α_A, α_{AH} , and α_{AB} . It is thus crucial to check that the sign of α_{ABH} is not correlated with the latter parameters. That this is indeed the case is easily seen by recalling that all the inequalities are required to be valid $\forall A, B, H$ in the field space, and noting that α_A, α_{AH} , and α_{AB} remain unchanged, while α_{ABH} flips sign, at the two field space points A and $-A$ (or equivalently at B and $-B$, or H and iH); see Eqs. (3.21) and (3.22). It follows that one can change freely the sign of α_{ABH} for any given configuration of α_A, α_{AH} , and α_{AB} . (As we will see in the next subsection, Figs. 3–5, this translates into domains symmetrical around

⁸Somewhat surprisingly, corresponding theorems, when the variable does not span the full $(-\infty, +\infty)$ interval, seem not to have been referenced in the mathematics literature before the 1970s (see [52]).

$\alpha_{ABH} = 0$.) The variable $Z \in (-\infty, +\infty)$ is thus genuinely uncorrelated with the other field dependent reduced parameters.

D. Global correlations among the α parameters

In this section we first determine the allowed domains of the α parameters taken two by two, and then combine the resulting six global correlations to obtain an analytical approximation of the full four-dimensional (4D) domain. Since the α parameters are ratios of gauge invariant quantities [cf. Eqs. (3.21) and (3.22)], it is convenient to choose a gauge that reduces the dependence on the set of components fields of the A , B , and H multiplets. Apart from the treatment of α_A versus α_{AH} , we carry all the discussion in this section assuming a gauge that diagonalizes the (Hermitian and traceless) B multiplet as defined in Eq. (3.1), which then takes the form

$$B = \begin{pmatrix} b & 0 \\ 0 & -b \end{pmatrix}. \quad (3.39)$$

It follows that the dependence on b cancels out in α_{AB} and, up to a global sign, in α_{ABH} .

1. α_A versus α_{AH}

These parameters are identical, respectively, to ζ and ξ that were defined and studied in detail in Sec. II A and Appendix A 3. We just recall here the corresponding domain:

$$(i): 0 \leq \alpha_{AH} \leq 1, \quad (3.40)$$

$$(ii): 2\alpha_{AH}(\alpha_{AH} - 1) + 1 \leq \alpha_A \leq 1, \quad (3.41)$$

illustrated in Fig. 1.

2. α_A versus α_{AB}

With no particular gauge choice but using the fact that the parameter α_A is a ratio, one can recast it in terms of reduced parameters in the following form:

$$\alpha_A = \frac{1}{4}(2\cos^4\theta + (3 + \cos 4\varphi)\sin^4\theta + (2 + \cos\rho \sin 2\varphi)\sin^2 2\theta), \quad (3.42)$$

where we defined

$$\begin{aligned} |a^0| &= a \cos\varphi \sin\theta, \\ |a^+| &= a \cos\theta, \\ |a^{++}| &= a \sin\varphi \sin\theta, \\ \rho &= \arg(a^0) - 2\arg(a^+) + \arg(a^{++}), \end{aligned} \quad (3.43)$$

with

FIG. 1. Projection of the α -parameters domain onto the (α_{AH}, α_A) plane.

$$0 \leq \varphi \leq \frac{\pi}{2}, \quad 0 \leq \theta \leq \frac{\pi}{2}, \quad 0 \leq \rho \leq 2\pi, \quad \text{and} \quad a = \sqrt{|a^0|^2 + |a^+|^2 + |a^{++}|^2}. \quad (3.44)$$

Furthermore, choosing a gauge for which Eq. (3.39) is valid, the α_{AB} parameter takes the very simple form,

$$\alpha_{AB} = \cos^2\theta. \quad (3.45)$$

Equations (3.42) and (3.45) lead straightforwardly to

$$\alpha_A = \frac{1}{4}(3 + (2 - 3\alpha_{AB})\alpha_{AB} + (\alpha_{AB} - 1)^2 \cos 4\varphi + (1 - \alpha_{AB})\alpha_{AB} \cos\rho \sin 2\varphi). \quad (3.46)$$

To determine the boundary of the allowed domain one can, for instance, study the variation of α_A in Eq. (3.46) as a quadratic function of $x \equiv \sin 2\varphi$ in the domain $0 \leq x \leq 1$ to identify the set of maximal and minimal possible values of α_A for a given α_{AB} depending on $\cos\rho$. The maximum is reached for $x = \frac{\alpha_{AB}}{1-\alpha_{AB}}\cos\rho$ which lies in the allowed domain only if $\cos\rho \geq 0$ and $\alpha_{AB} \leq \frac{1}{2}$. Otherwise, the maximum is reached at one of the boundary values $x = 0$ or $x = 1$. We find that the boundary of the domain is given by the following four curves:

$$\begin{aligned} (I): & \alpha_{AB} = 0 \quad \text{and} \quad \frac{1}{2} \leq \alpha_A \leq 1, \\ (II): & \alpha_A = \frac{1}{2} \quad \text{and} \quad 0 \leq \alpha_{AB} \leq 1, \\ (III): & \alpha_A = 1 \quad \text{and} \quad 0 \leq \alpha_{AB} \leq \frac{1}{2}, \\ (IV): & \frac{1}{2} \leq \alpha_{AB} \leq 1 \quad \text{and} \quad \alpha_A = \frac{1}{2} + 2(1 - \alpha_{AB})\alpha_{AB}; \end{aligned} \quad (3.47)$$

see also Fig. 2.

FIG. 2. The upper figure blue contour indicates the projection of the α -parameters domain onto the (α_{AB}, α_A) plane. The middle and lower figures are related to Sec. III E 1 to which the reader is referred for more details. The dashed straight lines illustrate arbitrary partitions defined by Eq. (3.35)-(I); the black solid parabolae illustrate arbitrary partitions defined by Eq. (3.35)-(II).

3. α_{AH} versus α_{ABH}

Similar to the preceding case, we recast α_{AH} and α_{ABH} in terms of reduced parameters and in the gauge where Eq. (3.39) holds,

$$\begin{aligned} \alpha_{AH} = & \frac{1}{2}(1 + \cos 2\varphi \cos 2\psi \sin^2 \theta) \\ & + \frac{1}{2\sqrt{2}}(\cos \varphi \cos \theta_3 + \sin \varphi \cos \theta_4) \sin 2\psi \sin 2\theta, \end{aligned} \quad (3.48)$$

$$\begin{aligned} \alpha_{ABH} = & \sqrt{2} \text{sgn}(b)(\sin \varphi \sin^2 \psi \cos \theta_2 \\ & - \cos \varphi \cos^2 \psi \cos \theta_1) \sin \theta, \end{aligned} \quad (3.49)$$

where θ and φ are as previously defined and

$$\begin{aligned} \theta_1 &= \arg(a^0) - 2\arg(\phi^0), \\ \theta_2 &= \arg(a^{++}) - 2\arg(\phi^+), \\ \theta_3 &= \arg(a^0) - \arg(a^+) - \arg(\phi^0) + \arg(\phi^+), \\ \theta_4 &= \arg(a^+) - \arg(a^{++}) - \arg(\phi^0) + \arg(\phi^+), \\ \cos \psi &= \frac{|\phi^0|}{\sqrt{|\phi^0|^2 + |\phi^+|^2}}, \end{aligned} \quad (3.50)$$

with $0 \leq \psi \leq \frac{\pi}{2}$ and $0 \leq \theta_i \leq 2\pi$. [Note that $\theta_1 = \theta_2 + \theta_3 + \theta_4$ (modulo 2π).]

FIG. 3. Projection of the α -parameters domain onto the $(\alpha_{ABH}, \alpha_{AH})$ plane.

A numerical parametric scan over the various angles allows one to guess the boundary of the α_{AH} versus α_{ABH} domain. The result turns out to be very simple given by the two curves:

$$(V): \alpha_{AH} = 1, \quad \forall \alpha_{ABH} \in [-\sqrt{2}, +\sqrt{2}], \quad (3.51)$$

$$(VI): \alpha_{AH} = \frac{1}{2} \alpha_{ABH}^2, \quad (3.52)$$

illustrated in Fig. 3. The proof for the upper boundary (3.51) is simple: It suffices to exhibit particular configurations of the various angles for which α_{AH} saturates its upper bound while α_{ABH} scans all its allowed domain. An example is $\varphi = \psi = \theta = \frac{\pi}{2}$, keeping all the θ_i 's free. This gives $\alpha_{AH} = 1$ and $\alpha_{ABH} = \sqrt{2} \cos \theta_2$, which proves the above statement. The lower boundary (3.52) is much more difficult to establish analytically. The proof is somewhat involved and will be relegated to Appendix D 3.

4. α_A versus α_{ABH}

Here again a numerical parametric scan over the various angles helps guess the boundary of the α_A versus α_{ABH} domain. However, one still needs for that to admit *ad hoc* that the whole boundary is obtained when $\sin \theta = 1$. The analytical proof is quite involved and is given in Appendix D 4 for completeness. We find that the boundary is determined by the following:

FIG. 4. Projection of the α -parameters domain onto the (α_{ABH}, α_A) plane.

$$(VII): \alpha_A = 1, \quad \text{for } \alpha_{ABH} \in [-\sqrt{2}, +\sqrt{2}],$$

$$(VIII): \alpha_A = \frac{1}{2}, \quad \text{for } \alpha_{ABH} \in [-1, +1],$$

$$(IX): \alpha_A = 1 - \alpha_{ABH}^2 + \frac{1}{2} \alpha_{ABH}^4,$$

$$\text{for } \alpha_{ABH} \in [-\sqrt{2}, -1] \cup [+1, +\sqrt{2}]. \quad (3.53)$$

5. α_{AB} versus α_{ABH}

From Eqs. (3.45), (3.49), (3.44), and (3.29), one obtains readily

$$\alpha_{ABH}^2 = 2Y^2(1 - \alpha_{AB}), \quad (3.54)$$

where Y [defined in Eq. (D35)] and α_{AB} vary independently in the domain $[0, 1]$. It is then clear that for each given value of α_{ABH} , α_{AB} reaches its maximal value compatible with Eq. (3.54) when $Y^2 = 1$. Also the minimal value $\alpha_{AB} = 0$ is reached for any value of α_{ABH}^2 . The boundary of the allowed domain in the plane α_{AB} versus α_{ABH} is thus delimited by the two curves:

$$(X): \alpha_{AB} = 0, \quad \forall \alpha_{ABH} \in [-\sqrt{2}, +\sqrt{2}], \quad (3.55)$$

$$(XI): \alpha_{AB} = 1 - \frac{1}{2} \alpha_{ABH}^2, \quad (3.56)$$

as shown on Fig. 5.

6. α_{AH} versus α_{AB}

The boundary of the allowed domain in the $(\alpha_{AB}, \alpha_{AH})$ plane is given by

$$(XII): \alpha_{AB} = 0, \quad \forall \alpha_{AH} \in [0, 1], \quad (3.57)$$

$$(XIII): \alpha_{AH} = 0, \quad \forall \alpha_{AB} \in \left[0, \frac{1}{2}\right], \quad (3.58)$$

$$(XIV): \alpha_{AH} = 1, \quad \forall \alpha_{AB} \in \left[0, \frac{1}{2}\right], \quad (3.59)$$

FIG. 5. Projection of the α -parameters domain onto the $(\alpha_{ABH}, \alpha_{AB})$ plane.

FIG. 6. Projection of the α -parameters domain onto the $(\alpha_{AB}, \alpha_{AH})$ plane.

$$(XV): \left(\alpha_{AB} - \frac{1}{2}\right)^2 + \left(\alpha_{AH} - \frac{1}{2}\right)^2 = \frac{1}{4},$$

$$\text{for } \alpha_{AB} \in \left[\frac{1}{2}, 1\right] \text{ and } \alpha_{AH} \in [0, 1]; \quad (3.60)$$

$$\begin{aligned} & \alpha_A \leq 1 \wedge \alpha_{AB} \geq 0 \wedge \alpha_{AB} \leq 1 - \frac{1}{2}\alpha_{ABH}^2 \\ & \wedge \\ & \alpha_{AH} \geq \frac{1}{2}\alpha_{ABH}^2 \wedge \alpha_A \geq 1 + 2(\alpha_{AH} - 1)\alpha_{AH} \\ & \wedge \\ & \left\{ \alpha_{AB} \leq \frac{1}{2} \vee \left\{ \alpha_A \leq \frac{1}{2} + 2(1 - \alpha_{AB})\alpha_{AB} \wedge \left(\alpha_{AB} - \frac{1}{2}\right)^2 + \left(\alpha_{AH} - \frac{1}{2}\right)^2 \leq \frac{1}{4} \right\} \right\}. \end{aligned} \quad (3.61)$$

This form is nontrivial in that it does not display explicitly all six correlations among the four α parameters; in particular, the correlation between α_A and α_{ABH} does not appear explicitly and, depending on α_{AB} , either only three or five of the six correlations are explicitly needed. These features will prove useful when resolving the constraints in Sec. III E 2. It is also informative to partially visualize the 4D potatoid by considering its 3D projections along each of the four directions. This amounts to combining the domains three by three which leads after some simplifications to

$$\{\alpha_A, \alpha_{AB}, \alpha_{AH}\} = \begin{cases} \frac{1}{2} \leq \alpha_A \leq 1 \wedge 0 \leq \alpha_{AB} \leq \frac{1}{2} + \sqrt{\frac{1-\alpha_A}{2}} \\ \wedge \\ \frac{1}{2}(1 - \sqrt{2\alpha_A - 1}) \leq \alpha_{AH} \leq \frac{1}{2}(1 + \sqrt{2\alpha_A - 1}), \end{cases} \quad (3.62)$$

see Fig. 6. The proof strategy is similar to the one in Sec. III D 2 albeit somewhat more involved, the convenient variable here to study the variation of α_{AH} being $x \equiv \cos 2\psi$. (See Appendix D 5 for details.)

7. The 4D α potatoid

The 2D projections of the α -parameters domain determined analytically in the previous subsections will allow, in some cases, a fully analytical resolving of the BFB conditions on the λ 's. Obviously generalizing beyond 2D along the same lines becomes nontractable analytically. In principle one can then proceed numerically, scanning over part or all of the seven angles entering Eqs. (3.42), (3.45), (3.48), and (3.49), to determine the 3D projections as well as the true 4D allowed domain of the α parameters. However, this would cut short the possibility of further analytical resolving for the conditions on the λ 's.

We will proceed differently here by constructing an analytical approximation of the true α -parameters domain from a back projection using only six planes. Obviously any point in the true domain should have its projections on the six planes lying within the six domains determined above. This necessary condition can be characterized by the interior of a four-dimensional convex domain that we will refer to as the 4D potatoid. To determine explicitly this 4D potatoid we first express separately in the form of a logical (inclusive) disjunction each of the six domains of Figs. 1–6, and then form the logical conjunction of these disjunctions. The resulting Boolean expression is somewhat involved but, interestingly enough, it eventually simplifies to the following form:

FIG. 7. Projection of the α potatoid along: (a) the α_A direction, (b) the α_{AB} direction, (c) the α_{ABH} direction, and (d) the α_{AH} direction.

$$\{\alpha_A, \alpha_{AH}, \alpha_{ABH}\} = 1 + 2(\alpha_{AH} - 1)\alpha_{AH} \leq \alpha_A \leq 1 \wedge \alpha_{ABH}^2 \leq 2\alpha_{AH}, \quad (3.63)$$

$$\{\alpha_A, \alpha_{AB}, \alpha_{ABH}\} = \begin{cases} \frac{1}{2} \leq \alpha_A \leq 1 \wedge 0 \leq \alpha_{AB} \leq 1 - \frac{\alpha_{ABH}^2}{2} \\ \wedge \\ \left\{ \alpha_{AB} \leq \frac{1}{2} \vee \frac{1}{2} - \sqrt{\frac{1-\alpha_A}{2}} \leq \alpha_{AB} \leq \frac{1}{2} + \sqrt{\frac{1-\alpha_A}{2}} \right\} \\ \wedge \\ \left\{ \alpha_{ABH}^2 \leq 1 \vee 1 - \sqrt{2\alpha_A - 1} \leq \alpha_{ABH}^2 \leq 1 + \sqrt{2\alpha_A - 1} \right\}, \end{cases} \quad (3.64)$$

$$\{\alpha_{AB}, \alpha_{AH}, \alpha_{ABH}\} = \begin{cases} 0 \leq \alpha_{AB} \leq 1 - \frac{\alpha_{ABH}^2}{2} \wedge \frac{\alpha_{ABH}^2}{2} \leq \alpha_{AH} \leq 1 \\ \wedge \\ \left\{ \alpha_{AB} \leq \frac{1}{2} \vee (1 - 2\alpha_{AB})^2 + (1 - 2\alpha_{AH})^2 \leq 1 \right\}. \end{cases} \quad (3.65)$$

Figure 7 shows these 3D projections. It is easy to check by eye from this figure that further projection on the various planes reproduces the domains shown in Figs. 1–6.

However, the rounded (and even nonsmooth) edges featured in Fig. 7 hint at the fact that looking at projections is necessary but not sufficient to determine the true 4D

domain of the α parameters. For instance, a point lying just outside the chopped edge in Fig. 7(a), which is a point excluded for sure, would still project on the interior of the domains of Figs. 3, 5, and 6. Obviously this is not yet fully a counterexample as the considered point might still project outside one of the three remaining 2D domains. But on general grounds the potatoid determined by Eq. (3.61), even though enclosing the true 4D domain, is not necessarily identical to it. Since relying on continuity arguments one does not expect holes in the interior of the true domain that would leave no imprint in the projections on the six planes, one concludes that differences between the potatoid and the true domain should be located on the boundaries of the former. We defer a detailed study showing that this is indeed the case till Sec. IV. There we will make use of interesting feedback on the issue from the more constrained Georgi-Machacek model.

E. Resolved forms of the precustodial BFB conditions

For now we ignore the above subtleties and exploit in the present section the domains of the α parameters, as determined so far, to push as much as possible an explicit resolving of the conditions given by Eqs. (3.32) and (3.33) for the λ parameters themselves.

1. Resolving $\alpha_o > 0 \wedge c_o > 0$

Resolving conditions (3.32) with respect to t , they become equivalent to Eqs. (3.34) and (3.35). As stressed at the end of Sec. III C 1, in order to fulfill the or structure of Eq. (3.35) one should in principle consider all possible partitions into subsets of the domain depicted in Fig. 2. Since obviously the two inequalities could be simultaneously satisfied in some parts of the domain, the subsets should be allowed to overlap. So, strictly speaking, we should consider *coverings* rather than *partitions*. More precisely,

A set of values $(\lambda_A^{(1)}, \lambda_A^{(2)}, \lambda_{AB}^{(1)}, \lambda_{AB}^{(2)})$ will satisfy Eq. (3.35) $\forall \alpha_A, \alpha_{AB}$, if and only if there exists a covering of the (α_A, α_{AB}) domain consisting of a family of subsets of this domain for which Eq. (3.35)-(I) is satisfied on a collection of these subsets, and Eq. (3.35)-(II) is satisfied on the complementary collection.

The task can seem daunting since there are *a priori* infinitely many ways of forming a covering of the domain. However, one can identify a clear procedure. Note first the obvious fact that, for a given $(\lambda_A^{(1)}, \lambda_A^{(2)}, \lambda_{AB}^{(1)}, \lambda_{AB}^{(2)})$ in the λ space, each of the two inequalities in Eq. (3.35) defines separately natural *partitions* of the (α_A, α_{AB}) domain, namely partitions formed by a collection made of subsets where the inequality is satisfied and subsets where it is not. Moreover, among all these natural partitions, one can show that a *minimal* partition, made of the smallest possible

number of subsets, is actually unique and made of at most two subsets.⁹ A clear strategy follows: For each given point in the λ space, determine the two minimal partitions defined, respectively, by Eq. (3.35)-(I) and Eq. (3.35)-(II), call them $\{s_{\text{yes}}^I, s_{\text{no}}^I\}$ and $\{s_{\text{yes}}^{II}, s_{\text{no}}^{II}\}$; then check whether their union forms a covering that satisfies the required property stated above in italics; that is, check whether

$$s_{\text{no}}^I \subset s_{\text{yes}}^{II}, \quad \text{or equivalently,} \quad s_{\text{no}}^{II} \subset s_{\text{yes}}^I, \quad (3.66)$$

to select or reject the considered point in λ space.

Given the linear dependence on α_{AB} in Eq. (3.35)-(I), the associated minimal partition corresponds simply to cutting the (α_A, α_{AB}) domain into regions by a straight line going vertically across the domain, at $\alpha_{AB} = \alpha_{AB}^* \equiv -\frac{\lambda_{AB}^{(1)}}{\lambda_{AB}^{(2)}}$, as illustrated by the dashed line in Fig. 2. The inequality (3.35)-(I) is then true for any α_{AB} in an entire interval of the form $[\alpha_{AB}^*, 1]$ or $[0, \alpha_{AB}^*]$ and false on their respective complement. This corresponds, respectively, to the two minimal partitions where the “yes” assignment holds for the right side or the left side region. Moreover, since α_{AB} lives in $[0, 1]$, the minimal partition reduces trivially to either $\{s_{\text{yes}}^I, \emptyset\}$ or $\{\emptyset, s_{\text{no}}^I\}$ if $\alpha_{AB}^* \notin [0, 1]$. It is thus convenient to consider separately the NAS conditions on $\lambda_{AB}^{(1)}$ and $\lambda_{AB}^{(2)}$ that correspond to each of these four configurations of the minimal partition. These NAS conditions are easy to write down given the monotonic dependence on $\alpha_{AB} \in [0, 1]$ in Eq. (3.35)-(I). They are given in Fig. 8 with the labels (i), (ii), (iii), and (iv). Note that (i) and (iv) correspond to the two extreme configurations, respectively, $\{s_{\text{yes}}^I, \emptyset\}$ [cf. Eq. (3.37)] and $\{\emptyset, s_{\text{no}}^I\}$, while (ii) and (iii) are the two intermediate generic partitions.

On the other hand, as can easily be seen from the dependence on α_A, α_{AB} in Eq. (3.35)-(II), the corresponding minimal partitions are determined by convex parabolae in the (α_{AB}, α_A) plane, illustrated by the black curves in Fig. 2. The middle and bottom figures in Fig. 2 show several possible configurations when $\alpha_{AB}^* \in [0, 1]$. The middle-left

⁹This is an immediate consequence of the binary “yes/no” characterization of the subsets of the natural partitions defined above. Indeed, starting from a given natural partition and taking the union of all the “yes” subsets and the union of all the “no” subsets form two subsets (including possibly an empty one) defining a minimal partition. The uniqueness proof then follows easily: if $\{s_1, s_2\}$ and $\{s'_1, s'_2\}$ are two minimal partitions, then at least one set s_i and one set s'_j should have a nonempty intersection $s_i \cap s'_j$, since the partitions cover the same domain. This implies the whole of s_i and s'_j to have the same yes/no characterization. But this contradicts the fact that the complementary subset of s_i has by definition the opposite characterization, unless $s_i \cap s'_j = s_i = s'_j$. The two remaining subsets should thus be identical, too, whence the uniqueness of the minimal partition.

FIG. 8. Boolean flowchart of the fully resolved form, i.e., with no dependence on the fields, of the NAS conditions on $\lambda_B, \lambda_H, \lambda_A^{(1)}, \lambda_A^{(2)}, \lambda_{AB}^{(1)}, \lambda_{AB}^{(2)}$ satisfying the inequalities given by Eqs. (3.34) and (3.35).

illustrates a generic case where Eq. (3.66) can never be satisfied irrespective of the yes/no configurations. The middle-right and bottom-left figures, and more generally when the solid and dashed lines do not cross, illustrate the necessary configurations to allow for Eq. (3.66), yet one still needs to examine the yes/no configurations for sufficiency. Finally the bottom-right figure where the two branches of the parabola cut through the domain is another configuration for which Eq. (3.66) is impossible. Finally, when $\alpha_{AB}^* \notin [0, 1]$, not represented in Fig. 2, the entire (α_{AB}, α_A) domain is contained either in the nonempty subset of $\{s_{\text{yes}}^I, \emptyset\}$ or in the nonempty subset of $\{\emptyset, s_{\text{no}}^I\}$. In the latter case it is required to be entirely contained in the yes region determined by the parabola.

Putting everything together, the problem becomes equivalent to solving for the following complementary conditions:

- (i) Eq. (3.35)-(I) valid $\forall \alpha_{AB} \in [0, 1]$, partition $\{s_{\text{yes}}^I, \emptyset\}$;
- (ii) $\alpha_{AB}^* = -\frac{\lambda_{AB}^{(1)}}{\lambda_{AB}^{(2)}}$,
Eq. (3.35)-(I) valid only $\forall \alpha_{AB} \in [0, \alpha_{AB}^*]$,
Eq. (3.35)-(II) should be valid $\forall \alpha_{AB} \in [\alpha_{AB}^*, 1]$,
i.e., $s_{\text{no}}^I \subset s_{\text{yes}}^{\text{II}}$;
- (iii) $\alpha_{AB}^* = -\frac{\lambda_{AB}^{(1)}}{\lambda_{AB}^{(2)}}$,
Eq. (3.35)-(I) valid only $\forall \alpha_{AB} \in [\alpha_{AB}^*, 1]$,
Eq. (3.35)-(II) should be valid $\forall \alpha_{AB} \in [0, \alpha_{AB}^*]$,
i.e., $s_{\text{no}}^I \subset s_{\text{yes}}^{\text{II}}$;
- (iv) Eq. (3.35)-(I) false $\forall \alpha_{AB} \in [0, 1]$, partition $\{\emptyset, s_{\text{no}}^I\}$,
Eq. (3.35)-(II) should be valid $\forall \alpha_{AB} \in [0, 1]$, partition $\{s_{\text{yes}}^{\text{II}}, \emptyset\}$,

where the numbering corresponds to that of Fig. 8.

We can now derive in a fully analytical way the resolved form of Eqs. (3.34) and (3.35), or equivalently of Eqs. (3.35) and (3.36) in conjunction with $\lambda_H > 0 \wedge \lambda_B > 0$. The NAS conditions thus obtained on the λ 's have no residual dependence on α_{AB} and α_A . To retrieve these NAS conditions we followed step-by-step the partitions described above and analyzed the nonmonotonic dependence on α_{AB} in Eq. (3.35)-(II) when applicable.

The details are very technical and will not be described here. We give the final result in Fig. 8 where we have defined the following Boolean expressions:

$$\mathcal{B}_3 \Leftrightarrow (2\lambda_A^{(1)} + \lambda_A^{(2)})\lambda_B > 12(\lambda_{AB}^{(1)} + \lambda_{AB}^{(2)})^2, \quad (3.67)$$

$$\begin{aligned} \mathcal{B}_4 \Leftrightarrow & 3(2\lambda_A^{(1)} + \lambda_A^{(2)})\lambda_{AB}^{(2)2} + 2\lambda_A^{(2)}(\lambda_A^{(1)} + \lambda_A^{(2)})\lambda_B \\ & < 12\lambda_A^{(2)}\lambda_{AB}^{(1)}(\lambda_{AB}^{(1)} + \lambda_{AB}^{(2)}) \\ & \vee 6\lambda_{AB}^{(2)}(\lambda_{AB}^{(1)} + \lambda_{AB}^{(2)}) + \lambda_A^{(2)}\lambda_B > 0, \end{aligned} \quad (3.68)$$

$$\mathcal{B}_5 \Leftrightarrow 2(\lambda_A^{(1)} + \lambda_A^{(2)})\lambda_B > 3(2\lambda_{AB}^{(1)} + \lambda_{AB}^{(2)})^2, \quad (3.69)$$

$$\begin{aligned} \mathcal{B}_6 \Leftrightarrow & (\lambda_A^{(1)} + \lambda_A^{(2)})\lambda_{AB}^{(2)2} > 4\lambda_A^{(2)}\lambda_{AB}^{(1)}(\lambda_{AB}^{(1)} + \lambda_{AB}^{(2)}) \\ & \vee 3\lambda_{AB}^{(2)2} + \lambda_A^{(2)}\lambda_B < 0, \end{aligned} \quad (3.70)$$

$$\mathcal{B}_7 \Leftrightarrow \lambda_B \min \{2(\lambda_A^{(1)} + \lambda_A^{(2)}), (2\lambda_A^{(1)} + \lambda_A^{(2)})\} > 12\lambda_{AB}^{(1)2}. \quad (3.71)$$

In writing this final form we used occasionally the fact that $\lambda_B > 0$ to obtain compact expressions where Eqs. (3.36) are implicitly taken into account in Eqs. (3.67), (3.69), and (3.71).

For a cross-check of our results we have performed various numerical scans simultaneously on the λ 's, and on α_A, α_{AB} in the domain defined by Eqs. (3.47)-I–(3.47)-IV. This amounts to checking the validity of the conditions Eqs. (3.35) and (3.36), and comparing the Boolean output with that of the resolved conditions of Fig. 8.¹⁰ One can take advantage of the fact that α_{AB} and α_A are not correlated in the square $[0, \frac{1}{2}] \times [\frac{1}{2}, 1]$ to replace for this part of the domain, and without loss of information, α_{AB} and α_A by their edge values in Eqs. (3.35)-(I) and (3.35)-(II). The parabola-edged part of the domain (where $\alpha_{AB} \in [\frac{1}{2}, 1]$) is more tricky. If not sufficiently finely meshed, a numerical scan could miss some features depending on the configuration of the maximum/minimum of the parabola. As an illustration we show in Fig. 9 the allowed 3D domains, for subsets of the λ parameters, obtained from the resolved exact conditions of Fig. 8 and compare them with the approximate ones obtained from requiring Eqs. (3.35) and (3.36) to hold for just three sets of benchmark values of α_{AB} and α_A lying on the boundary of their allowed domain. As expected, one of the benchmark sets leads to an approximate domain that is much less restrictive [the pink colored regions in Figs. 9(a) and 9(c)] than the exact domain shown in Figs. 9(b) and 9(d). However, one finds that the other benchmark set [the brown colored regions in Figs. 9(a) and 9(c)] leads unexpectedly to an extremely good approximation of the exact domain. Obviously this accidental agreement could not have been guessed without the comparison and is not by itself a cross-check of the validity of the conditions given in Fig. 8 and Eqs. (3.67)–(3.71). For that we have performed large scans, 4×10^6 points on a regular grid in the λ space in the configurations of Fig. 9, or fixing only $\lambda_H = 1$ and taking $1.3 - 2 \times 10^6$ points in the λ space with a much larger number of benchmark points, 60 benchmark points within, or 30 benchmark points on, the boundary of the (α_{AB}, α_A) domain. Counting the hits where the Boolean values of the approximate and exact conditions are equal or different we found in all cases a difference of less than 2% between the approximate and exact conditions. Another significant feature of the check is that the Boolean yield of the difference is found in 100% of the cases to be “approximate = True, exact = False.” Only one hit with the reverse configuration would have meant the exact conditions are wrong.

¹⁰Throughout the paper we rely significantly on the *Mathematica* package [53] for symbolic and numerical computations as well as for the generation of the plots.

FIG. 9. Upper figures: the allowed domain in the $\lambda_B, \lambda_{AB}^{(1)}, \lambda_{AB}^{(2)}$ space for $\lambda_A^{(1)} = -0.1, \lambda_A^{(2)} = 1, \lambda_H = 1$; (a) the brown domain corresponds to enforcing Eqs. (3.35) and (3.36) for just the three sets of values, $(\alpha_{AB}, \alpha_A) = (0, \frac{1}{2}), (\frac{1}{2}, \frac{1}{2}), (1, \frac{1}{2})$; the light pink indicates the increased domain when replacing the last set by $(\frac{1}{2}, 1)$; (b) exact resolved conditions of Fig. 8. Lower figures: the allowed domain in the $\lambda_B, \lambda_A^{(1)}, \lambda_{AB}^{(2)}$ space, for $\lambda_{AB}^{(1)} = -0.1, \lambda_A^{(2)} = 1, \lambda_H = 1$; (c) as in (a); (d) as in (b).

In summary, we have derived the NAS conditions for $\mathbf{a}_o > 0 \wedge \mathbf{c}_o > 0$ in a fully analytical resolved form. They are thus *necessary* for the BFB of the general potential given by Eqs. (3.2)–(3.4) and can safely be applied irrespective of the A, B , and H field configurations.¹¹ Further comments on these conditions are deferred to Secs. IV and V.

¹¹Note that an alternative approach to obtain these results is to start from the third inequality in Eq. (3.30) with no OR structure rather than from Eq. (3.33). Its advantage is to avoid the use of partitions and coverings but necessitates the study of functions with square roots as in Appendix B leading though to more compact conditions. We have checked the agreement of the two approaches. The partitions/coverings approach we developed will nevertheless be unavoidable for the all-field-directions full analytical resolving of the precustodial model in the case $\lambda_{ABH} = 0$, not treated in the present paper.

2. Partial resolving of $\mathbf{b}_o > 0 \vee 4\mathbf{a}_o\mathbf{c}_o - \mathbf{b}_o^2 > 0$

We investigate now Eq. (3.33) that should be valid $\forall Z, \alpha_A, \alpha_{AB}, \alpha_{AH}, \alpha_{ABH}$ in their allowed domains. [We use here the variable Z defined in Eq. (3.38) instead of t , and refer the reader to Sec. III C 2 for a discussion on the relevance of Z .] As argued repeatedly in Secs. III C 1 and III C 2 and discussed in detail in the previous subsection, the “or” structure in Eq. (3.33) implies that the validity of the inequalities should be required for all possible coverings of the $(Z, \alpha\text{-parameters})$ space. However, the situation is more complex here than in the previous subsection, since $4\mathbf{a}_o\mathbf{c}_o - \mathbf{b}_o^2$ [cf. Eq. (3.24)] involves simultaneously all four α ’s and is a complete quartic polynomial in Z . Given the particularly involved NAS conditions for quartic polynomials, Eqs. (G30a)–(G30d), we do not expect to resolve completely this case in an explicit form similar to that given

FIG. 10. Boolean flowchart for the resolving of $\mathfrak{b}_o > 0 \vee 4\mathfrak{a}_o \mathfrak{c}_o - \mathfrak{b}_o^2 > 0$. $z_{\pm}$ denote the two real-valued roots of $\mathfrak{b}_o(Z)$ when they exist. See text for the definitions of $\mathcal{B}_8$ and $\mathcal{B}_9$. We use the conventional notations $\wedge$, $\vee$ and $\neg$ for the Boolean operators “AND,” “OR” and “NOT,” respectively.

in Fig. 8. The aim here is to proceed as far as possible toward an explicitly resolved form, and then deal with the rest through mere numerical scans on the α parameters defined by Eq. (3.61), including some further refinements to be discussed in Sec. IV. To proceed let us first address the flowchart of the overall logic. This is sketched in Fig. 10, together with the following definitions:

- (i) $\mathcal{B}_8$ denotes the NAS conditions for $\mathfrak{b}_o$ to always have a constant sign;
- (ii) $\mathcal{B}_9^{(a,b)}$ denotes the NAS conditions for $4\mathfrak{a}_o \mathfrak{c}_o - \mathfrak{b}_o^2$ to be positive when Z is in the interval (a, b) and the α parameters satisfy Eq. (3.61).

The strategy underlying this flowchart is similar to the one adopted in the previous subsection (which the reader is referred to for definitions and notations), and should be clear by now. The upper left box of Fig. 10 corresponds to the λ -space points for which $\mathfrak{b}_o > 0$ defines two trivial minimal partitions, $\{s_{\text{yes}}^{\text{I}}, \emptyset\}$ or $\{\emptyset, s_{\text{no}}^{\text{I}}\}$, corresponding, respectively, to $\lambda_{BH} > 0$ and $\lambda_{BH} < 0$, while the lower left box corresponds to the λ -space points where $\mathfrak{b}_o > 0$ defines a generic minimal partition $\{s_{\text{yes}}^{\text{I}}, s_{\text{no}}^{\text{I}}\}$. The boxes to the right indicate the Boolean structure including the minimal generic partition $\{s_{\text{yes}}^{\text{II}}, s_{\text{no}}^{\text{II}}\}$ defined by $4\mathfrak{a}_o \mathfrak{c}_o - \mathfrak{b}_o^2 > 0$ to satisfy Eq. (3.66). We now investigate how far the

Boolean expressions $\mathcal{B}_8$ and $\mathcal{B}_9^{(\cdots)}$ can be resolved analytically.

• $\mathcal{B}_8$: Viewing $\mathfrak{b}_o$, Eq. (3.24), as a quadratic polynomial in Z , we denote by $z_{\pm}$ its two roots. Thus $\mathcal{B}_8$ corresponds to the NAS condition for which $z_{\pm}$ are not real valued, that is, to require the discriminant of this polynomial to be negative,

$$\mathcal{B}_8 \equiv (\alpha_{ABH} \lambda_{ABH})^2 - 8(\lambda_{AH}^{(1)} + \alpha_{AH} \lambda_{AH}^{(2)}) \lambda_{BH} \leq 0 \quad (3.72)$$

for all α_{AH} , α_{ABH} in the domain given by Eqs. (3.51) and (3.52). Taking into account the correlations at the boundary of this domain one can obtain condition $\mathcal{B}_8$ in a fully resolved analytical form. After some nontrivial Boolean simplifications we find

$$\begin{aligned} \{(\lambda_{ABH})^2 \leq 4(\lambda_{AH}^{(1)} + \lambda_{AH}^{(2)}) \lambda_{BH} \wedge ((\lambda_{ABH})^2 \\ \geq 4\lambda_{AH}^{(2)} \lambda_{BH} \vee \lambda_{AH}^{(1)} \lambda_{BH} \geq 0)\} \Leftrightarrow \mathcal{B}_8. \end{aligned} \quad (3.73)$$

Clearly then, the NAS conditions for the sufficient condition $\mathfrak{b}_o > 0$ read (see Fig. 10)

$$\mathcal{B}_8 \wedge \lambda_{BH} > 0. \quad (3.74)$$

However, as will be discussed later on in Sec. IV, the condition on the left-hand side of Eq. (3.73) is, in fact, only sufficient to yield $\mathcal{B}_8$.

• $\mathcal{B}_9^{(-\infty, +\infty)}$: To obtain $\mathcal{B}_9^{(-\infty, +\infty)}$ we consider $4\mathbf{a}_0\mathbf{c}_0 - \mathbf{b}_0^2$ as a quartic polynomial in Z and thus require all the conditions given by Eqs. (G30a)–(G30d). The coefficients $a_{i=0,\dots,4}$ are straightforwardly read from the combination $4\mathbf{a}_0\mathbf{c}_0 - \mathbf{b}_0^2$ upon use of Eqs. (3.24) and (3.38):

$$\begin{aligned} a_0 &= \gamma_0 - \delta_0^2, & a_1 &= -2\delta_0\delta_1, & a_2 &= \gamma_1 - \delta_1^2 - 2\delta_0\delta_2, \\ a_3 &= -2\delta_1\delta_2, & a_4 &= \gamma_2 - \delta_2^2, \end{aligned} \quad (3.75)$$

where

$$\begin{aligned} \delta_0 &= \lambda_{AH}^{(1)} + \alpha_{AH}\lambda_{AH}^{(2)}, & \delta_1 &= \frac{1}{2}\lambda_{ABH}, & \delta_2 &= \frac{\lambda_{BH}}{2\alpha_{ABH}^2}, \\ \gamma_0 &= \frac{1}{4}(\lambda_A^{(1)} + \alpha_A\lambda_A^{(2)})\lambda_H, & \gamma_1 &= \frac{(\lambda_{AB}^{(1)} + \alpha_{AB}\lambda_{AB}^{(2)})\lambda_H}{2\alpha_{ABH}^2}, \\ \gamma_2 &= \frac{\lambda_B\lambda_H}{24\alpha_{ABH}^4}. \end{aligned} \quad (3.76)$$

We provide here explicitly the resulting first three conditions given by Eqs. (G30a):

$$\begin{aligned} \mathcal{B}_9^{(-\infty, +\infty)} &\supset \lambda_B\lambda_H > 6\lambda_{BH}^2 \\ &\wedge \\ &(\lambda_A^{(1)} + \lambda_A^{(2)})\lambda_H > 4 \max \{(\lambda_{AH}^{(1)})^2, (\lambda_{AH}^{(1)} + \lambda_{AH}^{(2)})^2\} \\ &\wedge \\ &(\lambda_A^{(2)}\lambda_H < 2(\lambda_{AH}^{(2)})^2 \vee \lambda_A^{(2)}\lambda_H < 4 \max \{-\lambda_{AH}^{(1)}\lambda_{AH}^{(2)}, \lambda_{AH}^{(2)}(\lambda_{AH}^{(1)} + \lambda_{AH}^{(2)})\} \\ &\vee \lambda_A^{(2)}(2\lambda_A^{(1)} + \lambda_A^{(2)})\lambda_H > 4((\lambda_A^{(1)} + \lambda_A^{(2)})(\lambda_{AH}^{(2)})^2 + 2\lambda_A^{(2)}\lambda_{AH}^{(1)}(\lambda_{AH}^{(1)} + \lambda_{AH}^{(2)}))). \end{aligned} \quad (3.81)$$

Condition (3.79) appears much less amenable to a resolved form as it involves all four α parameters simultaneously. One can, however, still resolve it partially but this will not be pursued further here.¹² The remaining conditions corresponding to Eqs. (G30b), (G30c), and (G30d) will be treated numerically.

• $\mathcal{B}_9^{(-\infty, z_-), \mathcal{B}_9^{(z_+, +\infty)}$: To obtain these conditions one again considers $4\mathbf{a}_0\mathbf{c}_0 - \mathbf{b}_0^2$ as a quartic polynomial in Z . However, now the positivity is not required on all $Z \in (-\infty, +\infty)$, and one needs to rely on the results derived in Appendix H. Since the latter holds for $[0, +\infty)$, we first map one-to-one the domains $(-\infty, z_-]$

¹²For instance, since it is biquadratic in α_{ABH} with a positive definite coefficient of α_{ABH}^4 , a *sufficient* condition is then a negative discriminant. The latter has a simple form depending linearly on α_A , α_{AB} and quadratically on α_{AH} .

$$4a_0 = (\lambda_A^{(1)} + \alpha_A\lambda_A^{(2)})\lambda_H - 4(\lambda_{AH}^{(1)} + \alpha_{AH}\lambda_{AH}^{(2)})^2 > 0, \quad (3.77)$$

$$24\alpha_{ABH}^4 a_4 = \lambda_B\lambda_H - 6\lambda_{BH}^2 > 0, \quad (3.78)$$

$$\begin{aligned} 16\alpha_{ABH}^4 \Delta_0 &= \lambda_{ABH}^4 \alpha_{ABH}^4 - 4\lambda_{ABH}^2 (\lambda_H X_{AB} + 4\lambda_{BH} X_{AH}) \alpha_{ABH}^2 \\ &\quad + 4(\lambda_H X_{AB} - 2\lambda_{BH} X_{AH})^2 \\ &\quad + 8a_0(\lambda_B\lambda_H - 6\lambda_{BH}^2) > 0, \end{aligned} \quad (3.79)$$

where we defined

$$X_{AK} \equiv \lambda_{AK}^{(1)} + \alpha_{AK}\lambda_{AK}^{(2)} \quad (K = B, H). \quad (3.80)$$

Condition (3.77) can readily be resolved: Being linear in α_A , one requires it to hold simultaneously on the upper and lower boundary lines of the α_A domain given by Eq. (3.41). The resulting conditions depend only on α_{AH} quadratically and can be studied straightforwardly taking into account Eq. (3.40). After several Boolean simplifications we find the following resolved form of Eq. (3.77), adding also Eq. (3.78):

and $[z_+, +\infty)$ on $[0, +\infty)$ through the two changes of variable

$$Z = z_- - \xi \quad \text{and} \quad Z = z_+ + \xi, \quad (3.83)$$

respectively, with $\xi \in [0, +\infty)$, then search for the conditions on the quartic polynomial in ξ satisfying criterion (H13). We note, however, two simplifications due to the linear changes of variable: a_4 , the coefficient of Z^4 given by Eq. (3.78), is the same as that of ξ^4 . It follows that the *necessary* condition Eq. (3.81) remains valid. On the other hand, the coefficients a_0 are modified with respect to Eq. (3.77) to, respectively, a_0^- and a_0^+ given by

$$a_0^\mp = \lambda_H(6(\lambda_A^{(1)} + \alpha_A\lambda_A^{(2)}) + 12(\lambda_{AB}^{(1)} + \alpha_{AB}\lambda_{AB}^{(2)})z_\mp^2 + \lambda_B z_\mp^4). \quad (3.84)$$

Interestingly, one can show that when combined with Eqs. (3.34) and (3.35), the necessary constraints $a_0^\mp > 0$ as dictated by the first of Eqs. (G30a), will *always* be satisfied by Eq. (3.84) irrespective of the values of $z_\mp$. Indeed, given Eq. (3.34), when Eq. (3.35)-(I) is satisfied, then $a_0^\mp > 0$ follows trivially; and when Eq. (3.35)-(II) is satisfied, then $a_0^\mp$, taken as a quadratic equation in $z_\mp^2$, has no real-valued roots and thus again is always positive.

• $\mathcal{B}_9^{(z_-, z_+)}$: In this case a nonlinear change of variable

$$Z = z_- + (z_+ - z_-) \frac{\xi}{1 + \xi} \quad (3.85)$$

is used with $\xi \in [0, +\infty)$ before applying criterion (H13). Here, too, a simplification occurs for a_0 and a_4 after the change of variable. Up to a global positive definite denominator, they are expressed in terms of Eq. (3.84),

$$\begin{aligned} a_0 &= a_0^-, \\ a_4 &= a_0^+, \end{aligned} \quad (3.86)$$

and are thus *always* positive when combined with Eqs. (3.34) and (3.35), as explained above.

To summarize, we have identified a subset of analytically resolved necessary conditions in the various branches of the Fig. 10 flowchart. One now should combine these conditions with the other analytically resolved conditions given in Fig. 8 and Eqs. (3.67)–(3.71) and possibly also with those given by Eqs. (2.14)–(2.17). This allows a quick determination of necessary domains in the λ -space. Then adding the remaining necessary conditions that can be treated through numerical scans on the α parameters, one delineates the NAS BFB conditions. However, before doing so in Sec. V, we need to reexamine first the BFB conditions of the more constrained Georgi-Machacek model, as this will have some bearing on the general case.

F. The Georgi-Machacek BFB conditions

In [33] the authors provided a detailed study of the properties of the potential relying on a generalization of the parametrization used in [31]. They identified the two parameters

$$\hat{\omega} = \frac{\text{Tr}(\Phi^\dagger \tau^a \Phi \tau^b) \text{Tr}(X^\dagger t^a X t^b)}{\text{Tr}(\Phi^\dagger \Phi) \text{Tr}(X^\dagger X)}, \quad (3.87)$$

$$\hat{\xi} = \frac{\text{Tr}(X^\dagger X X^\dagger X)}{(\text{Tr}(X^\dagger X))^2}, \quad (3.88)$$

relevant to the study of the BFB conditions, writing $V_{\text{G-M}}^{(4)}$ in the form

$$V_{\text{G-M}}^{(4)} = \hat{r}^4 \cos^4 \hat{\gamma} (\hat{\lambda}_1 + (\hat{\lambda}_2 - \hat{\omega} \hat{\lambda}_5) \tan^2 \hat{\gamma} + (\hat{\xi} \hat{\lambda}_3 + \hat{\lambda}_4) \tan^4 \hat{\gamma}), \quad (3.89)$$

with

$$\hat{r}^2 \equiv \text{Tr}(\Phi^\dagger \Phi) + \text{Tr}(X^\dagger X), \quad (3.90)$$

$$\tan^2 \hat{\gamma} \equiv \frac{\text{Tr}(X^\dagger X)}{\text{Tr}(\Phi^\dagger \Phi)}. \quad (3.91)$$

Noting that $\text{Tr}(\Phi^\dagger \Phi) = 2H^\dagger H$ and $\text{Tr}(X^\dagger X) = 4\text{Tr}(AA^\dagger) + 2\text{Tr}(B^2)$ one can relate $\hat{r}$ and $\tan \hat{\gamma}$ to the parameters defined in Eqs. (3.16)–(3.19) to obtain

$$\tan^2 \hat{\gamma} = (1 + \cos^2 b) \tan^2 a \quad \text{and} \quad \hat{r}^2 \cos^2 \hat{\gamma} = 2r^2 \cos^2 a. \quad (3.92)$$

Then equating $V_{\text{G-M}}^{(4)}$, Eq. (3.89), with $V_{\text{p-c}}^{(4)}$, Eq. (3.4), and taking into account the above relations and Eqs. (3.11) and (3.12), one identifies $\hat{\omega}$ and $\hat{\xi}$ as the coefficients of $-\hat{\lambda}_5 \tan^2 \hat{\gamma}$ and $\hat{\lambda}_3 \tan^4 \hat{\gamma}$ which allows one to relate them to the parameters defined in the precustodial case, Eqs. (3.20)–(3.22), as follows:

$$\hat{\omega} = -\frac{1 - 2\alpha_{AH} - \sqrt{2}\alpha_{ABH}t}{2(2 + t^2)}, \quad (3.93)$$

$$\hat{\xi} = \frac{6 - 4\alpha_A + 4\alpha_{AB}t^2 + t^4}{(2 + t^2)^2}. \quad (3.94)$$

As a cross-check of the validity of these relations, one can indeed retrieve from the fact that $t \in [0, +\infty)$ and the exact knowledge of the two domains given by Eqs. (3.47)-I–(3.47)-IV and Eqs. (3.51) and (3.52) that $\hat{\omega} \in [-\frac{1}{4}, \frac{1}{2}]$ and $\hat{\xi} \in [\frac{1}{3}, 1]$ as already found in [33].

The allowed domain in the $(\hat{\omega}, \hat{\xi})$ plane has been given in [33]. This was done stating that the boundary of the domain is obtained from the real valued components of the neutral field directions, that is, keeping only $\text{Re}\chi^0$ and ξ^0 and zeroing all the others in Eqs. (3.87) and (3.88). However, no justification was given for this statement. The aim of the present section is to provide an explicit proof for the equation of the boundary of the $(\hat{\omega}, \hat{\xi})$ domain based on the symmetries of $V_{\text{G-M}}$. We choose to use $SU(2)_R$ to rotate away the lower as well as the imaginary parts of the upper components of H , so that

$$\frac{\text{Tr}(\Phi^\dagger \tau^a \Phi \tau^b)}{\text{Tr}(\Phi^\dagger \Phi)} = \frac{1}{4} \delta_{ab} \quad (3.95)$$

[note that Ref. [33] used $SU(2)_L$ instead], and use $SU(2)_L$ to rotate away, for instance, χ^{++} and the imaginary part of χ^+ , bringing the bi-triplet X in the form

$$X = \begin{pmatrix} \chi^{0*} & \xi^+ & 0 \\ -u & \xi^0 & u \\ 0 & -\xi^{+*} & \chi^0 \end{pmatrix}, \quad (3.96)$$

where $u(\equiv \text{Re}\chi^+)$ denotes a real-valued scalar field.¹³ With this choice of gauge $\hat{\omega}$ and $\hat{\xi}$ take the following form:

$$\hat{\omega} = \frac{1}{4}(2\sqrt{2}\cos\theta_0\cos(\arg(\chi^0)) + \sin\theta_0)\sin\theta_0\sin^2\theta_+ \quad (3.97)$$

$$+ \frac{1}{2}\cos(\arg(\xi^+))\cos\theta_+\cot\theta_u + \mathcal{O}(\cot^2\theta_u), \quad (3.98)$$

$$\hat{\xi} = 1 - \sin^2\theta_0\sin^2\theta_+ \left(1 + \frac{1}{4}(1 + 3\cos 2\theta_0)\sin^2\theta_+\right) \quad (3.99)$$

$$- \sqrt{2}\cos(\arg(\xi^+) + \arg(\chi^0))\cos\theta_+\sin^2\theta_+\sin 2\theta_0\cot\theta_u \\ + \mathcal{O}(\cot^2\theta_u), \quad (3.100)$$

where we defined the polar angles by

$$u = R\cos\theta_u, \quad (3.101)$$

$$|\xi^+| = R\cos\theta_+\sin\theta_u, \quad (3.102)$$

$$|\chi^0| = R\sin\theta_+\sin\theta_0\sin\theta_u, \quad (3.103)$$

$$\xi^0 = \sqrt{2}R\sin\theta_+\cos\theta_0\sin\theta_u, \quad (3.104)$$

$$R^2 = \frac{1}{2}\text{Tr}(X^\dagger X), \quad (3.105)$$

with

$$0 \leq \arg(\chi^0), \arg(\xi^+) \leq 2\pi, \quad (3.106)$$

$$0 \leq \theta_0, \theta_+ \leq \frac{\pi}{2}, \quad (3.107)$$

$$0 \leq \theta_u \leq \pi. \quad (3.108)$$

Note that due to the invariance of $V_{\text{G-M}}^{(4)}$ under $X \rightarrow -X$ one can always fix uniquely either the sign of ξ^0 or that of u . In our parametrization $\xi^0 > 0$ while u can take either sign. In Eqs. (3.98) and (3.100) we kept for simplicity only linear terms in u . We will come back to the exact contribution

¹³One could be tempted to zero, on top of χ^{++} , the (real-valued) ξ^0 entry rather than $\text{Im}\chi^+$. However, one can show that this is not possible through a noninfinitesimal $SU(2)$ rotation. More generally, one cannot zero more than two entries of X through $SU(2)_L \times SU(2)_R$ rotations.

later on. Here we first concentrate on the zeroth order u contributions to $\hat{\omega}$ and $\hat{\xi}$, i.e., Eqs. (3.97) and (3.99), which we dub $\hat{\omega}_0$ and $\hat{\xi}_0$. In Appendix E we give a detailed proof for the determination of the boundary in the $(\hat{\omega}_0, \hat{\xi}_0)$ domain, i.e., *under the working assumption that $u = 0 (= \cot\theta_u)$* . We find that this boundary is defined by the following upper and lower curves:

$$\hat{\xi}_0^{\max}(\hat{\omega}_0) = \frac{1}{3} + \frac{2}{27}(1 - 2\hat{\omega}_0 + 2\sqrt{(1 - 2\hat{\omega}_0)(1 + 4\hat{\omega}_0)})^2, \\ \text{for } \hat{\omega}_0 \in \left[-\frac{1}{4}, \frac{1}{2}\right], \quad (3.109)$$

$$\hat{\xi}_0^{\min}(\hat{\omega}_0) = \frac{1}{3} + \frac{2}{27}(1 - 2\hat{\omega}_0 - 2\sqrt{(1 - 2\hat{\omega}_0)(1 + 4\hat{\omega}_0)})^2, \\ \text{for } \hat{\omega}_0 \in \left[-\frac{1}{4}, -\frac{1}{6}\right], \quad (3.110a)$$

$$\hat{\xi}_0^{\min}(\hat{\omega}_0) = \frac{1}{3}, \quad \text{for } \hat{\omega}_0 \in \left[-\frac{1}{6}, \frac{1}{2}\right]. \quad (3.110b)$$

This reproduces exactly the boundary given in Ref. [33] as illustrated in Fig. 11 (note, however, that we deal with the inverse function with respect to Ref. [33]). As shown in Appendix E3 the condition $\sin^2\theta_+ = 1$, i.e., $\xi^+ = 0$, is sufficient and necessary for the determination of the $(\hat{\omega}_0, \hat{\xi}_0)$ boundary. In particular, the necessity of this condition is a nontrivial result. From Eq. (3.97) one sees that $\sin^2\theta_+ = 0$ could as well have defined a boundary. More importantly, the involved dependence on $\sin^2\theta_+$ in $\hat{\xi}_0$, Eq. (3.99), could in principle lead to portions of the boundary with $\sin^2\theta_+ < 1$, since we are interested in the projection on the $(\hat{\omega}_0, \hat{\xi}_0)$ plane. [This was, for instance, the case for the (α_A, α_{ABH}) domain studied in Sec. III D 4.] Moreover, this is not the end of the story because the boundary defined by Eqs. (3.109) and (3.110b) is obtained in the case $u = 0$. It remains to be seen whether $u \neq 0$ would possibly enlarge the allowed domain outside this boundary. We turn now to this point. The idea is to consider a subspace of the field space for which the boundary of $(\hat{\omega}_0, \hat{\xi}_0)$ is reached and determine within this subspace the boundary of $(\hat{\omega}, \hat{\xi})$ allowing for $u \neq 0$. As discussed above, such a subspace has necessarily $\xi^+ = 0$ ($\sin^2\theta_+ = 1$). The bi-triplet of Eq. (3.96) becomes

$$X = \begin{pmatrix} \chi^{0*} & 0 & 0 \\ -u & \xi^0 & u \\ 0 & 0 & \chi^0 \end{pmatrix}. \quad (3.111)$$

One then sees from Eqs. (3.98) and (3.100) that the first order u contributions vanish for any u in this subspace,

FIG. 11. The boundary in the $(\hat{\omega}_0, \hat{\xi}_0)$ plane delimiting the allowed inner domain, in the limit $u = 0$. This agrees with Ref. [33].

indicating that the boundary is indeed unchanged when $u \neq 0$ at least if u remains sufficiently small. In fact, this result remains true in general beyond the first order as a consequence of an accidental symmetry: $\text{Tr}(X^\dagger X)$ and $\text{Tr}(X^\dagger t^a X t^a)$ (summation over a) are invariant under the substitution $\chi^+ \leftrightarrow \xi^{++} (\chi^{++} \leftrightarrow \xi^+)$, and $\text{Tr}(X^\dagger X X^\dagger X)$ is invariant under the same substitution supplemented by $\chi^{++} \leftrightarrow \chi^{++*}$. Thus $\hat{\omega}$ and $\hat{\xi}$ are invariant under these substitutions, in which case X defined in Eq. (3.111) is replaced by

$$\tilde{X} = \begin{pmatrix} \chi^{0*} & u & 0 \\ 0 & \xi^0 & 0 \\ 0 & -u & \chi^0 \end{pmatrix}. \quad (3.112)$$

The key point is that the latter $\tilde{X}$ has the same form as X given by Eq. (3.96) with $u = 0$. We are then brought back to the same configuration that leads to the fact that the boundary is reached for $\xi^+ = 0$ and is given by Eqs. (3.109)–(3.110b); applied to the present case where X is replaced by $\tilde{X}$ implies similarly that the boundary is reached for $u = 0$ and is given by the same Eqs. (3.109)–(3.110b). This completes the proof that $u \neq 0$ in Eq. (3.96) remains within the boundary obtained for $u = 0$. Thus the full boundary in the $(\hat{\omega}, \hat{\xi})$ plane is given by Eqs. (3.109)–(3.110b):

$$\hat{\xi}_0^{\min}(\hat{\omega}) \leq \hat{\xi} \leq \hat{\xi}_0^{\max}(\hat{\omega}). \quad (3.113)$$

In the following we will refer to this domain as the $\omega - \xi$ chips.

IV. PEELING THE POTATOID WITH THE CHIPS

As already announced at the end of Sec. III D 7, the knowledge of the exact domain of the $\omega - \xi$ chips of the Georgi-Machacek model will have a spin-off on the refinement of the 4D α -parameters potatoid in the general precustodial model. That a model with an enlarged symmetry would backreact on a less symmetric and more general model is somewhat unusual. It can be understood as follows in the case at hand: The $SU(2)_L \times SU(2)_R$ symmetry of the Georgi-Machacek model has allowed one to regroup the four α parameters and the t parameter into just two relevant parameters $\hat{\omega}$ and $\hat{\xi}$ that are related to the former as given by Eqs. (3.93) and (3.94). However, the equations defining the $\omega - \xi$ chips, Eqs. (3.113), (3.109)–(3.110b), were arrived at thanks to the gauge and global symmetries, as well as to an accidental invariance of the quartic part of the Georgi-Machacek potential (see Sec. III F and Appendix E); in this, Eqs. (3.93) and (3.94) played no role. The latter, in conjunction with Eq. (3.113), will thus lead to a supplementary correlation among the α parameters and t that should be valid in the general precustodial model. It is in that sense that the Georgi-Machacek model informs about the more general model. Obviously, this information would have been redundant had we had beforehand a full knowledge of the exact 4D α -parameters domain. This is, however, not the case as pointed out in Sec. III D 7 regarding the α potatoid. Hence one can use the above information as a sufficient condition to exclude points in the α potatoid as follows: Each set of α parameters in the α potatoid defines, through Eqs. (3.93) and (3.94), a unique trajectory $(\hat{\omega}_\alpha(t), \hat{\xi}_\alpha(t))$ in the $(\hat{\omega}, \hat{\xi})$ plane, parametrized by $t \in [0, +\infty)$. If the trajectory goes out of the $\omega - \xi$ chips, then the corresponding set of α -parameters values should be excluded.

We show in Figs. 12 and 13 numerical scans taking into account this exclusion criterion.¹⁴ The red and blue dots delineate the somewhat convoluted regions of the α potatoid that are incompatible with the $\omega - \xi$ chips. As anticipated in Sec. III D 7 and visible from the different viewing angles in Fig. 12, the excluded portions lie only at the boundary of the α potatoid. Note that the domains (shown in pink) in Figs. 12 and 13 are 3D sections of the 4D α potatoid at fixed values of α_A or α_{AB} or α_{AH} , respectively; not to be confused with the 3D projections of the α potatoid shown in Figs. 7(a), 7(b), and 7(d), with which it would not be possible to disentangle boundaries

¹⁴In practice this is achieved by scanning over the four α parameters that satisfy Eq. (3.61) and following each trajectory $(\hat{\omega}_\alpha(t), \hat{\xi}_\alpha(t))$ scanning over $0 \leq u \leq \frac{\pi}{2}$ with $t = \sqrt{2} \tan u$. Alternatively, one can use the exact t -resolved form for Eqs. (3.93) and (3.94) (see Appendix F) and scan only on the α parameters. We used this latter alternative to cross-check our results.

FIG. 12. The $(\alpha_{AB}, \alpha_{AH}, \alpha_{ABH})$ 3D section of the 4D α potatoid at $\alpha_A = 1$, viewed from four different angles. The red and blue dots denote the regions excluded by the $\omega - \zeta$ chips. See text for further discussions.

unambiguously. Moreover, the choices of $\alpha_A = 1$, $\alpha_{AH} = \frac{1}{2}$, and $\alpha_{ABH} = 0$ made in Figs. 12 and 13 entail the inclusion, in the corresponding 3D sections, of the full 2D domains Eq. (3.47), Fig. 2, and Eqs. (3.51) and (3.52), Fig. 3, and Eqs. (3.40) and (3.41), Fig. 1, respectively. These scans will thus allow one to judge whether the resolved conditions on the λ 's given by Fig. 8, or those given by Eq. (3.82) or by Eq. (3.73), in which the pairs of parameters (α_A, α_{AB}) , (α_A, α_{AH}) , and $(\alpha_{AH}, \alpha_{ABH})$ have been eliminated, respectively, are indeed necessary and sufficient or not. The answer will be yes for the first two and no for the last:

- One sees from Fig. 13(a) that for $\alpha_{ABH} \gtrsim -0.27$ there are no exclusions by the $\omega - \zeta$ chips. In particular, the 2D section at $\alpha_{ABH} = 0$ corresponds to the full α_A, α_{AB} domain of Fig. 2 which is thus not reduced by the constraint from the $\omega - \zeta$ chips. In fact, this result could easily be retrieved once noted that the α_A, α_{AB} domain of Fig. 2 corresponds indeed to the 2D section of the α -potatoid Eq. (3.61) at $\alpha_{AH} = \frac{1}{2}$, $\alpha_{ABH} = 0$. These values imply $\omega = 0$ [cf. Eq. (3.93)]; and as seen

from Fig. 11, all points $(\omega = 0, \zeta)$ remain within the $\omega - \zeta$ chips $\forall \zeta \in [\frac{1}{3}, 1]$. It follows that the study in Sec. III E 1 that leads to the NAS conditions given by Fig. 8 remains valid, at least for the $\alpha_{AH} = \frac{1}{2}$, $\alpha_{ABH} = 0$ section. Moreover, since the domain of Fig. 2 is not only a projection but corresponds as well to the latter section of the α potatoid, then the above-mentioned NAS conditions are *sufficient* conditions for all other sections at fixed $\alpha_{AH}, \alpha_{ABH}$ since by construction they all fall in the interior of the α_A, α_{AB} domain of Fig. 2. Obviously this holds even if these sections have portions excluded by the $\omega - \zeta$ chips, e.g., when $\alpha_{ABH} < -0.27$ as seen from Fig. 13(a), since sufficiency is more constraining. We can thus safely conclude that the conditions given by Fig. 8 are NAS for the validity of Eq. (3.32) in all the α potatoid.

- Along a similar line of thought, one deduces from Fig. 13(b), where there are no exclusions by the $\omega - \zeta$ chips as soon as $\alpha_{ABH} \gtrsim -0.06$, and from the fact that

FIG. 13. Two 3D sections of the 4D α potatoid: (a) $(\alpha_A, \alpha_{AB}, \alpha_{ABH})$ at $\alpha_{AH} = \frac{1}{2}$; (b) $(\alpha_A, \alpha_{AH}, \alpha_{ABH})$ at $\alpha_{AB} = 0$. The red and blue dots denote the regions excluded by the $\omega - \zeta$ chips. See text for further discussions.

the projected domain shown in Fig. 1 is also retrieved as a 2D section at $\alpha_{AB} = \alpha_{ABH} = 0$, that the conditions given by Eq. (3.82) remain NAS for the validity of Eq. (3.77) in all the α potatoid.

- The case of Eq. (3.73) is more involved. This condition resulted from eliminating $(\alpha_{AH}, \alpha_{ABH})$ based on the full domain of Fig. 3. However, as seen from Fig. 12(b), a portion of this domain in the $-\sqrt{2} \leq \alpha_{ABH} \leq 0$ range is excluded by the $\omega - \zeta$ chips constraint. Equation (3.73) becomes thus only sufficient for the domain of Fig. 3 that corresponds furthermore to the 2D section at $\alpha_A = 1$, $\alpha_{AB} = 0$ on Figs. 12(a)–12(d). It is thus also only sufficient for the full α potatoid, again because the domain of Fig. 3 is the largest section. Note that one can do better by resolving the NAS conditions for this largest section, taking into account the actual $\omega - \zeta$ chips constraint which is simply defined by a straight line joining the points $(\alpha_{AH} = 0, \alpha_{ABH} = 0)$ and $(\alpha_{AH} = 1, \alpha_{ABH} = -\sqrt{2})$; see Fig. 12(b). The resulting truncated domain will, however, cease to be the largest section so that the obtained conditions are now only *necessary* for an extended fraction of the α potatoid. As seen from Fig. 12(d), the maximal section taking into account the $\omega - \zeta$ chips constraint does exist somewhere inside the 3D domain but would be difficult to determine analytically.

We end this section with a comment concerning α_{ABH} : as argued at the end of Sec. III C 2 the sign of α_{ABH} is not expected to be correlated with the three other α 's. If a given point $(\alpha_A, \alpha_{AB}, \alpha_{AH}, \alpha_{ABH})$ lies in the true 4D α -parameters domain, i.e., not just in the α potatoid, then the point $(\alpha_A, \alpha_{AB}, \alpha_{AH}, -\alpha_{ABH})$ lies also in this domain. This is best seen from Eq. (3.49) which is the only one that depends on the B field (in the chosen gauge), and only

through an arbitrary global sign. However, Eqs. (3.93) are not symmetrical under $\alpha_{ABH} \rightarrow -\alpha_{ABH}$, and as discussed above and shown in Figs. 12 and 13 the $\omega - \zeta$ chips peel the α potatoid asymmetrically with respect to α_{ABH} . This is not a contradiction because the $\omega - \zeta$ chips constraint is only sufficient but not necessary to exclude points. But given the general symmetry with respect to the sign flip of α_{ABH} , it follows that *for any domain excluded by the $\omega - \zeta$ chips one should also exclude the domain corresponding to the replacement $\alpha_{ABH} \rightarrow -\alpha_{ABH}$.*

V. PUTTING EVERYTHING TOGETHER: A USER'S GUIDE

It is time to recapitulate the various results we arrived at and then provide a roadmap for an optimal exploitation:

- While studying the general precustodial potential we were lead automatically in Secs. III C 1 and III E 1 to constraints that involved only the A and B multiplets for which we provided the fully resolved NAS BFB conditions in analytical form; see Fig. 8 and Eqs. (3.67)–(3.71). As such they thus correspond to the NAS conditions for a reduced model having only two triplets. Nonetheless, they do provide robust necessary BFB conditions for the full precustodial potential since they correspond to the potential in the $H = 0$ field direction.
- In Secs. III C 2 and III E 2 we addressed the parts of the constraints that involve simultaneously the three sectors H , A , and B . The sign of λ_{BH} turned out to be critical, but again the BFB conditions that we obtained in a fully resolved analytical form correspond to field subsectors, namely H , B or H , A [cf. Eqs. (3.81) and (3.82)] and are thus *necessary* for the full model. It is noteworthy that Eq. (3.82)

TABLE I. A roadmap for the complete NAS BFB conditions for the precustodial model. Check/cross marks following an equation number indicate that the equation should be satisfied/violated. See text for a detailed description.

①			
Eqs. (3.67)–(3.71), Fig. 8 ✓		Eqs. (2.14)–(2.17)	
②			
ⓐ Eq. (3.73) ✓	$\lambda_{BH} \geq 0$	Eq. (3.81) ✓	Eqs. (3.79), (G30b)–(G30d) ✓, with Eqs. (3.75) and (3.76) and $\forall \alpha$ -params, ω , ζ Eqs. (3.61), (3.113), (F1)
	$\lambda_{BH} < 0$		
ⓑ Eq. (3.73) ✗	$\lambda_{BH} \geq 0$	...	$4\mathbf{a}_o\mathbf{c}_o - \mathbf{b}_o^2$ Eqs. (3.24), (3.38), (3.85) $\rightarrow a_i$, (H13) ✓
	$\lambda_{BH} < 0$	Eq. (3.81) ✓	$4\mathbf{a}_o\mathbf{c}_o - \mathbf{b}_o^2$ Eqs. (3.24), (3.38), (3.83) $\rightarrow a_i$, (H13) ✓ and $\forall \alpha$ -params, ω , ζ Eqs. (3.61), (3.113), (F1)
Ⓒ			
ⓐ Eq. (3.73) ✓	$\lambda_{BH} \geq 0$	Eq. (3.81) ✓	Same as in ②
	$\lambda_{BH} < 0$		
ⓑ Eq. (3.73) ✓	$\lambda_{BH} \geq 0$	...	$\mathbf{b}_o + 2\sqrt{\mathbf{a}_o\mathbf{c}_o} > 0$ Eq. (3.30) ✓
	$\lambda_{BH} < 0$	Eq. (3.81) ✓	$\forall t \in [0, +\infty)$, α -params, ω , ζ Eqs. (3.61), (3.113), (F1)

reproduces Eqs. (2.14)–(2.17) of the Type-II seesaw model¹⁵ that we had arrived at following a different path in Sec. II A, a significant cross-check. Moreover, from the flowchart of Fig. 10 and the properties of $\mathcal{B}_9^{(a,b)}$ one finds that the constraint Eq. (3.81) should be applied whenever $\lambda_{BH} < 0$, thus retrieving the fully resolved NAS BFB conditions for the SM extended by one real $SU(2)$ triplet.

- (iii) We give in Table I a roadmap for a user’s implementation of the constraints following two alternative roads each made of two steps. Step ① is common and corresponds to the fully resolved necessary constraints that are also NAS if restricted to the A , B or H , A sectors. Note that these constraints are already stricter than the ones given in [34] under the assumption of two nonvanishing complex fields at once or the ones extended to the “custodial” direction in [35], as they are NAS in *all directions* within A , B or H , A . Also specifying to the Georgi-Machacek case we do retrieve the conditions found in [33]. Steps ② and Ⓒ are two technically different but theoretically equivalent ways to complete the NAS conditions. Note first that in both cases branches ⓐ and ⓑ approximate Eq. (3.73) as being necessary for the positivity of $\mathbf{b}_o$ (on top of it being sufficient). Despite the issue discussed in Sec. IV, this approximation is valid for all practical purposes, which we checked numerically by scanning over several tens of thousands of points in the α -parameters space and verified that Eqs. (3.72) and (3.73) delineated indeed the same

$(\lambda_{AH}^{(1,2)}, \lambda_{ABH})$ -space regions.¹⁶ Then the ⓐ branches with $\lambda_{BH} \geq 0$ are complete and provide fully resolved NAS BFB conditions. When $\lambda_{BH} < 0$, both ⓐ and ⓑ lead to the same fully resolved extra constraint involving the B , H sector, plus different sets of partially resolved constraints: In step ② as well as in step Ⓒ-ⓐ with $\lambda_{BH} < 0$, the latter constraints are resolved only with respect to the T and t parameters but still need a scan over the α parameters (including optionally the refinements of Sec. IV). In contrast, Ⓒ-ⓑ is resolved only with respect to T , and a supplementary scan is still required on t . Note also the different Boolean meanings in the last columns of ⓐ-ⓑ and Ⓒ-ⓑ. In the former one needs to find *at least one* set of values (u, v, c) satisfying a set of inequalities while the latter requires *all* values of t to satisfy one inequality.

We give in Fig. 14 an illustration of allowed λ domains following road ①-Ⓒ. A typical expectation is that the constraints are more stringent for negative values of the couplings associated with the positive definite operators that are present in the potential. This is indeed seen in Figs. 14(a), 14(c), and 14(d). In contrast, Fig. 14(b) shows that λ_{ABH} can be in equally sized negative or positive regions since this coupling corresponds to the only operators that are not positive definite [cf. Eq. (3.29)].

Let us close this section with an outlook on some issues related to the subject of the present paper but lying beyond its scope:

¹⁵With the correspondence $\Delta = A$, $\lambda = \lambda_H$, $\lambda_2 = \lambda_A^{(1)}/4$, $\lambda_3 = \lambda_A^{(2)}/4$, $\lambda_1 = \lambda_{AH}^{(1)}$, and $\lambda_4 = \lambda_{AH}^{(2)}$.

¹⁶This should not come as a surprise since the further refinement discussed in Sec. IV concerns only boundaries of the α potatoid that would require much finer scans as shown in Figs. 12 and 13.

FIG. 14. Necessary and sufficient 3D λ domains that ensure BFB of the precustodial potential Eq. (3.4) illustrated for fixed $\lambda_B = \lambda_H = \lambda_A^{(2)} = \lambda_{AB}^{(1)} = \lambda_{AB}^{(2)} = 1$, (a) the $(\lambda_A^{(1)}, \lambda_{AH}^{(1)}, \lambda_{AH}^{(2)})$ domain with $\lambda_{BH} = -\frac{1}{10}$, $\lambda_{ABH} = 1$; (b) the $(\lambda_{AH}^{(1)}, \lambda_{AH}^{(2)}, \lambda_{ABH})$ domain with $\lambda_{BH} = -\frac{1}{10}$, $\lambda_A^{(1)} = 1$; (c) the $(\lambda_{AH}^{(1)}, \lambda_{AH}^{(2)}, \lambda_{BH})$ domain with $\lambda_A^{(1)} = \lambda_{ABH} = 1$; and (d) the $(\lambda_A^{(1)}, \lambda_{AH}^{(1)}, \lambda_{BH})$ domain with $\lambda_A^{(2)} = \lambda_{ABH} = 1$.

A. Perturbative unitarity constraints

They typically bind the absolute magnitudes of the λ couplings and some of their combinations from above. These constraints should eventually be studied for the general precustodial model (see, however, [35]) and be combined with the NAS BFB conditions derived in this paper. Here we just note an interesting tension that might arise from such a combination, due to the form of conditions $\mathcal{B}_3$, $\mathcal{B}_5$, $\mathcal{B}_7$. The relatively large numerical factors appearing in these inequalities [see Eqs. (3.67), (3.69), (3.71)] can easily force $|\lambda_A^{(i=1,2)}|$ or λ_B to be (much) larger than one even for $|\lambda_{AB}^{(i=1,2)}|$, $|\lambda_H| \lesssim 1$. At least one among the conditions $\mathcal{B}_3$, $\mathcal{B}_5$, and $\mathcal{B}_7$ is active in cases (ii), (iii), or (iv) of the flowchart of Fig. 8. We illustrate a few

such configurations in Fig. 15. The domains shown in the figure are necessary but not sufficient; they can be reduced further when adding the rest of the NAS BFB conditions. Note that such a potential tension disappears in the limit of decoupling between the two triples ($\lambda_{AB}^{(i=1,2)} \rightarrow 0$) in accordance with the unitarity/BFB conditions found in [31].

B. Quantum corrections

They affect the tree-level constraints in various ways: they modify the form of the constraints, introduce a notion of scale at which they should be satisfied, and provide criteria for the validity of perturbativity, as treated, for instance, in [54], [35,55]. However, it is not often appreciated that combining perturbative unitarity and stability

FIG. 15. Trend of the $(\lambda_A^{(1)}, \lambda_A^{(2)})$ necessary domains in yellow, as dictated by the necessary BFB conditions of Fig. 8 and Eqs (3.67)–(3.71). The allowed domain lies to the right of each line, illustrated for $\lambda_B = 1$, $\lambda_H = \frac{1}{2}$ and (a) $\lambda_{AB}^{(1)} = -\frac{1}{5}$, $\lambda_{AB}^{(2)} = \frac{1}{2}$, (b) $\lambda_{AB}^{(1)} = \frac{1}{5}$, $\lambda_{AB}^{(2)} = -\frac{1}{2}$, (c) $\lambda_{AB}^{(1)} = -\frac{1}{5}$, $\lambda_{AB}^{(2)} = -\frac{3}{10}$, and (d) $\lambda_B = \frac{1}{2}$, $\lambda_H = \frac{1}{2}$, $\lambda_{AB}^{(1)} = -\frac{1}{5}$, $\lambda_{AB}^{(2)} = -\frac{3}{10}$.

requirements beyond the tree level needs some further care because the physical meaning of the running couplings becomes different in these two classes of constraints. Since unitarity is related to scattering processes, the proper objects are Green's functions. The scale appearing in the running couplings (and masses) of the renormalization group improved Green's functions encodes the way the scattering amplitudes scale with energy. In contrast, stability issues are expressed in terms of the renormalization group improved effective potential where now the scale on which depend the running couplings, masses, and fields is, in fact, a combination of the fields themselves and encode the modification of the shape of the potential (see, for instance, [56,57]¹⁷). It thus appears that, in so far as replacing the tree-level couplings by their runnings in the tree-level conditions is a good approximation, the potential stability conditions need not be required *at all* “scales,” from the electroweak scale all the way up to some very high cutoff Λ (e.g., M_{GUT} or M_{Planck}) as often done in the literature [17,32,34], but only at that scale Λ which represents the largest value of the fields. Barring Landau poles, there is indeed no physical reason to require the improved quartic part of the potential to remain positive for intermediate values of the fields. (Obviously this is at

¹⁷Where it was also stressed that even an additive constant becomes field dependent beyond tree level.

variance with the unitarity constraints that should be satisfied already at the energy scale of a given scattering experiment.) Furthermore, a long-standing issue is how to improve the effective potential in the presence of several scalar fields (see [58] for a recent reappraisal, and references therein). As concerns the NAS BFB conditions of Table I, they can be used beyond the tree level in two different ways: (i) The quartic part, $V_{\text{p-c}}^{(4)}$, Eq. (3.4), of the precustodial potential has the same form as the general counterterms needed to renormalize the Georgi-Machacek model accounting for a deviation from the tree-level correlations Eq. (3.12) due to the custodial symmetry breaking loop effect of the $U(1)_Y$ gauge couplings [34,48]. One is thus guaranteed that the ten λ couplings of $V_{\text{p-c}}^{(4)}$ will absorb the one-loop corrections of the Georgi-Machacek effective potential up to field dependent factors of the form $\log(\mathcal{M}(\phi_i)^2/Q^2) - c$, where $\mathcal{M}$ is typically a binomial function of the fields, Q is some renormalization scale, and c is a renormalization scheme dependent constant. It follows that satisfying the conditions of Table I on the λ 's that absorb the one-loop induced quartic couplings will also guarantee the stability of the full one-loop Georgi-Machacek effective potential at large field values with $\mathcal{M}(\phi_i)^2 \gg Q^2$. (ii) Table I can also obviously be used as a seed for the loop corrected stability conditions of the precustodial model itself, relying on whatever renormalization group improvement approaches are quoted above. The main difference with (i) will reside essentially in the renormalization conditions not enforcing the custodial symmetry of the potential at a given scale.

C. Comparison with other methods

As already recalled in the Introduction, several methods have been considered to address the boundedness-from-below conditions. Depending on the extended scalar sectors, the symmetries of the potential and the multiplet representations, the different methods can have varying practical applicability. An important case is when discrete symmetries are not imposed, thus allowing for odd powers of the scalar fields to occur in four-dimensional gauge invariant operators of the (most general) renormalizable scalar potential. This was considered for the two-Higgs-doublet model [36], for extensions with two real scalar singlets [42], or for the most general potential in left-right symmetric models [59]. It is also the case for the precustodial model with $\lambda_{ABH} \neq 0$ studied in this paper. The copositivity method ceases to be efficient in this case since one cannot write the potential in a bilinear form with vectors of dimension three or more having positive definite independent components. (The fact that such a two-dimensional form is still possible is not helpful as one hides the complexity of the conditions in the dependence on angles). In contrast, our approach remains applicable albeit with an extended set of the α parameters. It would provide further

insight into the all-directions NAS BFB conditions, for example in the study of left-right symmetric models, unlike in [60] which relied on [41,59] where specific directions were considered. A distinctive feature in this case is the appearance of positivity conditions for full quartic polynomials as was found for the model with two singlets discussed in [42] and in the present study with two triplets.

VI. CONCLUSION

We carried out in this paper a comprehensive study of tree-level necessary and sufficient conditions for a bounded from below potential in extensions of the SM with one or two $SU(2)_L$ triplet scalar fields. We derived for the first time the complete set of such conditions in the case of the general precustodial model having one complex and one real triplets. A fully resolved analytical form involving only the couplings was obtained for parts of these conditions. This could be achieved thanks to a parametrization of the 13-dimensional field space reducing the degrees of freedom to a small set of relevant gauge invariant variables. We determined precisely the compact domains in which most of these variables live, thus allowing a well-defined procedure for the other parts of the conditions that remained in a partially resolved form. It would be interesting to see how the more general methods quoted in the Introduction would perform in the presence of triplets. In particular, the fully resolved form we found in the purely two triplets sector may lend itself to a generalization to multiple fields. In the course of the study we were led to review some of the known results for the Type-II seesaw and Georgi-Machacek models providing complete proofs that were missing in the literature for key properties. The latter were important to settle on a firm basis in relation with an unexpected feedback of the Georgi-Machacek reduced variables on those of the precustodial model. Furthermore, we demonstrated the existence of simplified criteria for the positivity of a general quartic polynomial that can be used for any model with a renormalizable potential. The precustodial BFB conditions on the couplings have to be fulfilled for any consistent tree-level phenomenological analysis of the model. They find also their motivation as a pattern for the one-loop BFB conditions in the Georgi-Machacek model.

ACKNOWLEDGMENTS

We thank Abdesslam Arhrib for his fruitful collaboration at an early stage of this work, and Michele Frigerio as well as Michel Talon for profitable discussions. We also acknowledge an inspiring exchange with Kaladi Babu.

APPENDIX A: PROOF OF THE PROPERTIES OF ξ AND ζ

In the following we give the proof of Eq. (2.10), and then establish Eq. (2.11) and the ensuing correlations.

1. $0 \leq \xi \leq 1$

First we note that Δ being traceless implies the identity

$$\Delta\Delta^\dagger + \Delta^\dagger\Delta = \mathbf{1} \times \text{Tr}\Delta\Delta^\dagger \quad (\text{A1})$$

[see also Eq. (C2)], from which follows immediately

$$H^\dagger\Delta\Delta^\dagger H + H^\dagger\Delta^\dagger\Delta H = H^\dagger H \text{Tr}\Delta\Delta^\dagger. \quad (\text{A2})$$

Since $H^\dagger\Delta^\dagger\Delta H$ is positive definite, one then has

$$H^\dagger H \text{Tr}\Delta\Delta^\dagger - H^\dagger\Delta\Delta^\dagger H \geq 0, \quad (\text{A3})$$

and thus

$$\xi \equiv \frac{H^\dagger\Delta\Delta^\dagger H}{H^\dagger H \text{Tr}\Delta\Delta^\dagger} \leq 1. \quad (\text{A4})$$

Furthermore ξ is trivially greater than zero since it is the ratio of two positive definite quantities. Finally the two values 0 and 1 are effectively reached, respectively, when $H^\dagger\Delta = 0$ and $\Delta H = 0$, which is always possible for some given configurations of the H and Δ field components provided that $\text{Det}\Delta = 0$ when $H \neq 0$. Thus

$$0 \leq \xi \leq 1. \quad (\text{A5})$$

2. $\frac{1}{2} \leq \zeta \leq 1$

$\Delta\Delta^\dagger$ being a 2×2 matrix one has

$$\frac{1}{2}(\text{Tr}\Delta\Delta^\dagger)^2 - \frac{1}{2}\text{Tr}(\Delta\Delta^\dagger)^2 = \text{Det}\Delta\Delta^\dagger. \quad (\text{A6})$$

Then, using $\text{Det}\Delta\Delta^\dagger \equiv |\text{Det}\Delta|^2 \geq 0$ implies straightforwardly from Eq. (A6) that

$$\zeta \equiv \frac{\text{Tr}(\Delta\Delta^\dagger)^2}{(\text{Tr}\Delta\Delta^\dagger)^2} \leq 1. \quad (\text{A7})$$

Note that the value 1 is indeed reached when $\Delta\Delta^\dagger$ has one zero and one nonzero eigenvalues, which is always possible to find for some configurations of the Δ field components.

Also, we trivially have $\zeta \geq 0$ since it is the ratio of two positive definite quantities. However, the value 0 cannot be trivially reached, since if the numerator of ζ vanishes, then the denominator should vanish as well. In fact, ζ cannot go below $1/2$. To see this we rewrite ζ in terms of M_1^2 and M_2^2 the two (real and positive) eigenvalues of $\Delta\Delta^\dagger$,

$$\zeta = \frac{M_1^4 + M_2^4}{(M_1^2 + M_2^2)^2}. \quad (\text{A8})$$

It is now easy to study the function $\zeta(x) = (1+x^2)/(1+x)^2$ where $x \equiv M_1^2/M_2^2 \geq 0$ to show that it has a minimum of

$\zeta = 1/2$ at $x = 1$, that is, when $\Delta\Delta^\dagger$ has degenerate eigenvalues. One also retrieves the fact that $\zeta(x) \leq 1$ and reaches 1 for $x \rightarrow 0$ or $x \rightarrow \infty$. Thus

$$\frac{1}{2} \leq \zeta \leq 1. \quad (\text{A9})$$

3. Correlation between ξ and ζ

Since from Eqs. (2.6) and (2.7) ζ depends solely on Δ while ξ depends on both H and Δ , one could be tempted to assume that ζ and ξ can reach independently their extrema given by Eqs. (A5) and (A9), by varying independently H and Δ . This is, however, not true as one can see easily from the fact that ξ reaches its two extrema under the generic condition $\text{Det}\Delta = 0$ as discussed above Eq. (A5). But then $\text{Det}\Delta = 0$ together with Eq. (A6) imply necessarily $\zeta = 1$ so that $\zeta = \frac{1}{2}$ can never be reached when ξ takes its extremal values 0 or 1.

We use now the invariance under the general gauge transformation $H \rightarrow \mathcal{U}(x)H$, $\Delta \rightarrow \mathcal{U}(x)\Delta\mathcal{U}^\dagger(x)$, where $\mathcal{U}(x)$ denotes any element of $SU(2)_L \times U(1)_Y$, of the potential Eq. (2.1) and of the parameters defined in Eqs. (2.3)–(2.7). Since $\mathcal{U}(x)$ is unitary and $\Delta\Delta^\dagger$ Hermitian, we can always find, for any given field configuration Δ , a gauge transformation $\mathcal{U}_\Delta(x)$ that diagonalizes $\Delta\Delta^\dagger$. Then ζ takes the form given in Eq. (A8) and ξ reads

$$\xi = \frac{(M_2^2|\tilde{\phi}^0|^2 + M_1^2|\tilde{\phi}^+|^2)}{(M_1^2 + M_2^2)(|\tilde{\phi}^0|^2 + |\tilde{\phi}^+|^2)}, \quad (\text{A10})$$

where the tilde denotes the components of the transformed doublet $\tilde{H} = \mathcal{U}_\Delta(x)H$. It is then natural to define

$$c_\Delta^2 \equiv \frac{M_1^2}{M_1^2 + M_2^2}, \quad s_\Delta^2 \equiv 1 - c_\Delta^2, \quad (\text{A11})$$

$$c_H^2 \equiv \frac{|\tilde{\phi}^+|^2}{|\tilde{\phi}^0|^2 + |\tilde{\phi}^+|^2}, \quad s_H^2 \equiv 1 - c_H^2, \quad (\text{A12})$$

with their obvious range of variation $c_\Delta^2, c_H^2 \in [0, 1]$. Equations (2.12) and (2.13) follow then straightforwardly from Eqs. (A8), (A10)–(A12):

$$\xi = c_\Delta^2 c_H^2 + s_\Delta^2 s_H^2 = \frac{1}{2}(1 + c_{2H} c_{2\Delta}), \quad (\text{A13})$$

$$\zeta = c_\Delta^4 + s_\Delta^4 = \frac{1}{2}(1 + c_{2\Delta}^2), \quad (\text{A14})$$

where we have defined $c_{2H} = c_H^2 - s_H^2$, $c_{2\Delta} = c_\Delta^2 - s_\Delta^2$. It is crucial that these cosines vary independently from each other in their allowed domains $c_{2H} \in [-1, 1]$, $c_{2\Delta} \in [-1, 1]$. That they indeed scan independently all

their allowed domain is obvious from the definitions Eqs. (A11) and (A12) and the fact that $\mathcal{U}_\Delta(x)$ is invertible: Indeed, one can always choose the magnitudes of $M_1^2, M_2^2, |\tilde{\phi}^0|, |\tilde{\phi}^+|$ to reach any value of $c_H^2, c_\Delta^2 \in [0, 1]$; this will correspond to the domain of all field configurations obtained by gauge transforming $\tilde{H} \equiv (\tilde{\phi}^+, \tilde{\phi}^0)^T$ and $\tilde{\Delta} \equiv \text{diag}(e^{i\theta_1} M_1, e^{i\theta_2} M_2)$ with an arbitrary $\mathcal{U} \equiv \mathcal{U}_\Delta^{-1}$.

Eliminating $c_{2\Delta}^2$ in Eqs. (A13) and (A14) one finds

$$2\xi^2 - 2\xi + 1 + \frac{(c_{2H}^2 - 1)}{2} = c_{2H}^2 \zeta. \quad (\text{A15})$$

This allows one to determine the lower envelope in the ξ, ζ plane, i.e., when saturating the inequality in Eq. (2.11) as discussed in [32]. We will, however, rely directly on Eqs. (A13) and (A14) when determining the BFB conditions in the next section.

APPENDIX B: THE BFB CONDITIONS FOR THE TYPE-II SEESAW MODEL

1. The new necessary and sufficient BFB conditions

We give here a detailed proof of the NAS BFB conditions Eqs. (2.14) and (2.17). The condition $\lambda_2 + \zeta\lambda_3 \geq 0$ of Eq. (2.9) depends only on ζ so that the correlations given by Eqs. (A13) and (A14) are not relevant here. It is thus equivalent to replacing ζ by its two extreme values due to the monotonic dependence on ζ . Thus the first two conditions of Eq. (2.9) become

$$\lambda > 0 \wedge \lambda_2 + \lambda_3 \geq 0 \wedge \lambda_2 + \frac{\lambda_3}{2} \geq 0 \quad (\text{B1})$$

as was initially found in [31]. As for $\lambda_1 + \xi\lambda_4 + \sqrt{\lambda(\lambda_2 + \zeta\lambda_3)} > 0$ of Eq. (2.9), we first rewrite it in terms of c_{2H} and $c_{2\Delta}$ according to Eqs. (A13) and (A14), as

$$F(c_{2\Delta}, c_{2H}) > 0, \quad \forall c_{2\Delta}, c_{2H} \in [-1, 1], \quad (\text{B2})$$

where we defined

$$F(c_{2\Delta}, c_{2H}) \equiv \lambda_1 + (1 + c_{2H} c_{2\Delta}) \frac{\lambda_4}{2} + \sqrt{\lambda \left(\lambda_2 + (1 + c_{2\Delta}^2) \frac{\lambda_3}{2} \right)}. \quad (\text{B3})$$

However, since c_{2H} and $c_{2\Delta}$ are mutually independent, the monotonic dependence on c_{2H} in $F(c_{2\Delta}, c_{2H})$ again allows one to replace the above positivity condition equivalently by two positivity conditions corresponding to the two extreme values $c_{2H} = \pm 1$. We will thus replace once and for all the condition $\lambda_1 + \xi\lambda_4 + \sqrt{\lambda(\lambda_2 + \zeta\lambda_3)} > 0$ by

$$F(c_{2\Delta}, +) > 0 \quad \text{and} \quad F(c_{2\Delta}, -) > 0, \quad \forall c_{2\Delta} \in [-1, 1], \quad (\text{B4})$$

where we use the shorthand notation $F(c_{2\Delta}, \pm) \equiv F(c_{2\Delta}, \pm 1)$.

At this point a careful study is needed, as the dependence on $c_{2\Delta}$ is not trivially monotonic so that *a priori* one does not necessarily have

$$\text{Eq. (B4)} \Leftrightarrow \{F(+1, \pm) > 0 \wedge F(-1, \pm) > 0\}. \quad (\text{B5})$$

It is nonetheless noteworthy that this equivalence does hold in half of the parameter space region where $\lambda_3 < 0$ despite the nonmonotonicity of F in $c_{2\Delta}$, as we will see in a moment. Irrespective of the sign of λ_3 the first and second derivatives of F read

$$F'(c_{2\Delta}, \pm) = \frac{1}{2} \left(\frac{c_{2\Delta} \lambda \lambda_3}{\sqrt{\lambda(\lambda_2 + \frac{1}{2}(1 + c_{2\Delta}^2)\lambda_3)}} \pm \lambda_4 \right), \quad (\text{B6})$$

$$F''(c_{2\Delta}, \pm) = \frac{\lambda_3(2\lambda_2 + \lambda_3)}{(2\lambda_2 + \lambda_3(1 + c_{2\Delta}^2))^{3/2}}. \quad (\text{B7})$$

In the sequel we will assume without further reference the conditions given in Eq. (B1). It then immediately follows from Eq. (B7) that $F'' < 0$, $\forall c_{2\Delta} \in [-1, 1]$, whenever $\lambda_3 < 0$. This implies that if F admits an extremum, it will be necessarily a *maximum* so that Eq. (B5) is valid, since in this case the value of F at one of the two boundaries of $[-1, 1]$ is necessarily the smallest value it can take. On the other hand, if F does not admit an extremum, then Eq. (B5) is obviously valid as well, and one retrieves Eqs. (B15) and (B16).

We thus conclude that the BFB conditions Eqs. (B15) and (B16) initially found in [31] are necessary and sufficient, and thus complete, when $\lambda_3 < 0$.

The situation is quite different when $\lambda_3 > 0$. In this case F'' is non-negative over the full domain $[-1, 1]$. Thus if $F(c_{2\Delta}, \pm)$ admit extrema in the domain, they will be necessarily minima. On the other hand, $F'(c_{2\Delta}, +)$ and $F'(c_{2\Delta}, -)$ cannot vanish simultaneously (except in the special cases where $\lambda_4 = 0$ or $2\lambda_2 + \lambda_3 = 0$), but rather at two opposite values of $c_{2\Delta}$, as can be seen from Eq. (B6). This occurs for

$$c_{2\Delta}^{(\pm)} = \pm |\lambda_4| \sqrt{\frac{(2\lambda_2 + \lambda_3)}{\lambda_3(2\lambda\lambda_3 - \lambda_4^2)}} \quad (\text{B8})$$

with the consistency condition $0 \leq (c_{2\Delta}^{(\pm)})^2 \leq 1$. The latter condition reads

$$2\lambda\lambda_3 - \lambda_4^2 > 0 \wedge \lambda_4^2(2\lambda_2 + \lambda_3) \leq \lambda_3(2\lambda\lambda_3 - \lambda_4^2). \quad (\text{B9})$$

Note that the second of these two inequalities always implies the first due to the case assumption $\lambda_3 > 0$ and the validity of Eq. (B1). Moreover, this second inequality can be rewritten equivalently as

$$\sqrt{\lambda}\lambda_3 \geq \sqrt{(\lambda_2 + \lambda_3)\lambda_4^2}, \quad (\text{B10})$$

where we again relied on the case assumption $\lambda_3 > 0$. Thus Eq. (B10) is *necessary and sufficient for the existence of minima within the domain* $[-1, 1]$. In this case one of the two functions $F(c_{2\Delta}, +)$ and $F(c_{2\Delta}, -)$ will have a minimum at $c_{2\Delta}^{(+)}$ and the other at $c_{2\Delta}^{(-)}$. Moreover, the values of the two F functions at these minima turn out to be the *same*, given by

$$F_{\min} = \lambda_1 + \frac{\lambda_4}{2} + \sqrt{\lambda \left(\lambda_2 + \frac{\lambda_3}{2} \right) \left(1 - \frac{\lambda_4^2}{2\lambda\lambda_3} \right)}. \quad (\text{B11})$$

[To determine $F_{\min}$ some care should be taken by considering the sign of λ_4 and noting that the $\pm$ in Eq. (B8) refer neither to the sign of λ_4 nor to the two functions F .] In fact, the uniqueness of $F_{\min}$ is a direct consequence of the symmetry property $F(c_{2\Delta}, -c_{2H}) = F(-c_{2\Delta}, c_{2H})$; cf. Eq. (B3). This symmetry is also responsible for the fact that $c_{2\Delta}^{(+)}$ and $c_{2\Delta}^{(-)}$ are the opposite of each other so that when Eq. (B10) is satisfied they both remain in the domain $[-1, 1]$.

It follows that even though the two functions $F(c_{2\Delta}, +)$ and $F(c_{2\Delta}, -)$ do not reach their minimum for the same value of $c_{2\Delta}$, requiring

$$F_{\min} > 0 \quad (\text{B12})$$

when Eq. (B10) is satisfied will be equivalent to Eq. (B4). Note in particular that Eq. (B12) should imply $F(\pm 1, \pm) > 0$ and $F(\mp 1, \pm) > 0$, that is,

$$\begin{aligned} \lambda_1 + \frac{\lambda_4}{2} + \sqrt{\lambda \left(\lambda_2 + \frac{\lambda_3}{2} \right) \left(1 - \frac{\lambda_4^2}{2\lambda\lambda_3} \right)} &> 0 \Rightarrow \lambda_1 \\ &+ \sqrt{\lambda(\lambda_2 + \lambda_3)} > 0 \wedge \lambda_1 + \lambda_4 + \sqrt{\lambda(\lambda_2 + \lambda_3)} > 0, \end{aligned} \quad (\text{B13})$$

which is indeed the case.¹⁸ Finally, when Eq. (B10) is not satisfied, but still $\lambda_3 > 0$, then either $c_{2\Delta}^{(\pm)}$ are not real valued or they lie outside of the $[-1, 1]$ domain. In both cases the two functions $F(c_{2\Delta}, \pm)$ are monotonic on $[-1, 1]$ and Eq. (B5) applies, which is similar to the previously

¹⁸This is due to the inequality $\sqrt{\lambda(\lambda_2 + \lambda_3)} - \sqrt{\lambda(\lambda_2 + \frac{\lambda_3}{2})(1 - \frac{\lambda_4^2}{2\lambda\lambda_3})} > \pm \frac{\lambda_4}{2}$ being valid whenever $\lambda_3 > 0$ and Eq. (B1) are valid, and thus consistently also $2\lambda\lambda_3 - \lambda_4^2 > 0$.

discussed case of $\lambda_3 < 0$. Putting everything together one can summarize the conditions that are equivalent to $\lambda_1 + \xi\lambda_4 + \sqrt{\lambda(\lambda_2 + \xi\lambda_3)} > 0$ [or Eq. (B4)], as follows:

- (i) if $\sqrt{\lambda\lambda_3} < \sqrt{(\lambda_2 + \lambda_3)\lambda_4^2}$, then $F(\pm 1, \pm) > 0$ and $F(\mp 1, \pm) > 0$.
- (ii) if $\sqrt{\lambda\lambda_3} \geq \sqrt{(\lambda_2 + \lambda_3)\lambda_4^2}$, then $F_{\min} > 0$.

Adding Eq. (B1) to these conditions, we obtain the Boolean form of the necessary and sufficient BFB conditions as given by Eqs. (2.14) and (2.17).

2. The old conditions

We recall here for further reference the sufficient and almost necessary BFB conditions [31]:

$$\lambda > 0 \wedge \lambda_2 + \lambda_3 > 0 \wedge \lambda_2 + \frac{\lambda_3}{2} > 0, \quad (\text{B14})$$

$$\wedge \lambda_1 + \sqrt{\lambda(\lambda_2 + \lambda_3)} > 0 \wedge \lambda_1 + \sqrt{\lambda\left(\lambda_2 + \frac{\lambda_3}{2}\right)} > 0, \quad (\text{B15})$$

$$\wedge \lambda_1 + \lambda_4 + \sqrt{\lambda(\lambda_2 + \lambda_3)} > 0 \wedge \lambda_1 + \lambda_4 + \sqrt{\lambda\left(\lambda_2 + \frac{\lambda_3}{2}\right)} > 0. \quad (\text{B16})$$

APPENDIX C: THE PRECUSTODIAL POTENTIAL

We give hereafter some elements that can help define a systematic procedure to construct the precustodial potential Eqs. (3.3) and (3.4) from a minimal set of independent operators. They can be useful as well for the construction of extended models with several $SU(2)$ triplet and singlet fields.

Note first the following two general identities, valid for any 2×2 matrices M and N :

$$M + \sigma^2 M^\top \sigma^2 = \mathbf{1} \text{Tr} M, \quad (\text{C1})$$

$$MN + NM = \mathbf{1}(\text{Tr} MN - \text{Tr} M \text{Tr} N) + M \text{Tr} N + N \text{Tr} M. \quad (\text{C2})$$

The fundamental representation of $SU(2)$ is pseudoreal. In particular, any of its elements $\mathcal{U}$ satisfy

$$\sigma^2 \mathcal{U} \sigma^2 = \mathcal{U}^*, \quad (\text{C3})$$

where σ^2 denotes the second Pauli matrix. From this and Eq. (3.5) it follows that

$$\sigma^2 H^* \sim H, \quad (\text{C4})$$

$$H^\top \sigma^2 \sim H^\dagger, \quad (\text{C5})$$

$$\sigma^2 A^\top \sigma^2 \sim \sigma^2 A^* \sigma^2 \sim A \sim A^\dagger, \quad (\text{C6})$$

$$\sigma^2 B^\top \sigma^2 \sim B, \quad (\text{C7})$$

where the symbol $\sim$ stands for “... transforms like... under $SU(2)$.” To systematize further the discussion it is useful to define the 2×2 matrices

$$\mathbb{H}_0 = HH^\dagger, \quad (\text{C8})$$

$$\mathbb{H}_2 = HH^\top \sigma^2. \quad (\text{C9})$$

$\mathbb{H}_0$ is Hermitian and transforms as B under $SU(2)_L \times U(1)_Y$ but has a nonvanishing trace, while $\mathbb{H}_2$ is traceless and transforms as A under $SU(2)_L \times U(1)_Y$. From the tracelessness of $\mathbb{H}_2, A$, and B , Eq. (C1) implies

$$\begin{aligned} \sigma^2 \mathbb{H}_2^\top \sigma^2 &= -\mathbb{H}_2, & \sigma^2 \mathbb{H}_2^* \sigma^2 &= -\mathbb{H}_2^\dagger, & \sigma^2 A^\top \sigma^2 &= -A, \\ \sigma^2 A^* \sigma^2 &= -A^\dagger, & \sigma^2 B^\top \sigma^2 &= -B, \end{aligned} \quad (\text{C10})$$

thus trivializing Eqs. (C6) and (C7).

Similarly, Eq. (C2) leads to

$$(\mathbb{H}_2)^2 = \frac{1}{2} \mathbf{1} \text{Tr}(\mathbb{H}_2)^2, \quad A^2 = \frac{1}{2} \mathbf{1} \text{Tr} A^2, \quad B^2 = \frac{1}{2} \mathbf{1} \text{Tr} B^2. \quad (\text{C11})$$

For the sake of conciseness we do not write here other useful relations resulting from Eqs. (C1) and (C2), involving $\mathbb{H}_0$ or products involving A, B [generalizing Eq. (A1)]. We have now all the ingredients to show that the $SU(2)_L \times U(1)_Y$ invariant operators in Eqs. (3.3) and (3.4) form a complete and independent set: Any such operator is necessarily either in the form of a trace of a 2×2 matrix operator that is neutral under $U(1)_Y$ and constructed from a product of fields that transform similarly under $SU(2)_L$ or in the form of a product of such traces that are separately either neutral or charged under $U(1)_Y$. (Recall that the other invariant quantity, the determinant, is always expressible in terms of traces.) We sketch hereafter the main steps with some examples.

dim-2: the list of all $U(1)_Y$ neutral operators is $\mathbb{H}_0, B^2, AA^\dagger, A^\dagger A$; recall that $\text{Tr} \mathbb{H}_0 = H^\dagger H$.

dim-3: the exhaustive list of representative $U(1)_Y$ neutral operators is $B^3, \mathbb{H}_0 B, \mathbb{H}_2 A^\dagger, AA^\dagger B$. Only the first one drops out after tracing, since $\text{Tr} B^3 = 0$ as an immediate consequence of Eq. (C11) and $\text{Tr} B = 0$. All other neutral dim-3 operators obtained from the above list by arbitrary permutations of the fields or by substituting a field by its transpose or complex conjugate are, upon tracing, related to this list. This is obtained by successive use of Eqs. (C10) and (C11) and the like. For example, $\text{Tr} AA^\dagger B^*$ is forbidden since B^* does not transform as A , while $\text{Tr} A^* A^\top B^*$ is allowed but redundant: $\text{Tr} A^* A^\top B^* = (\text{Tr} AA^\dagger B)^* = -(\text{Tr} A^\dagger AB)^* = -\text{Tr} A^\top A^* B^* = +\text{Tr} A^\top A^* \sigma^2 B \sigma^2 = +\text{Tr} AA^\dagger B$, where

we used Eq. (C2) and the tracelessness of A, B for the second equality, and Eq. (C10) for the last two equalities.

dim-4: the exhaustive list of representative $U(1)_Y$ neutral operators is $\mathbb{H}_0\mathbb{H}_0, \mathbb{H}_2\mathbb{H}_2^\dagger, \mathbb{H}_0B^2, \mathbb{H}_0AA^\dagger, \mathbb{H}_2A^\dagger B, B^4, ABA^\dagger B, AA^\dagger AA^\dagger$. Note that products of two dim-2 traced operators should also be added. Thus a systematic strategy would be to reduce in the above list the traces of the product of four matrices to products of two traces, whenever possible. This is done using the same tricks as illustrated for dim-3. For example, $\text{Tr}B^4 = \text{Tr}(\frac{1}{2}\mathbf{1}\text{Tr}B^2)B^2 = \frac{1}{2}(\text{Tr}B^2)^2$ as a consequence of Eq. (C11), or $\text{Tr}ABA^\dagger B = -\text{Tr}AA^\dagger B^2 + \text{Tr}A(\text{Tr}A^\dagger B)B = -\frac{1}{2}(\text{Tr}AA^\dagger)(\text{Tr}B^2) + (\text{Tr}AB)(\text{Tr}A^\dagger B)$. Note that $\text{Tr}AA^\dagger AA^\dagger$ can be transformed similarly but we chose not to do so in Eq. (3.4) so as to keep close to the notations in the literature. Finally, it is immediate from the list above that there exists only one operator containing σ^2 up to complex conjugation, $\text{Tr}\mathbb{H}_2A^\dagger B$.

APPENDIX D: PROOFS OF PROPERTIES OF THE α PARAMETERS

1. $\alpha_{AB} \in [0, 1]$

Thanks to gauge invariance and to the fact that B is self-adjoint one can always find, for each given value of α_{AB} , an $SU(2)_L$ transformation $\mathcal{U}_L$ that diagonalizes B leading to

$$\alpha_{AB} \equiv \frac{\text{Tr}\tilde{A}B_d\text{Tr}\tilde{A}^\dagger B_d}{\text{Tr}\tilde{A}^\dagger \tilde{A}\text{Tr}(B_d^2)}, \quad (\text{D1})$$

where

$$\tilde{A} = \mathcal{U}_L A \mathcal{U}_L^\dagger \quad \text{and} \quad B_d = \mathcal{U}_L B \mathcal{U}_L^\dagger \equiv b_d \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (\text{D2})$$

Then all dependence on B drops out from α_{AB} , and one is then left with

$$\alpha_{AB} = \frac{|\tilde{a}^+|^2}{|\tilde{a}^0|^2 + |\tilde{a}^+|^2 + |\tilde{a}^{++}|^2}, \quad (\text{D3})$$

from which Eq. (3.28) follows immediately when $\tilde{A}$ scans all its field space values. It is to be stressed that appealing to gauge invariance is essential for the proof; indeed, without gauge invariance, one would still be at liberty to choose the B -field space direction such as $b^+ = 0$, leading through Eq. (D3) to the same result; however, this would be no proof that α_{AB} remains in the $[0, 1]$ domain in other field directions. This is similar to the reason why we believe the determination of the BFB conditions in [33] for the Georgi-Machacek model lacks a complete proof, a version of which we give in Sec. III F.

2. $\alpha_{ABH} \in [-\sqrt{2}, +\sqrt{2}]$

Again, in the gauge where the real field $B = B_d$ (and denoting the components of the gauge transformed H and A fields with a tilde), α_{ABH} defined in Eq. (3.22) takes the form

$$\alpha_{ABH} = \sqrt{2} \text{sgn}(b) \frac{\text{Re}(\tilde{a}^{++}(\tilde{\phi}^{++})^2 - \tilde{a}^0(\tilde{\phi}^{0*})^2)}{\sqrt{|\tilde{a}^0|^2 + |\tilde{a}^+|^2 + |\tilde{a}^{++}|^2}(|\tilde{\phi}^0|^2 + |\tilde{\phi}^+|^2)}. \quad (\text{D4})$$

Using the fact that $-|z| \leq \text{Re}(z) \leq |z|$ for any complex number z , one immediately finds

$$-\frac{|\tilde{a}^{++}||\tilde{\phi}^+|^2 + |\tilde{a}^0||\tilde{\phi}^0|^2}{\sqrt{|\tilde{a}^0|^2 + |\tilde{a}^+|^2 + |\tilde{a}^{++}|^2}(|\tilde{\phi}^0|^2 + |\tilde{\phi}^+|^2)} \leq \frac{\alpha_{ABH}}{\sqrt{2}} \leq \frac{|\tilde{a}^{++}||\tilde{\phi}^+|^2 + |\tilde{a}^0||\tilde{\phi}^0|^2}{\sqrt{|\tilde{a}^0|^2 + |\tilde{a}^+|^2 + |\tilde{a}^{++}|^2}(|\tilde{\phi}^0|^2 + |\tilde{\phi}^+|^2)}, \quad (\text{D5})$$

where the upper (lower) bound is effectively reached in the field directions where $\arg(\tilde{a}^{++}(\tilde{\phi}^{++})^2) = 0(\pi)$, $\arg(\tilde{a}^0(\tilde{\phi}^{0*})^2) = \pi(0)$. Moreover, since $\sqrt{|\tilde{a}^0|^2 + |\tilde{a}^+|^2 + |\tilde{a}^{++}|^2} \geq \sqrt{|\tilde{a}^0|^2 + |\tilde{a}^{++}|^2}$, α_{ABH} scans a larger domain in the direction $\tilde{a}^+ = 0$, namely

$$-\sqrt{2} \frac{|\tilde{a}^{++}||\tilde{\phi}^+|^2 + |\tilde{a}^0||\tilde{\phi}^0|^2}{\sqrt{|\tilde{a}^0|^2 + |\tilde{a}^{++}|^2}(|\tilde{\phi}^0|^2 + |\tilde{\phi}^+|^2)} \leq \alpha_{ABH} \leq \sqrt{2} \frac{|\tilde{a}^{++}||\tilde{\phi}^+|^2 + |\tilde{a}^0||\tilde{\phi}^0|^2}{\sqrt{|\tilde{a}^0|^2 + |\tilde{a}^{++}|^2}(|\tilde{\phi}^0|^2 + |\tilde{\phi}^+|^2)}. \quad (\text{D6})$$

Defining $x = |\tilde{\phi}^+|/|\tilde{\phi}^0|$ and $y = |\tilde{a}^{++}|/|\tilde{a}^0|$, the above domain is rewritten as

$$-f(x, y) \leq \alpha_{ABH} \leq +f(x, y), \quad (\text{D7})$$

with

$$f(x, y) = \sqrt{2} \frac{1 + yx^2}{(1 + x^2)\sqrt{(1 + y^2)}}. \quad (\text{D8})$$

Noting that $f(x, y) = f(1/x, 1/y)$, a straightforward study of the function $f(x, y)$ in the domain $x, y \in [0, +\infty)$ shows that it possesses a saddle point at $x = y = 1$ and reaches a global maximum at $x = y = 0$ and at $x = y \rightarrow +\infty$ given by $f(0, 0) = f_{\max} = \sqrt{2}$, whence Eq. (3.29). Incidentally, we note that the ill-defined point $H = 0, A = 0$ in α_{ABH} is automatically accounted for through the behavior of f .

3. Boundary of the $(\alpha_{AH}, \alpha_{ABH})$ domain

To prove that Eq. (3.52) gives the lower boundary we show hereafter that

$\delta \equiv \alpha_{ABH}^2 - 2\alpha_{AH}$ is either negative or vanishing. This combination is of the form

$$\delta(x) = -1 + ax^2 + bx\sqrt{1 - x^2}, \quad (\text{D9})$$

with $x \equiv \sin \theta$ and a, b easily read from Eqs. (3.49) and (3.48),

$$a = -\cos 2\varphi \cos 2\psi + 2(c_1 \cos \varphi \cos^2 \psi - c_2 \sin \varphi \sin^2 \psi)^2, \quad (\text{D10})$$

$$b = -\sqrt{2}(c_3 \cos \varphi + c_4 \sin \varphi) \sin 2\psi, \quad (\text{D11})$$

and $c_i \equiv \cos \theta_i$. The study of the $\delta(x)$ function shows that it reaches only one stationary point

$$\delta^{\text{stationary}} = \delta(x_0) = -1 + \frac{b}{2}(r + \sqrt{1 + r^2}), \quad (\text{D12})$$

for

$$x = x_0 = \frac{1}{\sqrt{2}} \sqrt{1 + \frac{r}{\sqrt{1 + r^2}}} \in [0, 1], \quad (\text{D13})$$

where we took into account the fact that $0 \leq x \leq 1$ [cf. Eq. (3.44)] and defined $r = a/b$.

One also finds

$$\left. \frac{d^2 \delta(x)}{dx^2} \right|_{x=x_0} = -4b(1 + r^2)(r + \sqrt{1 + r^2}), \quad (\text{D14})$$

so that it is only the sign of b that dictates whether this stationary point is a maximum ($b > 0$) or a minimum ($b < 0$). Note that b as defined in Eq. (D11) can take either sign since $c_3, c_4 \in [-1, 1]$. To proceed we consider the two cases:

- (i) $b < 0$, $\delta(x_0)$ is a minimum: In this case one has $\delta(x) \leq 0 \forall x \in [0, 1]$, if and only if $\delta(0) \leq 0$ and $\delta(1) \leq 0$. The first condition is trivially satisfied. The second is equivalent to $a \leq 1$. Since $\varphi, \psi \in [0, \frac{\pi}{2}]$ and c_1, c_2 vary independently in $[-1, 1]$, it

follows from Eq. (D10) that, for fixed φ and ψ , a reaches a maximum when $c_1 = -c_2 = \pm 1$. One thus has

$$\begin{aligned} a &\leq -\cos 2\varphi \cos 2\psi + 2(\cos \varphi \cos^2 \psi + \sin \varphi \sin^2 \psi)^2 \\ &= \frac{1}{4}(3 + \cos 4\psi + 2 \sin 2\varphi \sin^2 2\psi) \\ &\leq \frac{1}{4}(3 + \cos 4\psi + 2 \sin^2 2\psi) = 1, \end{aligned} \quad (\text{D15})$$

and $\delta(1) \leq 0$ as required.

- (ii) $b > 0$, $\delta(x_0)$ is a maximum: In this case one has $\delta(x) \leq 0 \forall x \in [0, 1]$, if and only if $\delta(x_0) \leq 0$. From Eq. (D12) and the fact that $b > 0$, the condition $\delta(x_0) \leq 0$ can be rewritten as

$$\sqrt{a^2 + b^2} \leq 2 - a. \quad (\text{D16})$$

This in turn is equivalent to

$$4(1 - a) - b^2 \geq 0, \quad (\text{D17})$$

since $a \leq 1$, Eq. (D15) being valid independently of the sign of b . Expressing $\sin \psi$ and $\cos \psi$ in terms of $T \equiv \tan \psi$, the above inequality is equivalently rewritten as

$$a_0 + a_2 T^2 + a_4 T^4 \geq 0, \quad (\text{D18})$$

with $T \in [0, +\infty)$ and

$$a_0 = (1 - c_1^2) \cos^2 \varphi, \quad (\text{D19})$$

$$a_2 = 1 + c_1 c_2 \sin 2\varphi - (c_3 \cos \varphi + c_4 \sin \varphi)^2, \quad (\text{D20})$$

$$a_4 = (1 - c_2^2) \sin^2 \varphi. \quad (\text{D21})$$

We can now examine the NAS positivity conditions for the biquadratic polynomial in T , namely

$$a_0 \geq 0 \wedge a_4 \geq 0 \wedge a_2 + 2\sqrt{a_0 a_4} \geq 0. \quad (\text{D22})$$

The first two are trivially satisfied. To prove the third we should take into account the correlation $\theta_1 = \theta_2 + \theta_3 + \theta_4$ (modulo multiples of 2π); see Eq. (3.50). Moreover, since $\sin \theta_1$ and $\sin \theta_2$ can take either sign, we can include the two cases by simply using the inequality $\sqrt{(1 - c_1^2)(1 - c_2^2)} \geq \sin \theta_1 \sin \theta_2$ to write

$$\begin{aligned} a_2 + 2\sqrt{a_0 a_4} &\geq 1 + \sin 2\varphi \cos(\theta_1 - \theta_2) \\ &\quad - (\cos \theta_3 \cos \varphi + \cos \theta_4 \sin \varphi)^2. \end{aligned} \quad (\text{D23})$$

Using $\theta_1 = \theta_2 + \theta_3 + \theta_4$ the right-hand side of the above inequality simplifies to

$$\begin{aligned} & 1 + \sin 2\varphi \cos(\theta_3 + \theta_4) \\ & - (\cos \theta_3 \cos \varphi + \cos \theta_4 \sin \varphi)^2 \\ & = (\sin \theta_3 \cos \varphi - \sin \theta_4 \sin \varphi)^2. \end{aligned} \quad (\text{D24})$$

Thus the third NAS positivity condition in Eq. (D22) is valid for all values of the angles φ and θ_i . This implies that Eq. (D18) is satisfied for all $T \geq 0$ and thus for all values of ψ . Equation (D16) then holds for all the values of the angles, in particular those compatible with $b > 0$; we have thus proven that $\delta(x) \leq 0 \forall x \in [0, 1]$ in this case, too.

This ends the proof that $\alpha_{ABH}^2 - 2\alpha_{AH} \leq 0$ holds for all field directions and that Eq. (3.52) gives the lower boundary in the $(\alpha_{ABH}, \alpha_{AH})$ plane.

4. Boundary of the (α_A, α_{ABH}) domain

We rewrite Eq. (3.49) as

$$-c_1 \sqrt{1 - x^2} (1 - y^2) + c_2 x y^2 = \alpha_{ABH} \quad (\text{D25})$$

with the obvious notations, $x = \sin \varphi$, $y = \sin \psi$, $c_1 = \sqrt{2} \cos \theta_1 \sin \theta$, $c_2 = \sqrt{2} \cos \theta_2 \sin \theta$. We seek the conditions on α_{ABH} , c_1 , and c_2 that ensure the existence of at least one value for $x \in [0, 1]$ for each value of $y^2 \in [0, 1]$ and vice versa. This can be worked out by solving for y^2 and considering the (relative) signs of c_1 and c_2 . One finds

- (i) When $c_1 \times c_2 \geq 0$, α_{ABH} can be of any sign, with x and α_{ABH} satisfying

$$\begin{aligned} 0 \leq \frac{\alpha_{ABH}}{c_2} \leq x, \quad \text{or} \quad x^2 \leq 1 - \frac{\alpha_{ABH}^2}{c_1^2} \\ \left(\text{when } \frac{\alpha_{ABH}}{c_2} \leq 0 \right). \end{aligned} \quad (\text{D26})$$

Thus upper and/or lower parts of the $[0, 1]$ domain for x will not be reached $\forall y^2 \in [0, 1]$, unless

$$\alpha_{ABH} = 0. \quad (\text{D27})$$

- (ii) When $c_1 \times c_2 \leq 0$, α_{ABH} and c_2 should have the same sign, with x and α_{ABH} satisfying

$$\begin{aligned} 0 \leq x^2 \leq \min \left\{ \frac{\alpha_{ABH}^2}{c_2^2}, 1 - \frac{\alpha_{ABH}^2}{c_1^2}, \frac{c_1^2}{c_1^2 + c_2^2} \right\} \\ \text{or} \quad \max \left\{ \frac{\alpha_{ABH}^2}{c_2^2}, 1 - \frac{\alpha_{ABH}^2}{c_1^2}, \frac{c_1^2}{c_1^2 + c_2^2} \right\} \leq x^2 \leq 1. \end{aligned} \quad (\text{D28})$$

Thus intermediate parts of the $[0, 1]$ domain for x will not be reached $\forall y^2 \in [0, 1]$, unless

$$\alpha_{ABH} = (\alpha_{ABH})_{\text{crit}} \equiv \frac{|c_1| |c_2|}{\sqrt{c_1^2 + c_2^2}}. \quad (\text{D29})$$

Since $|c_1|, |c_2| \in [0, \sqrt{2}]$, the maximum value for $|\alpha_{ABH}|$ from Eq. (D29) is obtained when $|c_1| = |c_2| = \sqrt{2}$ and corresponds to the maximal critical value $|(\alpha_{ABH})_{\text{crit}}^{\text{max}}| = 1$, not $\sqrt{2}$. A direct consequence is the absence of correlations between α_{ABH} and x or α_{ABH} and y^2 in the domain $\alpha_{ABH} \in [-1, 1]$; i.e., the square $[-1, 1] \times [0, 1]$ is totally filled in both cases. Indeed, for any given $\alpha_{ABH} \in [-1, 1]$ one can always find c_1 and c_2 of opposite signs satisfying Eq. (D29) so that to any x corresponds at least one y^2 and vice versa, thus varying freely in $[0, 1]$. Note also that $|c_1| = |c_2| = \sqrt{2}$ entails maximizing $|\cos \theta_1|$, $|\cos \theta_2|$, and $\sin \theta$ to 1.

We can study now the allowed domain in the plane (α_{ABH}, α_A) . We first determine the allowed (α_{ABH}, α_A) subdomain corresponding to $\sin \theta = 1$, and then show that all subdomains that correspond to $\sin \theta < 1$ are necessarily within that subdomain, which thus turns out to be the full (α_{ABH}, α_A) domain.

When $\sin \theta = 1$, the dependence on $\cos \rho$ drops out from Eq. (3.42), and one can easily solve for $x (= \sin \varphi)$ as a function of α_A ,

$$x_{\pm} = \sqrt{\frac{1}{2} (1 \pm \sqrt{2\alpha_A - 1})}. \quad (\text{D30})$$

The two $\pm$ solutions should be kept in the discussion as their union scans the full $[0, 1]$ domain of x allowing α_A to scan all its allowed domain $[\frac{1}{2}, 1]$. Similarly, since $\sin \theta = 1$, α_{ABH} will scan all its allowed domain $[-\sqrt{2}, +\sqrt{2}]$ by varying x , y , c_1 , and c_2 . Let us choose a couple of values (α_{ABH}, α_A) in their respective domains.

If $|\alpha_{ABH}| \leq 1$, then, relying on what was demonstrated after Eq. (D29), one can always find c_1 and c_2 (or equivalently $\cos \theta_1$ and $\cos \theta_2$) with opposite signs and the sign of c_2 being that of α_{ABH} , in such a way that $\alpha_{ABH} = (\alpha_{ABH})_{\text{crit}}$. It follows that for any $x \in [0, 1]$ there exists $y^2 \in [0, 1]$ consistent with the given value of α_{ABH} . In particular, this is true for the values of x corresponding, through Eq. (D30), to any given value of α_A . There is thus no obstruction on the independent choice of the values of α_{ABH} and α_A as long as $|\alpha_{ABH}| \leq 1$. It follows that the entire square $[-1, 1] \times [\frac{1}{2}, 1]$ is allowed in the (α_{ABH}, α_A) plane.

If $|\alpha_{ABH}| > 1$, one has to examine separately the conditions given by Eqs. (D26) and (D28). Note also that since $|\alpha_{ABH}| > 1$ the min and max in Eqs. (D28)

become uniquely defined, equaling, respectively, $1 - \frac{\alpha_{ABH}^2}{c_1^2}$ and $\frac{\alpha_{ABH}^2}{c_2^2}$. Plugging x as given by Eq. (D30) into the four inequalities, it is clear that a necessary condition in each case obtains when c_1 and c_2 take their extreme values $\pm\sqrt{2}$. Taking consistently into account the various sign conditions in each case as well as the $\pm$ in Eq. (D30) one determines the necessary condition relating α_A and α_{ABH} . One finds exactly the same inequality in the four cases, namely $\alpha_A \geq 1 - \alpha_{ABH}^2 + \frac{1}{2}\alpha_{ABH}^4$. Moreover, this condition is also sufficient since it allows at least the extremal values of c_1 and c_2 . Thus

$$\alpha_A = 1 - \alpha_{ABH}^2 + \frac{1}{2}\alpha_{ABH}^4 = \frac{1}{2}(1 + (\alpha_{ABH}^2 - 1)^2) \quad (\text{D31})$$

gives the lower boundary for α_A when $\alpha_{ABH} > 1$.

This completes the proof that when $\sin \theta = 1$, the allowed domain in the (α_{ABH}, α_A) plane is as defined by Eqs. (3.53) and illustrated in Fig. 4.

Since θ_1 and θ_2 appear only in α_{ABH} , they can be safely chosen without biasing the correlations between α_A and α_{ABH} , as long as they maximize the allowed domain of the latter. The angle θ is, however, common to α_A and α_{ABH} . One should then be careful that the value $\sin \theta = 1$ does not miss points in the allowed domain. A necessary condition for this not to happen is that $\sin \theta = 1$ still allows α_A and α_{ABH} to take any value in their respective domains as given by Eqs. (3.26) and (3.29). This is indeed the case as one can check from Eqs. (3.42) and (3.49) by varying all the other angles at fixed $\sin \theta = 1$.

However, this is not sufficient. One should still check that for $\sin \theta$ strictly smaller than one there exists no set of values for the remaining angle variables giving a point in the (α_{ABH}, α_A) plane that is *outside* the domain defined by Eqs. (3.53). To show this it suffices to prove [cf. Eq. (D31)] that

$$2\alpha_A - 1 - (\alpha_{ABH}^2 - 1)^2 \geq 0, \quad \forall \sin \theta, \quad (\text{D32})$$

whenever

$$|\alpha_{ABH}| > 1. \quad (\text{D33})$$

Rewriting Eq. (3.49) as

$$\alpha_{ABH} = \sqrt{2}Y \sin \theta, \quad (\text{D34})$$

where

$$Y = \sin \varphi \sin^2 \psi \cos \theta_2 - \cos \varphi \cos^2 \psi \cos \theta_1 \quad (\text{D35})$$

and $Y \in [-1, 1]$, condition (D33) implies

$$|Y| \geq \frac{1}{\sqrt{2}} \quad \text{and} \quad \sin \theta \geq \frac{1}{\sqrt{2}}, \quad (\text{D36})$$

since none of $|Y|$ and $\sin \theta$ can exceed one. We can thus replace Eq. (D33) by

$$1 \geq |Y| \geq \frac{1}{\sqrt{2}} \quad (\text{D37})$$

and

$$1 \geq \sin \theta \geq \frac{1}{\sqrt{2}|Y|}. \quad (\text{D38})$$

On the other hand, as seen from Eqs. (3.42) and (3.49), the only dependence on the angle ρ in Eq. (D32) is linear in $\cos \rho$ and with a positive coefficient:

$$\frac{1}{4} \sin 2\varphi \sin^2 2\theta \cos \rho + \dots \geq 0. \quad (\text{D39})$$

Condition (D32) is thus equivalent to the one where $\cos \rho$ takes its minimal value $\cos \rho = -1$, in which case (D32) can be recast in the form

$$a_4 \tau^4 + a_2 \tau^2 - 1 \geq 0 \quad (\text{D40})$$

with

$$a_2 = 2(2Y^2 - \sin 2\varphi), \quad (\text{D41})$$

$$a_4 = -4(Y^2 - \cos^2 \varphi)(Y^2 - \sin^2 \varphi), \quad (\text{D42})$$

where we defined $\tau = \tan \theta$ and dropped out a positive denominator. The coefficients of τ^2 and τ^4 in Eq. (D40) both satisfy

$$a_2 \geq 0, \quad (\text{D43})$$

$$a_4 \geq 0, \quad (\text{D44})$$

as a consequence of the lower bound in Eq. (D37). The first is immediate to establish. The positivity of a_4 is less obvious. Rewriting $|Y| \geq 1/\sqrt{2}$ and a_4 , respectively, as

$$0 \leq \left(Y - \frac{1}{\sqrt{2}}\right) \left(Y + \frac{1}{\sqrt{2}}\right), \quad (\text{D45})$$

$$a_4 = -4(Y - \cos \varphi)(Y + \cos \varphi)(Y - \sin \varphi)(Y + \sin \varphi), \quad (\text{D46})$$

and noting that Y is linear in $\cos \theta_1$ and $\cos \theta_2$ [cf. Eq. (D35)], one can easily study the sign of a_4 when

Eq. (D45) is satisfied, in terms of a bundle of six parallel straight lines with slope $\cot \varphi \cot^2 \psi$ in the $(\cos \theta_1, \cos \theta_2)$ plane; the sign alternates each time one of these lines is crossed. Moreover, since they are all parallel, it suffices to study the change of sign along a given axis in the $(\cos \theta_1, \cos \theta_2)$ plane, say the axis defined by $\cos \theta_2 = 0$. On this axis the inequality Eq. (D45) is satisfied if and only if

$$\frac{1}{\sqrt{2} \cos \varphi \cos^2 \psi} \leq \cos \theta_1 \leq 1 \quad \text{or} \quad -1 \leq \cos \theta_1 \leq -\frac{1}{\sqrt{2} \cos \varphi \cos^2 \psi}. \quad (\text{D47})$$

This implies

$$\cos \varphi \geq \frac{1}{\sqrt{2} \cos^2 \psi} \geq \frac{1}{\sqrt{2}}, \quad (\text{D48})$$

and thus

$$\sin \varphi \leq \frac{1}{\sqrt{2}} \quad \text{and} \quad \tan \varphi \leq 1. \quad (\text{D49})$$

On the other hand, it is easily seen from Eqs. (D35) and (D46) (with $\cos \theta_2 = 0$) that a_4 is positive if and only if $\cos \theta_1$ is between $1/\cos^2 \psi$ and $\tan \varphi/\cos^2 \psi$ or between $-1/\cos^2 \psi$ and $-\tan \varphi/\cos^2 \psi$, and negative otherwise. Using Eq. (D49) these conditions read

$$\frac{\tan \varphi}{\cos^2 \psi} \leq \cos \theta_1 \leq \frac{1}{\cos^2 \psi} \quad \text{or} \quad -\frac{1}{\cos^2 \psi} \leq \cos \theta_1 \leq -\frac{\tan \varphi}{\cos^2 \psi}. \quad (\text{D50})$$

And, again from Eq. (D49),

$$\frac{\tan \varphi}{\cos^2 \psi} \leq \frac{1}{\sqrt{2} \cos \varphi \cos^2 \psi}, \quad (\text{D51})$$

which shows that Eq. (D50) is satisfied whenever Eq. (D47) [or equivalently Eq. (D37)] is satisfied. Thus condition (D37) implies $a_4 \geq 0$. It is easy to see that this property remains true even when $\cos \theta_2 \neq 0$. Indeed, if (D37) is satisfied for a given point $(\cos \theta_1, \cos \theta_2)$, then it remains true on the entire straight line with slope $\cot \varphi \cot^2 \psi$ going through this point, in particular for the point intersecting the axis $\cos \theta_2 = 0$, and we are brought back to the known case.

Now back to Eq. (D40): The domain of variation of τ^2 corresponding to Eq. (D38) is given by

$$\frac{1}{2Y^2 - 1} \leq \tau^2 < +\infty. \quad (\text{D52})$$

Moreover, the quadratic function in τ^2 is a monotonically increasing function as can be seen from its derivative and Eqs. (D43) and (D44). Its minimum is thus reached for $\tau_{\min}^2 = \frac{1}{2Y^2 - 1}$ and is given by

$$a_4 \tau_{\min}^4 + a_2 \tau_{\min}^2 - 1 = \frac{2(1 - \sin 2\varphi)(4Y^2 + \sin 2\varphi - 1)}{(1 - 2Y^2)^2}, \quad (\text{D53})$$

which is obviously positive when Eq. (D37) is satisfied. Thus Eq. (D40) is always satisfied whenever Eqs. (D37) and (D38) are. This completes the proof that Eq. (D32) is satisfied whenever Eq. (D33) holds and that the full allowed domain in the (α_{ABH}, α_A) plane is given by Eqs. (3.53).

5. Boundary of the $(\alpha_{AB}, \alpha_{AH})$ domain

From Eq. (3.48) one sees that α_{AH} is of the form

$$\alpha_{AH}(x) = \frac{1}{2} + ax + b\sqrt{1 - x^2}, \quad \text{with } x \in [-1, +1], \quad (\text{D54})$$

where we defined $x \equiv \cos 2\psi$ and a and b are readily obtained from Eqs. (3.45) and (3.48),

$$a = \frac{1}{2}(1 - \alpha_{AB}) \cos 2\varphi, \quad (\text{D55})$$

$$b = \frac{1}{\sqrt{2}}(\cos \theta_3 \cos \varphi + \cos \theta_4 \sin \varphi) \sqrt{(1 - \alpha_{AB})\alpha_{AB}}. \quad (\text{D56})$$

It is easy to study the structure of maxima and minima of $\alpha_{AH}(x)$ at fixed a, b . One finds that it always has only one stationary point, at $x = \frac{a \operatorname{sgn}(b)}{\sqrt{a^2 + b^2}} \in [-1, +1]$, given by

$$\alpha_{AH}^{\text{stationary}} = \frac{1}{2} + \sqrt{a^2 + b^2} \operatorname{sgn}(b). \quad (\text{D57})$$

Moreover, this stationary point is found to be a minimum (respectively, maximum) when $b < 0$ (respectively, $b > 0$), and thus with a corresponding maximum (respectively, minimum) of α_{AH} given by $\max\{\alpha_{AH}(\pm 1)\}$ (respectively, $\min\{\alpha_{AH}(\pm 1)\}$). This leads to

$$\frac{1}{2} - \sqrt{a^2 + b^2} \leq \alpha_{AH} \leq \frac{1}{2} + |a|, \quad \text{iff } b \leq 0, \quad (\text{D58})$$

$$\frac{1}{2} - |a| \leq \alpha_{AH} \leq \frac{1}{2} + \sqrt{a^2 + b^2}, \quad \text{iff } b \geq 0. \quad (\text{D59})$$

The parameter b as defined by Eq. (D56) can take either sign when all the angles are varied [since $\cos \theta_3, \cos \theta_4 \in [-1, 1]$ and $\varphi \in [0, \frac{\pi}{2}]$; cf. Eq. (3.44)]. It is thus more

relevant to combine the α_{AH} domains given above, reducing them for fixed a and $|b|$ to

$$\frac{1}{2} - \sqrt{a^2 + b^2} \leq \alpha_{AH} \leq \frac{1}{2} + \sqrt{a^2 + b^2}, \quad (\text{D60})$$

or equivalently to

$$\left(\alpha_{AH} - \frac{1}{2}\right)^2 \leq a^2 + b^2. \quad (\text{D61})$$

Given Eq. (D56), the domain in Eq. (D61) is obviously maximized for $\cos \theta_3 = \cos \theta_4 = \pm 1$. Assuming these values we now show that α_{AH} will scan its full allowed domain $[0, 1]$, i.e., that $a^2 + b^2$ will reach $\frac{1}{4}$, only when $0 \leq \alpha_{AB} \leq \frac{1}{2}$. We first note from Eqs. (D55) and (D56) that $a^2 + b^2$ can be recast in the form

$$a^2 + b^2 = -\frac{1}{4}(1 + \sin 2\varphi)^2 \left(\alpha_{AB} - \frac{\sin 2\varphi}{1 + \sin 2\varphi}\right)^2 + \frac{1}{4}. \quad (\text{D62})$$

Since $\sin 2\varphi \in [0, 1]$, it is clear that $a^2 + b^2$ reaches $\frac{1}{4}$ iff $\alpha_{AB} = \frac{\sin 2\varphi}{1 + \sin 2\varphi} \in [0, \frac{1}{2}]$. It then follows from Eq. (D60) that all the α_{AH} domain $[0, 1]$ is allowed when $\alpha_{AB} \in [0, \frac{1}{2}]$, whence the boundaries given in Eqs. (3.57)–(3.59).

Finally, when $\frac{1}{2} \leq \alpha_{AB} \leq 1$, the study of $a^2 + b^2$ as a function of $\sin 2\varphi$ in Eq. (D62) shows that $a^2 + b^2$ reaches its maximum for $\sin 2\varphi = 1$, given by

$$a^2 + b^2|_{\max} = \frac{1}{4} - \left(\alpha_{AB} - \frac{1}{2}\right)^2. \quad (\text{D63})$$

Plugging this back into Eq. (D61) gives the largest allowed domain

$$\left(\alpha_{AH} - \frac{1}{2}\right)^2 + \left(\alpha_{AB} - \frac{1}{2}\right)^2 \leq \frac{1}{4}, \quad (\text{D64})$$

whence the half-circle boundary Eq. (3.60).

APPENDIX E: THE $(\hat{\omega}_0, \hat{\xi}_0)$ DOMAIN

To simplify the presentation we define

$$x = \cos 2\theta_0, \quad y = \cos \arg(\chi^0), \quad (\text{E1})$$

so that $\sin 2\theta_0 = +\sqrt{1-x^2}$ and $-1 \leq x, y \leq 1$; cf. Eqs. (3.106) and (3.107). We also define

$$w(x, y) = \frac{1}{8}(1 - x + 2\sqrt{2(1-x^2)}y) \quad (\text{E2})$$

so that the zeroth order u contribution to $\hat{\omega}$, Eq. (3.97), reads

$$\hat{\omega}_0(x, y, \theta_+) = w(x, y) \sin^2 \theta_+. \quad (\text{E3})$$

For later use we also denote by $x_y^>$ and $x_y^<$, respectively, the *largest* and *smallest* values of x satisfying the equation

$$w(x_y^>, y) = w(x_y^<, y) = \tilde{\omega}_0, \quad (\text{E4})$$

where $\tilde{\omega}_0$ is a given value of $\hat{\omega}_0 \in [-\frac{1}{4}, \frac{1}{2}]$. These two values of x are easily determined to be

$$x_y^{\gtrless} = \frac{1 - 8\tilde{\omega}_0 \pm 8\sqrt{y^2(2(1 - 4\tilde{\omega}_0)\tilde{\omega}_0 + y^2)}}{1 + 8y^2}. \quad (\text{E5})$$

Note also that they are reached if and only if $\sin^2 \theta_+ = 1$.

Using Eq. (E3) to eliminate $\sin^2 \theta_+$ from Eq. (3.99) one obtains straightforwardly a relation between $\hat{\omega}_0$ and the zeroth order u contribution to $\hat{\xi}$,

$$\hat{\xi}_0(x, y, \hat{\omega}_0) = 1 + c_1 \hat{\omega}_0 + c_2 \hat{\omega}_0^2 \quad (\text{E6})$$

with

$$c_1 = -\frac{(1-x)}{2w(x, y)}, \quad (\text{E7})$$

$$c_2 = -\frac{(1-x)(1+3x)}{8w(x, y)^2}. \quad (\text{E8})$$

x, y , and $\hat{\omega}_0$ can be varied independently of each other only locally, but they have global correlations due to Eq. (E3): From $0 \leq \sin^2 \theta_+ \leq 1$, one must require

$$w(x, y) \leq \hat{\omega}_0 \leq 0 \quad (\text{E9})$$

or

$$0 \leq \hat{\omega}_0 \leq w(x, y). \quad (\text{E10})$$

Apart from the special cases $\{x = 1, \hat{\omega}_0 = 0\}$ and $\{x = -1, \hat{\omega}_0 = (1/4) \sin^2 \theta_+\}$ where y varies freely in $[-1, +1]$, the above constraints dictate in general that the allowed ranges for y depend on $x (\neq -1, +1)$ and $\hat{\omega}_0$ as follows:

$$\text{if } \hat{\omega}_0 \geq 0, \quad \text{then } \max \left\{ -1, -\frac{1}{2\sqrt{2}} \sqrt{\frac{1-x}{1+x}} + \frac{2\sqrt{2}\hat{\omega}_0}{\sqrt{1-x^2}} \right\} \leq y \leq 1; \quad (\text{E11})$$

$$\text{if } \hat{\omega}_0 \leq 0, \quad \text{then } -1 \leq y \leq -\frac{1}{2\sqrt{2}} \sqrt{\frac{1-x}{1+x}} + \frac{2\sqrt{2}\hat{\omega}_0}{\sqrt{1-x^2}} \leq 0. \quad (\text{E12})$$

We now show the following key property:

$\hat{\xi}_0(x, y, \hat{\omega}_0)$, taken as a function of y , is increasing for

$$\hat{\omega}_0 \geq 0 \text{ and increasing for } \hat{\omega}_0 \leq 0. \quad (\text{E13})$$

The derivative of $\hat{\xi}_0$ reads

$$\frac{\partial \hat{\xi}_0}{\partial y} = \kappa^2 \hat{\omega}_0 \left(1 + \frac{\hat{\omega}_0}{2w(x, y)} (1 + 3x) \right), \quad (\text{E14})$$

where κ^2 is a positive definite x - and y -dependent prefactor. Using Eq. (E3), one finds that the last factor to the right is also positive, since $(1 + \frac{(1+3x)}{2} \sin^2 \theta_+) \geq \cos^2 \theta_+$ for $x \in [-1, +1]$.

1. Upper boundary

It follows from (E13) that the maximum of $\hat{\xi}_0$ for fixed x and $\hat{\omega}_0$ is given by $\hat{\xi}_0(x, +1, \hat{\omega}_0)$ (respectively, $\hat{\xi}_0(x, -1, \hat{\omega}_0)$) when $\hat{\omega}_0 \geq 0$ (respectively, $\hat{\omega}_0 \leq 0$). This suggests the study of these two functions in the corresponding negative and positive ranges of $\hat{\omega}_0$, which we will treat as families of functions of $\hat{\omega}_0$ parametrized by x :

$$\hat{\xi}_0^{(x)}(\hat{\omega}_0) = \hat{\xi}_0(x, +1, \hat{\omega}_0), \quad \text{for } 0 \leq \hat{\omega}_0 \leq w(x, +1) \quad \text{and} \\ x \in [-1, +1], \quad (\text{E15a})$$

$$\hat{\xi}_0^{(x)}(\hat{\omega}_0) = \hat{\xi}_0(x, -1, \hat{\omega}_0), \quad \text{for } w(x, -1) \leq \hat{\omega}_0 \leq 0 \quad \text{and} \\ x \in \left[-\frac{7}{9}, +1 \right], \quad (\text{E15b})$$

$$\hat{\xi}_0^{(x)}(\hat{\omega}_0) = 1, \quad \text{for } \hat{\omega}_0 = 0 \quad \text{and} \quad x \in \left[-1, -\frac{7}{9} \right]. \quad (\text{E15c})$$

In writing the above we took into account the consistency conditions Eqs. (E11) and (E12) and noted that $\hat{\omega}_0 < 0$ cannot be satisfied when $x \in [-1, -\frac{7}{9}]$. Obviously the upper boundary in the $(\hat{\omega}_0, \hat{\xi}_0)$ plane, that is, the function $\hat{\xi}_0^{\max}(\hat{\omega}_0)$ giving the maximal allowed value of $\hat{\xi}_0$ for a given $\hat{\omega}_0$, is obtained by determining the upper envelope of the family of functions $\hat{\xi}_0^{(x)}(\hat{\omega}_0)$ defined in Eqs. (E15a) and (E15b). We will show below that this envelope is given by

$$\hat{\xi}_0^{\max}(\hat{\omega}_0) = \hat{\xi}_0^{(x_{+1}^>)}(\hat{\omega}_0)|_{\hat{\omega}_0=w(x_{+1}^>, +1)}, \quad \text{for } \hat{\omega}_0 \geq 0, \quad (\text{E16a})$$

$$\hat{\xi}_0^{\max}(\hat{\omega}_0) = \hat{\xi}_0^{(x_{-1}^>)}(\hat{\omega}_0)|_{\hat{\omega}_0=w(x_{-1}^>, -1)}, \quad \text{for } \hat{\omega}_0 \leq 0, \quad (\text{E16b})$$

where $x_{\pm 1}^>$ have been defined in Eqs. (E4).

In other terms, the upper boundary is traced when $\hat{\omega}_0$ sits at the nonvanishing end points of its allowed domains given in Eqs. (E15a) and (E15b), thus corresponding to $\sin^2 \theta_+ = 1$ as noted after Eq. (E5), and for the largest value

of x that allows one to reach each end point. This result is a consequence of certain properties that can easily be shown by direct analytical (as well as numerical) inspection of the relevant functions and their first derivative, summarized hereafter without proof:

- (1) $w(x, -1)$ is ≤ 0 if and only if $x \in [-\frac{7}{9}, 1]$, and
 - (a) $w(x, -1)$ is *strictly decreasing* for $x \in [-\frac{7}{9}, \frac{1}{3}]$, spanning the full negative $\hat{\omega}_0$ domain $[-\frac{1}{4}, 0]$,
 - (b) $w(x, -1)$ is *strictly increasing* for $x \in [\frac{1}{3}, 1]$, spanning the full negative $\hat{\omega}_0$ domain $[-\frac{1}{4}, 0]$.

It follows that $x_{-1}^< \in [-\frac{7}{9}, \frac{1}{3}]$ and $x_{-1}^> \in [\frac{1}{3}, 1]$ [cf. Eq. (E4)].

- (2) $w(x, +1)$ is ≥ 0 in the entire x domain $[-1, +1]$, and
 - (a) $w(x, +1)$ is *strictly increasing* for $x \in [-1, -\frac{1}{3}]$, spanning partially the positive $\hat{\omega}_0$ domain $[\frac{1}{4}, \frac{1}{2}]$,
 - (b) $w(x, +1)$ is *strictly decreasing* for $x \in [-\frac{1}{3}, 1]$, spanning the full positive $\hat{\omega}_0$ domain $[0, \frac{1}{2}]$.

It follows that $x_{+1}^< \in [-1, -\frac{1}{3}]$ and $x_{+1}^> \in [-\frac{1}{3}, 1]$ [cf. Eq. (E4)].

- (3) in the domains of $x_{-1}^>$ and $x_{+1}^>$, that is, respectively, for $x \in [\frac{1}{3}, 1]$, $\hat{\omega}_0 \in [-\frac{1}{4}, 0]$, and $x \in [-\frac{1}{3}, 1]$, $\hat{\omega}_0 \in [0, \frac{1}{2}]$, $\hat{\xi}_0^{(x)}(\hat{\omega}_0)$ is a *strictly increasing* function of x . $[\frac{\partial}{\partial x} \hat{\xi}_0^{(x)}(\hat{\omega}_0)]$ vanishes only at the two isolated points $\{\hat{\omega}_0 = 0, \forall x\}$ and $\{\hat{\omega}_0 = \frac{1}{2}, x = -\frac{1}{3}\}$, where $\hat{\xi}_0$ takes its two extreme values 1 and $\frac{1}{3}$.

- (4) in the domains of $x_{-1}^<$ and $x_{+1}^<$, that is, respectively, for $x \in [-\frac{7}{9}, \frac{1}{3}]$, $\hat{\omega}_0 \in [-\frac{1}{4}, 0]$, and $x \in [-1, -\frac{1}{3}]$, $\hat{\omega}_0 \in [\frac{1}{4}, \frac{1}{2}]$, $\hat{\xi}_0^{(x)}(\hat{\omega}_0)$ taken as a function of x can be either *strictly increasing* or *strictly decreasing*, but it changes its monotonicity *at most* once depending on the value of $\hat{\omega}_0$.

- (5) $\hat{\xi}_0^{(x_{\pm 1}^>)}(\hat{\omega}_0) - \hat{\xi}_0^{(x_{\pm 1}^<)}(\hat{\omega}_0) = \frac{16}{27} (1 - 2\hat{\omega}_0)^{3/2} \sqrt{1 + 4\hat{\omega}_0} \geq 0$, valid for all $\hat{\omega}_0 \in [-\frac{1}{4}, \frac{1}{2}]$.

Consider the value of $\hat{\xi}_0 = \hat{\xi}_0^{(x_{-1}^>)}(\hat{\omega}_0)|_{\hat{\omega}_0=w(x_{-1}^>, -1)}$ for a given $\hat{\omega}_0 \leq 0$ [cf. Eq. (E16b)]. We now show that varying x in the vicinity of $x_{-1}^>$ does not allow one to find for the same $\hat{\omega}_0$ a larger value for $\hat{\xi}_0$: To find another value of $\hat{\xi}_0$ for the same $\hat{\omega}_0$ one should, according to Eq. (E15b), choose an x such that

$$w(x, -1) < \hat{\omega}_0 = w(x_{-1}^>, -1) = w(x_{-1}^<, -1). \quad (\text{E17})$$

(Note that the last equality is simply due to the definition of $x_{-1}^>$ and $x_{-1}^<$.) If x is taken sufficiently close to $x_{-1}^>$ so that $x \in [\frac{1}{3}, 1]$, then the above inequality is satisfied only if x is *strictly smaller* than $x_{-1}^>$ since by property (i) w is a strictly increasing function in the considered domain. It then follows from property (iii) that the new value of $\hat{\xi}_0$ is necessarily *strictly smaller* than the initial $\hat{\xi}_0^{(x_{-1}^>)}(\hat{\omega}_0)|_{\hat{\omega}_0=w(x_{-1}^>, -1)}$. Thus the latter is indeed a local maximum. But x can also be in the domain $[-\frac{7}{9}, \frac{1}{3}]$. In this case property (i) implies that x should be *strictly greater*

than $x_{-1}^<$ since w is a strictly decreasing function of x in the considered domain. Then according to property (iv):

Either $\hat{\zeta}_0^{(x)}(\hat{\omega}_0)$ did not change its monotonicity for the given value of $\hat{\omega}_0$ and the considered range for x within the $[-\frac{7}{9}, \frac{1}{3}]$ domain, which means it is still a strictly increasing function of x [cf. property (iii)]. In this case one has $\hat{\zeta}_0^{(x)}(\hat{\omega}_0) < \hat{\zeta}_0^{(\frac{1}{3})}(\hat{\omega}_0) < \hat{\zeta}_0^{(x_{-1}^<)}(\hat{\omega}_0)$ with $\hat{\omega}_0 = w(x_{-1}^<, -1)$.

Or $\hat{\zeta}_0^{(x)}(\hat{\omega}_0)$ changed once its monotonicity becoming a strictly decreasing function of x . In this case one has $\hat{\zeta}_0^{(x)}(\hat{\omega}_0) < \hat{\zeta}_0^{(x_{-1}^<)}(\hat{\omega}_0)$ because $x > x_{-1}^<$ as shown above. But then property (v) implies $\hat{\zeta}_0^{(x)}(\hat{\omega}_0) < \hat{\zeta}_0^{(x_{-1}^>)}(\hat{\omega}_0)$, which holds for any $\hat{\omega}_0$ including $\hat{\omega}_0 = w(x_{-1}^>, -1)$.

It follows that in all cases $\hat{\zeta}_0^{(x_{-1}^>)}(\hat{\omega}_0)|_{\hat{\omega}_0=w(x_{-1}^>, -1)}$ is indeed a global maximum.

A similar proof holds for the branch $\hat{\omega}_0 \geq 0$ noting though the reversed inequality in Eq. (E15a) as compared to Eq. (E15b), and the reversed behavior of w in property (ii) as compared to property (i). More specifically, one should look for an x such that $w(x, +1) > \hat{\omega}_0 = w(x_{+1}^>, +1) = w(x_{+1}^<, +1)$ [cf. Eqs. (E16a) and (E15a)] and consider separately the cases $x \in [-\frac{1}{3}, 1]$ and $x \in [-1, -\frac{1}{3}]$. In the first case the above inequality implies, using properties (ii) and (iii), that $x < x_{+1}^>$ and $\hat{\zeta}_0^{(x_{+1}^>)}(\hat{\omega}_0)|_{\hat{\omega}_0=w(x_{+1}^>, +1)}$ is a local maximum. In the second case property (ii) and the above inequality imply $x > x_{+1}^<$, and the result that $\hat{\zeta}_0^{(x_{+1}^>)}(\hat{\omega}_0)|_{\hat{\omega}_0=w(x_{+1}^>, +1)}$ is a maximum is again obtained upon the use of properties (iv) and (v).

We can now write explicitly $\hat{\zeta}_0^{\max}(\hat{\omega}_0)$. First, from Eqs. (E6), (E16b) and (E16a) and properties (i) and (ii) one finds the simple form

$$\hat{\zeta}_0^{\max}(\hat{\omega}_0) = Z(x_{+1}^>), \quad \text{for } 0 \leq \hat{\omega}_0 \leq \frac{1}{2}, \quad (\text{E18a})$$

$$\hat{\zeta}_0^{\max}(\hat{\omega}_0) = Z(x_{-1}^>), \quad \text{for } -\frac{1}{4} \leq \hat{\omega}_0 \leq 0, \quad (\text{E18b})$$

where

$$Z(x) = \frac{1}{8}(3 + x(2 + 3x)). \quad (\text{E19})$$

The explicit dependence on $\hat{\omega}_0$ is obtained by solving $\hat{\omega}_0 = w(x_{+1}^>, +1)$ for $x_{+1}^>$ and $\hat{\omega}_0 = w(x_{-1}^>, -1)$ for $x_{-1}^>$ and plugging the result back in Eqs. (E18a) and (E18b). In fact, a further simplification occurs because the two solutions are found to have exactly the same functional dependence on $\hat{\omega}_0$ [cf. Eq. (E5)], even though they correspond to different ranges of the latter:

$$x_{\pm 1}^> = \frac{1}{9}(1 - 8\hat{\omega}_0 + 8\sqrt{(1 - 2\hat{\omega}_0)(1 + 4\hat{\omega}_0)}). \quad (\text{E20})$$

Equations (E18a) and (E18b) can thus be merged into one single form for the full $\hat{\omega}_0$ range $[-\frac{1}{4}, \frac{1}{2}]$,

$$\begin{aligned} \hat{\zeta}_0^{\max}(\hat{\omega}_0) &= \frac{1}{3} + \frac{2}{27}(1 - 2\hat{\omega}_0 + 2\sqrt{(1 - 2\hat{\omega}_0)(1 + 4\hat{\omega}_0)})^2, \\ \hat{\omega}_0 &\in \left[-\frac{1}{4}, \frac{1}{2}\right], \end{aligned} \quad (\text{E21})$$

which reproduces the upper boundary given in [33] (note, however, that we deal with the inverse function with respect to the function considered in Ref. [33]); see also Fig. 11.

2. Lower boundary

We turn now to the determination of the lower boundary of the domain. In contrast with the previous case we cannot just study $\hat{\zeta}_0(x, +1, \hat{\omega}_0)$ and $\hat{\zeta}_0(x, -1, \hat{\omega}_0)$ as being the minima in the y domain, respectively, for $\hat{\omega}_0 \leq 0$ and $\hat{\omega}_0 \geq 0$ as suggested by the property (E13). Indeed, it is obvious from Eqs. (E3) and (E2) [see also Eq. (E15)] that $\hat{\zeta}_0(x, +1, \hat{\omega}_0)$ and more generally $\hat{\zeta}_0(x, y \geq 0, \hat{\omega}_0)$ are never compatible with $\hat{\omega}_0 < 0$. Moreover, $\hat{\zeta}_0(x, -1, \hat{\omega}_0)$ is compatible with $\hat{\omega}_0 \geq 0$ only in the reduced domain of $x \in [-1, -\frac{7}{9}]$ as already discussed after Eq. (E15c). This means that there could exist $y > -1$ and x outside this reduced domain for which values of $\hat{\zeta}_0$ smaller than $\hat{\zeta}_0(x, -1, \hat{\omega}_0)$ could be reached. Thus for both domains, $\hat{\omega}_0 \geq 0$ and $\hat{\omega}_0 \leq 0$, y should be varied away from $y = +1$ or -1 to determine the lower boundary function $\hat{\zeta}_0^{\min}(\hat{\omega}_0)$ that gives for each $\hat{\omega}_0$ the minimal allowed value for $\hat{\zeta}_0$. It is easy to see that for given $\hat{\omega}_0$ and x , the minimal value of $\hat{\zeta}_0$ is reached only when $\hat{\omega}_0 = w(x, y)$. This is a consequence of combining property (E13) with Eqs. (E9) and (E10) and the fact that $w(x, y)$ is an increasing function of y . For example, for a given positive $\hat{\omega}_0$ that should satisfy Eq. (E10) for say $y = +1$, decreasing y will monotonically decrease simultaneously $\hat{\zeta}_0$ [cf. (E13)] and $w(x, y)$. Since values of y such that $\hat{\omega}_0 > w(x, y)$ are forbidden by Eq. (E10), the minimum of $\hat{\zeta}_0$ is indeed reached when $\hat{\omega}_0 = w(x, y)$. A similar reasoning holds for negative $\hat{\omega}_0$ satisfying Eq. (E9) so that $\hat{\zeta}_0$ is reached when and only when $\hat{\omega}_0 = w(x, y)$. Thus in both cases the relevant functions are obtained for $x = x_y^>$ or $x_y^<$. Denoting by $\hat{\zeta}_0^{\pm}(y, \hat{\omega}_0)$ the two functions $\hat{\zeta}_0(x = x_y^{\pm}, y, \hat{\omega}_0)$ and using Eqs. (E19) and (E4), we find after some algebra

$$\begin{aligned} \hat{\zeta}_0^{\pm}(y, \hat{\omega}_0) &= \frac{1}{3} + \frac{2}{3} \left(\frac{1 - 6\hat{\omega}_0 + 2(y^2 \pm 3\sqrt{y^2(2\hat{\omega}_0 - 8\hat{\omega}_0^2 + y^2)})}{1 + 8y^2} \right)^2, \\ &\text{with } y^2 \in [-2\hat{\omega}_0 + 8\hat{\omega}_0^2, 1]. \end{aligned} \quad (\text{E22})$$

We note that these functions do not depend on the sign of y . Starting from Eq. (E22) it is straightforward to determine the configurations where $\hat{\zeta}_0$ reaches its absolute minimum value $\frac{1}{3}$. One finds

$$\hat{\zeta}_0^+ = \frac{1}{3} \quad \text{iff } \hat{\omega}_0 = \frac{1}{6} + \frac{\sqrt{y^2}}{3}, \quad (\text{E23})$$

$$\hat{\zeta}_0^- = \frac{1}{3} \quad \text{iff } \hat{\omega}_0 = \frac{1}{6} - \frac{\sqrt{y^2}}{3}, \quad (\text{E24})$$

where these values of $\hat{\omega}_0$ always lie within the validity domain of Eq. (E22). Varying y^2 in $[0, 1]$ we see that $\hat{\zeta}_0$ reaches the value of $\frac{1}{3}$ through either $\hat{\zeta}_0^+$ or $\hat{\zeta}_0^-$ for any value of $\hat{\omega}_0$ in $[-\frac{1}{6}, +\frac{1}{2}]$, while $\frac{1}{3}$ is never reached when $\hat{\omega}_0 \in [-\frac{1}{4}, -\frac{1}{6}]$. Thus for the $[-\frac{1}{6}, +\frac{1}{2}]$ subdomain, the lower boundary $\hat{\zeta}_0^{\min}(\hat{\omega}_0)$ is simply given by

$$\hat{\zeta}_0^{\min}(\hat{\omega}_0) = \frac{1}{3}, \quad \hat{\omega}_0 \in \left[-\frac{1}{6}, +\frac{1}{2}\right]. \quad (\text{E25})$$

To treat the $[-\frac{1}{4}, -\frac{1}{6}]$ subdomain we first note from Eq. (E22) the obvious inequality,

$$\hat{\zeta}_0^-(y, \hat{\omega}_0) < \hat{\zeta}_0^+(y, \hat{\omega}_0), \quad \text{for all } \hat{\omega}_0 < 0. \quad (\text{E26})$$

The lower boundary for the portion $\hat{\omega}_0 \in [-\frac{1}{4}, -\frac{1}{6}]$ is thus to be found within the $\hat{\zeta}_0^-$ branch. A straightforward analytical study shows that $\hat{\zeta}_0^-(y, \hat{\omega}_0)$ is a strictly decreasing function of y^2 for any $\hat{\omega}_0 \in [-\frac{1}{4}, -\frac{1}{6}]$.¹⁹ It follows that the lower boundary $\hat{\zeta}_0^{\min}(\hat{\omega}_0)$ is given by $\hat{\zeta}_0^-(y, \hat{\omega}_0)$ at $y^2 = 1$ (strictly speaking at $y = -1$ since $\hat{\omega}_0 < 0$),

$$\hat{\zeta}_0^{\min}(\hat{\omega}_0) = \frac{1}{3} + \frac{2}{27} \left(1 - 2\hat{\omega}_0 - 2\sqrt{1 + 2\hat{\omega}_0 - 8\hat{\omega}_0^2}\right)^2, \quad \hat{\omega}_0 \in \left[-\frac{1}{4}, -\frac{1}{6}\right]. \quad (\text{E27})$$

3. Comments

The functions given in Eqs. (E21), (E25) and (E27) provide the full boundary in the $(\hat{\omega}_0, \hat{\zeta}_0)$ domain. Given that χ^{++} and $\text{Im}\chi^+$ are put to zero by a gauge choice, i.e., Eq. (3.96), we have proven *under the working assumption* $\text{Re}\chi^+ \equiv u = 0$ in Eq. (3.96) that this boundary is obtained when $y = \pm 1$ and $\sin^2 \theta_+ = 1$, that is, for $\text{Im}\chi^0 = \xi^+ = 0$ [cf. Eqs. (E1) and (3.102)]. This agrees with [33] where the domain was determined by a numerical scan. There is, however, more to the proofs we provided: $\sin^2 \theta_+ = 1$ is not only sufficient but also necessary; indeed, as one can see from the various steps of the proofs given above, all the inequalities and monotonicity are strict.

It is important to stress that there is *a priori* no simple reason to believe that the domain $(\hat{\omega}_0, \hat{\zeta}_0)$ will be identical to the full domain of $(\hat{\omega}, \hat{\zeta})$, i.e., when relaxing the working assumption $u = 0$. The necessity of $\sin^2 \theta_+ = 1$ proved instrumental while completing the determination of the domain when $u \neq 0$; see Sec. III F.

APPENDIX F: RESOLVED NAS CONDITIONS FOR EQS. (3.93) AND (3.94)

Here we give without proof the necessary and sufficient conditions on the α parameters in order for the trajectories $(\hat{\omega}(t), \hat{\zeta}(t))$ given by Eqs. (3.93) and (3.94) to go through a given point $(\hat{\omega}, \hat{\zeta})$:

$$\begin{aligned} & \left\{ \hat{\zeta} \geq \alpha_{AB} \vee \hat{\zeta} \geq \frac{3}{2} - \alpha_A \right\} \wedge \hat{\zeta} \geq \frac{2\alpha_A + 2\alpha_{AB}^2 - 3}{2\alpha_A + 4\alpha_{AB} - 5} \\ & \wedge \\ & \left\{ \hat{\omega} \times \alpha_{ABH} \geq 0 \vee \min\left\{0, \frac{1}{4}\beta_{AH}\right\} \leq \hat{\omega} \leq \max\left\{0, \frac{1}{4}\beta_{AH}\right\} \right\} \\ & \wedge \\ & \frac{1}{8} \left(\beta_{AH} - \sqrt{4\alpha_{ABH}^2 + \beta_{AH}^2} \right) \leq \hat{\omega} \leq \frac{1}{8} \left(\beta_{AH} + \sqrt{4\alpha_{ABH}^2 + \beta_{AH}^2} \right) \\ & \wedge \\ & (r_1 \hat{\omega}^2 + r_2 \hat{\omega} + r_3 \hat{\zeta} + r_4) \hat{\omega}^2 + (r_5 + r_6(\hat{\zeta} + 1) + r_7 \hat{\omega})(\hat{\zeta} - 1) = 0, \end{aligned} \quad (\text{F1})$$

¹⁹More specifically, we find that the derivative $\frac{\partial}{\partial y^2} \hat{\zeta}_0^-(y, \hat{\omega}_0)$ vanishes only when $y^2 = (1/4)(1 - 6\hat{\omega}_0)^2$, a value ≥ 1 for $\hat{\omega}_0 \in [-\frac{1}{4}, -\frac{1}{6}]$ that is outside the y^2 domain. Thus $\frac{\partial}{\partial y^2} \hat{\zeta}_0^-(y, \hat{\omega}_0)$ does not change sign in the considered domain of $\hat{\omega}_0$. This sign is determined by choosing any value of $y^2 \in [2(-\hat{\omega}_0 + 4\hat{\omega}_0^2), 1]$; e.g., for $y^2 = 2(-\hat{\omega}_0 + 4\hat{\omega}_0^2)$ it is given by $\text{sgn}\{-8\hat{\omega}_0(-1 + 8\hat{\omega}_0)^3(1 - 6\hat{\omega}_0 + 8\hat{\omega}_0^2) = -$ for $\hat{\omega}_0 \in [-\frac{1}{4}, -\frac{1}{6}]$.

with

$$\begin{aligned}
r_1 &= 4(\beta_A + 2\beta_{AB} - 2)^2, \\
r_2 &= 4(1 - \beta_{AB})(\beta_A + 2\beta_{AB} - 2)\beta_{AH}, \\
r_3 &= (\beta_A + 2\beta_{AB} - 2)(\beta_{AH}^2 - 4\alpha_{ABH}^2), \\
r_4 &= 2\alpha_{ABH}^2(8 - (\beta_A - 4)(\beta_{AB} - 3)) + ((\beta_{AB} - 2)^2 - \beta_A - 1)\beta_{AH}^2, \\
r_5 &= \frac{1}{8}(4\alpha_{ABH}^4(\beta_A - 4) - 2\alpha_{ABH}^2(3 + \beta_{AB})\beta_{AH}^2 - \beta_{AH}^4), \\
r_6 &= \frac{1}{16}(\beta_{AH}^2 + 4\alpha_{ABH}^2)^2, \\
r_7 &= \frac{1}{2}\beta_{AH}(4\alpha_{ABH}^2(\beta_A + \beta_{AB} - 1) + (1 - \beta_{AB})\beta_{AH}^2),
\end{aligned}$$

where we defined

$$\beta_X \equiv 2\alpha_X - 1, \quad X = A, AB, AH. \quad (\text{F2})$$

The first three lines in Eq. (F1) are the NAS conditions that ensure the existence of at least one real-valued t solution to Eq. (3.94) and at least one real-valued positive t solution to Eq. (3.93). The last condition in Eq. (F1) guarantees a common t solution to both Eqs. (3.93) and (3.94). Note that Eq. (F1) is always satisfied for $\hat{\omega} = 0$, $\hat{\zeta} = 1$ for all α parameters in the α potatoid, which can be seen in particular from Eq. (3.47). This corresponds to the fact that the point $(\hat{\omega} = 0, \hat{\zeta} = 1)$ is always reached when $t \rightarrow \infty$, as evident from Eqs. (3.93) and (3.94).

The α -parameters sets that are excluded by the $\omega - \zeta$ chips (see the discussion in Sec. IV and footnote 14) correspond to those that satisfy Eq. (F1) when substituting therein $\hat{\zeta}$ by $\hat{\zeta}_0^{\max}(\hat{\omega}) + \epsilon$ or by $\hat{\zeta}_0^{\min}(\hat{\omega}) - \epsilon$, with ϵ an arbitrarily small positive number [cf. Eqs. (3.113), (3.109)–(3.110b)].

APPENDIX G: NEW NAS POSITIVITY CONDITIONS FOR QUARTIC POLYNOMIALS ON $\mathbb{R}$

In this section we consider the general conditions on the set of *real* coefficients $a_{i=0,1,2,3,4}$ that are necessary and sufficient to ensure

$$P(\xi) > 0, \quad \forall \xi \in (-\infty, +\infty), \quad (\text{G1})$$

where $P(\xi)$ is a quartic polynomial:

$$P(\xi) \equiv a_0 + a_1\xi + a_2\xi^2 + a_3\xi^3 + a_4\xi^4. \quad (\text{G2})$$

Our derivation does not rely on the known form of the four roots of $P(\xi) = 0$, and will actually allow one to cast the conditions in a simpler and more compact form than the ones usually relied upon in the literature [42,61].

To achieve this we take a different path than just writing down the well-known expressions of the four roots of $P(\xi)$.

We are interested in determining the exact $\{a_i\}$ space region for which $P(\xi)$ is positive valued for any ξ in $(-\infty, +\infty)$. Recalling a classic theorem on positive definiteness of even degree polynomials defined on $\mathbb{R}$ and having all their coefficients real valued, if $P(\xi)$ satisfies Eq. (G1), then it can be written in the form

$$P(\xi) \equiv Q(\xi)^2 + R(\xi)^2, \quad \forall \xi \in (-\infty, +\infty) \quad (\text{G3})$$

with

$$Q(\xi) = x_1 + y_1\xi + z_1\xi^2 \quad \text{and} \quad R(\xi) = x_2 + y_2\xi + z_2\xi^2, \quad (\text{G4})$$

where x_i , y_i , and z_i denote real numbers.²⁰

The exact $\{a_i\}$ space is then defined by the NAS conditions on the a_i coefficients such that there exist real numbers x_i , y_i , and z_i satisfying Eq. (G3). To determine these conditions we find it useful to geometrize this statement. Introducing the vectors,

$$\mathbf{x} = (x_1, x_2), \quad \mathbf{y} = (y_1, y_2), \quad \mathbf{z} = (z_1, z_2), \quad (\text{G5})$$

the identification of the coefficients of each ξ monomial in Eq. (G3) leads to

²⁰Note that taking Q and R as in Eq. (G4) is more general than actually needed. Indeed, $P(\xi)$ will satisfy Eq. (G1) if and only if its four roots are nonreal complex valued; that is, $P(\xi)$ of the form $P(\xi) = r(\xi - s)(\xi - \bar{s})(\xi - t)(\xi - \bar{t}) = r|(\xi - s)(\xi - t)|^2$, with $\text{Im}(s), \text{Im}(t) \neq 0$, s, t and their complex conjugates $\bar{s}, \bar{t}$ being the four roots, and r a positive real number. Expanding this form as the squared modulus of a complex number leads to Eq. (G4) but with one of the two polynomials Q and R being only linear in ξ . The symmetric choice made in Eq. (G4) lends itself, however, to a more convenient geometric discussion. Its equivalence with the more specific case above results from the invariance of Eq. (G3) under any rigid rotation of the three vectors $\mathbf{x}$, $\mathbf{y}$, and $\mathbf{z}$ defined in Eq. (G5).

$$\|\mathbf{x}\|^2 = a_0, \quad (\text{G6})$$

$$\|\mathbf{z}\|^2 = a_4, \quad (\text{G7})$$

$$2\mathbf{x} \cdot \mathbf{y} = a_1, \quad (\text{G8})$$

$$2\mathbf{y} \cdot \mathbf{z} = a_3, \quad (\text{G9})$$

$$\|\mathbf{y}\|^2 = a_2 - 2\mathbf{x} \cdot \mathbf{z}, \quad (\text{G10})$$

so that the problem is equivalent to determining three vectors knowing some of their moduli and scalar products and relations among them. The NAS conditions on the a_i will thus be determined by requiring consistent moduli of and angles between the three vectors $\mathbf{x}$, $\mathbf{y}$, $\mathbf{z}$. Equations (G6) and (G7) imply trivially the NAS conditions for the existence of the moduli of $\mathbf{x}$ and $\mathbf{z}$, namely $a_0 \geq 0 \wedge a_4 \geq 0$. However, the strict inequality Eq. (G1) forbids $a_0 = 0$ and $a_4 = 0$ [in the first case $P(\xi = 0) = 0$ and in the second P is cubic and possesses at least one real root]. The conditions should thus read

$$a_0 > 0 \wedge a_4 > 0. \quad (\text{G11})$$

Rewriting Eq. (G10) as

$$\|\mathbf{y}\|^2 = a_2 - 2\sqrt{a_0 a_4} \cos(\widehat{\mathbf{x}, \mathbf{z}}) \quad (\text{G12})$$

and using the boundedness of the cosine one finds the *necessary* condition for the existence of the modulus of $\mathbf{y}$:

$$a_2 + 2\sqrt{a_0 a_4} \geq 0. \quad (\text{G13})$$

It should be stressed that while this condition is necessary to ensure the existence of at least one choice of the angle $(\widehat{\mathbf{x}, \mathbf{z}})$, not knowing the sign of a_2 , for which the modulus of $\mathbf{y}$ exists, Eqs. (G11) and (G13) are in general not sufficient to guarantee the existence of the vectors themselves (apart from the special case $a_1 = a_3 = 0$); one still has to check for the consistency of the three scalar products: Eqs. (G8), (G6), and (G10) lead to

$$a_1 = 2\sqrt{a_0} \sqrt{a_2 - 2\sqrt{a_0 a_4} \cos(\widehat{\mathbf{x}, \mathbf{z}}) \cos(\widehat{\mathbf{y}, \mathbf{x}})}, \quad (\text{G14})$$

and Eqs. (G9), (G7), and (G10) to

$$a_3 = 2\sqrt{a_4} \sqrt{a_2 - 2\sqrt{a_0 a_4} \cos(\widehat{\mathbf{x}, \mathbf{z}}) \cos(\widehat{\mathbf{y}, \mathbf{z}})}. \quad (\text{G15})$$

Again, using $-1 \leq \cos \leq 1$, one retrieves two necessary conditions from these two equations that can be summarized as

$$a_2 + 2\sqrt{a_0 a_4} \geq \max\left\{\frac{a_1^2}{4a_0}, \frac{a_3^2}{4a_4}\right\}. \quad (\text{G16})$$

These conditions are stronger than condition (G13). There is, however, a further constraint that correlates Eqs. (G14) and (G15), namely $(\widehat{\mathbf{y}, \mathbf{z}}) = (\widehat{\mathbf{y}, \mathbf{x}}) + (\widehat{\mathbf{x}, \mathbf{z}})$. This transforms Eqs. (G14) and (G15) into

$$\frac{a_1}{\sqrt{a_0}} \eta - \frac{a_3}{\sqrt{a_4}} = 2\epsilon_{yx} \sqrt{1 - \eta^2} \left(a_2 - \frac{a_1^2}{4a_0} - 2\eta \sqrt{a_0 a_4} \right)^{\frac{1}{2}}, \quad (\text{G17})$$

$$\frac{a_3}{\sqrt{a_4}} \eta - \frac{a_1}{\sqrt{a_0}} = 2\epsilon_{yz} \sqrt{1 - \eta^2} \left(a_2 - \frac{a_3^2}{4a_4} - 2\eta \sqrt{a_0 a_4} \right)^{\frac{1}{2}}, \quad (\text{G18})$$

where $\eta \equiv \cos(\widehat{\mathbf{x}, \mathbf{z}})$, and ϵ_{yx} (respectively, ϵ_{yz}) indicates the *relative* sign between $\sin(\widehat{\mathbf{y}, \mathbf{z}})$ and $\sin(\widehat{\mathbf{z}, \mathbf{x}})$ (respectively, between $\sin(\widehat{\mathbf{y}, \mathbf{x}})$ and $\sin(\widehat{\mathbf{x}, \mathbf{z}})$). Note also that Eqs. (G17) and (G18) are obtained from one another under the exchange $a_1 \leftrightarrow a_3$ and $a_0 \leftrightarrow a_4$. The invariance of these conditions as well as any other positivity condition such as Eq. (G16), under $(a_1 \leftrightarrow a_3, a_0 \leftrightarrow a_4)$, corresponds to the invariance of the positivity condition under the duality transformation $\xi \rightarrow \xi^{-1}$:

$$P(\xi) > 0, \quad \forall \xi \in (-\infty, +\infty) \Leftrightarrow \xi^4 P(\xi^{-1}) > 0, \\ \forall \xi \in (-\infty, +\infty).$$

When the necessary conditions (G11) and (G16) are verified, one still has to check for the existence of at least one η satisfying Eqs. (G17) and (G18). Moreover, η has to satisfy

$$\eta \in [-1, \min\{1, \eta^*\}], \quad (\text{G19})$$

where

$$\eta^* \equiv \frac{1}{2\sqrt{a_0 a_4}} \left(a_2 - \max\left\{\frac{a_1^2}{4a_0}, \frac{a_3^2}{4a_4}\right\} \right) \quad (\text{G20})$$

is the critical value above which at least one of the square roots in Eqs. (G17) and (G18) turns complex and thus becomes invalid.²¹ To study further the conditions for the existence of η we square both sides of Eq. (G17). This leads to a cubic equation in η ,

$$I(\eta) = \hat{I}, \quad (\text{G21})$$

²¹Note that a necessary condition for the existence of η is obviously $\eta^* \geq -1$, leading back to Eq. (G16).

where we define for later use

$$I(\eta) \equiv (2\sqrt{a_0 a_4}(2\sqrt{a_0 a_4}\eta - a_2)(\eta + 1) + a_1 a_3)(\eta - 1), \quad (\text{G22})$$

$$\hat{I} \equiv \frac{(\sqrt{a_0} a_3 - a_1 \sqrt{a_4})^2}{2\sqrt{a_0 a_4}}. \quad (\text{G23})$$

It is important to note that Eq. (G21) would equally result from squaring Eq. (G18) due to the invariance under the permutation $(a_1 \leftrightarrow a_3, a_0 \leftrightarrow a_4)$. It follows that (G21) encodes by itself the information contained in (G17) as well as that contained in (G18), except for the one that is lost by squaring, namely the signs $\epsilon_{yx}, \epsilon_{yz}$. This loss of information is, however, not problematic, as the signs can be retrieved by plugging back in Eqs. (G17) and (G18) whatever valid solutions for η are found by solving (G21). Moreover, the constraint that only the solutions satisfying Eqs. (G19) and (G20) are valid is implicitly embedded in Eq. (G21): Whenever a solution is found satisfying $\eta \in [-1, +1]$, it automatically satisfies (G19) and (G20). The reason is that squaring both sides of Eq. (G17) enforces the positivity of the term under the square root. We thus conclude that the sought-after NAS conditions are those which guarantee the existence of (at least one) real-valued η satisfying simultaneously (G21) and $\eta \in [-1, +1]$, together with Eq. (G11). The function $I(\eta)$ being a cubic polynomial in η , one can in principle solve $I(\eta) = \hat{I}$ which has at least one, and up to three, real-valued solutions. One could, of course, proceed numerically, but this is not our aim. On the other hand, extracting information from the analytical expressions of the three roots of this cubic equation is not particularly tractable. In fact, $I(\eta)$ has some interesting properties listed below that are straightforward to establish and will allow us to determine analytically the NAS conditions without solving the equation. A straightforward calculation shows that one *always* has the following:

- (a) $I(\eta = -1) = -2a_1 a_3 \leq \hat{I}$,
- (b) $I(\eta = +1) = 0 \leq \hat{I}$,
- (c) $I(\eta = \eta^*) \leq \hat{I}$.

[Property (c) is valid for the two configurations of the Max in Eq. (G20).] Being a cubic polynomial, $I(\eta)$ possesses at most two stationary points $\eta_{\pm}$ given by

$$\eta_{\pm} = \frac{a_2 \pm \sqrt{\Delta_0}}{6\sqrt{a_0 a_4}}, \quad (\text{G24})$$

where we define

$$\Delta_0 = a_2^2 + 12a_0 a_4 - 3a_1 a_3. \quad (\text{G25})$$

Moreover, the coefficient of η^3 in $I(\eta)$, Eq. (G22), being always positive [cf. Eq. (G11)], one also has that

- (d) when $\Delta_0 > 0$, i.e., $\eta_{\pm}$ exist and are distinct turning points, then $\eta_- < \eta_+$ and $I(\eta)$ increases monotonically

in $(-\infty, \eta_-] \cup [\eta_+, +\infty)$ and decreases monotonically in $]\eta_-, \eta_+[$; η_- and η_+ correspond to local maximum and minimum, respectively.

- (e) if it does not possess turning points ($\Delta_0 \leq 0$), $I(\eta)$ increases monotonically everywhere.

We can now write down the full NAS conditions. As clear from Eq. (G21), they correspond to ensuring all possible configurations in the (η, I) plane for which $I(\eta)$ crosses (at least once) the positive horizontal line $I = \hat{I}$ within the η domain given by Eq. (G19). We will refer to these configurations as existence configurations (EC). To proceed, we begin by identifying a set of four necessary conditions for the EC, then show that they form together with Eq. (G16) a set of sufficient conditions as well.

We note first that, when it exists, η_+ is always the position of the local minimum of $I(\eta)$. Properties (b) and (d) then imply that this minimum is necessarily negative whenever $\eta_+ \leq 1$. But since $\hat{I}$ is positive definite it follows that when η_+ lies in the relevant domain $[-1, +1]$, it never plays a role in the realization of the EC. We thus concentrate hereafter on η_- and η^* .

Properties (b) and (e) imply that the EC are never realized if $\Delta_0 \leq 0$, since in this case I would reach $\hat{I}$ only for $\eta > 1$, that is, outside its allowed domain [cf. Eq. (G19)] (except possibly for the nongeneric case where $a_3\sqrt{a_0} = a_1\sqrt{a_4}$);

$$\text{a necessary condition is thus } \Delta_0 > 0. \quad (\text{G26})$$

It follows that $\eta_{\pm}$ exist and are turning points. Similarly, properties (b) and (d) imply that the EC cannot be realized if $\eta_- > 1$ since again I cannot reach $\hat{I}$ within the allowed η domain Eq. (G19);

$$\text{a necessary condition is thus } \eta_- \leq 1. \quad (\text{G27})$$

Furthermore, the EC cannot be realized if $\eta_- > \eta^*$, since, according to property (d), $I(\eta)$ would in this case be monotonically increasing at η^* , and for it to reach $\hat{I}$ one would still have to increase η above η^* as implied by property (c), which is outside its allowed domain [cf. Eq. (G19)];

$$\text{a necessary condition is thus } \eta_- \leq \eta^*. \quad (\text{G28})$$

Since among the two turning points $\eta_{\pm}$, only η_- plays a role and is a local minimum; obviously if $I(\eta_-) < \hat{I}$, then EC are never realized in the relevant η domain. $I(\eta)$ still reaches $\hat{I}$ but outside this domain as a consequence of property (b);

$$\text{a necessary condition is thus } I(\eta_-) \geq \hat{I}. \quad (\text{G29})$$

It is now easy to see that the latter condition, in conjunction with the necessary conditions Eqs. (G11) and (G26)–(G28), would also form a set of sufficient conditions if and only if $\eta_- \geq -1$. Indeed, if $\eta_- < -1$, then to ensure that Eq. (G21) can be fulfilled for an η in the allowed domain would also require $I(\eta = -1) \geq \hat{I}$ which is generically in contradiction with property (a).

Fortunately, however, $\eta_- < -1$ is forbidden anyway by the necessary condition Eq. (G16). [This can be proven by showing, upon use of Eq. (G16) which implies in particular $a_2 + 6\sqrt{a_0 a_4} \geq 0$, that $\eta_- < -1$ would lead to $(a_2 + 2\sqrt{a_0 a_4})^2 < \frac{a_1^2 a_3^2}{16a_0 a_4}$ that contradicts Eq. (G16).] Thus η_- always satisfies $\eta_- \geq -1$.

We therefore conclude that adding the necessary condition Eq. (G16) to Eqs. (G11) and (G26)–(G29), one obtains a set of necessary and sufficient conditions. There is, however, more to it. One can show that (G28) actually implies Eq. (G16). The latter can hence be discarded without loss of generality.²² Putting everything together, the NAS conditions read finally

$$P(\xi) > 0, \forall \xi \in (-\infty, +\infty) \Leftrightarrow a_0 > 0 \wedge a_4 > 0 \wedge \Delta_0 > 0 \quad (\text{G30a})$$

$$\wedge$$

$$\sqrt{\Delta_0} + 2a_2 - \frac{3}{4} \max \left\{ \frac{a_1^2}{a_0}, \frac{a_3^2}{a_4} \right\} > 0 \quad (\text{G30b})$$

$$\wedge$$

$$\sqrt{\Delta_0} - a_2 + 6\sqrt{a_0 a_4} > 0 \quad (\text{G30c})$$

$$\wedge$$

$$2\Delta_0^{\frac{3}{2}} - \Delta_1 > 0, \quad (\text{G30d})$$

where we defined

$$\Delta_1 = 2a_2^3 + 27(a_0 a_3^2 + a_4 a_1^2) - 72a_0 a_2 a_4 - 9a_1 a_2 a_3. \quad (\text{G31})$$

Note that we have switched all the inequalities over to strict. The nongeneric equality cases can lead to different conditions. However, as argued at the beginning of Sec. III C, only strict positivity will be relevant. We have performed a numerical check of the above NAS conditions by scanning randomly over $a_0, a_4 \in [0, 100]$ and $a_1, a_2, a_3 \in [-100, 100]$ for 10^5 points, then solved numerically $P(\xi) = 0$ for each point and checked that whenever Eqs. (G30a)–(G30d) are satisfied, $P(\xi)$ has no real roots, and whenever one of the conditions is violated $P(\xi)$ has at least one real root. We also performed another nontrivial check based on the obvious fact that a translation of $P(\xi)$ to $P(\xi + \xi_0)$ for any $\xi_0 \in \mathbb{R}^*$ should not affect the positivity. It follows that the NAS conditions obtained after the translation, where the

modified coefficients $\tilde{a}_{0,1,2,3}$ depend explicitly on ξ_0 while $\tilde{a}_4 = a_4$, should be equivalent to the initial ones. Incidentally we find that ξ_0 cancels out in the modified Δ_0 and Δ_1 , which means that these two quantities can be reexpressed as functions of *differences* of the four roots of $P(\xi)$, and lead to the same conditions as before. In contrast, $\tilde{a}_0$ and the modified Eqs. (G30b) and (G30c) still depend on ξ_0 . That $\tilde{a}_0 > 0$ is valid when the initial NAS conditions Eqs. (G30a)–(G30d) are satisfied follows immediately from the fact that $\tilde{a}_0 = P(\xi_0)$. It remains to be checked that the involved dependence on ξ_0 in the modified Eqs. (G30b) and (G30c) does not lead to further NAS conditions. We verified that this is indeed the case through a numerical scan over 5×10^3 points in the a_i space satisfying Eqs. (G30a)–(G30c) followed by a scan over 2×10^3 values of ξ_0 for each of these points; the modified Eqs. (G30b) and (G30c) were found to be automatically satisfied for all values of ξ_0 .

In order to appreciate the simplification arrived at with Eqs. (G30a) and (G30d), one can compare with common knowledge [61,62]: Δ_0 and Δ_1 being defined as in [62], we note that the discriminant of $P(\xi)$ can be factorized as follows: $\Delta = (2\Delta_0^{\frac{3}{2}} - \Delta_1)(2\Delta_0^{\frac{3}{2}} + \Delta_1)/27$. Equation (G30d) requires the positivity of the first factor. It should then be clear that instead of relying on the signs of Δ , D , and P in

²²The proof consists in showing that (G28), more explicitly Eq. (G30b), together with Eq. (G30a), leads to Eq. (G16). We just sketch here the main steps: If $2a_2 - \frac{3}{4} \max \left\{ \frac{a_1^2}{a_0}, \frac{a_3^2}{a_4} \right\} > 0$, then obviously $a_2 - \frac{1}{4} \max \left\{ \frac{a_1^2}{a_0}, \frac{a_3^2}{a_4} \right\} > 0$ and Eq. (G16) is satisfied. If $2a_2 - \frac{3}{4} \max \left\{ \frac{a_1^2}{a_0}, \frac{a_3^2}{a_4} \right\} < 0$, then one can nonambiguously square the inequality in Eq. (G30b) and study it as a quadratic polynomial in a_2 . One then finds that it is satisfied only in a closed domain of a_2 for which Eq. (G16) is always satisfied whatever the configuration of the max.

the notations of [62], where the first two are complicated expressions, with an “and/or” structure as summarized in [62], we only need the signs of Δ_0 and just one of the two factors of Δ and two other simple relations involving Δ_0 with exclusively an “and” structure. Moreover, the “and” structure leads to unambiguous determination of necessary conditions. Another benefit of our approach is that it leads almost immediately to the conditions established in the following section.

APPENDIX H: NEW NAS POSITIVITY CONDITIONS FOR QUARTIC POLYNOMIALS ON $\mathbb{R}^+$

In this section we consider the NAS conditions on the parameters of the quartic polynomial Eq. (G2) that ensure its positivity for all non-negative ξ ,

$$P(\xi) > 0, \quad \forall \xi \in [0, +\infty). \quad (\text{H1})$$

Here, the form given by Eq. (G3), although sufficient, is no longer necessary. It should be replaced by the necessary and sufficient form [50–52]

$$P(\xi) \equiv Q(\xi)^2 + R(\xi)^2 + (A(\xi)^2 + B(\xi)^2)\xi, \quad (\text{H2})$$

where, since $P(\xi)$ is a quartic polynomial, Q and R keep the same form as in Eq. (G4), and

$$A(\xi) = u_1 + v_1\xi, \quad B(\xi) = u_2 + v_2\xi, \quad (\text{H3})$$

with u_i and v_i denoting real numbers. Equating the

coefficients of identical monomials in ξ on both sides of Eq. (H2), one finds that Eqs. (G6) and (G7) remain unchanged while Eqs. (G8)–(G10) are slightly modified:

$$\|\mathbf{x}\|^2 = a_0, \quad (\text{H4})$$

$$\|\mathbf{z}\|^2 = a_4, \quad (\text{H5})$$

$$2\mathbf{x} \cdot \mathbf{y} = a_1 - \|\mathbf{u}\|^2, \quad (\text{H6})$$

$$2\mathbf{y} \cdot \mathbf{z} = a_3 - \|\mathbf{v}\|^2, \quad (\text{H7})$$

$$\|\mathbf{y}\|^2 = a_2 - 2\mathbf{x} \cdot \mathbf{z} - 2\mathbf{u} \cdot \mathbf{v}, \quad (\text{H8})$$

where we introduced the vectors

$$\mathbf{u} = (u_1, u_2), \quad \mathbf{v} = (v_1, v_2). \quad (\text{H9})$$

The study carried out in Appendix G can thus be taken over unchanged to the present case with the following replacements:

$$a_1 \rightarrow a_1 - u^2, \quad (\text{H10})$$

$$a_3 \rightarrow a_3 - v^2, \quad (\text{H11})$$

$$a_2 \rightarrow a_2 - 2uvc, \quad (\text{H12})$$

where $u \equiv \|\mathbf{u}\|$, $v \equiv \|\mathbf{v}\|$, and $-1 \leq c \equiv \cos(\widehat{\mathbf{u}, \mathbf{v}}) \leq 1$ can be chosen arbitrarily in their domains. We thus reach the general solution to our problem:

The NAS conditions on $a_{i=0,1,2,3,4}$ for Eq. (H1) are obtained from Eqs. (G30a)–(G30d)

in which the replacements Eqs. (H10)–(H12) should lead to satisfied inequalities

$$\text{for at least one choice of } u \geq 0, v \geq 0, \text{ and } -1 \leq c \leq 1. \quad (\text{H13})$$

This shows in what sense Eq. (H1) is less constraining than Eq. (G1). Indeed, consider the domain $\mathcal{S}$ of all points in the $(a_0, a_1, a_2, a_3, a_4)$ space that satisfy conditions (G30a)–(G30d), and thus Eq. (G1). Obviously $\mathcal{S}$ will also satisfy Eq. (H1) since the latter is contained in Eq. (G1). But now, any point $(a_0, a'_1, a'_2, a'_3, a_4)$ lying outside of $\mathcal{S}$ and thus not satisfying Eq. (G1), will satisfy Eq. (H1) if it can be related to a point in $\mathcal{S}$ through the relations $a'_1 > a_1$

and $a'_3 > a_3$, and $a'_2 = a_2 + 2c\sqrt{a'_1 - a_1}\sqrt{a'_3 - a_3}$ with arbitrary $c \in [-1, +1]$. This is so because using Eqs. (H10)–(H12) will bring the point back into the $\mathcal{S}$ domain. The additional set of points $(a_0, a'_1, a'_2, a'_3, a_4)$ together with $\mathcal{S}$ lead obviously to a domain for which Eq. (H1) is satisfied larger than that for which Eq. (G1) is.

- [1] G. Aad *et al.* (ATLAS Collaboration), Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC, *Phys. Lett. B* **716**, 1 (2012).
- [2] S. Chatrchyan *et al.* (CMS Collaboration), Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC, *Phys. Lett. B* **716**, 30 (2012).
- [3] A. Pich, Flavour anomalies, *Proc. Sci. LHCP2019* (2019) 078 [arXiv:1911.06211].
- [4] J. Olsen, Experimental overview of Higgs physics, HIGGS2020, 2020, https://indico.cern.ch/event/900384/contributions/3795997/attachments/2129919/3586941/Experimental_Overview.pdf.
- [5] W. Konetschny and W. Kummer, Nonconservation of total lepton number with scalar bosons, *Phys. Lett. B* **70**, 433 (1977).
- [6] T. P. Cheng and L.-F. Li, Neutrino masses, mixings and oscillations in $SU(2) \times U(1)$ models of electroweak interactions, *Phys. Rev. D* **22**, 2860 (1980).
- [7] G. Lazarides, Q. Shafi, and C. Wetterich, Proton lifetime and fermion masses in an $SO(10)$ model, *Nucl. Phys. B* **181**, 287 (1981).
- [8] J. Schechter and J. W. F. Valle, Neutrino masses in $SU(2) \times U(1)$ theories, *Phys. Rev. D* **22**, 2227 (1980).
- [9] R. N. Mohapatra and G. Senjanovic, Neutrino Mass and Spontaneous Parity Nonconservation, *Phys. Rev. Lett.* **44**, 912 (1980).
- [10] R. N. Mohapatra and G. Senjanovic, Neutrino masses and mixings in gauge models with spontaneous parity violation, *Phys. Rev. D* **23**, 165 (1981).
- [11] H. Georgi and M. Machacek, Doubly charged Higgs bosons, *Nucl. Phys. B* **262**, 463 (1985).
- [12] M. S. Chanowitz and M. Golden, Higgs boson triplets with $M(W) = M(Z) \cos \theta_w$, *Phys. Lett.* **165B**, 105 (1985).
- [13] J. Gluza, M. Kordiaczynska, and T. Srivastava, Discriminating HTM and MLRSM models in colliders studies of a doubly charged Higgs boson pair production and its subsequent leptonic decays, arXiv:2006.04610.
- [14] R. Padhan, D. Das, M. Mitra, and A. Kumar Nayak, Probing doubly and singly charged Higgs bosons at the pp collider HE-LHC, *Phys. Rev. D* **101**, 075050 (2020).
- [15] R. Primulando, J. Julio, and P. Uttayarat, Scalar phenomenology in type-II seesaw model, *J. High Energy Phys.* **08** (2019) 024.
- [16] B. Fuks, M. Nemevšek, and R. Ruiz, Doubly charged Higgs boson production at hadron colliders, *Phys. Rev. D* **101**, 075022 (2020).
- [17] D. K. Ghosh, N. Ghosh, I. Saha, and A. Shaw, Revisiting the high-scale validity of the type II seesaw model with novel LHC signature, *Phys. Rev. D* **97**, 115022 (2018).
- [18] B. A. Ouazghour, A. Arhrib, R. Benbrik, M. Chabab, and L. Rahili, Theory and phenomenology of a two-Higgs-doublet type-II seesaw model at the LHC run 2, *Phys. Rev. D* **100**, 035031 (2019).
- [19] B. Ait-Ouazghour and M. Chabab, The Higgs Potential in 2HDM extended with a Real Triplet Scalar: A roadmap, arXiv:2006.12233.
- [20] P. Bhupal Dev, D. K. Ghosh, N. Okada, and I. Saha, 125 GeV Higgs boson and the type-II Seesaw model, *J. High Energy Phys.* **03** (2013) 150.
- [21] P. S. B. Dev, C. M. Vila, and W. Rodejohann, Naturalness in testable type II seesaw scenarios, *Nucl. Phys. B* **921**, 436 (2017).
- [22] P. Bhupal Dev and Y. Zhang, Displaced vertex signatures of doubly charged scalars in the type-II seesaw and its left-right extensions, *J. High Energy Phys.* **10** (2018) 199.
- [23] P. B. Dev, S. Khan, M. Mitra, and S. K. Rai, Doubly-charged Higgs boson at a future electron-proton collider, *Phys. Rev. D* **99**, 115015 (2019).
- [24] M. Frank, B. Fuks, K. Huitu, S. Mondal, S. K. Rai, and H. Waltari, Left-right supersymmetric option at a high-energy upgrade of the LHC, *Phys. Rev. D* **101**, 115014 (2020).
- [25] K. Huitu, A minimal supersymmetric left-right model, dark matter and signals at the LHC, *Eur. Phys. J. Special Topics* **229**, 3187 (2020).
- [26] M. Aaboud *et al.* (ATLAS Collaboration), Search for doubly charged Higgs boson production in multi-lepton final states with the ATLAS detector using proton–proton collisions at $\sqrt{s} = 13$ TeV, *Eur. Phys. J. C* **78**, 199 (2018).
- [27] ATLAS Collaboration, Search for doubly- and singly-charged Higgs bosons decaying into vector bosons in multi-leptons final states with the ATLAS detector using proton-proton collisions at $\sqrt{s} = 13$ TeV, ATLAS-CONF-2020-056, 2020.
- [28] S. Chatrchyan *et al.* (CMS Collaboration), A search for a doubly-charged Higgs boson in pp collisions at $\sqrt{s} = 7$ TeV, *Eur. Phys. J. C* **72**, 2189 (2012).
- [29] CMS Collaboration, A search for doubly-charged Higgs boson production in three and four lepton final states at $\sqrt{s} = 13$ TeV. CMS-PAS-HIG-16-036, 2017.
- [30] A. W. El Kaffas, W. Khater, O. M. Og Reid, and P. Osland, Consistency of the two Higgs doublet model and CP violation in top production at the LHC, *Nucl. Phys. B* **775**, 45 (2007).
- [31] A. Arhrib, R. Benbrik, M. Chabab, G. Moultaqa, M. Peyranere, J. Ramadan, and L. Rahili, The Higgs potential in the Type II Seesaw model, *Phys. Rev. D* **84**, 095005 (2011).
- [32] C. Bonilla, R. M. Fonseca, and J. W. F. Valle, Consistency of the triplet seesaw model revisited, *Phys. Rev. D* **92**, 075028 (2015).
- [33] K. Hartling, K. Kumar, and H. E. Logan, The decoupling limit in the Georgi-Machacek model, *Phys. Rev. D* **90**, 015007 (2014).
- [34] S. Blasi, S. De Curtis, and K. Yagyu, Effects of custodial symmetry breaking in the Georgi-Machacek model at high energies, *Phys. Rev. D* **96**, 015001 (2017).
- [35] M. E. Krauss and F. Staub, Perturbativity Constraints in BSM Models, *Eur. Phys. J. C* **78**, 185 (2018).
- [36] M. Maniatis, A. von Manteuffel, O. Nachtmann, and F. Nagel, Stability and symmetry breaking in the general two-Higgs-doublet model, *Eur. Phys. J. C* **48**, 805 (2006).
- [37] M. Maniatis and O. Nachtmann, Stability and symmetry breaking in the general three-Higgs-doublet model, *J. High Energy Phys.* **02** (2015) 058.
- [38] M. Maniatis and O. Nachtmann, Stability and symmetry breaking in the general n -Higgs-doublet model, *Phys. Rev. D* **92**, 075017 (2015).

- [39] I. Ivanov, Minkowski space structure of the Higgs potential in 2HDM, *Phys. Rev. D* **75**, 035001 (2007); Erratum, *Phys. Rev. D* **76**, 039902 (2007).
- [40] K. Kannike, Vacuum stability conditions from copositivity criteria, *Eur. Phys. J. C* **72**, 2093 (2012).
- [41] J. Chakraborty, P. Konar, and T. Mondal, Copositive criteria and boundedness of the scalar potential, *Phys. Rev. D* **89**, 095008 (2014).
- [42] K. Kannike, Vacuum stability of a general scalar potential of a few fields, *Eur. Phys. J. C* **76**, 324 (2016); Erratum, *Eur. Phys. J. C* **78**, 355 (2018).
- [43] I. P. Ivanov, M. Köpke, and M. Mühlleitner, Algorithmic boundedness-from-below conditions for generic scalar potentials, *Eur. Phys. J. C* **78**, 413 (2018).
- [44] M. Köpke, BFB, a Mathematica package to check boundedness of general Higgs potentials, Available at <https://git.io/vFQvi>, 2017.
- [45] K. Babu, I. Gogoladze, and S. Khan, Radiative electroweak symmetry breaking in Standard Model extensions, *Phys. Rev. D* **95**, 095013 (2017).
- [46] N. G. Deshpande and E. Ma, Pattern of symmetry breaking with two Higgs doublets, *Phys. Rev. D* **18**, 2574 (1978).
- [47] G. Chauhan, Vacuum stability and symmetry breaking in left-right symmetric model, *J. High Energy Phys.* **12** (2019) 137.
- [48] J. F. Gunion, R. Vega, and J. Wudka, Naturalness problems for $\rho = 1$ and other large one loop effects for a standard model Higgs sector containing triplet fields, *Phys. Rev. D* **43**, 2322 (1991).
- [49] M. Aoki and S. Kanemura, Unitarity bounds in the Higgs model including triplet fields with custodial symmetry, *Phys. Rev. D* **77**, 095009 (2008); **89**, 059902(E) (2014).
- [50] V. Powers and B. Reznick, Polynomials That are positive on an interval, *Trans. Am. Math. Soc.* **352**, 4677 (2000).
- [51] O. Benoist, Writing positive polynomials as sums of (few) squares, in *European Mathematical Society NewsLetter*, Issue 2017 (2017–9), pp. 8–13.
- [52] G. Pólya and G. Szegő, *Problems and Theorems in Analysis II. Classics in Mathematics* (Springer-Verlag, New York, 1976).
- [53] W. R. Inc., Mathematica, Version 11.3, Champaign, IL, 2018.
- [54] F. Staub, Reopen parameter regions in two-Higgs doublet models, *Phys. Lett. B* **776**, 407 (2018).
- [55] M. E. Krauss and F. Staub, Unitarity constraints in triplet extensions beyond the large s limit, *Phys. Rev. D* **98**, 015041 (2018).
- [56] M. Bando, T. Kugo, N. Maekawa, and H. Nakano, Improving the effective potential, *Phys. Lett. B* **301**, 83 (1993).
- [57] C. Ford, D. Jones, P. Stephenson, and M. Einhorn, The effective potential and the renormalization group, *Nucl. Phys. B* **395**, 17 (1993).
- [58] L. Chataignier, T. Prokopec, M. G. Schmidt, and B. Swiezewska, Single-scale renormalisation group improvement of multi-scale effective potentials, *J. High Energy Phys.* **03** (2018) 014.
- [59] J. Chakraborty, P. Konar, and T. Mondal, Constraining a class of B – L extended models from vacuum stability and perturbativity, *Phys. Rev. D* **89**, 056014 (2014).
- [60] J. Chakraborty, J. Gluza, T. Jelinski, and T. Srivastava, Theoretical constraints on masses of heavy particles in left-right symmetric models, *Phys. Lett. B* **759**, 361 (2016).
- [61] G. Ulrich and L. T. Watson, Positivity conditions for quartic polynomials, *SIAM J. Sci. Stat. Comput.* **15**, 528 (1994).
- [62] See for instance Quartic function, https://en.wikipedia.org/wiki/Quartic_function.