Messenger inflation in gauge mediation and super-WIMP dark matter

Shinsuke Kawai¹ and Nobuchika Okada²

¹*Department of Physics, Sungkyunkwan University, Suwon 440-746, Korea*

²*Department of Physics and Astronomy, University of Alabama, Tuscaloosa, Alabama 35487, USA*

(Received 26 March 2021; accepted 29 September 2021; published 26 October 2021)

We discuss phenomenological viability of a novel inflationary model in the minimal gauge mediated supersymmetry breaking scenario. In this model, cosmic inflation is realized in the flat direction along the messenger supermultiplets and a natural dark matter candidate is the gravitino from the out-of-equilibrium decay of the binolike neutralino at late times, which is called the super-WIMP scenario. The produced gravitino is warmish and can have a large free-streaming length; thus the cusp anomaly in the small scale structure formation may be mitigated. We show that the requirement of the Standard Model Higgs boson mass to be $m_{h^0} = 125.1$ GeV gives a relation between the spectrum of the cosmic microwave background and the messenger mass M . We find, for the e -folding number $N_e = 60$, the Planck 2018 constraints (TT, TE, EE + lowE + lensing + BK15 + BAO, 68% confidence level) give $M > 3.64 \times 10^7$ GeV. The gravitino dark matter mass is $m_{3/2} < 5.8$ GeV and the supersymmetry breaking scale Λ is found to be in the range $(1.28\text{--}1.33) \times 10^6$ GeV. Future cosmic microwave background observation is expected to give tighter constraints on these parameters.

DOI: [10.1103/PhysRevD.104.083539](https://doi.org/10.1103/PhysRevD.104.083539)

I. INTRODUCTION

The direct detection of gravitational waves by the Laser Interferometer Gravitational-wave Observatory and the discovery of the Higgs boson in the Large Hadron Collider have been major achievements in this century. For fundamental science, equally important are the null results, or the *absence* of discoveries. Today, we are confronted with several null results contrary to expectations of theorists. One of them is the nondetection of cosmic microwave background (CMB) B -mode polarization of primordial origin. Since many simple inflationary models predict a large tensor mode perturbation and hence a detectably large B -mode polarization, they are now strongly disfavored by the observation. Consequently, realistic inflationary model building requires a more sophisticated framework than, for example, the simple chaotic-type inflation models that were once considered standard. Another important null result we encountered concerns supersymmetric particles. It now seems unlikely that the hierarchy problem is solved by the low scale supersymmetry scenario, as was once hoped. Nevertheless, supersymmetry itself is an exquisite theoretical framework that resolves many of the shortcomings inherent in the

Standard Model (SM), such as the radiative stability of the scalar potential and the natural unification of the gauge couplings. It is also well known that supersymmetry is one of the few avenues that the local Lorentz-invariant quantum field theory can be consistently extended [1,2]. Furthermore, supersymmetry is an essential ingredient when constructing an ultraviolet complete theory, such as string theory. Thus from a theoretical perspective, it is reasonable to suppose that supersymmetry may well be realized at an energy scale beyond our current experimental reach, rather than to dismiss it as an outright irrelevance.

Motivated by the precision measurements of the CMB spectrum, we discuss, in this paper, a scenario of cosmic inflation within the minimal supersymmetric Standard Model (MSSM). In this scenario, inflation is implemented by known component fields, and its prediction is shown to be compatible with current CMB observations. A notable feature of this scenario is that it predicts gravitino dark matter produced nonthermally at late times, which is called the super-WIMP dark matter in the literature [3–6]. As supersymmetry is broken in the real world, any phenomenological model of supersymmetry must include a breaking mechanism. The gauge-mediated supersymmetry breaking (GMSB) [7–12] is a simple, elegant, and highly predictive mechanism of supersymmetry breaking, which is not afflicted by the conundrum of flavor-changing neutral currents. In GMSB, the breaking of supersymmetry is communicated from the hidden sector to the MSSM sector by a pair (or pairs) of the messenger supermultiplets Φ and $\bar{\Phi}$ that are charged under the SM gauge group and coupled

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

to a hidden sector field S , which is singlet under the SM gauge group. We investigate cosmic inflation along the flat direction for the messenger fields Φ and $\bar{\Phi}$. This possibility seems relatively unexplored. We use 125.1 GeV of the Higgs mass as a phenomenological input and assume that the present dark matter abundance consists of the super-WIMP. Then, this *messenger inflation* model predicts a relation between the CMB spectrum and the mass of the messengers.

In the next section we review the GMSB scenario and explain our notations. Super-WIMP dark matter, which is one of the key features of this scenario, is discussed in Sec. III. We describe our inflationary model and its predictions in Sec. IV, and we discuss constraints on the parameter space in Sec. V. In Sec. VI we conclude with some remarks.

II. MINIMAL GAUGE MEDIATED SUPERSYMMETRY BREAKING

In GMSB, the messenger fields are vectorlike pairs of chiral supermultiplets D, D^c and $L, \bar{L}$ transforming under the SM gauge group as in the following table:

	$SU(3)_c$	$SU(2)_L$	$U(1)_Y$
D^c	$\mathbf{3}^*$	$\mathbf{1}$	$+1/3$
$\bar{L}$	$\mathbf{1}$	$\mathbf{2}$	$-1/2$
D	$\mathbf{3}$	$\mathbf{1}$	$-1/3$
L	$\mathbf{1}$	$\mathbf{2}$	$+1/2$

In order to maintain the gauge coupling unification, it is customary to assume that the messengers come in multiplets

$$\bar{\Phi} = (D^c, \bar{L}), \quad \Phi = (D, L), \quad (1)$$

so that $\bar{\Phi}$ is in $\mathbf{5}^*$ and Φ is in $\mathbf{5}$ of a global $SU(5)$ symmetry group. More generally, the messenger fields may be treated as N_5 copies¹ of such pairs $(\bar{\Phi}_I, \Phi_I)$, $I = 1, 2, \dots, N_5$. The superpotential for the GMSB scenario is

$$W = (yS + M)\bar{\Phi}\Phi + W_{\text{hid}}(S), \quad (2)$$

where S is a SM gauge singlet that represents the dynamics of supersymmetry breaking, M is the mass of the messenger fermions, and y is a dimensionless coupling constant. The hidden sector superpotential $W_{\text{hid}}(S)$ is assumed to give a nonzero F -term for the S field so that the supersymmetry is broken in the vacuum,

$$\langle S \rangle = 0 + \theta^2 F_S, \quad \langle \Phi \rangle = \langle \bar{\Phi} \rangle = 0. \quad (3)$$

The messenger mass terms then become

¹ N_5 is the Dynkin index of the representation: $N_5 = 1$ for $\mathbf{5}^* + \mathbf{5}$, $N_5 = 3$ for $\mathbf{10}^* + \mathbf{10}$, $N_5 = 5$ for $\mathbf{24}^* + \mathbf{24}$, etc.

$$\mathcal{L} \supset -M\Psi_{\bar{\Phi}}\Psi_{\Phi} + \text{H.c.}$$

$$- (\bar{\Phi} \quad \Phi^*) \begin{pmatrix} M^2 & yF_S \\ yF_S & M^2 \end{pmatrix} \begin{pmatrix} \bar{\Phi}^* \\ \Phi \end{pmatrix}, \quad (4)$$

where $\bar{\Phi}$ and Φ represent the scalar components and $\Psi_{\bar{\Phi}}$ and Ψ_{Φ} are the corresponding fermionic counterparts. The effect of supersymmetry breaking is manifest as the eigenvalues for the scalar masses are seen to deviate from the fermionic masses. As the squared scalar masses need to be positive, we must have $M^2 > yF_S$. The leading contribution to the gaugino masses arises from one-loop,

$$M_a = N_5 \frac{\alpha_a}{4\pi} \Lambda, \quad (5)$$

whereas the soft masses for scalars are generated at two-loop,

$$m_i^2 = 2N_5\Lambda^2 \sum_{a=1}^3 C_a(R_i) \left(\frac{\alpha_a}{4\pi} \right)^2. \quad (6)$$

Here we included the multiplicities N_5 and introduced

$$\alpha_a \equiv \frac{g_a^2}{4\pi} \Lambda, \quad \Lambda \equiv \frac{yF_S}{M}. \quad (7)$$

The index $a = 1, 2, 3$ stands for $U(1)_Y, SU(2)_L, SU(3)_c$, and $C_a(R_i)$ are the quadratic Casimir for the representation R_i . The gravitino mass is

$$m_{3/2} = \frac{F_S}{\sqrt{3}M_P}, \quad (8)$$

where $M_P = 2.44 \times 10^{18}$ GeV is the reduced Planck mass.

The minimal GMSB scenario is highly predictive in the sense that the mass spectrum is determined by a handful of parameters

$$N_5, \quad M, \quad \Lambda, \quad \tan\beta, \quad \text{sign}\mu. \quad (9)$$

One may use the Higgs mass [13]

$$h_0 = 125.10 \pm 0.14 \text{ GeV} \quad (10)$$

to constrain the parameter space. In Table I we show the prediction of the minimal GMSB scenario for the mass parameters using the SOFTSUSY4.1.10 package [14], when N_5 , Λ , and $\tan\beta$ are given as the input parameters and M is fixed by the condition (10). The sign of μ is chosen to be positive. Furthermore, if the gravitino mass is known, from (7) and (8) we may find²

²This y represents the same degree of freedom as c_{grav} commonly used in the literature. In [15,16] which analyze the super-WIMP scenario after the discovery of the Higgs boson, y (n_{grav}) is set to unity.

TABLE I. The mass spectra in the units of GeV and the CMB parameters n_s and r . We used SOFTSUSY4.1.10 [14]. The three parameters N_5 , M , $\tan\beta$ are the input, and the remaining mGMSB parameter Λ is fixed by the Higgs mass $m_h = 125.1$ GeV. We assume $\mu > 0$. The gravitino mass $m_{3/2}$ is determined by (13) with $\Omega_{3/2}h^2 = 0.120$ in the super-WIMP scenario. The n_s and r values on the first column are for the minimally coupled case ($M = 1.87 \times 10^7$ for $N_e = 50$, $M = 1.46 \times 10^7$ for $N_e = 60$).

$\tan\beta$	10			45	
N_5	1			5	
M	10^7	10^8	10^9	10^{13}	10^{10}
Λ	1.262×10^6	1.300×10^6	1.318×10^6	1.287×10^6	3.157×10^5
h_0	125.1				
H_0	5924	6521	6919	7456	3680
A_0	5924	6521	6919	7456	3680
$H^\pm$	5924	6522	6920	7456	3681
$\tilde{g}$	8087	8263	8346	8087	9437
$\tilde{\chi}_{1,2}^0$	1764, 3290	1809, 3370	1832, 3407	1780, 3296	2176, 3893
$\tilde{\chi}_{3,4}^0$	4034, 4039	4648, 4650	5026, 5028	5372, 5374	3922, 4026
$\tilde{\chi}_{1,2}^\pm$	3290, 4039	3370, 4651	3407, 5028	3296, 5374	3893, 4027
$\tilde{u}, \tilde{c}_{1,2}$	$1.084 \times 10^4, 1.154 \times 10^4$	$1.080 \times 10^4, 1.157 \times 10^4$	$1.065 \times 10^4, 1.147 \times 10^4$	$9664, 1.060 \times 10^4$	8291, 8763
$\tilde{t}_{1,2}$	9721, 1.102×10^4	9441, 1.094×10^4	9097, 1.076×10^4	7622, 9702	7215, 8045
$\tilde{d}, \tilde{s}_{1,2}$	$1.076 \times 10^4, 1.154 \times 10^4$	$1.070 \times 10^4, 1.157 \times 10^4$	$1.054 \times 10^4, 1.147 \times 10^4$	$9460, 1.060 \times 10^4$	8237, 8763
$\tilde{b}_{1,2}$	$1.073 \times 10^4, 1.101 \times 10^4$	$1.067 \times 10^4, 1.093 \times 10^4$	$1.051 \times 10^4, 1.076 \times 10^4$	9416, 9700	7764, 8044
$\tilde{\nu}_{e,\mu}$	4386	4627	4812	5236	3224
$\tilde{\nu}_\tau$	4381	4621	4804	5222	3114
$\tilde{e}, \tilde{\mu}_{1,2}$	2344, 4387	2547, 4628	2734, 4813	3479, 5237	1766, 3226
$\tilde{\tau}_{1,2}$	2327, 4382	2525, 4622	2707, 4805	3438, 5223	1321, 3118
$m_{3/2}$	6.137	5.375	4.741	2.721	376.8
$\Omega_{3/2}h^2$	0.120				
ξ	(0)	0.06310	1.339	2.419×10^4	0.007754
n_s	(0.9418)	0.9606	0.9615	0.9616	0.9565
r ($N_e = 50$)	(0.3106)	0.01498	0.004713	0.004192	0.07526
ξ	(0)	0.08371	1.612	2.883×10^4	0.01218
n_s	(0.9512)	0.9673	0.9678	0.9678	0.9652
r ($N_e = 60$)	(0.2600)	0.008777	0.003270	0.002964	0.03858

$$y = \frac{\Lambda M}{\sqrt{3}M_P m_{3/2}}. \quad (11)$$

As we see in the next section, the requirement for the super-WIMP scenario that the nonthermally generated gravitino comprises the observed abundance of dark matter indeed determines the gravitino mass parameter. In the rest of this paper, for definiteness we choose $\mu > 0$ and consider the minimal GMSB with the super-WIMP dark matter scenario, where the prediction is entirely given by N_5 , M , and $\tan\beta$.

III. SUPER-WIMP DARK MATTER

In GMSB, the gauge mediation effects dominate over the gravity mediation effects. Since the latter contribution is of the order of the gravitino mass, we must have

$$m_{3/2} = \frac{F_S}{\sqrt{3}M_P} \ll \frac{\alpha_a}{4\pi} \Lambda. \quad (12)$$

The gaugino masses and the soft masses are both of the order of $\frac{\alpha_a}{4\pi} \Lambda$. Thus the gravitino is the lightest supersymmetric particle (LSP) in GMSB. The next lightest supersymmetric particle (NLSP) is either a neutralino or a stau. See Table I.

As the neutralinos are unstable, the freeze-out neutralino dark matter scenario that leads to the WIMP miracle in generic supersymmetry breaking models is not applicable to GMSB. However, the gravitino produced as a decay product of the NLSP neutralino is stable and hence is a good candidate of dark matter. In this gravitino dark matter scenario, dubbed the super-WIMP scenario [5,6], the number density of the gravitino dark matter particles is the same as that of the thermally produced long-lived neutralino NLSP. The relic abundance of the gravitino dark matter is then

$$\Omega_{3/2}h^2 = \Omega_{\tilde{\chi}_1^0}h^2 \times \left(\frac{m_{3/2}}{m_{\tilde{\chi}_1^0}} \right) = 0.120, \quad (13)$$

where we used the center value of the Planck 2018 result [17] $\Omega_c h^2 = 0.120 \pm 0.001$ for the dark matter abundance. For the binolike neutralino $\tilde{\chi}_1^0 \simeq \tilde{B}$, the abundance of the thermally produced relics is approximated by³

$$\Omega_{\tilde{\chi}_1^0} h^2 \simeq \frac{8.7 \times 10^{-11} \times (n+1) x_f^{n+1} \text{ GeV}^{-2}}{\sqrt{g_*} \sigma_{\tilde{B}}}, \quad (14)$$

where $n = 1$ for the p -wave and

$$x_f = X - \left(n + \frac{1}{2}\right) \ln X, \quad (15)$$

$$X \simeq \ln \left[0.19 \times (n+1) \frac{g}{\sqrt{g_*}} M_{\text{P}} m_{\tilde{\chi}_1^0} \sigma_{\tilde{B}} \right], \quad (16)$$

with $g = 2$ the helicity degrees of freedom for the NLSP neutralino $\tilde{\chi}_1^0$ (Majorana particle). We use the SM value $g_* = 106.75$ for the relativistic degrees of freedom, as the thermal production takes place well below the supersymmetry breaking scale. The pair annihilation cross section of the binolike neutralino $\sigma_{\tilde{B}}$ is dominated by the p -wave pair annihilation process through the exchange of the right-handed charged sleptons in the t -channel,

which, for a pure bino $\tilde{B}$ with the right-handed slepton exchange is evaluated as [21]

$$\sigma_{\tilde{B}} = \frac{3g_Y^4 \rho (1 + \rho^2)}{2\pi m_{\tilde{e}_R}^2 (1 + \rho)^4}, \quad (17)$$

where $\rho \equiv (m_{\tilde{\chi}_1^0}/m_{\tilde{e}_R})^2$. We evaluate the gauge coupling g_Y at the breaking scale using the SOFTSUSY4.1.10 package [14]; for the minimal GMSB parameters $\tan \beta = 10$, $N_5 = 1$, and $M = 10^{7-13}$, it is found to be $g_Y \simeq 0.367$.

The gravitino mass is then determined by the condition (13) for the super-WIMP scenario, and the Yukawa coupling parameter y is found from (11). The gravitino mass thus obtained is listed in Table I. Figure 1 shows the gravitino mass $m_{3/2}$ and the Yukawa coupling y as functions of the messenger mass M , when $N_5 = 1$, $\tan \beta = 10$, and $\mu > 0$. The gravitino mass stays within a few GeV range, whereas the Yukawa coupling is seen to vary exponentially. The perturbative limit $y \lesssim \mathcal{O}(1)$ corresponds to $M \lesssim 10^{13}$ GeV.

³This formula for the freeze-out relic abundance is well known [18]. We checked for our parameter choices that numerical results of the micrOMEGAS package [19,20] agree within a percent.

FIG. 1. The gravitino mass $m_{3/2}$ (black, scale on the left axis) and the Yukawa coupling y (red, scale on the right axis) as functions of the messenger fermion mass M . We use $N_5 = 1$, $\tan \beta = 10$. The Λ parameter is fixed by the condition that the Higgs mass is 125.10 GeV.

IV. MESSENGER INFLATION

The inflationary model is constructed by embedding the MSSM (including the messenger and the hidden sector) in supergravity. The part of the Lagrangian pertinent for the inflationary model is

$$\mathcal{L} \supset \int d^4\theta \phi^\dagger \phi \mathcal{K} + \left\{ \int d^2\theta \phi^3 W + \text{H.c.} \right\}, \quad (18)$$

where W is the superpotential given by (2) and the Kähler potential in the superconformal framework [22–28] is here assumed to be noncanonical,

$$\mathcal{K} = -3M_{\text{P}}^2 + |\tilde{\Phi}|^2 + |\Phi|^2 + |S|^2 - \frac{3}{2} \gamma (\tilde{\Phi} \Phi + \text{H.c.}) - \frac{\zeta}{M_{\text{P}}^2} |S|^4, \quad (19)$$

where γ and ζ are dimensionless parameters. The mechanism of GMSB discussed in Sec. II is essentially unaltered by this choice of the Kähler potential.

The supergravity embedding is similar to the one used in various supersymmetric extensions of the SM Higgs inflation model [29–39]. It was discussed in [40] that there is no direct analog of SM Higgs inflation in the MSSM, whereas in an extended model such as the next-to-minimal supersymmetric Standard Model (NMSSM), a supersymmetric version of the SM Higgs inflation model may be constructed. It would be interesting to point out that within GMSB, the MSSM alone can accommodate successful slow-roll inflation of the same type.

The scalar potential is computed once the superpotential (2) and the Kähler potential (19) are given. As discussed in [29,41], the instabilities that may exist in the direction of S are controlled by introducing a higher order term in the Kähler potential, that is, the last term of (19). The inflationary

FIG. 2. The scalar potential in the Einstein frame, for the $\tan\beta = 10$, $N_5 = 1$, $M = 10^8$ GeV case of Table I. The non-minimal coupling is chosen to be $\xi = 0.08371$ corresponding to $N_e = 60$ e -foldings, and the quartic parameter in the Kähler potential (19) is chosen to be $\zeta = 0.1$. The labels are in the $M_P = 1$ units. The left panel shows the potential in the S field direction, with the value of φ set at M_P , $5M_P$, and $20M_P$. The potential is seen to be stable at $S = 0$. The right panel shows the potential in the φ direction, with $S = 0$.

trajectory is then taken in the flat direction along the messenger multiplets, parametrized by

$$\bar{\Phi} = \Phi = \frac{1}{2}\varphi, \quad \langle S \rangle = 0. \quad (20)$$

In Fig. 2 we show an example of the scalar potential $V_E(\varphi, S)$ in the Einstein frame, for the $\tan\beta = 10$, $N_5 = 1$, $M = 10^8$ GeV, and $N_e = 60$ case of Table I. To be concrete, we use the Polonyi superpotential $W_{\text{hid}}(S) = \Lambda_S^2 S$, where $\Lambda_S \equiv \sqrt{F_S}$, for the hidden sector (2), and the quartic parameter is chosen to be $\zeta = 0.1$. When the hidden sector field is stabilized at $S = 0$, the scalar part of the action is

$$S_{\text{scalar}} = \int d^4x \sqrt{-g} \left[\frac{M_P^2 + \xi\varphi^2}{2} R - \frac{1}{2}(\partial\varphi)^2 - V_J(\varphi) \right], \quad (21)$$

where R is the scalar curvature, $\xi \equiv \frac{\gamma}{4} - \frac{1}{6}$, and

$$V_J(\varphi) = \frac{y^2}{16}\varphi^4 + M^2\varphi^2 \frac{4M_P^2 - (\frac{1}{6} + 2\gamma)\varphi^2}{8M_P^2 - \gamma(2 - 3\gamma)\varphi^2}. \quad (22)$$

The second term of (22) is negligible when $y\varphi \gg M$, which is the case of our interest since $M/y \sim 10^{13}$ GeV (see Fig. 1) and $\varphi \gtrsim 10^{-2}M_P$ during inflation. Thus the effective theory of messenger inflation is the nonminimally coupled φ^4 inflation model, that includes the minimally coupled φ^4 model and the SM Higgs inflation model as special cases.

The prediction of the nonminimal φ^4 model is understood in the framework of the standard slow roll paradigm [42]. Transforming the Jordan frame action (21) into the Einstein frame by Weyl rescaling of the metric, the field φ has a deformed potential

FIG. 3. The prediction of the messenger inflation model for the primordial tilt n_s and the tensor-to-scalar ratio r . The plots for the e -folding numbers $N_e = 50$ and 60 are shown, with the Planck 2018 1- and 2- σ constraints indicated on the background. The dashed line is $r = \frac{16}{3}(1 - n_s)$, corresponding to the minimally coupled ($\xi = 0$) case.

$$V_E(\varphi) = \frac{y^2}{16} \frac{\varphi^4}{(M_P^2 + \xi\varphi^2)^2}. \quad (23)$$

The field φ has a noncanonical kinetic term in the Einstein frame and is related to the canonically normalized field $\hat{\varphi}$ by

$$d\hat{\varphi} = \frac{\sqrt{M_P^2 + \xi\varphi^2 + 6\xi^2\varphi^2}}{M_P^2 + \xi\varphi^2} d\varphi. \quad (24)$$

One may now introduce the slow roll parameters to analyze the model and obtain the prediction for the cosmological parameters.

The model contains two real parameters ξ and y . We use the normalization of the scalar amplitude⁴ to determine the nonminimal coupling ξ . Then, apart from the e -folding number N_e , the spectrum is entirely given by a single parameter, which may be chosen to be the Yukawa coupling y .

In Sec. II and Sec. III we saw that the minimal GMSB with the successful super-WIMP dark matter scenario is controlled by a set of parameters

$$N_5, \quad M, \quad \tan\beta. \quad (25)$$

The Yukawa coupling (11) is also determined by them. Thus, for a given e -folding number N_e , the CMB spectrum of this inflationary model is uniquely determined by the parameter set (25). Figure 3 shows the primordial tilt n_s and the tensor-to-scalar ratio r , when $N_5 = 1$ and $\tan\beta = 10$, as M is varied. The scalar self-coupling for the minimally coupled φ^4 inflation corresponds to $M = 1.87 \times 10^7$ GeV

⁴In numerics we used the Planck 2018 [17] TT, TE, EE + lowE + lensing + BAO value $A_s = \exp(3.047) \times 10^{-10}$ at the pivot scale $k_0 = 0.05 \text{ Mpc}^{-1}$.

for $N_e = 50$ and $M = 1.46 \times 10^7$ GeV for $N_e = 60$, and thus, the messenger mass M giving the Planck-normalized scalar amplitude must be larger than these values.

The constraints on the CMB spectrum give the lower bound on the messenger mass M . We find, from the Planck 2018 results (TT + TE + EE + lowE + lensing + BK15 + BAO) [43],

$$M > 4.40 \times 10^8 \text{ GeV} \quad (68\% \text{ C.L.}), \quad (26)$$

$$M > 5.78 \times 10^7 \text{ GeV} \quad (95\% \text{ C.L.}) \quad (27)$$

for e -folding number $N_e = 50$, and

$$M > 3.64 \times 10^7 \text{ GeV} \quad (68\% \text{ C.L.}), \quad (28)$$

$$M > 3.01 \times 10^7 \text{ GeV} \quad (95\% \text{ C.L.}) \quad (29)$$

for $N_e = 60$. We saw, in Fig. 1, that perturbativity $y \lesssim \mathcal{O}(1)$ requires $M \lesssim 10^{13}$ GeV. In the context of nonminimally coupled φ^4 inflation, it is argued, e.g., in [44–47], that unitarity is violated for too large ξ (see also [48], however). In our case, this gives another upper bound for the messenger mass M . If the unitarity bound is given by $\xi \lesssim 100$, for example, we must have $M \lesssim 10^{11}$ GeV for both $N_e = 50$ and $N_e = 60$, which is stronger than the one given by the perturbativity requirement.

V. THEORETICAL AND OBSERVATIONAL CONSTRAINTS

Let us now discuss further details of the model and its phenomenological consistency. Our main focus is on the $N_5 = 1$ and $\tan \beta = 10$ case, except in Sec. V E 1 where the viability of the stau NLSP scenario is discussed.

A. Reheating temperature

Although the messenger fields are coupled to the SM matter fields by the SM gauge interactions, the decay channel of the messenger inflaton is through a Yukawa interaction, if such an interaction exists at all. Indeed, the $\bar{\Phi}$ field is in the $\mathbf{5}^*$ representation of $SU(5)$, and recalling that this is the same representation as that of the SM fermions (the lepton doublets and the down-type quark singlets) in the Georgi-Glashow $SU(5)$ grand unified theory, a gauge invariant $\mathbf{5}^* \cdot \mathbf{10} \cdot \mathbf{5}^*$ Yukawa interaction can exist for $\bar{\Phi}$. Using the component fields (1), the superpotential responsible for the decay of the $\bar{\Phi}$ would be

$$W \supset y_L \bar{L} H_d e^c + y_D D^c H_d Q, \quad (30)$$

where e^c and Q are the MSSM fields in $\mathbf{10}$ and H_d is the down-type Higgs in $\mathbf{5}^*$ of $SU(5)$. Although this Yukawa interaction breaks the Z_2 symmetry that exchanges Φ and $\bar{\Phi}$, it has negligible effects on the GMSB mechanism.

The reheating temperature may then be evaluated for the perturbative decay $\bar{L} \rightarrow \tilde{H}_d e^c$ through the y_L interaction.⁵ The condition that the decay width

$$\Gamma \simeq \frac{y_L^2}{8\pi} M \quad (31)$$

becomes comparable to the Hubble parameter gives

$$T_{\text{rh}} \simeq \frac{y_L}{2\pi} \sqrt{MM_{\text{P}}} \left(\frac{45}{2g_*} \right)^{1/4}. \quad (32)$$

The reheating temperature is thus controlled by the coupling y_L .

B. Thermal gravitino production

The gravitino is produced also by the scattering of particles in the thermal plasma during the radiation dominated era after inflation. Its relic abundance is evaluated as [52–54]

$$\Omega_{3/2}^{\text{TP}} h^2 \sim 0.3 \left(\frac{T_{\text{rh}}}{10^8 \text{ GeV}} \right) \left(\frac{1 \text{ GeV}}{m_{3/2}} \right) \left(\frac{M_3}{1 \text{ TeV}} \right)^2, \quad (33)$$

where M_3 is the gluino mass. For the successful super-WIMP dark matter scenario, this contribution should be subdominant to the nonthermally produced relic abundance (13). For $N_5 = 1$ and $\tan \beta = 10$, the gluino mass is $M_3 \sim 10$ TeV. As the gravitino mass range of our interest is $m_{3/2} \sim 1$ GeV, the thermally produced gravitino is negligible when

$$T_{\text{rh}} \lesssim 10^6 \text{ GeV}. \quad (34)$$

On the other hand, the NLSP neutralinos of mass $m_{\tilde{\chi}_1^0} \sim$ a few TeV need to be produced from the thermal plasma and thus

$$T_{\text{rh}} \gtrsim \text{a few TeV}. \quad (35)$$

The reheating temperature of this scenario must satisfy both (35) and (34). For a given messenger mass M , the constraints on the reheating temperature give bounds on the Yukawa coupling y_L in (30). For example, $T_{\text{rh}} \sim 10^5$ GeV is a reasonable value of reheating temperature in our scenario with $M \sim 10^8$ GeV and $y_L \sim 10^{-8}$.

C. Big bang nucleosynthesis

Another important requirement for the success of the super-WIMP scenario is that the big bang nucleosynthesis is not disturbed by the late time decay of the neutralino NLSP [55]. The lifetime of the binolike neutralino is evaluated for the decay modes $\tilde{B} \rightarrow \gamma \tilde{G}$ as [5,6]

⁵We neglect nonlinear effects [49–51] for simplicity.

$$\tau_{\tilde{B}} \simeq 0.74 \times \left(\frac{m_{3/2}}{1 \text{ GeV}} \right)^2 \left(\frac{1 \text{ TeV}}{m_{\tilde{B}}} \right)^5 \text{ s}. \quad (36)$$

In our scenario with $N_5 = 1$ and $\tan \beta = 10$, the lifetime of the NLSP neutralino is $\tau_{\tilde{B}} < 1$ s. Thus the NLSP neutralino decays before the big bang nucleosynthesis commences ($\simeq 1$ s), leaving the big bang nucleosynthesis intact.

D. Gravitino free streaming length

As the gravitino is much lighter than the NLSP neutralino, the gravitino is energetic when it is produced. The production occurs at late times, when the Hubble expansion rate is already small and the effect of redshift on the gravitino is not significant. Thus the nonthermally produced gravitino has a relatively long free streaming length [56,57],

$$\lambda_{\text{FS}} = \int_{\tau_{\tilde{B}}}^{t_{\text{eq}}} \frac{v(t)}{a(t)} dt \simeq 0.18 \times \left(\frac{1 \text{ TeV}}{m_{\tilde{B}}} \right)^{3/2} \text{ Mpc}. \quad (37)$$

Although the cold dark matter scenario is known to be successful in explaining the structure formation at large scales $\gtrsim 1$ Mpc, recent studies of N -body simulation suggest that the structure formation at small scales favors warm dark matter, with the free streaming length $\lambda_{\text{FS}} \simeq 0.1$ Mpc [58]. It can be seen from Table I and (37) that our model indeed gives $\lambda_{\text{FS}} \simeq 0.1$ Mpc, preferred by the small scale structure formation.⁶

E. Competing dark matter scenarios

It is well known that in some parameter region of the minimal GMSB scenario, a stau, rather than a neutralino, can be the NLSP. An example is shown on the rightmost column in Table I. Since we are studying the supersymmetric model, some charged configuration in a flat direction, called a Q -ball, can also be stable and behave as dark matter. Let us briefly comment on these two possibilities below.

1. Stau NLSP scenario

As the gaugino mass (5) is proportional to N_5 and the soft scalar mass (6) is proportional to $\sqrt{N_5}$, the scalar leptons become relatively lighter than neutralinos when N_5 is large. The lightest stau $\tilde{\tau}_1$, instead of the lightest neutralino χ_1^0 , may then become the NLSP.⁷ An example of the parameters and the mass spectrum giving the 125.1 GeV Higgs boson mass is shown in Table I. In this case, the late time decay of the stau NLSP produces the gravitino, which may be considered as super-WIMP dark matter.

Similar to the neutralino NLSP case (13), the relic abundance of the nonthermally produced gravitino is related to that of the stau NLSP $\Omega_{\tilde{\tau}_1} h^2$ by

$$\Omega_{3/2} h^2 = \Omega_{\tilde{\tau}_1} h^2 \times \left(\frac{m_{3/2}}{m_{\tilde{\tau}_1}} \right). \quad (38)$$

For the (purely right handed) stau, the dominant channels of annihilation are through the electromagnetic interaction into two photons and the t -channel annihilation into two taus with bino exchange:

of which the latter contribution is subdominant except when the neutralino mass $m_{\tilde{B}}$ is very close to the stau mass $m_{\tilde{\tau}_1}$ [62]. For the first diagram, the annihilation cross section is $\sigma_{\tilde{\tau}_1} = 4\pi\alpha_{\text{em}}^2/m_{\tilde{\tau}_1}^2$, where $\alpha_{\text{em}} = g_{\text{em}}^2/4\pi$ and g_{em} is the electromagnetic coupling constant. The stau NLSP abundance $\Omega_{\tilde{\tau}_1} h^2$ is evaluated using the contribution from the first diagram, by equations similar to (14), (15), (16) with $n = 0$ (s -wave annihilation), $g = 2$, and $\sigma_{\tilde{B}}$, $m_{\tilde{\chi}_1^0}$ replaced by $\sigma_{\tilde{\tau}_1}$, $m_{\tilde{\tau}_1}$.

As the gravitino relic abundance is given by (38), the gravitino mass is determined by the condition $\Omega_{3/2} h^2 = 0.120$; that is, the gravitino produced by the late time decay of the stau NLSP comprises the total abundance of the present dark matter. For example, we find $m_{3/2} = 376.8$ GeV when $\tan \beta = 45$, $N_5 = 5$, and $M = 10^{10}$ GeV, as listed in Table I. The gravitino mass of a few hundred GeV and the stau mass of $\sim \text{TeV}$, however, are excluded by the constraints from the big bang nucleosynthesis [63]. Although the parameters of the minimal GMSB may be adjusted, the gravitino mass and the stau mass stay in the same order in the stau NLSP case. It thus seems difficult to construct a stau NLSP version of this inflationary scenario satisfying phenomenological requirements.

2. Q -ball dark matter

The reheating temperature (34) is lower than what is required for generic (nonresonant) leptogenesis. In the supersymmetric Standard Model, it is then natural to suppose that the baryon asymmetry of the Universe is generated along a charged flat direction, i.e., by the Affleck-Dine mechanism [64]. It has been pointed out [65] that some charged configurations along a flat direction, called Q -balls, can be stable and hence considered as dark matter. The conditions for Q -ball dark matter, namely the stability and the correct relic abundance, are studied, for example, in [66]. There are known to be two types of Q -balls, called the gauge mediation type and the “new” type. For both types, the parameter ranges for the Q -balls to be dark matter are very different from the parameter range of our scenario; for the reheating temperature (34) and the gravitino mass of a few GeV, the Q -balls are actually unstable, and they cannot remain as dark matter today.

⁶The effects of the gravitino on the Hubble tension are discussed recently in [59].

⁷The stau NLSP scenario may also arise from the effects of hidden sector renormalization [60,61].

VI. FINAL REMARKS

In this paper we discussed the possibilities that the messenger fields in gauge mediation may be identified as the inflaton field for cosmic inflation. The requirements that the Higgs boson mass is 125.1 GeV and the gravitino dark matter has the correct relic abundance make the minimal GMSB super-WIMP scenario extremely predictive. In particular, we have shown that there appears a direct relation between the prediction of the CMB spectrum and the messenger mass scale, as indicated in Fig. 3.

The theoretical framework of this inflationary scenario is minimalistic. It is the MSSM with the minimal GMSB. It is amusing to observe that the phenomenologically viable and observationally compatible inflationary scenario can be constructed in such a simple and well-known framework. In a sense, the model presented in this paper may be considered as the simplest supersymmetric analog of the SM Higgs inflation model. Of course, the relatively large

125.1 GeV Higgs boson mass indicates that the supersymmetric particles (except the gravitino) are heavy, so that testing this scenario by collider experiments is difficult. In cosmology, there are many projects for CMB observation; for example, the CMB B -mode observation by the LiteBIRD satellite experiments [67] targets the tensor-to-scalar ratio of order $\Delta r \lesssim 0.001$. Figure 3 suggests that our scenario may be falsified by such experiments; otherwise the value of the tensor-to-scalar ratio r will predict the scale of supersymmetry breaking.

ACKNOWLEDGMENTS

This work was supported in part by the National Research Foundation of Korea Grant-in-Aid for Scientific Research Grant No. NRF-2018R1D1A1B07051127, the NRF-JSPS Collaboration “String Axion Cosmology” (S. K.) and by the United States Department of Energy Grant No. DE-SC0012447 (N. O.).

-
- [1] S. R. Coleman and J. Mandula, All possible symmetries of the S matrix, *Phys. Rev.* **159**, 1251 (1967).
 - [2] R. Haag, J. T. Lopuszanski, and M. Sohnius, All possible generators of supersymmetries of the s matrix, *Nucl. Phys.* **B88**, 257 (1975).
 - [3] L. Covi, J. E. Kim, and L. Roszkowski, Axinos as Cold Dark Matter, *Phys. Rev. Lett.* **82**, 4180 (1999).
 - [4] L. Covi, H.-B. Kim, J. E. Kim, and L. Roszkowski, Axinos as dark matter, *J. High Energy Phys.* **05** (2001) 033.
 - [5] J. L. Feng, A. Rajaraman, and F. Takayama, Superweakly Interacting Massive Particles, *Phys. Rev. Lett.* **91**, 011302 (2003).
 - [6] J. L. Feng, S. Su, and F. Takayama, Supergravity with a gravitino LSP, *Phys. Rev. D* **70**, 075019 (2004).
 - [7] M. Dine and W. Fischler, A phenomenological model of particle physics based on supersymmetry, *Phys. Lett. B* **110**, 227 (1982).
 - [8] C. R. Nappi and B. A. Ovrut, Supersymmetric Extension of the $SU(3) \times SU(2) \times U(1)$ Model, *Phys. Lett.* **113B**, 175 (1982).
 - [9] L. Alvarez-Gaume, M. Claudson, and M. B. Wise, Low-energy supersymmetry, *Nucl. Phys.* **B207**, 96 (1982).
 - [10] M. Dine and A. E. Nelson, Dynamical supersymmetry breaking at low-energies, *Phys. Rev. D* **48**, 1277 (1993).
 - [11] M. Dine, A. E. Nelson, and Y. Shirman, Low-energy dynamical supersymmetry breaking simplified, *Phys. Rev. D* **51**, 1362 (1995).
 - [12] M. Dine, A. E. Nelson, Y. Nir, and Y. Shirman, New tools for low-energy dynamical supersymmetry breaking, *Phys. Rev. D* **53**, 2658 (1996).
 - [13] P. A. Zyla *et al.* (Particle Data Group), Review of particle physics, *Prog. Theor. Exp. Phys.* **2020**, 083C01 (2020).
 - [14] B. C. Allanach, SOFTSUSY: a program for calculating supersymmetric spectra, *Comput. Phys. Commun.* **143**, 305 (2002).
 - [15] N. Okada, SuperWIMP dark matter and 125 GeV Higgs boson in the minimal GMSB, [arXiv:1205.5826](https://arxiv.org/abs/1205.5826).
 - [16] J. L. Feng, Z. Surujon, and H.-B. Yu, Confluence of constraints in gauge mediation: The 125 GeV Higgs boson and goldilocks cosmology, *Phys. Rev. D* **86**, 035003 (2012).
 - [17] N. Aghanim *et al.* (Planck Collaboration), Planck 2018 results. VI. Cosmological parameters, *Astron. Astrophys.* **641**, A6 (2020).
 - [18] E. W. Kolb and M. S. Turner, *The Early Universe*, Frontiers in Physics Vol. 69 (Addison-Wesley, Redwood City, 1990).
 - [19] G. Belanger, F. Boudjema, A. Pukhov, and A. Semenov, MicroOMEGAs: A program for calculating the relic density in the MSSM, *Comput. Phys. Commun.* **149**, 103 (2002).
 - [20] G. Belanger, F. Boudjema, A. Pukhov, and A. Semenov, MicroOMEGAs:: Version 1.3, *Comput. Phys. Commun.* **174**, 577 (2006).
 - [21] N. Arkani-Hamed, A. Delgado, and G. F. Giudice, The well-tempered neutralino, *Nucl. Phys.* **B741**, 108 (2006).
 - [22] M. Kaku, P. Townsend, and P. van Nieuwenhuizen, Properties of conformal supergravity, *Phys. Rev. D* **17**, 3179 (1978).
 - [23] W. Siegel and S. J. Gates Jr, Superfield supergravity, *Nucl. Phys.* **B147**, 77 (1979).
 - [24] E. Cremmer, S. Ferrara, L. Girardello, and A. Van Proeyen, Yang-Mills theories with local supersymmetry: Lagrangian, transformation laws and SuperHiggs effect, *Nucl. Phys.* **B212**, 413 (1983).
 - [25] S. Ferrara, L. Girardello, T. Kugo, and A. Van Proeyen, Relation between different auxiliary field formulations of

- $N = 1$ supergravity coupled to matter, *Nucl. Phys.* **B223**, 191 (1983).
- [26] T. Kugo and S. Uehara, Improved superconformal gauge conditions in the $N = 1$ supergravity {Yang-Mills} matter system, *Nucl. Phys.* **B222**, 125 (1983).
- [27] T. Kugo and S. Uehara, Conformal and poincare tensor calculi in $N = 1$ supergravity, *Nucl. Phys.* **B226**, 49 (1983).
- [28] T. Kugo and S. Uehara, $N = 1$ superconformal tensor calculus: Multiplets with external lorentz indices and spinor derivative operators, *Prog. Theor. Phys.* **73**, 235 (1985).
- [29] S. Ferrara, R. Kallosh, A. Linde, A. Marrani, and A. Van Proeyen, Superconformal symmetry, NMSSM, and inflation, *Phys. Rev. D* **83**, 025008 (2011).
- [30] M. Arai, S. Kawai, and N. Okada, Higgs inflation in minimal supersymmetric SU(5) GUT, *Phys. Rev. D* **84**, 123515 (2011).
- [31] C. Pallis and N. Toumbas, Non-minimal Higgs inflation and non-thermal leptogenesis in a supersymmetric pati-salam model, *J. Cosmol. Astropart. Phys.* **12** (2011) 002.
- [32] M. Arai, S. Kawai, and N. Okada, Supersymmetric standard model inflation in the Planck era, *Phys. Rev. D* **86**, 063507 (2012).
- [33] M. Arai, S. Kawai, and N. Okada, Higgs-lepton inflation in the supersymmetric minimal seesaw model, *Phys. Rev. D* **87**, 065009 (2013).
- [34] M. Arai, S. Kawai, and N. Okada, Supersymmetric B-L inflation near the conformal coupling, *Phys. Lett. B* **734**, 100 (2014).
- [35] S. Kawai and N. Okada, TeV scale seesaw from supersymmetric Higgs-lepton inflation and BICEP2, *Phys. Lett. B* **735**, 186 (2014).
- [36] S. Kawai and J. Kim, Testing supersymmetric Higgs inflation with non-Gaussianity, *Phys. Rev. D* **91**, 045021 (2015).
- [37] S. Kawai and J. Kim, Multifield dynamics of supersymmetric Higgs inflation in SU(5) GUT, *Phys. Rev. D* **93**, 065023 (2016).
- [38] G. K. Leontaris, N. Okada, and Q. Shafi, Non-minimal quartic inflation in supersymmetric $SO(10)$, *Phys. Lett. B* **765**, 256 (2017).
- [39] N. Okada and Q. Shafi, Gravity waves and gravitino dark matter in μ -hybrid inflation, *Phys. Lett. B* **787**, 141 (2018).
- [40] M. B. Einhorn and D. T. Jones, Inflation with non-minimal gravitational couplings in supergravity, *J. High Energy Phys.* **03** (2010) 026.
- [41] S. Ferrara, R. Kallosh, A. Linde, A. Marrani, and A. Van Proeyen, Jordan frame supergravity and inflation in NMSSM, *Phys. Rev. D* **82**, 045003 (2010).
- [42] N. Okada, M. U. Rehman, and Q. Shafi, Tensor to scalar ratio in non-minimal ϕ^4 inflation, *Phys. Rev. D* **82**, 043502 (2010).
- [43] Y. Akrami *et al.* (Planck Collaboration), Planck 2018 results. X. Constraints on inflation, *Astron. Astrophys.* **641**, A10 (2020).
- [44] J. Barbon and J. Espinosa, On the naturalness of Higgs inflation, *Phys. Rev. D* **79**, 081302 (2009).
- [45] C. Burgess, H. M. Lee, and M. Trott, Power-counting and the validity of the classical approximation during inflation, *J. High Energy Phys.* **09** (2009) 103.
- [46] M. P. Hertzberg, On inflation with Non-minimal coupling, *J. High Energy Phys.* **11** (2010) 023.
- [47] R. N. Lerner and J. McDonald, Higgs inflation and naturalness, *J. Cosmol. Astropart. Phys.* **04** (2010) 015.
- [48] F. Bezrukov, A. Magnin, M. Shaposhnikov, and S. Sibiryakov, Higgs inflation: Consistency and generalisations, *J. High Energy Phys.* **01** (2011) 016.
- [49] A. D. Dolgov and D. P. Kirilova, On particle creation by a time-dependent scalar field, *Yad. Fiz.* **51**, 273 (1990) [*Sov. J. Nucl. Phys.* **51**, 172 (1990)].
- [50] J. H. Traschen and R. H. Brandenberger, Particle production during out-of-equilibrium phase transitions, *Phys. Rev. D* **42**, 2491 (1990).
- [51] S. Kawai and Y. Nakayama, Reheating of the Universe as holographic thermalization, *Phys. Lett. B* **759**, 546 (2016).
- [52] M. Bolz, W. Buchmuller, and M. Plumacher, Baryon asymmetry and dark matter, *Phys. Lett. B* **443**, 209 (1998).
- [53] M. Bolz, A. Brandenburg, and W. Buchmuller, Thermal production of gravitinos, *Nucl. Phys.* **B606**, 518 (2001).
- [54] H. Eberl, I. D. Gialamas, and V. C. Spanos, Gravitino thermal production revisited, *Phys. Rev. D* **103**, 075025 (2021).
- [55] M. Kawasaki, K. Kohri, and T. Moroi, Big-Bang nucleosynthesis and hadronic decay of long-lived massive particles, *Phys. Rev. D* **71**, 083502 (2005).
- [56] S. Borgani, A. Masiero, and M. Yamaguchi, Light gravitinos as mixed dark matter, *Phys. Lett. B* **386**, 189 (1996).
- [57] E. Pierpaoli, S. Borgani, A. Masiero, and M. Yamaguchi, The formation of cosmic structures in a light gravitino dominated Universe, *Phys. Rev. D* **57**, 2089 (1998).
- [58] W. B. Lin, D. H. Huang, X. Zhang, and R. H. Brandenberger, Nonthermal Production of WIMPs and the Subgalactic Structure of the Universe, *Phys. Rev. Lett.* **86**, 954 (2001).
- [59] Y. Gu, M. Khlopov, L. Wu, J. M. Yang, and B. Zhu, Light gravitino dark matter: LHC searches and the Hubble tension, *Phys. Rev. D* **102**, 115005 (2020).
- [60] M. Arai, S. Kawai, and N. Okada, A gauge mediation scenario with hidden sector renormalization in MSSM, *Phys. Rev. D* **81**, 035022 (2010).
- [61] M. Arai, S. Kawai, and N. Okada, Renormalization effects on the MSSM from a calculable model of a strongly coupled hidden sector, *Phys. Rev. D* **84**, 075002 (2011).
- [62] T. Asaka, K. Hamaguchi, and K. Suzuki, Cosmological gravitino problem in gauge mediated supersymmetry breaking models, *Phys. Lett. B* **490**, 136 (2000).
- [63] M. Kawasaki, K. Kohri, T. Moroi, and A. Yotsuyanagi, Big-Bang Nucleosynthesis and Gravitino, *Phys. Rev. D* **78**, 065011 (2008).
- [64] I. Affleck and M. Dine, A new mechanism for baryogenesis, *Nucl. Phys.* **B249**, 361 (1985).
- [65] A. Kusenko and M. E. Shaposhnikov, Supersymmetric Q balls as dark matter, *Phys. Lett. B* **418**, 46 (1998).
- [66] S. Kasuya, M. Kawasaki, and T. T. Yanagida, IceCube potential for detecting Q-ball dark matter in gauge mediation, *Prog. Theor. Exp. Phys.* **(2015)**, 053B02.
- [67] M. Hazumi *et al.*, LiteBIRD: A satellite for the studies of B-mode polarization and inflation from cosmic background radiation detection, *J. Low Temp. Phys.* **194**, 443 (2019).