**Erratum: Analysis of the diffractive production of W 's and dijets at the DESY HERA
and Fermilab Tevatron colliders
[Phys. Rev. D 60, 054005 (1999)]**

R. J. M. Covolan and M. S. Soares
(Published 24 November 1999)

PACS number(s): 12.40.Nn, 13.60.Hb, 13.85.Qk, 13.87.-a, 99.10.+g

The theoretical predictions given in Table I of the original paper are compared to the experimental measurement in an inappropriate way. These predictions were obtained by computing only the interaction of pomerons (emitted by protons) with antiprotons (that means events with rapidity gaps on the side from which antiprotons come from). However, the experimental rate is for both sides, i.e., events with a rapidity gap on the proton or the antiproton side.

Because of the symmetry of the problem, in order to obtain the correct comparison, one has just to multiply the theoretical predictions by a factor of 2. By doing so one obtains the new Table I given here. As can be seen from the table, this change implies that the results obtained with the renormalized flux factor (fits 3 and 4) agree very well with the experimental rate. In spite of this correction, all the discussions and conclusions remain unchanged.

TABLE I. Diffractive productions rates of W 's (all values are given in percentages).

Experiment (W 's)	Rate	STD Fit 1 (hard-hard)	STD Fit 2 (free-delta)	REN Fit 3 (hard-hard)	REN Fit 4 (free-zero)
CDF (rap gap)	1.15 ± 0.55	6.24	7.08	1.06	1.16