Soft-photon corrections to the Bethe-Heitler process in the $\gamma p \rightarrow l^+ l^- p$ reaction

Matthias Heller, Oleksandr Tomalak, and Marc Vanderhaeghen Institut für Kernphysik and PRISMA Cluster of Excellence, Johannes Gutenberg Universität, D-55099 Mainz, Germany

(Received 5 March 2018; published 30 April 2018)

We report on the calculation of first-order quantum electrodynamics (QED) corrections for the $\gamma p \rightarrow l^+ l^- p'$ process. An upcoming experiment at MAMI (Mainz) aims to compare the cross sections of muon- and electron-pair production in this reaction to test lepton universality. Precise knowledge of the electromagnetic radiative corrections is needed for these measurements. As a first step, we present the leading QED radiative corrections in the soft-photon approximation when accounting for the finite lepton mass. For the kinematics at MAMI, we find corrections of the percent level for muons, and of order 10% for electrons.

DOI: 10.1103/PhysRevD.97.076012

I. INTRODUCTION

Recent experiments found a significant difference in the proton charge radius, comparing measurements with electrons and muons. Currently, the most precise measurements with electron scattering were performed by the A1 Collaboration in Mainz [1,2]. The proton radius extracted from these measurements is $R_E = 0.879(8)$ fm. For muonic measurements, there is currently only the proton radius extraction by muonic spectroscopy [3,4], yielding a significantly smaller value than the extraction by electron scattering experiments. The reported value of the muonic hydrogen experiments is $R_F = 0.84087(39)$ fm [4].

This discrepancy, often referred to as the proton radius puzzle, has triggered a lot of activity in recent years. Explanations for this puzzle reach from systematic errors in the extraction, see Refs. [5-12], to physics beyond the standard model, see for example in Refs. [13-22]. If one tries to explain this puzzle by new physics, one has to give up lepton universality as a consequence, since this requires the same, universal coupling for all leptons.

To shed further light on this puzzle and test lepton universality, the MUSE experiment has been proposed, which aims at comparing the scattering of muons and electrons on a proton target [23,24]. In Ref. [25], the authors suggested another test of lepton universality by comparing the cross section of lepton-pair production for muons and electrons in the process $\gamma p \rightarrow l^+ l^- p'$. Such experiment only requires a relative measurement through the ratio of electron- and muon-pair production cross sections slightly above dimuon production threshold. According to the finding of Ref. [25], the measurement of this ratio with absolute precision of around 7×10^{-4} can test lepton universality at 3σ significance level. An upcoming experiment at MAMI is planned to perform such measurements [26].

For a precise theoretical prediction, it is, however, necessary to include higher-order corrections to the process. In this article, we report as a first step on the calculation of the first-order QED corrections in the soft-photon limit when accounting for the finite lepton mass.

The outline of the paper is as follows. In Sec. II, we introduce the kinematical notations for the process $\gamma p \rightarrow l^+ l^- p'$ and give the formulas for the cross section at tree level. In Sec. III, we evaluate the first-order QED corrections to the cross section in the soft-photon approximation. This limit is defined by a soft scaling of the loop momenta. We give the analytic expressions for the real and virtual corrections. We show that they factorize in terms of the tree level cross section, and explicitly check the cancellation of infrared divergences. In Sec. IV, we present the results of this work. We quantify how the ratio of cross sections of muon- and electron-pair production to electron-pair production is affected by radiative corrections. We give our conclusions and an outlook in Sec. V.

II. LEPTON-PAIR PRODUCTION AT TREE LEVEL

The Bethe-Heitler process at tree level is described by two graphs, see Fig. 1. We use p(p') for the momenta of the initial (final) proton, and $p_3(p_4)$ for the momenta of

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.



FIG. 1. The Bethe-Heitler process at tree level.

leptons $l^-(l^+)$ respectively. The initial photon has momentum p_1 , and the virtual photon momentum in the one-photon exchange graphs of Fig. 1 is defined as $p_2 = p - p'$. The Mandelstam variables for this process are defined as

$$(p_3 + p_4)^2 = s_{ll},\tag{1}$$

$$(p_3 - p_1)^2 = t_{ll}, (2)$$

$$p_2^2 = (p - p')^2 = t.$$
 (3)

The on-shell condition for external particles implies:

$$p_3^2 = p_4^2 = m^2, (4)$$

$$p^2 = p'^2 = M^2,$$
 (5)

$$p_1^2 = 0.$$
 (6)

At leading order, the scattering amplitude \mathcal{M}_{0} is given by

$$\mathcal{M}_{0} = \bar{u}(p_{3})(ie) \left[\gamma^{\nu} \frac{i(\not p_{3} - \not p_{1} + m)}{(p_{3} - p_{1})^{2} - m^{2}} \gamma^{\mu} + \gamma^{\mu} \frac{i(\not p_{1} - \not p_{4} + m)}{(p_{1} - p_{4})^{2} - m^{2}} \gamma^{\nu} \right] (ie) v(p_{4}) \frac{-i}{t} \varepsilon_{\nu}(p_{1}) \\ \times \bar{u}(p')(-ie) \Gamma_{\mu}(t) u(p),$$
(7)

where the electromagnetic vertex Γ^{μ} for the proton is expressed as

$$\Gamma^{\mu}(t) = F_D(t)\gamma^{\mu} - iF_P(t)\frac{\sigma^{\mu\nu}(p_2)_{\nu}}{2M},$$
(8)

with the proton's Dirac and Pauli form factors F_D and F_P , respectively.

The corresponding unpolarized differential cross section $d\sigma_0$ is given by

$$\left(\frac{d\sigma}{dtds_{ll}d\Omega_{ll}^{CM_{l+l^{-}}}}\right)_{0} = \frac{1}{(2\pi)^{4}} \frac{1}{64} \frac{\beta}{(2ME_{\gamma})^{2}} \left[\sum_{i} \sum_{f} \left(\mathcal{M}_{0}^{*}\mathcal{M}_{0}\right)\right], \quad (9)$$

where E_{γ} is the lab energy of the initial photon and $\Omega_{ll}^{CM_{l+l^{-}}}$ is the solid angle of the lepton pair in their center-of-mass frame, in which the lepton velocity is denoted by

$$\beta = \sqrt{1 - \frac{4m^2}{s_{ll}}}.$$
 (10)

In Eq. (9), we average over all polarizations in the initial state and sum over the polarizations in the final state. We express the cross section as a product of hadronic and leptonic parts as

$$\left(\frac{d\sigma}{dtds_{ll}d\Omega_{ll}^{CM_{l+l-}}}\right)_0 = \frac{\alpha^3\beta}{16\pi(2ME_{\gamma})^2 t^2} L_{\mu\nu} H^{\mu\nu},\qquad(11)$$

where the fine-structure constant is defined as $\alpha \equiv e^2/4\pi \approx 1/137$. Furthermore, the unpolarized leptonic tensor $L_{\mu\nu}$ (including the average over the initial photon polarization) is given by

$$L^{\mu\nu} = -\frac{1}{2} \operatorname{Tr} \left[(\not p_3 + m) \left(\gamma^{\alpha} \frac{(\not p_3 - \not p_1 + m)}{(p_3 - p_1)^2 - m^2} \gamma^{\mu} + \gamma^{\mu} \frac{(\not p_1 - \not p_4 + m)}{(p_1 - p_4)^2 - m^2} \gamma^{\alpha} \right) (\not p_4 - m) \times \left(\gamma^{\nu} \frac{(\not p_3 - \not p_1 + m)}{(p_3 - p_1)^2 - m^2} \gamma_{\alpha} + \gamma_{\alpha} \frac{(\not p_1 - \not p_4 + m)}{(p_1 - p_4)^2 - m^2} \gamma^{\nu} \right) \right],$$
(12)

and the unpolarized hadronic tensor $H^{\mu\nu}$ by

$$H^{\mu\nu} = \frac{1}{2} \operatorname{Tr}[(\not\!\!p' + M)\Gamma^{\mu}(\not\!\!p + M)(\Gamma^{\dagger})^{\nu}].$$
(13)

Using (8), the unpolarized hadronic tensor can be expressed as

$$H^{\mu\nu} = \left(-g^{\mu\nu} + \frac{p_2^{\mu}p_2^{\nu}}{p_2^2}\right) [4M^2 \tau G_M^2(t)] + \tilde{p}^{\mu} \tilde{p}^{\nu} \frac{4}{1+\tau} [G_E^2(t) + \tau G_M^2(t)], \qquad (14)$$

where $\tilde{p} \equiv (p + p')/2$, $\tau \equiv -t/(4M^2)$, and where we conveniently express the hadronic tensor in terms of electric (*G_E*) and magnetic (*G_M*) form factors defined as

$$G_E = F_D - \tau F_P, \tag{15}$$

$$G_M = F_D + F_P, \tag{16}$$

which are functions of the spacelike momentum transfer t.

For the electric and magnetic proton form factors, which enter the total cross sections for lepton-pair production, we exploit the fit of Ref. [2], which is based on a global analysis of the electron-proton scattering data at $Q^2 < 10 \text{ GeV}^2$ with an empirical account of TPE corrections.

In the experimental setup, when only the recoil proton is measured, one has to integrate (11) over the lepton angles:

$$\left(\frac{d\sigma}{dtds_{ll}}\right)_0 = \frac{\alpha^3\beta}{16\pi(2ME_\gamma)^2t^2} \cdot \int d\Omega_{ll}^{CM_{l+l-}} L_{\mu\nu} H^{\mu\nu}.$$
 (17)

The kinematical invariant t is in one-to-one relation with the recoiling proton lab momentum \vec{p}' (or energy E'):

$$|\vec{p}'| = 2M\sqrt{\tau(1+\tau)},\tag{18}$$

$$E' = M(1+2\tau),$$
 (19)

whereas the invariant s_{ll} is then determined from the recoiling proton lab scattering angle:

$$\cos \theta_{p'} = \frac{s_{ll} + 2(s + M^2)\tau}{2(s - M^2)\sqrt{\tau(1 + \tau)}},$$
(20)

where s is the squared center-of-mass energy, which can be expressed in terms of the initial photon-beam energy E_{γ} :

$$s = 2E_{\gamma}M + M^2. \tag{21}$$

In Ref. [25], the authors calculated the ratio *R* of cross sections between electron- and muon-pair production:

$$R(s_{ll}, s_{ll}^{0}) \equiv \frac{[\sigma_0(\mu^+\mu^-)](s_{ll}) + [\sigma_0(e^+e^-)](s_{ll})}{[\sigma_0(e^+e^-)](s_{ll}^{0})}, \quad (22)$$

which depends on the invariant mass of the lepton pair s_{ll} , and a reference point s_{ll}^0 to which the measurement is normalized.

The corresponding plot for the kinematical range accessible at MAMI is shown in Fig. 2. The normalization is shown for the choice $s_{II}^0 = s_{II}$, i.e., at each point above the muon-pair production threshold the sum of the cross sections for muon- and electron-pair production is divided by the corresponding cross section for electron-pair production. In this plot, the blue curve describes the scenario, when lepton universality holds, i.e., $G_E^\mu = G_E^e$, while the red curve corresponds to a case when lepton universality is broken by an amount of 1%. The blue band describes the 3σ deviation if this observable is measured with an absolute



FIG. 2. Ratio of the cross sections in $\gamma p \rightarrow (e^+e^- + \mu^+\mu^-)p$ vs $\gamma p \rightarrow (e^+e^-)p$. The blue band corresponds to a 3σ band, where $\sigma = 7 \times 10^{-4}$.

accuracy of 7×10^{-4} . We will show in this work that radiative corrections shift this curve by more than 3σ , making their inclusion indispensable for a comparison with experiment.

III. LEADING-ORDER RADIATIVE CORRECTIONS IN THE SOFT-PHOTON LIMIT

We evaluate the first-order QED corrections to the $\gamma p \rightarrow l^+ l^- p$ process in the soft-photon limit. This limit is defined by a scaling of the momenta *k* of virtual photons in the loops and real photon momenta in the bremsstrahlung process, with respect to external scales, as

$$k \sim \lambda,$$
 (23)

where λ is a small parameter. We calculate the diagrams at leading order in λ . This procedure reproduces all infrareddivergent contributions and results in a finite, gaugeinvariant piece. The resulting cross section correction factorizes in terms of the tree-level cross section given by Eq. (11).

A. Virtual corrections

We start by calculating the one-loop virtual radiative corrections. In the soft-photon approximation, only box diagrams contribute. We list all propagators and their scaling with λ in Table I:

The integral measure d^4k scales as

$$d^4k \sim \lambda^4. \tag{24}$$

Therefore, to obtain a contribution of order 1, we need a denominator of order λ^4 . This is only possible for the box diagrams when the first 3 propagators of Table I are present in a Feynman integral.

TABLE I. Scaling of the propagator denominators with the expansion parameter λ . Only integrals with the first 3 propagators contribute, since these integrals have a denominator scaling as λ^4 , which is the scaling of the integral measure in the numerator.

| Propagator denominator | Scaling at least as |
|---------------------------|---------------------|
| $(k+p_3)^2 - m^2$ | λ |
| $(k - p_4)^2 - m^2$ | λ |
| k^2 | λ^2 |
| $(p_3 - p_1 + k)^2 - m^2$ | 1 |

For the box diagrams shown in Fig. 3, we obtain the following leading contribution:

$$\mathcal{M}^{\text{box}} = (ie^2) 4 \cdot (p_3 p_4) \cdot \mathcal{M}_0 \mu^{4-d} \\ \times \int \frac{d^d k}{(2\pi)^d} \frac{1}{(p_3 + k)^2 - m^2} \frac{1}{(k - p_4)^2 - m^2} \frac{1}{k^2} + \mathcal{O}(\lambda) \\ = -\frac{e^2}{8\pi^2} (s_{ll} - 2m^2) \cdot \mathcal{M}_0 \cdot C_0(m^2, s_{ll}, m^2, 0, m^2, m^2),$$
(25)

with the 3-point function C_0 in dimensional regularization, see Ref. [27]¹:

$$C_{0}(m^{2}, s_{ll}, m^{2}, 0, m^{2}, m^{2})$$

$$= \frac{1}{s_{ll}\beta} \left\{ \left[\frac{1}{\epsilon_{IR}} - \gamma_{E} + \ln\left(\frac{4\pi\mu^{2}}{m^{2}}\right) \right] \ln\left(\frac{\beta - 1}{\beta + 1}\right) + 2\mathrm{Li}_{2}\left(\frac{\beta - 1}{2\beta}\right) + \ln^{2}\left(\frac{\beta - 1}{2\beta}\right) - \frac{1}{2}\ln^{2}\left(\frac{\beta - 1}{\beta + 1}\right) - \frac{\pi^{2}}{6} \right\}.$$

$$(26)$$

In Eqs. (25), (26), μ is a scale introduced to account for the correct energy dimension of the integral. Physical quantities have to be independent of this scale, as well as of the infrared regulator $\epsilon_{IR} \equiv 2 - d/2 < 0$. All other diagrams are infrared finite and scale at least as λ . Therefore, the other graphs do not contribute in the softphoton limit.

Although the box diagrams are UV finite, we have to include counterterm corrections, shown in Fig. 4, since they contain infrared-divergent parts in the on-shell subtraction scheme, which we follow here. We describe these contributions according to Ref. [28].

In the on-shell subtraction scheme, the vertex counterterm is defined to fix the electron charge e at $q^2 = 0$. Considering the vertex function in Fig. 5, one can decompose the diagram into two tensor structures with corresponding form factors F and G:



FIG. 3. QED box diagrams contributing to the radiative corrections calculation in the soft-photon approximation.

$$\bar{u}(p')\Gamma^{\mu}u(p) = \bar{u}(p') \left[(1 + F(q^2))\gamma^{\mu} + iG(q^2)\sigma^{\mu\nu}\frac{q_{\nu}}{2m} \right] u(p),$$
(27)

with

$$q = p' - p. \tag{28}$$

Only $F(q^2)$ is UV divergent, and one finds at $q^2 = 0$ the renormalization constant:

$$Z_{1} = 1 - F(0)$$

$$= 1 - \frac{e^{2}}{(4\pi)^{2}} \left\{ \left[\frac{1}{\epsilon_{\rm UV}} - \gamma_{E} + \ln\left(\frac{4\pi\mu^{2}}{m^{2}}\right) \right] + 2 \left[\frac{1}{\epsilon_{\rm IR}} - \gamma_{E} + \ln\left(\frac{4\pi\mu^{2}}{m^{2}}\right) \right] + 4 \right\}.$$
(29)

This leads to the renormalized vertex:

$$\tilde{\Gamma}^{\mu} = \Gamma^{\mu} + (Z_1 - 1)\gamma^{\mu}, \qquad (30)$$

that in the soft-photon limit $(\tilde{\Gamma}_s^{\mu})$, which corresponds to taking only the infrared-divergent part, is expressed as

$$\tilde{\Gamma}_{\rm s}^{\mu} = -\frac{\alpha}{2\pi} \gamma^{\mu} \left[\frac{1}{\epsilon_{\rm IR}} - \gamma_E + \ln\left(\frac{4\pi\mu^2}{m^2}\right) \right]. \tag{31}$$

The contribution of the two vertex counterterms in Fig. 4 is then given by

$$\mathcal{M}_{\text{vertex}}^{\text{ct}} = -\frac{\alpha}{\pi} \left[\frac{1}{\epsilon_{\text{IR}}} - \gamma_E + \ln\left(\frac{4\pi\mu^2}{m^2}\right) \right] \mathcal{M}_0. \quad (32)$$

The self-energy counterterm is defined from the lepton self-energy correction $\Sigma(p)$, which is expressed in terms of the lepton propagator *S*:

¹We use the same notation for this function as in http://qcdloop .fnal.gov/



FIG. 4. Counterterm diagrams, which contribute to the $\gamma p \rightarrow l^+ l^- p$ process. These give rise to infrared-divergent contributions in the on-shell subtraction scheme and have therefore to be accounted for when calculating the radiative corrections in the soft-photon approximation.



FIG. 5. Diagrams for the calculation of the counterterms. The upper diagram defines the vertex counterterm, the lower diagram corresponds to the lepton self-energy.

$$iS = iS^0 + iS^0(-i)\Sigma(p)iS,$$
(33)

with free fermion propagator given by

$$S^{0}(p) = \frac{\not p + m}{p^{2} - m^{2}}.$$
(34)

Calculating up to first order, we have to include the one-loop correction:

$$-i\Sigma(\not\!\!\!p) = -e^2\mu^{4-d} \int \frac{d^dk}{(2\pi)^d} \frac{\gamma^{\alpha}(\not\!\!\!p + k\!\!\!/ + m)\gamma_{\alpha}}{((p+k)^2 - m^2)k^2}.$$
 (35)

The on-shell renormalization condition fixes the pole at $p^2 = m^2$ with residue equal to one. This gives the renormalization constants Z_2 and Z_m :

$$Z_2 = 1 + \frac{d\Sigma(\not p)}{d\not p} \bigg|_{\not p=m},\tag{36}$$

$$(1-Z_m)Z_2m = \Sigma(m). \tag{37}$$

The evaluation of $\Sigma(p)$ and its derivative, results in the renormalization constants:

$$Z_{2} = 1 - \frac{e^{2}}{(4\pi)^{2}} \left\{ \left[\frac{1}{\epsilon_{\rm UV}} - \gamma_{E} + \ln\left(\frac{4\pi\mu^{2}}{m^{2}}\right) \right] + 2 \left[\frac{1}{\epsilon_{\rm IR}} - \gamma_{E} + \ln\left(\frac{4\pi\mu^{2}}{m^{2}}\right) \right] + 4 \right\},$$
(38)

$$Z_2 Z_m = 1 - \frac{e^2}{(4\pi)^2} \left\{ 4 \left[\frac{1}{\epsilon_{\rm UV}} - \gamma_E + \ln\left(\frac{4\pi\mu^2}{m^2}\right) \right] + 2 \left[\frac{1}{\epsilon_{\rm IR}} - \gamma_E + \ln\left(\frac{4\pi\mu^2}{m^2}\right) \right] + 8 \right\}.$$
(39)

The renormalized self-energy is then given by

$$\tilde{\Sigma}(p) = \Sigma(p) - (Z_2 - 1)\not p + (Z_2 Z_m - 1)m.$$
(40)

Taking only the infrared-divergent piece in the soft-photon limit $(\tilde{\Sigma}_s)$, we obtain:

$$\tilde{\Sigma}_{\rm s}(p) = \frac{\alpha}{2\pi} (\not\!\!p - m) \left[\frac{1}{\epsilon_{\rm IR}} - \gamma_E + \ln \left(\frac{4\pi\mu^2}{m^2} \right) \right].$$
(41)

The contribution of the self-energy counterterm M_{se}^{ct} in Fig. 4 is therefore given by

$$\mathcal{M}_{\rm se}^{\rm ct} = \frac{\alpha}{2\pi} \left[\frac{1}{\epsilon_{\rm IR}} - \gamma_E + \ln\left(\frac{4\pi\mu^2}{m^2}\right) \right] \mathcal{M}_0. \tag{42}$$

Adding virtual corrections of Eq. (25) and counterterms of Eqs. (32) and (42), we obtain the virtual one-loop correction in the soft-photon limit $\mathcal{M}_{s;V}$:

$$\mathcal{M}_{s;V} = -\frac{\alpha}{2\pi} \left\{ (s_{ll} - 2m^2) C_0(m^2, s_{ll}, m^2, 0, m^2, m^2) + \left[\frac{1}{\epsilon_{IR}} - \gamma_E + \ln\left(\frac{4\pi\mu^2}{m^2}\right) \right] \right\} \mathcal{M}^0.$$
(43)

The resulting virtual correction to the cross section is then given, to first order in α , by

$$\left(\frac{d\sigma}{dt_{ll}ds_{ll}}\right)_{s;V} = 2\operatorname{Re}[\mathcal{M}_0^* \times \mathcal{M}_{s;V}].$$
(44)

It can be expressed as

$$\left(\frac{d\sigma}{dt_{ll}ds_{ll}}\right)_{s;V} = \left(\frac{d\sigma}{dt_{ll}ds_{ll}}\right)_0 (\delta_{s;V}^{IR} + \delta_{s;V}), \quad (45)$$

with the infrared-divergent part:

$$\delta_{s;V}^{IR} = \left(\frac{-\alpha}{\pi}\right) \left[\left(\frac{1+\beta^2}{2\beta}\right) \ln\left(\frac{1-\beta}{1+\beta}\right) + 1 \right] \\ \times \left[\frac{1}{\epsilon_{IR}} - \gamma_E + \ln\left(\frac{4\pi\mu^2}{m^2}\right)\right], \tag{46}$$

and the finite part:

$$\delta_{s;V} = \left(\frac{-\alpha}{\pi}\right) \left(\frac{1+\beta^2}{2\beta}\right) \\ \times \left\{ 2\text{Li}_2\left(\frac{2\beta}{\beta+1}\right) + \frac{1}{2}\ln^2\left(\frac{1-\beta}{1+\beta}\right) - \pi^2 \right\}.$$
(47)

B. Soft-photon bremsstrahlung

Besides the QED virtual radiative corrections, one has to account for processes with radiation of undetected photons.

The diagrams contributing to the soft bremsstrahlung from the lepton side are shown in Fig. 6. Note that the diagram, where the photon is attached to the internal lepton line, vanishes for $\lambda \to 0$ and does not contribute in the soft-photon limit. Denoting the momentum of the photon by *k*, we find the squared matrix element for this process in the form:

$$|\mathcal{M}(\gamma p \to \gamma_{s} l^{+} l^{-} p)|^{2} = |\mathcal{M}(\gamma p \to l^{+} l^{-} p)|^{2} (-e^{2}) \left[\frac{p_{3}^{\mu}}{p_{3} \cdot k} - \frac{p_{4}^{\mu}}{p_{4} \cdot k} \right] \cdot \left[\frac{p_{3\mu}}{p_{3} \cdot k} - \frac{p_{4\mu}}{p_{4} \cdot k} \right].$$

$$(48)$$

To calculate the contribution to the cross section, one then has to integrate over the undetected soft-photon energy up to a small value ΔE_s , determined by the experimental resolution.



FIG. 6. Diagrams with real photon emission from the lepton lines for the Bethe-Heitler process. In the soft-photon limit, the diagram with the photon attached to the internal (off-shell) fermion line does not contribute.

Due to the energy-momentum conserving δ -function, $\delta^4(p_1 + p - p_3 - p_4 - p' - k)$, the integration domain has a complicated shape in the lab system. The integration can be carried out in the rest frame S of the real (p_1) and virtual (p_2) photons, which is also the rest frame of the dilepton pair and soft photon, defined by

$$\vec{p}_1 + \vec{p}_2 = \vec{p}_3 + \vec{p}_4 + \vec{k} = 0.$$
⁽⁴⁹⁾

In such frame, the dependence of the integral with respect to the soft-photon momentum becomes isotropic. For the differential cross section, we then need to evaluate:

$$\left(\frac{d\sigma}{dtds_{ll}}\right)_{s;R} = -\left(\frac{d\sigma}{dtds_{ll}}\right)_0 \frac{e^2}{(2\pi)^3} \int_{|\vec{k}| < \Delta E_s} \frac{d^3\vec{k}}{2k^0} \\ \times \left[\frac{m^2}{(p_3k)^2} + \frac{m^2}{(p_4k)^2} - \frac{2(p_3p_4)}{(p_3k)(p_4k)}\right], \quad (50)$$

where the integration is performed in the frame S.

The integrals are infrared divergent and can be carried out analytically after dimensional regularization. They have been worked out, e.g., in Ref. [29]. For the kinematics in system S, where the soft-photon momentum:

$$|\vec{k}| \ll |\vec{p}_3|, \ |\vec{p}_4|,$$
 (51)

with the lepton momenta:

$$p_3^0 = p_4^0 = \frac{\sqrt{s_{ll}}}{2}, \qquad \vec{p}_3 = -\vec{p}_4,$$
 (52)

we obtain:

$$\left(\frac{d\sigma}{dtds_{ll}}\right)_{s;R} = \left(\frac{d\sigma}{dtds_{ll}}\right)_0 (\delta_{s;R}^{IR} + \delta_{s;R}), \qquad (53)$$

where $\delta_{s;R}^{IR}$ is the infrared-divergent contribution due to real photon emission:

$$\delta_{s;R}^{IR} = \left(\frac{-\alpha}{\pi}\right) \left[\left(\frac{1+\beta^2}{2\beta}\right) \ln\left(\frac{1+\beta}{1-\beta}\right) - 1 \right] \\ \times \left[\frac{1}{\epsilon_{IR}} - \gamma_E + \ln\left(\frac{4\pi\mu^2}{m^2}\right) \right],$$
(54)

and $\delta_{s;R}$ is the corresponding finite part:

$$\delta_{s;R} = \left(\frac{-\alpha}{\pi}\right) \left\{ \ln\left(\frac{4\Delta E_s^2}{m^2}\right) \left[1 + \left(\frac{1+\beta^2}{2\beta}\right) \ln\left(\frac{1-\beta}{1+\beta}\right)\right] + \frac{1}{\beta} \ln\left(\frac{1-\beta}{1+\beta}\right) + \left(\frac{1+\beta^2}{2\beta}\right) \left[2Li_2\left(\frac{2\beta}{1+\beta}\right) + \frac{1}{2}\ln^2\left(\frac{1-\beta}{1+\beta}\right)\right] \right\}.$$
 (55)

The maximum value of the undetected soft-photon energy ΔE_s is defined in the system S. One can reexpress it in terms of the detector resolutions. We consider the case of detecting the recoil proton only. The energy E' and angle $\theta_{p'}$ of the scattered proton are measured in the lab frame. The missing mass M_{miss} of the system is defined by

$$M_{\rm miss}^2 = (p_3 + p_4 + k)^2 = s_{ll} + 2M_{\rm miss}E_s, \quad (56)$$

$$E_s = \frac{M_{\rm miss}^2 - s_{ll}}{2M_{\rm miss}},\tag{57}$$

where E_s denotes the soft-photon energy.

The missing mass M_{miss} is experimentally determined from the quantity:

$$M_{\text{miss}}^2 = (p_1 + p - p')^2$$
$$= 4M\tau \left(E_{\gamma} \sqrt{\frac{1+\tau}{\tau}} \cos \theta_{p'} - E_{\gamma} - M \right), \quad (58)$$

where τ is determined from the lab proton momentum by Eq. (18), and $\theta_{p'}$ is the experimentally measured recoil proton scattering angle in the laboratory frame.

For the process without radiation, this angle is given by Eq. (20), which can be equivalently obtained from Eq. (58) by the replacement $M_{\text{miss}}^2 \rightarrow s_{ll}$:

$$s_{ll} = 4M\tau \left(E_{\gamma} \sqrt{\frac{1+\tau}{\tau}} \cos \theta_{p'} |_{\text{no rad}} - E_{\gamma} - M \right).$$
 (59)

Combining Eqs. (58) and (59), we can express the softphoton energy of Eq. (57) approximately as:

$$E_{\rm s} = \frac{2ME_{\gamma}\sqrt{\tau(1+\tau)}}{\sqrt{s_{ll}}} \left[\cos\theta_{p'} - \cos\theta_{p'}\right]_{\rm no\,rad}.$$
 (60)

Consequently, the experimental recoiling proton angular resolution, denoted as $\Delta \theta_{p'}$, determines the maximum value ΔE_s of the undetected soft-photon energy, which enters the radiative correction of Eq. (55), as

$$\Delta E_s = \frac{2ME_{\gamma}\sqrt{\tau(1+\tau)}}{\sqrt{s_{ll}}}\sin\theta_{p'}\Delta\theta_{p'}.$$
 (61)

C. Total result and exponentiation

Adding the real and virtual contributions of Eqs. (54) and (46), we find a cancellation of all infrared divergences on the level of the cross section:

$$\delta_{\rm s;R}^{\rm IR} + \delta_{\rm s;V}^{\rm IR} = 0. \tag{62}$$

For the finite part of the first-order QED corrections in the soft-photon approximation:

$$\delta = \delta_{\rm s:R} + \delta_{\rm s:V},\tag{63}$$

we find the result:

$$\delta = -\left(\frac{\alpha}{\pi}\right) \left\{ \left[\ln\left(\frac{4\Delta E_s^2}{m^2}\right) + \ln\left(\frac{1-\beta}{1+\beta}\right) \right] \times \left[1 + \left(\frac{1+\beta^2}{2\beta}\right) \ln\left(\frac{1-\beta}{1+\beta}\right) \right] + \left(\frac{1-\beta}{\beta}\right) \ln\left(\frac{1-\beta}{1+\beta}\right) + \left(\frac{1+\beta^2}{2\beta}\right) \left[4\text{Li}_2\left(\frac{2\beta}{1+\beta}\right) - \pi^2 \right] \right\}, \quad (64)$$

which reduces in the limit $s_{ll} \gg 4m^2$ to:

$$\delta = -\left(\frac{\alpha}{\pi}\right) \left\{ \ln\left(\frac{4\Delta E_s^2}{s_{ll}}\right) \left[1 + \ln\left(\frac{m^2}{s_{ll}}\right)\right] - \frac{\pi^2}{3} \right\}.$$
 (65)

To account for the emission of a higher amount of soft photons or higher-order virtual corrections due to soft photons in the loop, we follow Ref. [30] and exponentiate the terms leading to double logarithmic enhancements as

$$\left(\frac{d\sigma}{dtds_{ll}}\right)_{s,tot} = \left(\frac{d\sigma}{dtds_{ll}}\right)_{0} \cdot F \exp\left\{-\frac{\alpha}{\pi} \left[\ln\left(\frac{4\Delta E_{s}^{2}}{m^{2}}\right) + \ln\left(\frac{1-\beta}{1+\beta}\right)\right] \left[1 + \left(\frac{1+\beta^{2}}{2\beta}\right) \ln\left(\frac{1-\beta}{1+\beta}\right)\right]\right\} \\ \times \left\{1 - \frac{\alpha}{\pi} \left[\left(\frac{1-\beta}{\beta}\right) \ln\left(\frac{1-\beta}{1+\beta}\right) + \left(\frac{1+\beta^{2}}{2\beta}\right) \left[4\text{Li}_{2}\left(\frac{2\beta}{1+\beta}\right) - \pi^{2}\right]\right]\right\} \\ \equiv \left(\frac{d\sigma}{dtds_{ll}}\right)_{0} (1+\delta_{\exp}).$$
(66)

Note that in Eq. (66) terms of single logarithmic nature of order α are still missing, and require a full one-loop calculation. The normalization factor *F* in Eq. (66) is due to the physical assumption that in an experiment the sum of all soft-photon energies is smaller than ΔE_s , instead of requiring that each soft-photon energy is individually smaller than ΔE_s . It was shown in Ref. [30] that when including the leading correction from unity, the normalization factor *F* is given by:

$$F = 1 - \frac{\alpha^2}{3} \left[1 + \left(\frac{1+\beta^2}{2\beta}\right) \ln\left(\frac{1-\beta}{1+\beta}\right) \right]^2 + \cdots$$
 (67)

Although we account for the factor *F* explicitly, its deviation from unity is quite small: for $s_{ll} = 0.077 \text{ GeV}^2$ approximately -2.4×10^{-3} for electrons and -8.5×10^{-6} for muons.

IV. RESULTS AND DISCUSSION

In Fig. 7, we show the corrections at fixed $s_{ll} = 0.077 \text{ GeV}^2$ as a function of the soft-photon energy. We observe a logarithmic behavior of the correction factor δ which gives rise to the so-called radiative tail. We also show the exponentiated form, δ_{exp} , given by Eq. (66), which estimates higher-order effects of soft-photon corrections. Assuming a value $\Delta E_s = 0.01 \text{ GeV}$, δ at first order differs by about 0.006 for electron-pair production and is indistinguishable at the level of precision for muon-pair production (the difference is around -1.0×10^{-4}).

In Fig. 8, we show the radiative corrections to the cross section in the kinematical range of s_{ll} between 0 and 0.08 GeV². The muon threshold is at $s_{ll} = 4m_{\mu}^2 \approx 0.045$ GeV² (vertical dashed red line in Fig 8). We observe that the corrections for electrons are negative of order 10%, while the corrections for muons are positive of order 1%.

Taking radiative corrections into account, the ratio of Eq. (22) is now given by

$$R(s_{ll}, s_{ll}^{0}) = \frac{[\sigma_0(\mu^+\mu^-)(1+\delta^\mu)](s_{ll}) + [\sigma_0(e^+e^-)(1+\delta^e)](s_{ll})}{[\sigma_0(e^+e^-)(1+\delta^e)](s_{ll}^{0})},$$
(68)





FIG. 7. QED corrections to the cross section in the soft-photon limit as a function of the soft-photon energy ΔE_s , which corresponds to the integrated over angular bins $\Delta \theta_{p'}$ according to Eq. (61). This variation stems from the integrated over radiative tail. The external kinematics and the di-lepton invariant mass $s_{II} = 0.077 \text{ GeV}^2$ are indicated on the plot.

FIG. 8. First-order QED corrections to the cross section in the soft-photon limit, using $\Delta E_s = 0.01$ GeV. The vertical dashed red line indicates the muon-pair production threshold at $s_{ll} \approx 0.045$ GeV².



FIG. 9. Ratio of cross sections between electron- and muon-pair production at tree level (blue curve) and with account of firstorder QED corrections estimated using $\Delta E_s = 0.01$ GeV (orange curve) with 3σ error bands. The red curve denotes the scenario when lepton universality is broken with $G_E^{\mu}/G_E^e = 1.01$, including the radiative corrections in the soft-photon approximation.

which depends on the measured invariant lepton mass s_{ll} and the reference point s_{ll}^0 , to which the cross section is normalized. δ^e and δ^{μ} are given by Eq. (66). One chooses $s_{ll}^0 < 4m_{\mu}^2$, such that the reference measurement is below the muon-pair-production threshold, and only electron pairs are created.

In Fig. 9, we show the differential cross section ratio *R* of Eq. (22), including first-order QED corrections in the softphoton approximation with $\Delta E_s = 0.01$ GeV. One sees from this plot, that the inclusion of radiative corrections is indispensable, since the ratio of cross sections, defined in Eq. (22), is shifted to higher values by more than the 3σ band. The radiative corrections to *R* are of the order of a few percent. The red curve in Fig. 9 shows the scenario when lepton universality is violated by $G_E^{\mu}/G_E^e = 1.01$. Following Ref. [25], we use 3σ bands around the curves,



FIG. 10. Ratio of cross sections between electron- and muonpair production at tree level (blue curve) and with account of firstorder QED corrections estimated using $\Delta E_s = 0.01$ GeV (orange curve), normalized to the electron-pair production cross section at $s_{II}^0 = 0.02$ GeV².

with the experimental resolution $\sigma = 7 \times 10^{-4}$. The statement that lepton universality can be tested with a 3σ confidence level remains true if one adds radiative corrections as can be seen in Fig. 9.

In Fig. 10, we show the corresponding ratio between the cross sections normalized to a value below the muon-pair production threshold. As a reference point, we choose $s_{ll}^0 = 0.02 \text{ GeV}^2$. The bands now correspond to the renormalized 3σ bands, i.e.,

$$\sigma = 7 \times 10^{-4} \cdot \frac{\sigma(e^+e^-)(s_{ll})}{\sigma(e^+e^-)(s_{ll}^0)}.$$

V. CONCLUSIONS AND OUTLOOK

In this work, we have calculated QED radiative corrections to the photoproduction of electron and muon pairs on a proton target in the soft-photon approximation. Only radiation from the produced pair and box diagrams with photon and lepton legs contribute in this approximation when accounting for the finite lepton mass. The resulting correction to the cross section factorizes in terms of the tree-level contribution. We expressed the proportionality factor in a compact analytical form. With account of radiative corrections, the ratio of photoproduction cross sections of $e^+e^- + \mu^+\mu^-$ to e^+e^- pairs at the same beam energy (as well when compared to the e^+e^- cross section at an energy below the muon-production threshold) increases by a percent amount comparing to the tree level result. Such changes are significantly larger than the precision needed to distinguish between the proton charge radii extractions from experiments with muons and electrons. It makes a correct inclusion of radiative corrections paramount for the experimental realization. As a next step, we plan to extend the radiative correction result in the soft-photon approximation presented in this work to a full one-loop QED calculation on the lepton side and to include the box diagrams resulting from the two-photon exchange between lepton and proton with an intermediate proton state using the techniques developed in Refs. [31,32] for elastic l^-p scattering. For the leading corrections resulting from the hadronic side we expect, from the corresponding results for the elastic l^-p scattering, to receive cross section corrections at the percent level for the electron case. Such anticipated corrections would translate in a change of the ratio of $e^+e^- + \mu^+\mu^-$ to e^+e^- cross sections at the per mille level, corresponding with the 1σ accuracy goal discussed above for this quantity.

ACKNOWLEDGMENTS

We would like to thank Dr. Aleksandrs Aleksejevs and Shihao Wu for useful discussions. This work was supported by the Deutsche Forschungsgemeinschaft DFG in part through the Collaborative Research Center [The Low-Energy Frontier of the Standard Model (SFB 1044)], and in part through the Cluster of Excellence [Precision Physics, Fundamental Interactions and Structure of Matter (PRISMA)]. Matthias Heller is supported in part by GRK Symmetry Breaking (DFG/GRK 1581). Our figures were generated using JAXODRAW [33], based on AXODRAW [34]. For our *Mathematica* plots, we use the package MATEX.²

²https://github.com/szhorvat/MaTeX.

- J. C. Bernauer *et al.* (A1 Collaboration), High-Precision Determination of the Electric and Magnetic Form Factors of the Proton, Phys. Rev. Lett. **105**, 242001 (2010).
- [2] J. C. Bernauer *et al.* (A1 Collaboration), Electric and magnetic form factors of the proton, Phys. Rev. C 90, 015206 (2014).
- [3] R. Pohl *et al.*, The size of the proton, Nature (London) **466**, 213 (2010).
- [4] A. Antognini *et al.*, Proton structure from the measurement of 2S - 2P transition frequencies of muonic hydrogen, Science **339**, 417 (2013).
- [5] I. T. Lorenz, H.-W. Hammer, and U. G. Meissner, The size of the proton—closing in on the radius puzzle, Eur. Phys. J. A 48, 151 (2012).
- [6] I. T. Lorenz and U. G. Meiner, Reduction of the proton radius discrepancy by 3, Phys. Lett. B 737, 57 (2014).
- [7] I. T. Lorenz, U. G. Meiner, H.-W. Hammer, and Y.-B. Dong, Theoretical constraints and systematic effects in the determination of the proton form factors, Phys. Rev. D 91, 014023 (2015).
- [8] G. Lee, J. R. Arrington, and R. J. Hill, Extraction of the proton radius from electron-proton scattering data, Phys. Rev. D 92, 013013 (2015).
- [9] J. Arrington and I. Sick, Evaluation of the proton charge radius from e-p scattering, J. Phys. Chem. Ref. Data 44, 031204 (2015).
- [10] J. Arrington, An examination of proton charge radius extractions from e-p scattering data, J. Phys. Chem. Ref. Data 44, 031203 (2015).
- [11] K. Griffioen, C. Carlson, and S. Maddox, Consistency of electron scattering data with a small proton radius, Phys. Rev. C 93, 065207 (2016).
- [12] D. W. Higinbotham, A. A. Kabir, V. Lin, D. Meekins, B. Norum, and B. Sawatzky, Proton radius from electron scattering data, Phys. Rev. C 93, 055207 (2016).
- [13] D. Tucker-Smith and I. Yavin, Muonic hydrogen and MeV forces, Phys. Rev. D 83, 101702 (2011).
- [14] V. Barger, C. W. Chiang, W. Y. Keung, and D. Marfatia, Proton Size Anomaly, Phys. Rev. Lett. 106, 153001 (2011).
- [15] V. Barger, C. W. Chiang, W. Y. Keung, and D. Marfatia, Constraint on Parity-Violating Muonic Forces, Phys. Rev. Lett. **108**, 081802 (2012).
- [16] B. Batell, D. McKeen, and M. Pospelov, New Parity-Violating Muonic Forces and the Proton Charge Radius, Phys. Rev. Lett. **107**, 011803 (2011).
- [17] P. Brax and C. Burrage, Atomic precision tests and light scalar couplings, Phys. Rev. D 83, 035020 (2011).

- [18] U. D. Jentschura, Lamb shift in muonic hydrogen. II. Analysis of the discrepancy of theory and experiment, Ann. Phys. (Amsterdam) **326**, 516 (2011).
- [19] C. E. Carlson and B. C. Rislow, New physics and the proton radius problem, Phys. Rev. D **86**, 035013 (2012).
- [20] L. B. Wang and W. T. Ni, Proton radius puzzle and large extra dimensions, Mod. Phys. Lett. A 28, 1350094 (2013).
- [21] R. Onofrio, Proton radius puzzle and quantum gravity at the Fermi scale, Europhys. Lett. **104**, 20002 (2013).
- [22] S. G. Karshenboim, D. McKeen, and M. Pospelov, Constraints on muon-specific dark forces, Phys. Rev. D 90, 073004 (2014); Publisher's Note: Constraints on muonspecific dark forces, Phys. Rev. D90, 079905 (2014).
- [23] R. Gilman *et al.* (MUSE Collaboration), Studying the proton "radius" puzzle with μp elastic scattering, arXiv:1303.2160.
- [24] R. Gilman *et al.* (MUSE Collaboration), Technical design report for the Paul Scherrer institute experiment R-12-01.1: Studying the proton "Radius" puzzle with μp elastic scattering, arXiv:1709.09753.
- [25] V. Pauk and M. Vanderhaeghen, Lepton Universality Test in the Photoproduction of e^-e^+ versus $\mu^-\mu^+$ Pairs on a Proton Target, Phys. Rev. Lett. **115**, 221804 (2015).
- [26] A2 Coll. at MAMI, Letter of intent, https://www.blogs .uni-mainz.de/fb08-mami-experiments/files/2016/07/A2-LOI2016-1.pdf.
- [27] W. Beenakker, S. Dittmaier, M. Kramer, B. Plumper, M. Spira, and P. M. Zerwas, NLO QCD corrections to t anti-t H production in hadron collisions, Nucl. Phys. B653, 151 (2003).
- [28] M. Vanderhaeghen, J. M. Friedrich, D. Lhuillier, D. Marchand, L. Van Hoorebeke, and J. Van de Wiele, QED radiative corrections to virtual Compton scattering, Phys. Rev. C 62, 025501 (2000).
- [29] G. 't Hooft and M. J. G. Veltman, Scalar one loop integrals, Nucl. Phys. B153, 365 (1979).
- [30] D. R. Yennie, S. C. Frautschi, and H. Suura, The infrared divergence phenomena and high-energy processes, Ann. Phys. (N.Y.) 13, 379 (1961).
- [31] O. Tomalak and M. Vanderhaeghen, Subtracted dispersion relation formalism for the two-photon exchange correction to elastic electron-proton scattering: comparison with data, Eur. Phys. J. A 51, 24 (2015).
- [32] O. Tomalak and M. Vanderhaeghen, Two-photon exchange corrections in elastic muon-proton scattering, Phys. Rev. D 90, 013006 (2014).
- [33] D. Binosi and L. Theussl, JaxoDraw: A graphical user interface for drawing Feynman diagrams, Comput. Phys. Commun. 161, 76 (2004).
- [34] J. A. M. Vermaseren, Axodraw, Comput. Phys. Commun. 83, 45 (1994).