$D = 10$ dyonic black holes in string inspired models

Gaston Giribet,¹ Marcela Lagos,² Julio Oliva,² and Aldo Vera²

¹*Physics Department, New York University, 726 Broadway, New York City, New York 10003, USA*

²*Departamento de Física, Universidad de Concepción, Casilla 160C, Concepción 4030000, Chile*

(Received 14 August 2018; published 13 September 2018)

We consider string-inspired models in $D = 10$ spacetime dimensions, which include couplings with 1- and 3-form fields as well as R^4 higher-curvature corrections to the gravitational action. For such models, we explicitly construct a family of black hole solutions with both electric and magnetic charges, and with different horizon topologies. The solutions exhibit some features similar to those of self-gravitating monopoles in Einstein-Yang-Mills theory, which we discuss. When higher-curvature corrections are switched off, our solutions reduce to charged p -brane solutions previously studied in the literature. Novel qualitative features appear due to the R^4 terms, though. Such is the case for the emergence of branch singularities for charged solutions that, nonetheless, can be shielded by the event horizon.

DOI: [10.1103/PhysRevD.98.064022](https://doi.org/10.1103/PhysRevD.98.064022)

I. INTRODUCTION

String theory induces modifications to Einstein gravity which are represented by the presence of higher-curvature terms in the low-energy effective action. Type II string theories, for example, contain quartic (R^4) terms which appear at cubic order in the α' expansion. The theory also contains additional fields; in the case of type IIA strings, apart from the fields that are already present in the bosonic sector, the theory contains 1- and 3-form fields, together with their magnetic duals.

The quartic modification to Einstein-Hilbert action in the type II theories at tree level has the form

$$\int (c_1 t_8 + c_2 \epsilon_{10} \cdot \epsilon_{10}) R^4, \quad (1)$$

where $c_1 = 8c_2$ is a positive coupling, ϵ_{10} is the Levi-Civita pseudotensor in ten dimensions (which here appears with two of its indices contracted with those of a second copy of itself), and t_8 is a rank-8 tensor introduced in Ref. [1]; see also Ref. [2]. In Eq. (1), R^4 represents the tensor product of four Riemann tensors. This yields

$$12 \int d^{10}x \sqrt{-g} ((c_1 - 8c_2) (R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma})^2 + \dots), \quad (2)$$

where the ellipsis stands for other contractions of four Riemann tensors. There are, in addition, a dilaton envelope

factor and couplings between the curvature and the Kalb-Ramond field, which we omit here. Notice that for $c_1 = 8c_2$ the squared Kretschmann scalar disappears from the action, which allows Einstein spaces to persist as solutions.

The condition $c_1 \geq 0$ is crucial for a theory like (1) to avoid superluminal behavior [3]. For simplicity, we will focus on the case $c_1 = 0 \neq c_2$. Remarkably, for a particular choice of R^4 terms we will be able to write down black hole solutions that are charged under different p -form fluxes analytically. The simplification with respect to the case $c_1 \neq 0$ comes from the fact that the terms $\epsilon_{10} \cdot \epsilon_{10} R^4$ correspond to the dimensional extension of the Pfaffian whose integral, in eight dimensions, computes the Euler characteristic in virtue of the Chern-Weil-Gauss-Bonnet theorem. The topological origin of this term is what makes the field equations be of second order and, thus, analytically tractable. Being of second order, the theory is, in addition, free of Ostrogradsky instabilities; the perturbative theory results are free of ghosts around flat space [4]. Besides, a theory with such R^4 terms (provided neither quadratic nor cubic terms are present) does not exhibit the causality problems of the type discussed in Ref. [5]. All this makes the case $c_1 = 0$ an excellent arena to investigate the qualitative features that the presence of R^4 terms may introduce.

In $D = 10$ dimensions we thus consider the action

$$I = \frac{1}{16\pi\ell_p^8} \int \left(R - F_{(2)} \wedge *F_{(2)} - F_{(4)} \wedge *F_{(4)} + \frac{\alpha^3}{24} \epsilon_8 \epsilon_8 R^4 \right), \quad (3)$$

where ℓ_p is the Planck length. $F_{(2)}$ and $F_{(4)}$ are the field strengths associated with the 1-form $A_{(1)}$ and the 3-form

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

$A_{(3)}$, respectively $(F_{(p+1)} = dA_{(p)})$. Conventions are such that $\int F_{(p)} \wedge *F_{(p)} = (1/p) \int d^{10}x \sqrt{-g} F_{\mu_1 \dots \mu_p} F^{\mu_1 \dots \mu_p}$. One can also add a cosmological constant term $-1/(8\pi\ell_p^8) \times \int d^{10}x \sqrt{-g} \Lambda$. The coupling constant α has mass dimension -2 . The tensor structure in the quartic terms is given by $\epsilon_8 \epsilon_8 = (1/2) \epsilon_{10} \cdot \epsilon_{10}$, which can be conveniently written as

$$\epsilon_8 \epsilon_8 R^4 = \delta_{\nu_1 \nu_2 \dots \nu_8}^{\mu_1 \mu_2 \dots \mu_8} R^{\nu_1 \nu_2}_{\mu_1 \mu_2} R^{\nu_3 \nu_4}_{\mu_3 \mu_4} R^{\nu_5 \nu_6}_{\mu_5 \mu_6} R^{\nu_7 \nu_8}_{\mu_7 \mu_8}, \quad (4)$$

where δ is the totally antisymmetric Kronecker symbol [6]. The sign $\alpha^3 > 0$ is the one that yields a consistent model; see Ref. [3].

For the theory above, we will write down analytic dyonic black hole solutions charged under both the 1- and 3-form fields, and with different horizon topologies. The local geometry of the base manifolds (i.e., of the constant-time sections of the horizons) of these solutions will be the direct product of 2^m copies of (2^{3-m}) -dimensional constant-curvature manifolds, with $m = \{0, 1, 2, 3\}$. For example, in the case of such constant-curvature manifolds being (2^{3-m}) -spheres, the case $m = 0$ would correspond to the ten-dimensional, R^4 -corrected Reissner-Nordström black hole, whose base manifold is an 8-sphere. The other extreme of the list, the case $m = 3$, would correspond to a topological black hole with flat a horizon, namely, a black brane. Higher-genus topological black holes with compact horizons require to consider as a base manifold the product of Fuchsian quotients of hyperbolic spaces. The two most interesting and less-simple examples are black hole solutions whose horizons are either products of four 2-spheres or of two 4-spheres. These solutions are supported by magnetic fluxes on the spheres and admit net electric charges. The latter are the cases we will study in this paper.

II. BLACK HOLE SOLUTIONS

A. The solution with $S^2 \times S^2 \times S^2 \times S^2$

Let us start with the case $m = 2$, that is, solutions whose base manifolds are a direct product of four constant-curvature 2-manifolds. We begin by considering 2-spheres. The configuration of the 3-form for such a solution is given by

$$F_{(4)} = \frac{Q_2}{r^4} dt \wedge dr \wedge \sum_{i=1}^4 \text{vol}(S_i^2) + P_2 \sum_{i < j}^4 \text{vol}(S_i^2) \wedge \text{vol}(S_j^2), \quad (5)$$

where $\text{vol}(S_i^2)$ stands for the volume form of the i th 2-sphere S_i^2 , $i = \{1, 2, 3, 4\}$. Q_2 and P_2 are two integration constants associated with the electric and magnetic charges under the 3-form field $A_{(3)}$. Similarly, the configuration of the 1-form is given by

$$F_{(2)} = \frac{Q_0}{r^8} dt \wedge dr, \quad (6)$$

which corresponds to the Coulombian potential $A_{(1)} = Q_0/(7r^7)dt$.

The metric takes the form

$$ds^2 = -H(r)dt^2 + \frac{dr^2}{H(r)} + r^2 \sum_{i=1}^4 \frac{dz_i d\bar{z}_i}{(1 + \frac{\sigma}{4} z_i \bar{z}_i)^2}, \quad (7)$$

where $(z_i, \bar{z}_i)$ are complex projective variables on the i th 2-(pseudo)sphere. $\sigma = 1$ corresponds to the metric of the base manifold being a product of four unit 2-spheres. The cases $\sigma = 0, -1$ correspond to the locally flat spaces and locally hyperbolic spaces, respectively. H is a function of the radial coordinate r , which is given by the fourth-order polynomial equation

$$\alpha^3 r^8 (g_0 H^4 - g_1 \sigma H^3 + g_2 |\sigma| H^2 - g_3 \sigma H + g_4 |\sigma|) - g_5 r^{14} H = T, \quad (8)$$

with the matter content given by

$$T = -t_1 Q_0^2 - t_2 Q_2^2 r^4 + t_3 \ell_p^8 M r^7 + t_4 P_2^2 r^8 - t_5 \sigma r^{14},$$

where the coefficients g_i and t_i in their prime factorization forms are

$$\begin{aligned} g_0 &= 2^7 \cdot 3^4 \cdot 5^2 \cdot 7^2, & g_1 &= 2^9 \cdot 3^4 \cdot 5^2 \cdot 7, \\ g_2 &= 2^8 \cdot 3^5 \cdot 5 \cdot 7, & g_3 &= 2^9 \cdot 3^3 \cdot 5 \cdot 7, \\ g_4 &= 2^7 \cdot 3^3 \cdot 5 \cdot 7, & g_5 &= 2^3 \cdot 3^2 \cdot 5 \cdot 7, \end{aligned}$$

and

$$\begin{aligned} t_1 &= 3^2 \cdot 5, & t_2 &= 2^3 \cdot 3^2 \cdot 5 \cdot 7, \\ t_3 &= 2^5 \cdot 3^2 \cdot 5 \cdot 7\pi, & t_4 &= 2^2 \cdot 3^4 \cdot 5 \cdot 7, \\ t_5 &= 2^3 \cdot 3^2 \cdot 5. \end{aligned}$$

If $(\hat{\lambda}\alpha/64\pi\ell_p^8) \int d^{10}x \sqrt{-g} \delta_{\nu_1 \nu_2 \nu_3 \nu_4}^{\mu_1 \mu_2 \mu_3 \mu_4} R^{\nu_1 \nu_2}_{\mu_1 \mu_2} R^{\nu_3 \nu_4}_{\mu_3 \mu_4}$ is also included in the action, then the left-hand side of Eq. (8) receives the piece $\hat{\lambda}\alpha r^{12}(g_6 H^2 - g_7 \sigma H + g_8 |\sigma|)$ with $g_6 = 2^4 \cdot 3^3 \cdot 5 \cdot 7^2$, $g_7 = 2^5 \cdot 3^3 \cdot 5 \cdot 7$, and $g_8 = 2^4 \cdot 3^3 \cdot 7$.

In the case of a nonvanishing cosmological constant ($\Lambda \neq 0$), the matter content T receives an additional term $70\Lambda r^{16}$. The general relativity (GR) limit (or, equivalently, the large- r^2/α limit) of the solution yields

$$H \simeq -\frac{r^2 \Lambda}{36} + \frac{\sigma}{7} - \frac{4\pi\ell_p^8 M}{r^7} - \frac{9P_2^2}{2r^6} + \frac{Q_2^2}{r^{10}} + \frac{Q_0^2}{56r^{14}} + \dots, \quad (9)$$

where the ellipsis stands for subleading terms in orders of α/r^2 . This is consistent with the results for charged p -brane

solutions with nonspherical horizons in Einstein theory.¹ Solutions with base manifolds that are not constant-curvature manifolds were studied, for instance, in Refs. [9,10] (see also references therein).

Notice that, in Eq. (9), the term with the charge P_2 presents a dampening that is weaker than the Newtonian potential in ten dimensions, which typically leads to a divergent gravitational energy. This is reminiscent of what happens with self-gravitating monopoles in Einstein-Yang-Mills theory [11]. In addition, the sign of such an energy contribution to the gravitational potential seems to introduce instabilities.

The electrically charged solution for finite α also presents curious features. In particular, there exists a branch singularity that occurs at a finite distance that, in principle, can be smaller than the horizon location r_+ . Such a branch singularity happens when radicands in the solution to a polynomial equation such as Eq. (8) take negative values. This happens because of the relative signs between the Newtonian term and the terms with the charges $Q_{0,2}$, which makes the radicands vanish for finite r . This results in nonreal components of the metric and divergences in the curvature invariants, which are typical features of charged solutions in Lovelock theory (see Fig. 1).

It is worth mentioning that, if we add to the action the term $(\lambda\alpha/64\pi\ell_p^8) \int d^{10}x \sqrt{-g} (F_{\mu_1\mu_2\mu_3\mu_4} F^{\mu_1\mu_2\mu_3\mu_4})^2$ and consider the case $P_2 = Q_0 = 0$, then Eq. (8) receives the following contribution on the right-hand side:

$$2^3 \cdot 3^3 \cdot 5 \cdot 7 r^7 \int_{r_0}^r dr X^2(r) (2^5 3^2 \alpha \lambda + r^4 \lambda), \quad (10)$$

where $X(r)$ is the solution to the equation

$$X(r)r^4 + 2^6 3 \alpha \lambda X^3(r) = Q_2. \quad (11)$$

The ultraviolet cutoff r_0 above contributes to the mass.

B. The solution with $S^4 \times S^4$

Solutions of a similar type exist for base manifolds that are the direct product of two constant-curvature 4-manifolds. This corresponds to $m = 1$. In this case, the magnetic configuration of the 3-form is given by

$$F_{(4)} = \hat{P}_2 \sum_{i=1}^2 \text{vol}(S_i^4),$$

$$\text{vol}(S_i^4) = \frac{dz_i \wedge d\bar{z}_i \wedge dw_i \wedge d\bar{w}_i}{(1 + \frac{1}{4} z_i \bar{z}_i + \frac{1}{4} w_i \bar{w}_i)^4}, \quad (12)$$

where $\text{vol}(S_i^4)$ stands for the volume form of the 4-sphere S_i^4 , $i = \{1, 2\}$. Again, the integration constant $\hat{P}_2$ is

¹We thank Marcello Ortaggio for calling our attention to Refs. [7,8] which contain black hole solutions to which ours tend in the limit $\alpha^3 \rightarrow 0$.

FIG. 1. Lapse functions for the case $(S^2)^4$ charged under a Maxwell field and uncharged under $F_{(4)}$. We use $M = 10^{-3}$ and the doublets (Q_0, α) equal to $(0, 0)$ (blue), $(0, 8 \times 10^{-2})$ (green), $(10^{-2}, 8 \times 10^{-2})$ (red), and $(1.55 \times 10^{-2}, 8 \times 10^{-2})$ (purple).

associated with the magnetic charge. The metric for this solution (with $\sigma = 1$) takes the form

$$ds^2 = -\hat{H}(r)dt^2 + \frac{dr^2}{\hat{H}(r)} + r^2 \sum_{i=1}^2 \frac{dz_i d\bar{z}_i + dw_i d\bar{w}_i}{(1 + \frac{1}{4} z_i \bar{z}_i + \frac{1}{4} w_i \bar{w}_i)^2}. \quad (13)$$

We have that the local geometry of the base manifold is now given by a product of two unit 4-spheres (or, more generally, of a pair of identical 4-manifolds of constant-curvature $\sigma = \{0, \pm 1\}$). For this case, $\hat{H}$ is given by the following fourth-order polynomial equation:

$$\alpha^3 r (\hat{g}_0 \hat{H}^4 - \hat{g}_1 \sigma \hat{H}^3 + \hat{g}_2 |\sigma| \hat{H}^2 - \hat{g}_3 \sigma \hat{H} + \hat{g}_4 |\sigma|) + (-\hat{g}_5 \hat{H} + \hat{g}_6 \sigma) r^7 = T, \quad (14)$$

with the matter contribution

$$T = \hat{t}_1 \ell_p^8 M + \hat{t}_2 \hat{P}_2^2 r + \hat{t}_3 \Lambda r^9, \quad (15)$$

where the coefficients $\hat{g}_i$ are given by

$$\begin{aligned} \hat{g}_0 &= 2^6 \cdot 3^4 \cdot 5^2 \cdot 7^2, & \hat{g}_1 &= 2^8 \cdot 3^5 \cdot 5^2 \cdot 7, \\ \hat{g}_2 &= 2^7 \cdot 3^5 \cdot 5 \cdot 7^2, & \hat{g}_3 &= 2^8 \cdot 3^5 \cdot 5 \cdot 7, \\ \hat{g}_4 &= 2^6 \cdot 3^5 \cdot 5 \cdot 7, & \hat{g}_5 &= 2^2 \cdot 3^2 \cdot 5 \cdot 7, \\ \hat{g}_6 &= 2^2 \cdot 3^3 \cdot 5, \end{aligned}$$

FIG. 2. Lapse functions for the case $(S^2)^4$ dyonic- $F_{(4)}$ black holes uncharged under a Maxwell field. We use $\alpha = 0.08$, $Q_0 = 0$, and $M = 10^{-3}$ as well as the doublets (Q_2, P_2) with $(1.45 \times 10^{-2}, 2 \times 10^{-2})$ (purple), $(10^{-2}, 2 \times 10^{-2})$ (green), and $(10^{-2}, 8 \times 10^{-2})$ (red).

and the coefficients $\hat{t}_i$ are given by

$$\hat{t}_1 = 2^4 \cdot 3^2 \cdot 5 \cdot 7\pi, \quad \hat{t}_2 = 2 \cdot 3^3 \cdot 5 \cdot 7, \quad \hat{t}_3 = 5 \cdot 7.$$

The GR limit of this solution yields

$$\hat{H} \simeq -\frac{\Lambda r^2}{36} + \frac{3\sigma}{7} - \frac{3\hat{P}_2^2}{2r^6} - \frac{4\pi\ell_P^8 M}{r^7} + \dots, \quad (16)$$

where the ellipsis stands for subleading terms in orders of α/r^2 . Again, we observe the weakened dampening of the term with charge $\hat{P}_2$. The fact that the GR solution (16) is recovered when $\alpha = 0$ shows that we are considering a perturbative solution of the higher-curvature theory. These theories typically present up to four different branches of static solutions, some of which are nonexistent (divergent) in the $\alpha \rightarrow 0$ limit (see Fig. 2).

If $(\hat{\lambda}\alpha/64\pi\ell_P^8) \int d^{10}x \sqrt{-g} \delta_{\nu_1\nu_2\nu_3\nu_4}^{\mu_1\mu_2\mu_3\mu_4} R^{\nu_1\nu_2}_{\mu_1\mu_2} R^{\nu_3\nu_4}_{\mu_3\mu_4}$ is included in the action, then the left-hand side of Eq. (14) receives the piece $\hat{\lambda}\alpha r^5(\hat{g}_7|\sigma| - \hat{g}_8\sigma\hat{H} + \hat{g}_9\hat{H}^2)$ with $\hat{g}_7 = 2^3 \cdot 3^3 \cdot 7^2$, $\hat{g}_8 = 2^4 \cdot 3^4 \cdot 5 \cdot 7$, and $\hat{g}_9 = 2^3 \cdot 3^3 \cdot 5 \cdot 7^2$, and it also leads to a solvable equation.

C. Thermodynamics

The new qualitative features introduced by the higher-curvature terms in the action can also be analyzed by taking a glance at the thermodynamics of the solution. For instance, consider the black hole solution with $m = 1$ and

$P_2 = 0 = \Lambda$, whose Hawking temperature, including both R^2 and R^4 terms, is given by

$$T_H = \frac{8\sigma r_+^{14} + 48\hat{\lambda}|\sigma|r_+^{12} + 384\alpha^3|\sigma|r_+^8 - 24Q_2^2 r_+^4 - Q_0^2}{32\pi r_+^9 (192\alpha^3\sigma + 12\hat{\lambda}\sigma r_+^4 + r_+^6)}. \quad (17)$$

This is obtained by computing $T_H = H'(r_+)/ (4\pi)$. From Eq. (17) we can see the short-distance modifications to the GR behavior. At large r_+^2/α , Eq. (17) clearly reproduces the thermodynamical behavior of asymptotically flat black hole solutions to Einstein's equations. In contrast, provided $\alpha \neq 0$, the small- r_+ behavior deviates from that of GR.

Short-distance contributions are also seen in the black hole entropy formula. The entropy, which can be computed by using the Wald formula, takes the form

$$S_{\text{BH}} = \frac{(4\pi)^4 r_+^8}{4\ell_P^8} + \frac{4\sigma\hat{\lambda}\alpha r_+^6}{\ell_P^8} + \frac{192(4\pi)^4 \alpha^3 \sigma r_+^2}{\ell_P^8}, \quad (18)$$

from which we observe that the Bekenstein-Hawking area-law term gets supplemented by contributions of higher-curvature terms that vanish in the limit $\alpha \rightarrow 0$. Such terms scale with lower powers of the volume, as expected from dimensional analysis. Unlike the term proportional to the horizon area, the sign of the other two terms is sensitive to the curvature of the base manifold (σ) as well as to the sign of the coupling constant of the higher-curvature terms (α). Different choices of those signs lead to different qualitative behaviors, including some pathological ones. Such a dependence on the signs of the couplings is also observed in the expression for the temperature, although, as expected, it becomes relevant only at short distances.

III. CONCLUSIONS

We have derived analytic dyonic black hole solutions for string-inspired models that include R^4 terms together with $F_{(2)}$ and $F_{(4)}$ fluxes, with different horizon topologies. We focused our attention on those ten-dimensional solutions whose base manifolds are given by the tensor product of constant-curvature manifolds. We have shown that magnetically charged solutions of this type exhibit weakened asymptotics that are reminiscent of those of self-gravitating monopoles in Einstein-Yang-Mills theory. The solutions with no magnetic charges, in contrast, have finite gravitational energy.

We have also shown that the higher-curvature terms make the electrically charged solutions develop a branch singularity at a finite distance, which can be well shielded by the event horizon. It is remarkable that, even in the case in which both R^2 and $F_{(4)}$ terms are included in the action, the particular Lovelock-type R^4 tensor structure we considered here allowed us to write down dyonic black hole solutions in a closed form. This provides a tractable model to study the physics of higher-curvature effects analytically.

ACKNOWLEDGMENTS

The authors thank Marcello Ortaggio for discussions. The work of G. G. is supported by the NSF through Grant No. PHY-1214302. M. L. and A. V. appreciate the support

of CONICYT Fellowships No. 21141229 and No. 21151067, respectively. J. O. is supported by CONICYT Grant No. 1181047.

-
- [1] D.J. Gross and E. Witten, Superstring modifications of Einstein's equations, *Nucl. Phys.* **B277**, 1 (1986).
 - [2] S. Giusto and S. D. Mathur, Fuzzball geometries and higher derivative corrections for extremal holes, *Nucl. Phys.* **B738**, 48 (2006).
 - [3] A. Gruzinov and M. Kleban, Causality constrains higher curvature corrections to gravity, *Classical Quantum Gravity* **24**, 3521 (2007).
 - [4] B. Zwiebach, Curvature squared terms and string theories, *Phys. Lett. B* **156**, 315 (1985).
 - [5] X.O. Camanho, J.D. Edelstein, J. Maldacena, and A. Zhiboedov, Causality constraints on corrections to the graviton three-point coupling, *J. High Energy Phys.* **02** (2016) 020.
 - [6] D. Lovelock, The Einstein tensor and its generalizations, *J. Math. Phys.* **12**, 498 (1971).
 - [7] Y. Bardoux, M. M. Caldarelli, and C. Charmousis, Shaping black holes with free fields, *J. High Energy Phys.* **05** (2012) 054.
 - [8] Y. Bardoux, C. Charmousis, and T. Kolyvaris, Lovelock solutions in the presence of matter sources, *Phys. Rev. D* **83**, 104020 (2011).
 - [9] G. Gibbons and S.A. Hartnoll, A gravitational instability in higher dimensions, *Phys. Rev. D* **66**, 064024 (2002).
 - [10] G. Dotti and R.J. Gleiser, Obstructions on the horizon geometry from string theory corrections to Einstein gravity, *Phys. Lett. B* **627**, 174 (2005).
 - [11] G.W. Gibbons and P.K. Townsend, Self-gravitating Yang monopoles in all dimensions, *Classical Quantum Gravity* **23**, 4873 (2006).