Erratum: Refined gluino and squark pole masses beyond leading order **[Phys. Rev. D 74, 075009 (2006)]**

Stephen P. Martin

(Received 11 December 2018; published 27 December 2018)

DOI: [10.1103/PhysRevD.98.119901](https://doi.org/10.1103/PhysRevD.98.119901)

In Eq. (A11), $C_a(G)$ should be $C_a(A)$.

In Eqs. (A17) and (A18), there were several incorrect coefficients. These equations should read:

$$x^{(2)} = h_a^2 C_a(F) \{ C_a(A) [-1111/24 + 4\pi^2/3 - 4\pi^2 \ln 2 + 6\zeta_3] + S_a^L [71 + 8\pi^2]/12 + S_a^H [143 - 16\pi^2]/12 \} + h_a h_b C_a(F) C_b(F) [8\pi^2 \ln 2 - 5\pi^2 - 12\zeta_3 + 7/8], \quad (\text{A17})$$

and

$$x^{(3)} = 64 h_a C_a(F) [h_b h_c C_b(F) C_c(F) d_1^{(3)} + h_a h_b \{ C_a(A) C_b(F) d_2^{(3)} + S_{ab}^L d_{4A}^{(3)}/2 + S_a^L C_b(F) d_{4B}^{(3)}/2 + S_{ab}^H d_5^{(3)}/2 \} + h_a^2 \{ [C_a(A)]^2 d_3^{(3)} + C_a(A) S_a^L d_6^{(3)}/2 + C_a(A) S_a^H d_7^{(3)}/2 + S_a^L S_a^H d_8^{(3)}/4 + [S_a^H]^2 d_9^{(3)}/4 + [S_a^L]^2 d_{10}^{(3)}/4 \}]. \quad (\text{A18})$$