Low energy neutrinos from stopped muons in the Earth

Wan-Lei Guo*

Institute of High Energy Physics, Chinese Academy of Sciences, P.O. Box 918, Beijing 100049, China

(Received 17 December 2018; published 23 April 2019)

We explore the low energy neutrinos from stopped cosmic ray muons in the Earth. Based on the muon intensity at the sea level and the muon energy loss rate, the depth distributions of stopped muons in the rock and sea water can be derived. Then we estimate the μ^- decay and nuclear capture probabilities in the rock. Finally, we calculate the low energy neutrino fluxes and find that they depend heavily on the detector depth d . For $d = 1000$ m, the ν_e , $\bar{\nu}_e$, ν_μ , and $\bar{\nu}_\mu$ fluxes in the range of $13 \text{ MeV} \leq E_\nu \leq 53 \text{ MeV}$ are averagely 10.8%, 6.3%, 3.7%, and 6.2% of the corresponding atmospheric neutrino fluxes, respectively. The above results will be increased by a factor of 1.4 if the detector depth $d < 30$ m. In addition, we find that most neutrinos come from the region within 200 km and the near horizontal direction, and the $\bar{\nu}_e$ flux depends on the local rock and water distributions.

DOI: 10.1103/PhysRevD.99.073007

I. INTRODUCTION

Atmospheric neutrinos are a very important neutrino source to study the neutrino oscillation physics. In 1998, the Super-Kamiokande (Super-K) experiment reported the first evidence of neutrino oscillations based on a zenith angle dependent deficit of atmospheric muon neutrinos [1]. Atmospheric neutrinos are produced in the Earth's atmosphere as a result of cosmic ray interactions and the weak decays of secondary mesons, in particular pions and kaons [2]. At the same time, a large amount of muons are also produced and some of them can penetrate the rock and sea water of Earth's surface to significant depths. These penetrating muons are the important background source for some underground experiments [3]. It is well known that these muons will finally stop in the Earth and then produce the low energy neutrinos through decay or nuclear capture [4]. However, these neutrinos are not included in the previous literatures [5–7]. Here we shall focus on these neglected neutrinos from stopped muons in the Earth.

Muons are the most numerous charged particles at sea level [8]. After losing energy by ionization and radiative processes, the stopped μ^+ in the rock and sea water will generate two low energy neutrinos ν_e and $\bar{\nu}_\mu$ ($E_\nu \leq 53 \text{ MeV}$) through $\mu^+ \rightarrow e^+ + \nu_e + \bar{\nu}_\mu$. Unlike μ^+ , the stopped μ^- may undergo either decay or capture by the nucleus [4]. In the nuclear capture case, a stopped μ^- can only produce a

neutrino ν_μ with energy less than the muon mass. These low energy neutrinos will be the background source in the searches of some relevant physics, such as diffuse supernova relic neutrinos [9,10], dark matter annihilation in the Sun [11] and our galaxy [12], solar neutrino conversion [13], and proton decays catalyzed by grand unified theory (GUT) monopoles in the Sun [14]. So it is necessary for us to investigate the low energy neutrinos induced by stopped muons in the Earth.

In this paper, we shall calculate the neutrino fluxes from stopped cosmic ray muons in the Earth. In Sec. II, the stopped $\mu^\pm$ distributions in the rock and sea water will be given in terms of the muon intensity at the sea level and the muon energy loss rate. Section III is devoted to the $\bar{\nu}_e$ and ν_μ energy spectra from a stopped μ^- . Based on the atomic capture and nuclear capture abilities of 10 dominant elements in the upper continental crust, we estimate the μ^- decay and nuclear capture probabilities in the rock and sea water. In Sec. IV, we numerically calculate the low energy neutrino fluxes according to the stopped $\mu^\pm$ distributions and the μ^- decay probability, and discuss their features. In addition, an approximation formula to compute the neutrino fluxes has also been presented. Finally, our conclusions will be given in Sec. V.

II. DISTRIBUTIONS OF STOPPED MUONS

Muons are the most numerous charged particles at sea level [8]. For the energy and angular distribution of cosmic ray muons at the sea level, we use the following parametrization [15]

$$I(p_\mu, \theta) = I_v(\xi) \cos^3 \theta \quad \text{for } p_\mu > 1 \text{ GeV}, \quad (1)$$

*guowl@ihep.ac.cn

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](#). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

with $\xi = p_\mu \cos \theta$ and θ is the zenith angle. The vertical muon intensity is given by

$$I_v(p_\mu) = c_1 p_\mu^{-(c_2 + c_3 \log p_\mu + c_4 \log^2 p_\mu + c_5 \log^3 p_\mu)}, \quad (2)$$

where $c_1 = 0.00253 \text{ cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1} \text{ GeV}^{-1}$, $c_2 = 0.2455$, $c_3 = 1.288$, $c_4 = -0.2555$, and $c_5 = 0.0209$. It is found that Eq. (1) is valid for all zenith angles and the muon momentum $p_\mu > 1 \text{ GeV}$ [15]. For $p_\mu \leq 1 \text{ GeV}$, the muon energy spectrum is almost flat and the corresponding angular distribution is steeper than $\cos^2 \theta$ [8]. Therefore we assume

$$I(p_\mu, \theta) = 0.00389 \cos^3 \theta \quad \text{for } p_\mu \leq 1 \text{ GeV}, \quad (3)$$

in the units of $\text{cm}^{-2} \text{ s}^{-1} \text{ sr}^{-1} \text{ GeV}^{-1}$. In terms of Eqs. (1)–(3), the total muon flux for a horizontal detector can be derived from

$$J_\mu = 2\pi \int I(p_\mu, \theta) \cos \theta d\cos \theta dp_\mu = 1 \text{ cm}^{-2} \text{ min}^{-1}, \quad (4)$$

which is familiar to experimentalists for horizontal detectors [8]. The muons with $p_\mu > 1 \text{ GeV}$ make a 71% contribution to J_μ . With the help of the muon charge ratio in Fig. 6 of Ref. [16], we calculate the μ^+ flux $J_{\mu^+} = 0.0093 \text{ cm}^{-2} \text{ s}^{-1}$ and the μ^- flux $J_{\mu^-} = 0.00737 \text{ cm}^{-2} \text{ s}^{-1}$, and the corresponding muon charge ratio $J_{\mu^+}/J_{\mu^-} = 1.26$.

The long-lived muons can penetrate the rock and sea water of the Earth's surface to significant depths. The muon range $X(p_\mu)$ in the standard rock and water may be found in Ref. [17]. Then one can easily get the stop depth $x = X(p_\mu) \cos \theta$ for a muon with the momentum p_μ and incidence zenith angle θ at the sea level. Based on the muon distribution $I(p_\mu, \theta)$, muon charge ratio [16] and muon range $X(p_\mu)$ [17], we calculate the $\mu^\pm$ stop rate per unit volume $S_{\mu^\pm}(x)$ in the standard rock and water. In Fig. 1, the $\mu^\pm$ stop

rate $S_{\mu^\pm}(x)$ and the charge ratio of stopped muons S_{μ^+}/S_{μ^-} as a function of stop depth x have been plotted. It is clear that the depth x of most stopped muons is less than 30 m.

III. NEUTRINO ENERGY SPECTRA FROM A STOPPED MUON

It is well known that a stopped μ^+ will quickly decay into a positron and two neutrinos through $\mu^+ \rightarrow e^+ + \nu_e + \bar{\nu}_\mu$. The ν_e and $\bar{\nu}_\mu$ energy spectra (normalized to 1) can be written as [18]

$$f_{\nu_e} = \frac{192}{m_\mu} \left[\left(\frac{E_\nu}{m_\mu} \right)^2 \left(\frac{1}{2} - \frac{E_\nu}{m_\mu} \right) \right], \quad (5)$$

$$f_{\bar{\nu}_\mu} = \frac{64}{m_\mu} \left[\left(\frac{E_\nu}{m_\mu} \right)^2 \left(\frac{3}{4} - \frac{E_\nu}{m_\mu} \right) \right], \quad (6)$$

where m_μ is the muon mass and $E_\nu \leq m_\mu/2$. Unlike μ^+ , a stopped μ^- can not only decay $\mu^- \rightarrow e^- + \bar{\nu}_e + \nu_\mu$, but also be captured by nucleus and produce a neutrino ν_μ with $E_\nu < m_\mu$, such as $\mu^- + {}^{16}\text{O} \rightarrow {}^{16}\text{N}^* + \nu_\mu$. In fact, the stopped μ^- will be quickly attached to an atom and form a muonic atom (atomic capture) [4] when it stops in the rock or sea water. Then it cascades down to the lowest 1s level in a timescale of the order of 10^{-13} s through emitting Auger electrons and muonic x-rays. In the following time, the bounded μ^- in a muonic atom has only two choices, to decay or to capture on the nucleus (nuclear capture) [4].

In order to estimate the μ^- decay and nuclear capture probabilities, we should first consider the relative abilities of atomic capture for different elements in the rock. Egidy and Hartmann [19] find a semiempirical approach and give the average atomic capture probability $P(Z)$ for 65 elements, normalized to 1 for ${}^{16}\text{O}$. For the rock chemical composition, we take the upper continental crust data from Ref. [20]. Then the mass and number percentages of 10 dominant elements in

FIG. 1. The muon stop rate per unit volume $S_{\mu^\pm}$ (left) and the charge ratio of stopped muons S_{μ^+}/S_{μ^-} (right) as a function of stop depth x in the standard rock and water.

TABLE I. The μ^- atomic capture percentages and decay probabilities D_{μ^-} in a muonic atom for 10 dominant elements of the upper continental crust [20]. The corresponding mass and number percentages, average atomic capture probability $P(Z)$, μ^- mean life τ_{μ^-} , and Huff factor Q have also been listed.

Elements	Mass (%)	Number (%)	$P(Z)$	Atomic capture (%)	τ_{μ^-} (ns)	Huff factor	D_{μ^-} (%)
O	47.51	62.13	1.00	60.26	1795.4	0.998	81.56
Si	31.13	23.89	0.84	19.46	756	0.992	34.14
Al	8.15	3.91	0.76	2.88	864	0.993	39.05
Fe	3.92	2.27	3.28	7.21	206	0.975	9.14
Ca	2.57	2.07	1.90	3.81	332.7	0.985	14.92
Na	2.43	2.27	1.00	2.21	1204	0.996	54.58
K	2.32	1.28	1.54	1.91	435	0.987	19.54
Mg	1.50	1.99	0.93	1.79	1067.2	0.995	48.33
Ti	0.38	0.17	2.66	0.45	329.3	0.981	14.70
P	0.07	0.02	1.04	0.02	611.2	0.991	27.57

the upper continental crust have been calculated and listed in Table I. Considering the corresponding atomic capture probability $P(Z)$, we derive the atomic capture percentages of 10 elements as shown in the fifth column of Table I. It is worthwhile to stress that the water is a ^{16}O target since μ^- p can easily penetrate nearby ^{16}O atoms [4].

The decay rate Λ_{decay} and nuclear capture rate Λ_{capture} of the bounded μ^- in a muonic atom have the following relation [4]

$$\Lambda_{\text{total}} = \Lambda_{\text{capture}} + Q\Lambda_{\text{decay}}, \quad (7)$$

where $\Lambda_{\text{total}} = 1/\tau_{\mu^-}$, $\Lambda_{\text{decay}} = 1/\tau_{\mu^+}$, and Q is the Huff factor [21]. Then the μ^- decay probability can be easily obtained by

$$D_{\mu^-} = Q \frac{\Lambda_{\text{decay}}}{\Lambda_{\text{total}}} = Q \frac{\tau_{\mu^-}}{\tau_{\mu^+}}. \quad (8)$$

With the help of $\tau_{\mu^+} = 2196.98$ ns [8] and the μ^- mean life in Ref. [21], we calculate the μ^- decay probabilities D_{μ^-} for 10 dominant elements as listed in the last column of Table I. Combining the atomic capture percentages and the corresponding D_{μ^-} in Table I, one can find that the averaged decay probability $D_{\mu^-} = 60.65\%$ and nuclear capture probability $C_{\mu^-} = 39.35\%$ for negative muons stopped in the rock. Since the water can be approximated as an ^{16}O target, the μ^- decay and nuclear capture probabilities in the water are $D_{\mu^-} = 81.56\%$ and $C_{\mu^-} = 18.44\%$, respectively. For the $\bar{\nu}_e$ and ν_μ energy spectra, we ignore the differences between the free μ^- decay and the bounded μ^- decay [4], and have

$$f_{\bar{\nu}_e} = f_{\nu_e} D_{\mu^-}, \quad (9)$$

$$f_{\nu_\mu} = f_{\bar{\nu}_\mu} D_{\mu^-} + \tilde{f}_{\nu_\mu} C_{\mu^-}, \quad (10)$$

where $\tilde{f}_{\nu_\mu}$ is the ν_μ energy spectrum (normalized to 1) from the μ^- nuclear capture. It is found that $\tilde{f}_{\nu_\mu}$ is fairly similar to

the γ spectrum in the reaction of the π^- capture on nucleus [4]. Therefore we use the γ spectrum from the $^{16}\text{O}(\pi^-, \gamma)^{16}\text{N}^*$ experimental results [22] and require that the maximal neutrino energy only reaches 95 MeV for $\tilde{f}_{\nu_\mu}$, because the muon mass is 34 MeV less than that of a pion.

IV. NEUTRINO FLUXES FROM STOPPED MUONS

Since the produced neutrinos from stopped muons are isotropic, the neutrino differential fluxes can be written as

$$\frac{d\phi_{\nu_i}}{dE_\nu} = f_{\nu_i} \int S_{\mu^\pm}(x) \frac{2\pi(R_\oplus - x)^2 \sin \vartheta}{4\pi r^2} d\vartheta dx, \quad (11)$$

where $R_\oplus = 6371$ km is the Earth's radius, and ϑ is the angle between the stopped $\mu^\pm$ and detector point seen from the Earth's center. The distance between the stopped $\mu^\pm$ and detector is given by

$$r^2 = (R_\oplus - x)^2 + (R_\oplus - d)^2 - 2 \cos \vartheta (R_\oplus - x)(R_\oplus - d), \quad (12)$$

where d is the detector depth. In fact, the neutrino oscillation should be considered [23], but it is beyond the scope of this paper. For a large neutrino detector, different detector parts will receive notably different neutrino fluxes from stopped muons within $r < 100$ m. Therefore we take a virtual spherical detector with a 25 m radius for the following analysis, which will increase about 1% flux for the $d < 30$ m case. The integral flux ϕ_{ν_i} can be obtained from Eq. (11). In the left panel of Fig. 2, we plot the cumulative percentage of ϕ_{ν_i} as a function of surface distance $L = R_\oplus \vartheta$ for three typical detector depths in the rock case. Different flavors have almost identical results even in the water case. It is worthwhile to stress that the stopped muons within the surface distance $L = 200$ km contribute 68.7% and 56.1% of ϕ_{ν_i} for $d = 0$ m and $d = 1000$ m, respectively. In addition, one may easily find

FIG. 2. The cumulative percentage of the neutrino integral flux ϕ_{ν_i} as a function of surface distance $L = R_\oplus \vartheta$ (left) and the ν_e integral flux ϕ_{ν_e} with $E_\nu < 53$ MeV as a function of zenith angle θ (right).

that most neutrinos come from the near horizontal direction. Since the recoil electrons can carry the directional information of incident neutrinos in the elastic scattering of neutrinos on electrons [14] and ν_e has the largest cross section, we calculate the zenith angular distribution of ν_e integral flux in the rock case, as shown in the right panel of Fig. 2. It is found that the smaller d case has the narrower peak at the horizontal direction. Note that the underground part of the virtual spherical detector contributes the $\cos \theta > 0$ angular distribution for the depth of detector center $d = 0$ m case. $\bar{\nu}_e$, ν_μ , and $\bar{\nu}_\mu$ have similar angular distributions even in the water case.

In Fig. 3, we show the neutrino integral fluxes ϕ_{ν_i} as a function of detector depth d for different flavors in the rock and water cases. It is clear that ϕ_{ν_i} depends heavily on the detector depth d . For ϕ_{ν_e} , $\phi_{\bar{\nu}_e}$, and ϕ_{ν_μ} , the differences between the rock and water cases are very small. However, the ν_μ differential fluxes in the rock and water cases have obvious differences due to different values of D_{μ^-} and C_{μ^-}

FIG. 3. The neutrino flux ϕ_{ν_i} as a function of detector depth d for different neutrino flavors.

in Eq. (10). Note that $\phi_{\bar{\nu}_e}$ in the water case is larger than that in the rock case because of the larger D_{μ^-} in the water case. Therefore, the $\bar{\nu}_e$ flux depends on the local rock and water distributions for a given detector.

Before presenting the differential neutrino fluxes, we here introduce an approximation formula to calculate the neutrino fluxes. Since the depth of most stopped muons is less than 30 m as shown in Fig. 1, one may assume $S_{\mu^\pm}(x) = J_{\mu^\pm} \delta(x)$ and subsequently simplify Eq. (11) to

$$\frac{d\phi_{\nu_i}}{dE_\nu} = f_{\nu_i} J_{\mu^\pm} A(d), \quad (13)$$

where $A(d) = 0.5 R_\oplus^2 \int \sin \vartheta r_{x=0}^{-2} d\vartheta$ expresses the conversion factor from the muon flux $J_{\mu^\pm}$ at the sea level to the neutrino flux ϕ_{ν_i} for a detector with a depth d , and $A(R_\oplus) = 1$. For $d < 10^4$ m, $A(d)$ can be approximated as

$$A(d) = \text{Min}[6.62, 8.39 - 1.21 \log d], \quad (14)$$

where d is in units of meter. It is found that Eqs. (13) and (14) can describe the exact results shown in Fig. 3 with an error of 2%.

By use of Eq. (11), we calculate the differential neutrino fluxes as shown in the left panel of Fig. 4. Here $D_{\mu^-} = 70\%$ ($C_{\mu^-} = 30\%$) and a Super-K detector depth $d = 1000$ m have been assumed. For comparison, the total atmospheric neutrino fluxes at the Super-K site from the Bartol [5], Honda [6], and Battistoni [7] groups have also been shown. It is found that the neutrino fluxes from stopped muons are much less than the atmospheric neutrino fluxes. In the right panel of Fig. 4, we plot the ratios of ϕ_{ν_i} to the corresponding atmospheric neutrino flux from the Battistoni group [7] for different flavors. For $13 \text{ MeV} \leq E_\nu \leq 53 \text{ MeV}$, the ν_e , $\bar{\nu}_e$, ν_μ , and $\bar{\nu}_\mu$ fluxes are averagely 10.8%, 6.3%, 3.7%, and 6.2% of the corresponding atmospheric neutrino fluxes,

FIG. 4. The differential neutrino fluxes $d\phi_{\nu_i}/dE_\nu$ (left) and the ratios of $d\phi_{\nu_i}/dE_\nu$ to the corresponding atmospheric neutrino flux from Ref. [7] (right) for different flavors and $d = 1000$ m. In the left panel, the total atmospheric neutrino fluxes at the Super-K site from the Bartol [5], Honda [6], and Battistoni [7] groups have also been shown.

respectively. It is worthwhile to stress that the above results will be increased by a factor of 1.4 if the detector depth $d < 30$ m.

V. CONCLUSIONS

In conclusion, we have investigated the low energy neutrinos from stopped cosmic ray muons in the Earth. The $\mu^\pm$ stop rates per unit volume $S_{\mu^\pm}(x)$ in the rock and sea water have been calculated in terms of the muon intensity $I(p_\mu, \theta)$ at the sea level and the muon range $X(p_\mu)$. Based on the atomic capture and nuclear capture abilities of 10 dominant elements in the upper continental crust, we estimate the μ^- decay and nuclear capture probabilities in the rock and sea water. Then the neutrino energy spectra f_{ν_e} , $f_{\bar{\nu}_e}$, f_{ν_μ} , and $f_{\bar{\nu}_\mu}$ from a stopped muon are given. Finally, we present the low energy neutrino fluxes and give simultaneously a good approximation to calculate them. It is found that most neutrinos come from the surface distance $L < 200$ km region and the near horizontal direction. For the integral fluxes ϕ_{ν_e} , $\phi_{\bar{\nu}_e}$, and ϕ_{ν_μ} , the differences between the rock and water cases

are very small. On the contrary, the $\phi_{\bar{\nu}_e}$ depends on the local rock and water distributions because of different μ^- decay probabilities. Note that all ϕ_{ν_i} depend heavily on the detector depth d . For the Super-K detector depth $d = 1000$ m, the ν_e , $\bar{\nu}_e$, ν_μ , and $\bar{\nu}_\mu$ fluxes in the range of $13 \text{ MeV} \leq E_\nu \leq 53 \text{ MeV}$ are averagely 10.8%, 6.3%, 3.7% and 6.2% of the corresponding atmospheric neutrino fluxes, respectively. The above results will be increased by a factor of 1.4 if the detector depth $d < 30$ m. These low energy neutrinos should be considered in searches of some related topics.

ACKNOWLEDGMENTS

We are grateful to Meng-Yun Guan and Ji-Lei Xu for their useful discussions and helps. This work is supported in part by the National Nature Science Foundation of China (NSFC) under Grants No. 11575201 and No. 11835013, and the Strategic Priority Research Program of the Chinese Academy of Sciences under Grant No. XDA10010100.

-
- [1] Y. Fukuda *et al.* (Super-Kamiokande Collaboration), *Phys. Rev. Lett.* **81**, 1562 (1998).
 - [2] M. Honda, T. Kajita, K. Kasahara, S. Midorikawa, and T. Sanuki, *Phys. Rev. D* **75**, 043006 (2007).
 - [3] S. W. Li and J. F. Beacom, *Phys. Rev. C* **89**, 045801 (2014).
 - [4] D. F. Measday, *Phys. Rep.* **354**, 243 (2001).
 - [5] T. K. Gaisser, T. Stanev, and G. Barr, *Phys. Rev. D* **38**, 85 (1988).
 - [6] M. Honda, T. Kajita, K. Kasahara, and S. Midorikawa, *Phys. Rev. D* **52**, 4985 (1995).
 - [7] G. Battistoni, A. Ferrari, T. Montaruli, and P. R. Sala, *Astropart. Phys.* **23**, 526 (2005).
 - [8] C. Patrignani *et al.* (Particle Data Group), *Chin. Phys. C* **40**, 100001 (2016).
 - [9] S. Ando and K. Sato, *New J. Phys.* **6**, 170 (2004).
 - [10] F. An *et al.* (JUNO Collaboration), *J. Phys. G* **43**, 030401 (2016).

- [11] C. Rott, J. Siegal-Gaskins, and J. F. Beacom, *Phys. Rev. D* **88**, 055005 (2013); N. Bernal, J. Marín-Albo, and S. Palomares-Ruiz, *J. Cosmol. Astropart. Phys.* **08** (2013) 011.
- [12] S. Palomares-Ruiz and S. Pascoli, *Phys. Rev. D* **77**, 025025 (2008).
- [13] A. Gando *et al.* (KamLAND Collaboration), *Astrophys. J.* **745**, 193 (2012).
- [14] K. Ueno *et al.* (Super-Kamiokande Collaboration), *Astropart. Phys.* **36**, 131 (2012).
- [15] D. Reyna, [arXiv:hep-ph/0604145](#).
- [16] V. A. Naumov, [arXiv:hep-ph/0201310](#).
- [17] D. E. Groom, N. V. Mokhov, and S. I. Striganov, *At. Data Nucl. Data Tables* **78**, 183 (2001).
- [18] P. Coloma, P. B. Denton, M. C. Gonzalez-Garcia, M. Maltoni, and T. Schwetz, *J. High Energy Phys.* **04** (2017) 116.
- [19] T. Von Egidy and F. J. Hartmann, *Phys. Rev. A* **26**, 2355 (1982).
- [20] R. L. Rudnick and S. Gao, Composition of the continental crust, in *The crust*, edited by H. D. Holland and K. K. Turekian, Treatise on Geochemistry Vol. 3 (Elsevier-Pergamon, Oxford, 2003), pp. 1–64.
- [21] T. Suzuki, D. F. Measday, and J. P. Roalsvig, *Phys. Rev. C* **35**, 2212 (1987).
- [22] G. Strassner *et al.*, *Phys. Rev. C* **20**, 248 (1979).
- [23] O. L. G. Peres and A. Y. Smirnov, *Phys. Rev. D* **79**, 113002 (2009).