Loops in Anti-de Sitter Space

Igor Bertan^{*} and Ivo Sachs[†]

Arnold Sommerfeld Center for Theoretical Physics, Ludwig Maximilian University of Munich,
Theresienstraße 37, D-80333 München, Germany

(Received 3 May 2018; published 4 September 2018)

We obtain analytic results for the four-point amplitude, at one loop, of an interacting scalar field theory in four-dimensional, Euclidean anti-de Sitter space without exerting any conformal field theory knowledge. For the two-point function, we provide analytic expressions up to two loops. In addition, we argue that the critical exponents of correlation functions near the conformal boundary of anti-de Sitter space provide the necessary data for the renormalization conditions, thus replacing the usual on-shell condition.

DOI: 10.1103/PhysRevLett.121.101601

Introduction.—Over the past sixty years, there has been tremendous progress in the calculation of scattering amplitudes in quantum field theory—in particular, concerning higher loop amplitudes in Yang-Mills theory and (super) gravity.

At the same time, we have very few analytic results for loop amplitudes in curved space-times. More precisely, while their short-distance expansion and, in particular, the structure of counterterms they give rise to is rather well known [1,2], we know little about their dependence on coordinates beyond that (e.g. Ref. [3] and references therein). Even in de Sitter (dS) or anti-de Sitter (AdS) space, which are maximally symmetric, admitting the same number of isometries as Minkowski space, little is known about such amplitudes; see Refs. [4–10] for recent progress. The reason for this is that, while in Minkowski space, the momentum representation leads to a hierarchy of elementary integrals; in dS or AdS, this is not the case, and the coordinate representation generally leads to integral expressions that are more manageable in (A)dS. Still, except for some special cases, we lack the technical tools for performing the integrations completely.

In this Letter we report on some progress considering the simplest interacting renormalizable field theory. Concretely, we compute the two- and four-point functions for $\lambda\phi^4$ theory [11] to the second order in the coupling λ on the Poincaré patch of Euclidean AdS_4 by explicitly evaluating the corresponding one- and two-loop integrals in coordinate representation. Working on AdS, we avoid complications that arise from IR effects on dS, for instance [3]. It turns out that even this simplified setting is beyond reach for external

legs at generic points in AdS, but for insertions on the conformal boundary we are able to get explicit expressions.

Being able to go beyond the short-distance expansion, we encounter an interesting complication concerning the renormalization conditions: For distances that are small compared to the curvature scale, the problem reduces to that in flat space, and the physical masses provide the right boundary conditions for the renormalized propagator, for instance. At scales of the order of the curvature radius, however, there is no meaningful definition of the mass of a scalar field, and one needs to identify a reasonable renormalization condition. In the present case, we will find that the critical exponents of correlation functions at the conformal boundary of AdS provide just that. Indeed, the bulk amplitudes on AdS with external legs inserted at the boundary define a crossing symmetric point correlation function of some hitherto unknown primary operator of a conformal field theory (CFT) on the conformal boundary by construction, and therefore a consistent CFT. Of course, we do not know what is the microscopic realization of this CFT, nor do we need it. What matters is that a primary operator has a well-defined dimension which is given by the critical exponent of its correlation function near the conformal boundary. This is what replaces the physical mass at large distances (see also Ref. [12]).

Concretely, let us consider a scalar field with classical action [13]

$$S = - \int d\mu_x \left(\frac{1}{2} (\partial\phi)^2 + \frac{m^2}{2} \phi^2 + \frac{\lambda}{4!} \phi^4 \right) \quad (1)$$

on the Poincaré patch of hyperbolic space of radius $1/a$ with the metric

$$ds^2 = \frac{1}{a^2 z^2} (dz^2 + dx^{i^2}), \quad (z, x^i) \in (\mathbb{R}_{\geq 0}, \mathbb{R}^3). \quad (2)$$

There are two admissible boundary conditions for the classical scalar field corresponding to the asymptotic

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](#). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

behavior $\phi(z, x^i) \sim z^\Delta \varphi(x^i)$ with $a^2 \Delta(\Delta - 3) = m^2$. Here we will focus on the conformally coupled scalar for which $m^2 = -2a^2$ and therefore $\Delta = 1, 2$. For $\Delta = 1$, there are further complications due to infrared divergencies. We thus focus on $\Delta = 2$ in this case. The scalar propagator is then given by (cf. Ref. [1] and references therein)

$$\Lambda(K) = \frac{a^2 K^2}{4\pi^2(1 - K^2)}, \quad (3)$$

where K is the invariant bilocal function

$$K \equiv K_{x,y} = \frac{2zw}{(x^i - y^i)^2 + z^2 + w^2} \quad (4)$$

with coordinates $x^\mu = (z, x^i)$ and $y^\mu = (w, y^i)$. Henceforth we denote $x^2 \equiv x^{i^2}$. Taking one point to the conformal boundary, $z \sim 0$, then

$$\frac{K}{z} \sim \bar{K} = \frac{2w}{(x - y)^2 + w^2} \quad (5)$$

reduces to the usual bulk-to-boundary propagator [14].

Two-point function.—To order λ^2 , the two-point function contains the following fundamental constituents:

$$\begin{array}{cccc} \text{---} \bullet \text{---} & \text{---} \bullet \text{---} \text{---} \text{---} & \text{---} \bullet \text{---} \text{---} \text{---} & \text{---} \bullet \text{---} \text{---} \text{---} \\ \mathcal{I}_2 & \mathcal{H}_2 & \mathcal{L}_2 & \mathcal{K}_2 \end{array} \quad (6)$$

where the first diagram is just a mass shift which will, however, play a prominent role in the following. It corresponds to the elementary integral (here $K = K_{x_1, x_2}$)

$$\begin{aligned} \mathcal{I}_2 &= \int d\mu_x \Lambda(x_1, x) \Lambda(x_2, x) \\ &= -\frac{K}{8\pi^2(1 - K^2)} \left(\log \frac{1 - K}{1 + K} + K \log \frac{4K^2}{1 - K^2} \right). \end{aligned} \quad (7)$$

The one-loop tadpole diagram

$$\mathcal{H}_2 = \int d\mu_x \Lambda(x, x_1) \Lambda(x, x_2) \Lambda(x, x) \quad (8)$$

requires regularization at short distances, $K \rightarrow 1$. We choose

$$K \rightarrow \frac{K}{1 + \epsilon}, \quad (9)$$

which cuts out a small ϵ -ball around the pole in the propagator and rescales it by $1/(1 + \epsilon)$. With this, the tadpole diagram reduces to a mass counterterm as expected:

$$\mathcal{H}_2 = \frac{a^2}{8\pi^2} \left(\frac{1}{\epsilon} - \frac{9}{2} \right) \mathcal{I}_2 + \mathcal{O}(\epsilon). \quad (10)$$

For the two-loop tadpole diagram

$$\mathcal{L}_2 = \int d\mu_{x,y} \Lambda(x_1, x) \Lambda(x_2, x) \Lambda(x, y)^2 \Lambda(y, y), \quad (11)$$

we adopt the same regularization as above. It is then possible to show that for $z \neq 0$, the y integral is well defined and is independent of x . Consequently, the nested integral (11) factorizes as

$$\mathcal{L}_2 = \frac{a^2}{(4\pi^2)^3} \frac{1}{\epsilon(2 + \epsilon)(1 + \epsilon)^4} \mathcal{M}_2 \times \mathcal{I}_2, \quad (12)$$

where $\mathcal{I}_2$ is again the mass shift. Using translation invariance to set $x = (1, 0)$, we find

$$\mathcal{M}_2 = 8 \int_{-\infty}^{\infty} d^4 y \frac{[y^2 + (w + 1)^2 + \epsilon Q]^{-2}}{[y^2 + (w - 1)^2 + \epsilon Q]^2},$$

with $Q = y^2 + w^2 + 1$. This integral is most easily computed using Schwinger parameters. For small ϵ , we then end up with

$$\mathcal{L}_2 = \frac{a^2 \pi^2}{2(4\pi^2)^3} \left(\frac{14 + 13 \log \frac{\epsilon}{2}}{2} - \frac{1 + \log \frac{\epsilon}{2}}{\epsilon} \right) \mathcal{I}_2 + \mathcal{O}(\epsilon).$$

Finally, we consider the sunset diagram

$$\mathcal{K}_2 = \int d\mu_{x,y} \Lambda(x_1, x) \Lambda(x, y)^3 \Lambda(x_2, y). \quad (13)$$

Let us first consider the subdiagram with only one external leg attached:

$$\mathcal{J}_2 = \int d\mu_y \Lambda(x, y)^3 \Lambda(x_2, y). \quad (14)$$

If we denote the restriction of $\mathcal{J}_2$ to the conformal boundary by J_2 , we have

$$J_2 = \frac{2^7 a^4 z_2^2 z''^6}{(4\pi^2)^4 (1 + \epsilon)^4} \int_{-\infty}^{\infty} d^4 y \frac{w^4 [Q + 2z''w + \epsilon Q]^{-3}}{[Q - 2z''w + \epsilon Q]^3},$$

where $Q = y^2 + z''^2 + w^2$. Here we use translation invariance, as above, to set $x'_2 = (0, 0)$, followed by an inversion with $z'' = z'/(x'^2 + z'^2)$ [15]. This integral is again evaluated using Schwinger, leading to

$$J_2 = \frac{a^4 \pi^2}{4(4\pi^2)^4} K_{x, x_2}^2 \left(\frac{1}{\epsilon} + 3 \log \frac{\epsilon}{2} - \frac{1}{2} \right), \quad (15)$$

where we use the fact that $2z'' = \bar{K}_{x, x_2}$ to recover the covariant form. The full sunset diagram can now be obtained by attaching the remaining leg to J_2 . This yields

$$\mathcal{K}_2 = \frac{a^2 \pi^2}{4(4\pi^2)^3} \left(\frac{1}{\epsilon} + 3 \log \frac{\epsilon}{2} - \frac{1}{2} \right) \mathcal{I}_2, \quad (16)$$

where $\mathcal{I}_2$ is the mass shift (7), and where we use the symmetry of $\mathcal{K}_2$ under permutation of the external legs to construct the unique extension of the above correlator in the bulk.

In conclusion, all diagrams that contribute to the two-point function, up to the second order in the coupling constant, reduce to the mass shift diagram which relates the Lagrangian mass to the conformal dimension of a primary operator, $\mathcal{O}$, when evaluated at the boundary. This suggests replacing the renormalization condition defining the physical mass, which itself is not well defined in AdS, with a renormalization condition on the conformal dimension of $\mathcal{O}$. Our choice sets Δ to 2 at all orders in the perturbation. Below, we will see that this is consistent with the four-point function.

Four-point function.—Up to second order in λ , the one-particle irreducible diagrams contributing to the four-point function are

$$\mathcal{I}_4 \quad \mathcal{K}_4 \quad (17)$$

The tree-level contribution of the quartic vertex to the four-point function, given by

$$\mathcal{I}_4 = \int d\mu_x \Lambda(x_1, x) \Lambda(x_2, x) \Lambda(x_3, x) \Lambda(x_4, x), \quad (18)$$

has already been calculated for external legs inserted on the boundary. Here we just quote the result [16]:

$$I_4 = \frac{4^2 a^4 (\prod_{i=1}^4 z_i)^2}{(4\pi^2)^3 (\eta \zeta \prod_{i<j} r_{ij})^{\frac{4}{3}}} \times \int_0^\infty dz_2 F_1 \left[2, 2; 4; 1 - \left(\frac{\eta + \zeta}{\eta \zeta} \right)^2 - \frac{4 \sinh^2 z}{\eta \zeta} \right], \quad (19)$$

where we introduce $r_{ij} = |x_i - x_j|$ and the conformal cross ratios of the coordinates on the boundary $\eta = r_{14} r_{23} / r_{12} r_{34}$, $\zeta = r_{14} r_{23} / r_{13} r_{24}$.

Let us calculate the one-loop correction given by the double integral

$$\mathcal{K}_4 = \int d\mu_{x,y} \Lambda(x_1, x) \Lambda(x_2, x) \Lambda(x, y)^2 \Lambda(x_3, y) \Lambda(x_4, y),$$

considering again first the simpler integral

$$\mathcal{J}_4 = \int d\mu_y \Lambda(x, y)^2 \Lambda(x_3, y) \Lambda(x_4, y). \quad (20)$$

By sending x_3 and x_4 to the boundary, it takes the form

$$J_4 = \frac{a^8 (z_3 z_4)^2}{(4\pi^2)^4 (1 + \epsilon)^8} \int d\mu_y \frac{\bar{K}_{x_3, y}^2 \bar{K}_{x_4, y}^2 (1 + K_{x, y} + \epsilon)^{-2}}{K_{x, y}^{-4} (1 - K_{x, y} + \epsilon)^2}.$$

We may safely set $\epsilon = 0$ in the prefactor, since the integral diverges logarithmically. As before, we translate the points x, x_3, x_4 by $(0, -x_4^i)$, which gives $x_4' = (0, 0)$, and we perform an inversion with the inverted points denoted by double primes so that

$$J_4 = \frac{a^4 z'^4 (z_3 z_4)^2}{2^{-7} (4\pi^2)^4 r_{34}^4} \int_{-\infty}^{\infty} \frac{d^4 y}{w^{-4}} \frac{[(Q_- + \epsilon Q)(Q_+ + \epsilon Q)]^{-2}}{[(x_3'' - x'' - y)^2 + w^2]^2},$$

where $Q_{\pm} = y^2 + (z'' \pm w)^2$ and $Q = y^2 + z''^2 + w^2$. Integrating yields

$$J_4 = \frac{a^4 \pi^2}{(4\pi^2)^4} K_{x, x_3}^2 K_{x, x_4}^2 \left(\log \frac{\alpha^2 + 1}{2\epsilon} - 2 \right), \quad (21)$$

where

$$\frac{4}{\alpha^2 + 1} \equiv r_{34}^2 \bar{K}_{x, x_3} \bar{K}_{x, x_4} = \frac{4z''^2}{(x_3'' - x'')^2 + z''^2}. \quad (22)$$

Finally, we attach the remaining two external legs to (20). Sending all x_i to the boundary, we have

$$K_4 = \frac{4^4 a^4 (\prod_{i=1}^4 z_i)^2 \pi^2}{2(4\pi^2)^6} \int_{-\infty}^{\infty} d^4 x \frac{z^4 (\log \frac{\alpha^2 + 1}{2\epsilon} - 2)}{\prod_{i=1}^4 [(x_i - x)^2 + z^2]^2}.$$

We then repeat the by-now-familiar procedure by translating x_k , $k = 1, \dots, 4$ by $(0, -x_4^i)$ (denoted by primes), inverting all points (denoted by double primes), and then making the substitution $(z'', x''^i) = (z, x^i + x_3''^i)$. The integration is then performed in the usual fashion, leading to

$$K_4 = \frac{1}{16\pi^2} \left[-I_4 \left(\frac{11}{3} + \log \frac{\epsilon}{8} \right) + L_4 \right], \quad (23)$$

where

$$L_4 = \frac{3 \times 4^2 a^4 (\prod_{i=1}^4 z_i)^2}{2(4\pi^2)^3 (\eta \zeta \prod_{i<j} r_{ij})^{\frac{4}{3}}} \times \int_0^\infty ds \int_0^1 dr \frac{[sr(1-r)] \log(1+s)}{(1+s)^2 \left[\frac{sr(1-r)}{\eta^2} + \frac{r}{\zeta^2} + 1 - r \right]^2}.$$

This is the main result of this Letter. To continue, L_4 can be evaluated numerically or, alternatively, order by order in an expansion in $1 - \zeta^{-2}$ and η^{-1} . Then the expansion coefficients of (23) contain important physical information, which can be extracted by comparing them with the operator product expansion (OPE) in CFT.

Comparison to conformal field theory.—In flat space, loop corrections to the tree-level amplitudes contain

information about the coupling dependence of the masses of resonances, for instance. In AdS, where there is no scattering, the role of physical masses is taken by the dimensions of operators of some CFT dual [14,17,18]. For a scalar field ϕ in AdS, this CFT is characterized by the existence of a scalar operator $\mathcal{O}$ dual to ϕ . Our renormalization scheme fixes its two-point function to

$$\langle \mathcal{O}(x_1) \mathcal{O}(x_2) \rangle = \text{circle with a dot in the middle} = \frac{N_\phi}{r_{12}^4}, \quad (24)$$

where $N_\phi = a^2(2z_1 z_2)^2/4\pi^2$. To continue, we take $z_i = z \sim 0$ for all external legs. Then, expanding the holographic four-point function in the variables η^{-1} and $Y = 1 - \zeta^{-2}$ yields

$$\begin{aligned} \langle \mathcal{O}(x_1) \mathcal{O}(x_2) \mathcal{O}(x_3) \mathcal{O}(x_4) \rangle &= 3 \times \text{circle with a dot in the middle} + \lambda \text{circle with four dots} + \frac{\lambda^2}{2} \left(3 \times \text{circle with four dots and a cross} \right) + \mathcal{O}(\lambda^3) \\ &= \frac{N_\phi^2}{(r_{12} r_{34})^4} \left[1 + \frac{1}{\eta^4} \sum_{l,m=0}^{\infty} F_{lm}(\log \eta, \lambda_R) \frac{Y^m}{\eta^{2l}} \right], \end{aligned} \quad (25)$$

where each F_{lm} can be derived from the results of the previous sections. Note that the factor “3” in the diagrams merely indicates that there are three diagrams of this type that contribute to the correlator. Here we furthermore introduce the renormalized coupling constant by a non-minimal subtraction of the form

$$\lambda = \lambda_R + \frac{\lambda_R^2}{32\pi^2} \left(5 + 3 \log \frac{\epsilon}{8} \right) + \mathcal{O}(\lambda_R^3). \quad (26)$$

At this point, we should emphasize that a nonvanishing beta function of the bulk theory does not spoil the conformal symmetry on the boundary, since scale transformations on the boundary correspond to translations in AdS.

Let us now consider a general four-point function of identical scalar operators $\mathcal{O}$ of conformal dimension ν in a three-dimensional CFT. Its decomposition in conformal blocks (CBs) G_s^ν is [19]

$$\langle \mathcal{O}(x_1) \mathcal{O}(x_2) \mathcal{O}(x_3) \mathcal{O}(x_4) \rangle = \sum_{s, \nu_s} \frac{C_s^{\nu_s} \times G_s^{\nu_s}(\eta, Y)}{(r_{12} r_{34})^4 \eta^{\nu_s - s}}, \quad (27)$$

where the only unknowns are the spectrum of the CFT—i.e., the spin s and conformal dimension ν_s of all primary operators, as well as the associated OPE coefficients $C_s^{\nu_s}$. These can, in turn, be determined by comparison with (25). Our recipe is the following: we first compute F_{lm} and identify the set of possibly contributing CBs at that order in η and Y . Afterwards, we expand (27) in the conformal dimensions

$\nu_s = \bar{\nu}_s + \lambda_R \gamma_s^{\bar{\nu}_s(1)} + \lambda_R^2 \gamma_s^{\bar{\nu}_s(2)} + \mathcal{O}(\lambda_R^3)$ and OPE coefficients $C_s^{\nu_s} = C_s^{\bar{\nu}_s(0)} + \lambda_R C_s^{\bar{\nu}_s(1)} + \lambda_R^2 C_s^{\bar{\nu}_s(2)} + \mathcal{O}(\lambda_R^3)$. This allows for a direct comparison with F_{lm} at different orders in λ_R .

For brevity, in this Letter we will focus on the terms with $(l, m) = (0, 0), (1, 0), (0, 1), (0, 2)$. The CBs which contribute at these orders are

$$\begin{aligned} G_0^{\nu_0} &= 1 + \frac{\nu_0}{4} Y + \frac{\nu_0^3(\nu_0 + 1)^{-1}}{8(2\nu_0 - 1)\eta^2} + \frac{\nu_0(\nu_0 + 2)^2}{32(\nu_0 + 1)} Y^2, \\ G_1^{\nu_1} &= -\frac{1}{2} Y - \frac{\nu_1 + 1}{8} Y^2, \\ G_2^{\nu_2} &= -\frac{1}{2\eta^2} + \frac{3}{8} Y^2. \end{aligned}$$

Note that the identity operator $\mathbb{1}$ appears in (25) in the form of the s -channel disconnected diagram, as its CB reads $G_0^0 = 1$. Thus, $C_0^0 = N_\phi^2$.

For $(l, m) = (0, 0)$, one finds

$$\begin{aligned} F_{00} &= 2 + \frac{\lambda_R}{48\pi^2} (-1 + 6 \log \eta) \\ &\quad + \frac{\lambda_R^2}{3 \times 2^8 \pi^4} (5 - 11 \log \eta + 3(\log \eta)^2). \end{aligned}$$

The only contribution comes from the primary operator $:\mathcal{O}^2:$ having conformal dimension $\bar{\nu}_0 = 4$. Then, comparing the corresponding CB expansion with F_{00} at lowest order in the coupling constant yields $C_0^{4(0)} = 2N_\phi^2$. At first order in λ_R , we get

$$\gamma_0^{4(1)} = -\frac{1}{16\pi^2}, \quad C_0^{4(1)} = -\frac{N_\phi^2}{48\pi^2};$$

whereas at second order in λ_R , we find

$$\begin{aligned} \gamma_0^{4(1)} &= \pm \frac{1}{16\pi^2}, \quad \gamma_0^{4(2)} = \frac{5}{3 \times 2^8 \pi^4}, \\ C_0^{4(2)} &= \frac{5N_\phi^2}{3 \times 2^8 \pi^4}. \end{aligned}$$

Note that $\gamma_0^{4(1)}$ agrees at different orders in λ_R . This provides an important consistency test for the AdS/CFT duality beyond tree level in the bulk. In previous work, this property was taken as part of the definition of loop diagrams in the bulk (e.g., Refs. [5–7]). It is reassuring to see that it is indeed compatible with an actual bulk calculation. Put differently, this supports the argument that CFT does indeed describe the structure underlying amplitudes of QFT in AdS rather than acting merely as a definition of some bulk theory specified by its correlation functions.

For $(l, m) = (0, 1)$, F_{01} reads

$$F_{01} = 2 + \frac{\lambda_R}{96\pi^2}(-5 + 12 \log \eta) + \frac{\lambda_R^2}{12 \times 2^8 \pi^4}(31 - 50 \log \eta + 12(\log \eta)^2).$$

In addition to $:\mathcal{O}^2:$, there might be a contribution of a vector operator of dimension $\bar{\nu}_1 = 5$. However, by comparing F_{01} with the expansions of the CBs, one can infer that the vector operator does not appear in the OPE. This agrees with our expectation based on general CFT arguments.

The term satisfying $(l, m) = (0, 2)$ reads

$$F_{02} = 3 + \frac{9\lambda_R}{80\pi^2}\left(-\frac{11}{20} + \log \eta\right) + \frac{\lambda_R^2}{500 \times 2^8 \pi^4}(1408 - 1965 \log \eta + 450(\log \eta)^2).$$

Bearing in mind that the vector operator of dimension 5 does not appear, the only new operator which contributes is the spin-2 primary of the schematic form $:\mathcal{O}\partial^i\partial^j\mathcal{O}:$ with $\bar{\nu}_2 = 6$. It follows that

$$C_2^{6(0)} = \frac{16N_\phi^2}{5}, \quad \gamma_2^{6(1)} = C_2^{6(1)} = 0, \\ \gamma_2^{6(2)} = -\frac{1}{20 \times 2^8 \pi^4}, \quad C_2^{6(2)} = -\frac{11N_\phi^2}{375 \times 2^8 \pi^4}.$$

Also, here $\gamma_2^{6(1)}$ agrees at both orders in λ_R . An interesting observation is that the spin-2 primary, in spite of not being conserved, does not acquire an anomalous dimension at first order in the coupling λ_R . However, it does get modified at second order.

The last term we consider here corresponds to $(l, m) = (1, 0)$ with

$$F_{10} = \frac{\lambda_R}{120\pi^2}\left(\frac{17}{5} + 12 \log \eta\right) + \frac{\lambda_R^2}{150 \times 2^8 \pi^4}\left(-\frac{491}{5} - 584 \log \eta + 120(\log \eta)^2\right),$$

which requires a new scalar operator $:\mathcal{O}\nabla^2\mathcal{O}:$ of conformal dimension $\bar{\nu}_0 = 6$ and

$$C_0^{6(0)} = \frac{8N_\phi^2}{7}, \quad \gamma_0^{6(1)} = -\frac{1}{16\pi^2}, \quad C_0^{6(1)} = \frac{239N_\phi^2}{14 \times 420\pi^2}, \\ \gamma_0^{6(2)} = \frac{23}{15 \times 2^7 \pi^4}, \quad C_0^{6(2)} = -\frac{1637N_\phi^2}{5 \times 2^6 \times 1029}.$$

Again, there is an agreement of $\gamma_0^{6(1)}$ at different orders in λ_R . Eventually, this gives a complete characterization of all operators of spin $s \leq 2$ entering the OPE.

Conclusions.—In this Letter, we computed quantum corrections to the two- and four-point correlation functions up to second order in the coupling constant for the simplest scalar field theory in AdS_4 . The obtained results for the two- and four-point functions are mutually consistent. Furthermore, the holographic four-point function can systematically be expanded in the conformal invariants to reveal the OPE structure of the dual CFT, along with the corrections to both the OPE coefficients and conformal dimensions of primary operators. This was carried out here at low orders, disclosing a mathematically consistent dual CFT. In particular, the absence of the stress tensor and of any conserved current becomes explicit.

To summarize, the conformally coupled bulk $\lambda\phi^4$ theory describes a one-parameter family of dual conformal field theories whose OPE coefficients and dimensions are parametrized by the renormalized coupling λ_R . Generalization of the result presented here to boundary conditions with $\Delta = 1$ is possible, although with some extra complications concerning infrared divergencies for the tadpole and sunset diagrams. For massless spin-1 and spin- $\frac{1}{2}$ particles in AdS, the propagator is again given by (3) modulo parallel transport of the polarization vectors. This means that spin-1 and spin- $\frac{1}{2}$ fields lead to similar integrals to those computed here, and hence QED and scalar QED in AdS can be quantized in the same way.

The authors would like to thank Evgeny Skvortsov for very substantial input and advice. This work was supported by the DFG Transregional Collaborative Research Centre TRR 33 and the DFG cluster of excellence “Origin and Structure of the Universe”.

*igor.bertan@physik.lmu.de

†ivo.sachs@physik.lmu.de

- [1] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, Cambridge, England, 1984).
- [2] J. Comellas, P. E. Haagensen, and J. I. Latorre, Differential renormalization for curved space and finite temperature, *Int. J. Mod. Phys. A* **10**, 2819 (1995).
- [3] E. T. Akhmedov, Lecture notes on interacting quantum fields in de sitter space, *Int. J. Mod. Phys. D* **23**, 1430001 (2014).
- [4] J. Penedones, Writing CFT correlation functions as AdS scattering amplitudes, *J. High Energy Phys.* **03** (2011) 025.
- [5] A. L. Fitzpatrick and J. Kaplan, Analyticity and the holographic S-matrix, *J. High Energy Phys.* **10** (2012) 127.
- [6] O. Aharony, L. F. Alday, A. Bissi, and E. Perlmutter, Loops in AdS from conformal field theory, *J. High Energy Phys.* **07** (2017) 036.
- [7] L. Rastelli and X. Zhou, Mellin Amplitudes for Supergravity on $\text{AdS}_5 \times S^5$, *Phys. Rev. Lett.* **118**, 091602 (2017).

- [8] C. Cardona, Mellin-(Schwinger) representation of One-loop Witten diagrams in AdS, [arXiv:1708.06339](#).
- [9] S. Giombi, C. Sleight, and M. Taronna, Spinning AdS loop diagrams: two point functions, *J. High Energy Phys.* **06** (2018) 030.
- [10] E. Y. Yuan, Simplicity in AdS Perturbative Dynamics, [arXiv:1801.07283](#).
- [11] The cubic coupling is known to be extremal, and the simplest λ diagram is divergent.
- [12] N. Beisert *et al.*, Review of AdS/CFT integrability: An overview, *Lett. Math. Phys.* **99**, 3 (2012).
- [13] $d\mu_{x_1, x_2, \dots} = d^4x_1 \sqrt{|g(x_1)|} \times d^4x_2 \sqrt{|g(x_2)|} \dots$
- [14] E. Witten, Anti de Sitter space and holography, *Adv. Theor. Math. Phys.* **2**, 253 (1998).
- [15] D. Z. Freedman, S. D. Mathur, A. Matusis, and L. Rastelli, Correlation functions in the CFT_d/AdS_{d+1} correspondence, *Nucl. Phys.* **B546**, 96 (1999).
- [16] W. Muck and K. S. Viswanathan, Conformal field theory correlators from classical scalar field theory on anti-de Sitter space, *Phys. Rev. D* **58**, 041901 (1998).
- [17] J. M. Maldacena, The large N limit of superconformal field theories and supergravity, *Adv. Theor. Math. Phys.* **2**, 231 (1998).
- [18] S. S. Gubser, I. R. Klebanov, and A. M. Polyakov, Gauge theory correlators from non-critical string theory, *Phys. Lett. B* **428**, 105 (1998).
- [19] F. A. Dolan and H. Osborn, Conformal four point functions and the operator product expansion, *Nucl. Phys.* **B599**, 459 (2001).