**Erratum: Extremely Low Electron-ion Temperature
Relaxation Rates in Warm Dense Hydrogen:
Interplay between Quantum Electrons and Coupled Ions
[Phys. Rev. Lett. 122, 015001 (2019)]**

Qian Ma, Jiayu Dai , Dongdong Kang, M. S. Murillo, Yong Hou, Zengxiu Zhao, and Jianmin Yuan

(Received 12 August 2019; published 28 August 2019)

DOI: [10.1103/PhysRevLett.123.099901](https://doi.org/10.1103/PhysRevLett.123.099901)

b_c in the coupled Gericke, Murillo, and Schlages (GMS) model in this Letter has a typographical error. The correct b_c is $Ze^2/\frac{1}{2}(k_B T_e + k_B T_i)$ and not $\frac{1}{2}Ze^2/(k_B T_e + k_B T_i)$. To introduce the coupling between electrons and ions, we define $b_c = Ze^2/\frac{1}{2}(k_B T_e + k_B T_i)$ instead of $b_c = b_0 = Ze^2/k_B T_e$, typically used in the GMS model $L_{\text{GMS}} = \frac{1}{2} \ln \{1 + [(\lambda_D^2 + R_i^2)/(\Lambda^2/8\pi + b_c^2)]\}$.