⁹The data shown in Fig. 2(b) do not extend to an energy high enough to show the decrease in efficiency expected from the decrease in absorption in the counter gas.

¹⁰For ease of construction and maintenance the polarimeter is built in modular form. Each module consists of four counters arranged in a cross and four

lithium blocks that fill in the corners of the cross.

¹¹The camera data show that the axis of the polarimeter stayed within a circle of 30' radius centered at R.A. 16^h8^m, Dec -16°5' during the Sco X-1 observations. This is 2°15' from Sco X-1. The background observations were centered at R.A. 15^h15.25^m, Dec -0°55'.

ERRATA

MUTUAL FRICTION IN He II NEAR THE SUPERFLUID TRANSITION. G. Ahlers [Phys. Rev. Letters 22, 54 (1969)].

The third line below Eq. (1) should read, "(q is in $W \text{ cm}^{-2}$),..."

MEASUREMENT OF THE K_1^0 CHARGED-TO-NEUTRAL DECAY BRANCHING RATIO. B. Gobbi, D. Green, W. Hakel, R. Moffett, and J. Rosen [Phys. Rev. Letters 22, 682 (1969)], and STUDY OF $K_1^0 \rightarrow \pi^0 \pi^0$ AND $K_2^0 \rightarrow \pi^0 \pi^0$ INTERFERENCE. B. Gobbi, D. Green, W. Hakel, R. Moffett, J. Rosen, B. Goz, and D. Tycko [Phys. Rev. Letters 22, 685 (1969)].

The figures for these two Letters were transposed: Those appearing with the first Letter pertained to the second, and vice versa. The captions are with the proper Letters and thus not with the figures they describe.