ERRATA

CRITERIA FOR THE ELIMINATION OF DISCRETE AMBIGUITIES IN NUCLEAR OPTICAL POTENTIALS. D. A. Goldberg and S. M. Smith [Phys. Rev. Lett. 29, 500 (1972)].

The formulas appearing in Ref. 14 should be corrected to read as follows:

$$\delta_l = k \int_{r_2}^R \{1 - [\hbar^2(l + \frac{1}{2})^2 / 2\mu r^2 - U(r) - V_c(r)] / E\}^{1/2} dr - k \int_{r_1}^R [1 - (l + \frac{1}{2})^2 / k^2 r^2 - 2\eta / kr]^{1/2} dr,$$

where r_2 and r_1 are defined by the equations

$$1 - [\hbar^2(l + \frac{1}{2})^2 / 2\mu r_2^2 + V(r_2) - V_c(r_2)] / E = 0, \quad 1 - (l + \frac{1}{2})^2 / k^2 r_1^2 - 2\eta / kr_1 = 0.$$

The calculations referred to in the text were performed using the correct formulas.

ANTICENTRIFUGAL STRETCHING IN ^{20}Ne .
H. C. Lee and R. Y. Cusson [Phys. Rev. Lett. 29, 1525 (1972)].

In Table I, the first number in column 3 should be 3.862 instead of 3.682, and the heading for columns 5 and 6 should be $\Delta\rho = \rho_0 - \rho_J$ instead of $\Delta\rho = \rho_J - \rho_0$.

PRECISE TEST OF SU(2) CHIRAL BREAKING.
A COLEMAN-GLASHOW FORMULA FOR PSEUDOSCALAR MESONS. G. Cicogna, F. Strocchi, and R. Vergara Caffarelli [Phys. Rev. Lett. 29, 1702 (1972)].

We would like to clarify Ref. 7 as follows: "It must be specified that in the first paper of Ref. 4, electromagnetic mass differences were assumed to arise almost entirely from conventional electromagnetic effects, and therefore the value given there for ϵ_3 was not considered as related directly to such mass differences."