Hyperfine structure of the 2^3P state in ^9Be and the nuclear quadrupole moment

Mariusz Puchalski and Jacek Komasa

Faculty of Chemistry, Adam Mickiewicz University, Uniwersytetu Poznańskiego 8, 61-614 Poznań, Poland

Krzysztof Pachucki

Faculty of Physics, University of Warsaw, Pasteura 5, 02-093 Warsaw, Poland

(Received 1 February 2021; accepted 22 February 2021; published 31 March 2021)

We have performed accurate calculations of the hyperfine structure of the 2^3P state in the ^9Be atom with the help of highly optimized, explicitly correlated functions, accounting for the leading finite nuclear mass, radiative, nuclear structure, and relativistic effects. By comparison with measurements, we have determined the ^9Be nuclear quadrupole moment to be $Q_N = 0.05350(14)$ barns, which is not only the most accurate result but also disagrees with previous determinations.

DOI: [10.1103/PhysRevResearch.3.013293](https://doi.org/10.1103/PhysRevResearch.3.013293)

The electric quadrupole moment Q_N of nuclei is a measure of the deformation of the nuclear charge distribution with respect to the spherical symmetry. This deformation comes from a strong dependence of nuclear forces on the orientation of the nucleon spin and is present in atomic nuclei with a spin equal to or greater than one, as it is for ^9Be with $I = 3/2$. In principle, Q_N could be determined from the nuclear structure theory, but such calculations with controlled accuracy have not yet been feasible. Only very recently and only for such a simple nucleus as the deuteron have theoretical predictions based on the chiral effective field theory reached the uncertainty of 1% [1], and this result agreed with the more precise value from the rotational spectroscopy of the deuterium hydride (HD) molecule [2].

Originally, Q_N for many nuclei was determined by the electron scattering of the nuclei, for example, for ^7Li [3]. However, for ^9Be the available experimental data are not only limited in accuracy but also differ noticeably from each other, indicating that the scattering results are not always conclusive [4,5]. Nowadays, reliable Q_N values can be derived from a combination of theoretical and atomic or molecular spectroscopic data. This has been realized for LiH, LiF, and LiCl molecules [6], leading to a $Q_N(^7\text{Li})$ consistent with the nuclear scattering value [3]. Although appropriate theoretical calculations were performed for BeH^+ [7,8] as well as for the $^7,^9\text{Be}^-$ anion [9] years ago, there have been no corresponding measurements reported so far. Therefore, the method of choice for finding $Q_N(^9\text{Be})$ is atomic spectroscopy.

In the accurate determination of Q_N from atomic spectroscopy, it is important to understand the electron-nucleus

interaction at the fundamental level. Recent advances in measurements of electronic [10] and muonic atoms [11], together with progress in the theoretical description of atomic spectra [12], indicate the importance of the nuclear structure effects. Therefore, accurate knowledge of the nuclear quadrupole moment will give access to details of the electron-nucleus interactions which, so far, have not been visible, for example, the nuclear quadrupole polarizability [13].

The ground state 2^1S_0 of Be is fully symmetric; thus, the nuclear quadrupole moment does not lead here to any observable effect. In contrast, in the lowest excited 2^3P level, which is metastable, the hyperfine interactions cause level splitting, and this splitting can be accurately measured [14]. So, it is atomic structure theory that must interpret the hyperfine structure in terms of the magnetic dipole and the electric quadrupole moments of the nucleus. Still, the same hyperfine interactions that lead to the hyperfine splitting also cause the hyperfine mixing of different 2^3P_J levels. This mixing had been accounted for in Ref. [14] but in a very simplified way. Although the accuracy of these approximations has been questioned [7], consecutive atomic structure calculations, using modern and sophisticated approaches, have relied on these original inaccurate calculations of the hyperfine mixing. An exception is the work by Beloy *et al.* [15], in which this mixing was calculated in the second order of perturbation theory using the relativistic configuration interaction combined with many body perturbation theory (CI + MBPT) approach, but its numerical precision turned out to be insufficient to obtain Q_N with a competitive accuracy. This hyperfine mixing is not specific just to Be atoms; it is certainly present in all the other atomic systems. Therefore, the determinations of quadrupole moments for many other nuclei might have lower accuracy than anticipated, which should be verified, as we have done here for the ^9Be nucleus.

In this work, we employ well-optimized explicitly correlated Gaussian wave functions to have good control over the numerical accuracy. With these wave functions, we perform calculations of the ^9Be hyperfine structure, with a com-

plete treatment of the hyperfine mixing of different 2^3P_J levels. Moreover, we fully account for the leading radiative (QED), finite nuclear mass, and nuclear structure effects, while dealing with relativistic and higher order corrections approximately. Our result for the electric quadrupole moment $Q_N(^9\text{Be})$ show that all of its previous determinations were not as accurate as claimed, due to a very approximate treatment of the hyperfine mixing and neglect of the radiative (QED) corrections.

I. FINE AND HYPERFINE STRUCTURE HAMILTONIAN

The most accurate approach for light atomic systems solves at first the nonrelativistic Hamiltonian

$$H = \frac{\vec{p}_N^2}{2m_N} + \sum_a \frac{\vec{p}_a^2}{2} - \sum_a \frac{Z}{r_a} + \sum_{a<b} \frac{1}{r_{ab}}, \quad (1)$$

and next includes relativistic effects as a perturbation. Relevant relativistic effects are split into the fine structure Hamiltonian

$$H_{\text{fs}} = Q_1 + Q_2, \quad (2)$$

$$Q_1 = \sum_a \frac{Z\alpha}{4} \vec{\sigma}_a \cdot \left[\frac{(g-1)\vec{r}_a}{m^2 r_a^3} \times \vec{p}_a - \frac{g}{m m_N} \frac{\vec{r}_a}{r_a^3} \times \vec{p}_N \right] + \sum_{a \neq b} \frac{\alpha}{4 m^2} \vec{\sigma}_a \cdot \left[g \frac{\vec{r}_{ab}}{r_{ab}^3} \times \vec{p}_b - (g-1) \frac{\vec{r}_{ab}}{r_{ab}^3} \times \vec{p}_a \right], \quad (3)$$

$$Q_2 = - \sum_{a<b} \frac{3\alpha g^2}{16 m^2} \sigma_a^i \sigma_b^j \left(\frac{r_{ab}^i r_{ab}^j}{r_{ab}^5} - \frac{\delta^{ij}}{3 r_{ab}^3} \right), \quad (4)$$

and the hyperfine structure Hamiltonian

$$H_{\text{hfs}} = \vec{I} \cdot \vec{Q} + \frac{3 I^i I^j}{I(2I-1)} \frac{Q^{ij}}{6}, \quad (5)$$

$$\vec{Q} = \vec{Q}_1 + \vec{Q}_2 + \vec{Q}_3, \quad (6)$$

$$\vec{Q}_1 = \sum_a \frac{1}{3} \frac{Z\alpha g_N}{m m_N} \vec{\sigma}_a \pi \delta^3(r_a), \quad (7)$$

$$\vec{Q}_2 = \sum_a \frac{Z\alpha g_N}{2 m m_N} \frac{\vec{r}_a}{r_a^3} \times \vec{p}_a - \frac{Z\alpha (g_N-1)}{2 m_N^2} \frac{\vec{r}_a}{r_a^3} \times \vec{p}_N, \quad (8)$$

$$Q_3^j = \sum_a \frac{Z\alpha g_N}{8 m m_N} \sigma_a^i \left(\frac{3 r_a^i r_a^j}{r_a^5} - \frac{\delta^{ij}}{r_a^3} \right), \quad (9)$$

$$Q^{ij} = - \sum_a \alpha Q_N \left(\frac{3 r_a^i r_a^j}{r_a^5} - \frac{\delta^{ij}}{r_a^3} \right), \quad (10)$$

where $\vec{p}_N = - \sum_a \vec{p}_a$ is the nuclear momentum, $\vec{I}$ is the nuclear spin, g is the free-electron g factor, Q_N is the electric quadrupole moment of the nucleus, and g_N is the nuclear g factor,

$$g_N = \frac{m_N}{Z m_p} \frac{\mu}{\mu_N} \frac{1}{I}. \quad (11)$$

Leading QED effects are included in the free electron g factor, while higher order corrections are accounted for as described later on.

Because the hyperfine splittings are only about 100 times smaller than the fine splitting, one cannot assume that hyperfine levels have a definite angular momentum $\vec{J} = \vec{L} + \vec{S}$, but only definite $\vec{L}$ and $\vec{S}$. It is convenient therefore to extend the original formulation of the hyperfine splitting by Hibbert in Ref. [16] and represent the fine and the hyperfine structure of the 2^3P state in terms of an effective Hamiltonian, instead of expectation values only,

$$H_{\text{eff}} = c_1 \vec{L} \cdot \vec{S} + c_2 (L^i L^j)^{(2)} (S^i S^j)^{(2)} + a_1 \vec{I} \cdot \vec{S} + a_2 \vec{I} \cdot \vec{L} + a_3 (L^i L^j)^{(2)} S^i I^j + \frac{b}{6} \frac{3(I^i I^j)^{(2)}}{I(2I-1)} \frac{3(L^i L^j)^{(2)}}{L(2L-1)}, \quad (12)$$

where the coefficients $a_1, a_2, a_3, b, c_1, c_2$ are independent of J but are specific to the particular state. These coefficients can be obtained, for example, from the matrix elements of H_{fs} and H_{hfs} in the decoupled $|M_L, M_S\rangle$ or $|J, M_J\rangle$ basis. Once these coefficients are known, the above Hamiltonian can be diagonalized, yielding the hyperfine levels. Alternatively, the hyperfine structure can be represented in terms of J -dependent A_J and B_J coefficients

$$H_{\text{hfs, eff}} = A_J \vec{I} \cdot \vec{J} + \frac{B_J}{6} \frac{3(I^i I^j)^{(2)}}{I(2I-1)} \frac{3(J^i J^j)^{(2)}}{J(2J-1)}, \quad (13)$$

which are conventionally used to represent the measured values, while the fine structure is given in terms of differences in the centroid energies.

If the atomic levels had a definite value J , the fine structure would be given by the expectation value of H_{eff} , namely

$$E_{\text{fs}}(J) = \begin{cases} c_1 + c_2/6 & \text{for } J = 2 \\ -c_1 - 5 c_2/6 & \text{for } J = 1, \\ -2 c_1 + 5 c_2/3 & \text{for } J = 0 \end{cases} \quad (14)$$

and the hyperfine structure by

$$A_J = \begin{cases} a_1/2 + a_2/2 + a_3/6 & \text{for } J = 2 \\ a_1/2 + a_2/2 - 5 a_3/6 & \text{for } J = 1 \end{cases}, \quad B_J = \begin{cases} b & \text{for } J = 2 \\ -b/2 & \text{for } J = 1 \end{cases}. \quad (15)$$

Because, in general, one cannot assume that the levels have a definite value of J , the effective hyperfine Hamiltonian in Eq. (12) has to be diagonalized numerically. Nonetheless, the hyperfine structure can still be represented in terms of A_J and B_J coefficients, and we use them for comparison of theoretical predictions with experimental results and for the determination of the nuclear quadrupole moment Q_N .

II. RELATIVISTIC, RADIATIVE, AND FINITE NUCLEAR SIZE CORRECTIONS

The hyperfine Hamiltonian in Eq. (5) represents the leading hyperfine interactions, but there are also many small corrections which are often overlooked in literature. These corrections contain terms with higher powers of the fine structure constant α . Because most of them are proportional to the Fermi contact interaction, i.e., to $\vec{I} \cdot \vec{Q}_1$ in Eq. (5), we account for them in terms of the following factor:

$$\tilde{a}_1 = a_1(1 + \epsilon). \quad (16)$$

Below, we briefly describe contributions included in the ϵ term.

The $\mathcal{O}(\alpha)$ correction is analogous to that in hydrogenic systems [17] and consists of two parts. The first part is due to the nuclear recoil

$$H_{\text{rec}}^{(5)} = \left[-\frac{3Z\alpha}{\pi} \frac{m}{m_N} \ln\left(\frac{m_N}{m}\right) \right] \vec{I} \cdot \sum_a \frac{2}{3} \frac{Z\alpha g_N}{m m_N} \vec{\sigma}_a \pi \delta^3(r_a) \quad (17)$$

and numerically is very small, almost negligible. The recoil contribution to ϵ amounts to $\epsilon_{\text{rec}} = -0.000011$. The second part of the $\mathcal{O}(\alpha)$ correction is due to the finite nuclear size and the nuclear polarizability, and is given by [17,18]

$$H_{\text{fs}}^{(5)} = [-2Z\alpha m r_Z] \vec{I} \cdot \sum_a \frac{2}{3} \frac{Z\alpha g_N}{m m_N} \vec{\sigma}_a \pi \delta^3(r_a), \quad (18)$$

where r_Z is a kind of effective nuclear radius called the Zemach radius. Disregarding the inelastic effects, this radius can be written down in terms of the electric charge ρ_E and magnetic-moment ρ_M densities as

$$r_Z = \int d^3r d^3r' \rho_E(r) \rho_M(r') |\vec{r} - \vec{r}'|. \quad (19)$$

Nevertheless, the inelastic, i.e., polarizability, corrections can be significant, but because they are very difficult to calculate, they are usually neglected. In this work, we account for possible inelastic effects by employing r_Z from a comparison of very accurate calculations of hfs in $^9\text{Be}^+$ with the experimental value, namely $r_Z = 4.07(5)$ fm [18]. Because this correction is also proportional to the contact Fermi interaction, we represent it in terms of $\epsilon_{\text{fs}} = -0.000615$.

Next, there are radiative and relativistic corrections of the relative order $\mathcal{O}(\alpha^2)$. The radiative correction, beyond that included by the free electron g factor, is [17]

$$H_{\text{rad}}^{(6)} = Z\alpha^2 \left(\ln 2 - \frac{5}{2} \right) \vec{I} \cdot \sum_a \frac{2}{3} \frac{Z\alpha g_N}{m m_N} \vec{\sigma}_a \pi \delta^3(r_a) \quad (20)$$

and the corresponding ϵ factor is $\epsilon_{\text{rad}} = -0.000384$. The $\mathcal{O}(\alpha^2)$ relativistic and higher order corrections are much more complicated. They have been calculated for the ground state of $^9\text{Be}^+$ [18]. Here we take this result, assume that it is proportional to the Fermi contact interaction, and obtain $\epsilon_{\text{rel}} = 0.001664$. This is the only approximation we assume in this work, and as a consequence we neglect the mixing of the 2^3P_1 state with the nearby lying 2^1P_1 . Exactly for this reason we will use only the hyperfine splitting of the 2^3P_2 state, which does not mix with 2^1P_1 for the determination of Q_N . The resulting total correction is

$$\epsilon = \epsilon_{\text{rec}} + \epsilon_{\text{fs}} + \epsilon_{\text{rad}} + \epsilon_{\text{rel}} = 0.654 \times 10^{-3}. \quad (21)$$

Some previous works present these multiplicative correction for all individual hyperfine contributions, but in our opinion this cannot be fully correct because higher order relativistic corrections may include additional terms, beyond that in H_{eff} in Eq. (12). These corrections are expected to be much smaller than the experimental uncertainty for the B_J coefficient, and therefore are neglected.

III. ECG WAVE FUNCTION AND EXPECTATION VALUES

Let us now move to the calculations of the a_i , b , and c_i coefficients. To obtain sufficiently high accuracy for these parameters, we express the four-electron atomic wave function as a linear combination of properly symmetrized explicitly correlated Gaussian (ECG) functions,

$$\Psi(\{\vec{r}_a\}) = \sum_{n=1}^K t_n \mathcal{A}[\phi_n(\{\vec{r}_a\}) \chi_{\{a\}}], \quad (22)$$

where t_n are linear coefficients, $\mathcal{A}$ is the antisymmetrization operator over electronic indices, and $\{a\}$ and $\{\vec{r}_a\}$ denote the sequence of electron indices and coordinates, respectively. The electronic P symmetry of the states was enforced using the following spatial functions:

$$\phi^i(\{\vec{r}_a\}) = r_a^i \exp \left[- \sum_b w_b r_b^2 - \sum_{c<d} u_{cd} r_{cd}^2 \right] \quad (23)$$

with w_b and u_{cd} being nonlinear variational parameters, while the spin functions corresponding to different spin $S = 1$ projections were, for $\{a\} = (1, 2, 3, 4)$,

$$\chi_{\{a\}}^{1-1} = \frac{1}{\sqrt{2}} (\alpha_1 \beta_2 - \beta_1 \alpha_2) \beta_3 \beta_4, \quad (24)$$

$$\chi_{\{a\}}^{10} = \frac{1}{2} (\alpha_1 \beta_2 - \beta_1 \alpha_2) (\alpha_3 \beta_4 - \beta_4 \alpha_3), \quad (25)$$

$$\chi_{\{a\}}^{11} = \frac{1}{\sqrt{2}} (\alpha_1 \beta_2 - \beta_1 \alpha_2) \alpha_3 \alpha_4, \quad (26)$$

where α_i and β_i are the one-electron spin-up and spin-down functions. To control the uncertainty of our results, we performed the calculations with several basis sets, successively increasing their size. The nonlinear parameters were optimized variationally with respect to the nonrelativistic energy E until the energy reached stability in a desired number of digits. From the analysis of convergence, we obtained the extrapolated nonrelativistic energies and mean values used in Table I. The largest wave functions optimized variationally were composed of 6144 terms, leading to the nonrelativistic energy in agreement with the result obtained by Kedzioriski *et al.* [19].

The optimized nonrelativistic wave functions were subsequently employed to evaluate the fine and hyperfine parameters from the expectation values of the corresponding quantum-mechanical operators (2)–(10). For this purpose, we chose $J = 2$, $L = 1$, and for an arbitrary M

$$c_1 = \langle J, M | Q_1 | J, M \rangle, \quad (27)$$

$$c_2 = 6 \langle J, M | Q_2 | J, M \rangle, \quad (28)$$

$$a_1 = \langle J, M | \vec{Q}_1 \cdot \vec{J} | J, M \rangle / 3, \quad (29)$$

$$a_2 = \langle J, M | \vec{Q}_2 \cdot \vec{J} | J, M \rangle / 3, \quad (30)$$

$$a_3 = \langle J, M | \vec{Q}_3 \cdot \vec{J} | J, M \rangle, \quad (31)$$

$$b = \langle L, M | Q^{ij} L^i L^j | L, M \rangle / 5. \quad (32)$$

The numerical values of these parameters and their convergence with increasing size of the ECG basis are shown in

TABLE I. Convergence of the nonrelativistic energy and theoretical fine and hyperfine structure parameters for the 2^3P state of ^9Be (in MHz). Mass $m_N = 9.012\,183\,07(8)$ u [20] and magnetic moment $\mu/\mu_N = -1.177\,432(3)$ [21] were used for the ^9Be nucleus.

K	$E/\text{a.u.}$	c_1	c_2	a_1	a_2	a_3	b/Q_N (MHz/barn)
1536	-14.566 340 608 5	32 986.385	5 399.032	-231.116 70	-22.698 781	14.788 494	27.150 476
2048	-14.566 341 144 7	32 987.849	5 399.186	-231.126 66	-22.698 923	14.788 389	27.149 896
3072	-14.566 341 380 0	32 989.401	5 399.240	-231.130 99	-22.698 986	14.788 467	27.149 112
4096	-14.566 341 441 5	32 989.854	5 399.190	-231.129 61	-22.699 124	14.788 579	27.148 954
6144	-14.566 341 466 0	32 990.027	5 399.172	-231.128 99	-22.699 186	14.788 630	27.148 898
∞	-14.566 341 474(8)	32 990.2(2)	5 399.15(2)	-231.128 4(6)	-22.699 24(5)	14.788 68(5)	27.148 87(3)

Table I. Their relative accuracy can be estimated collectively as better than 6×10^{-6} .

IV. FINE AND HYPERFINE STRUCTURE

The hyperfine transition frequencies $\nu_J(F; F+1)$ can be expressed in terms of the A_J and B_J parameters and vice versa. For this, one writes

$$\langle H_{\text{hfs,eff}} \rangle_F = A_J A_{IJF} + B_J B_{IJF} + C_J C_{IJF}, \quad (33)$$

where

$$A_{IJF} = \frac{1}{2} K, \quad (34)$$

$$B_{IJF} = \frac{3/4 K(K+1) - I(I+1)J(J+1)}{2I(2I-1)J(2J-1)}, \quad (35)$$

with $K = F(F+1) - I(I+1) - J(J+1)$ and the total angular momentum $\vec{F} = \vec{S} + \vec{L} + \vec{I}$. The octupole term C_{IJF} is given, e.g., by Schwartz [22] and Jaccarino [23].

For $J = 1$, we arrive at

$$A_1 = -\frac{1}{6} \nu_1 \left(\frac{1}{2}; \frac{3}{2} \right) - \frac{3}{10} \nu_1 \left(\frac{3}{2}; \frac{5}{2} \right), \quad (36)$$

$$B_1 = \frac{1}{3} \nu_1 \left(\frac{1}{2}; \frac{3}{2} \right) - \frac{1}{5} \nu_1 \left(\frac{3}{2}; \frac{5}{2} \right), \quad (37)$$

and for $J = 2$

$$A_2 = -\frac{3}{50} \nu_2 \left(\frac{1}{2}; \frac{3}{2} \right) - \frac{7}{50} \nu_2 \left(\frac{3}{2}; \frac{5}{2} \right) - \frac{4}{25} \nu_2 \left(\frac{5}{2}; \frac{7}{2} \right), \quad (38)$$

$$B_2 = \frac{2}{5} \nu_2 \left(\frac{1}{2}; \frac{3}{2} \right) + \frac{2}{5} \nu_2 \left(\frac{3}{2}; \frac{5}{2} \right) - \frac{16}{35} \nu_2 \left(\frac{5}{2}; \frac{7}{2} \right), \quad (39)$$

TABLE II. A_J , B_J , and C_J parameters determined using Eqs. (36)–(40) from experimental [14] and theoretical hfs frequencies ν_J (in MHz). The theoretical B_2 is by definition equal to the experimental one, which serves for determination of b in Eq. (41).

	Experimental	Theoretical	Difference
A_1	-140.1564(8)	-140.1091	-0.0473(8)
B_1	-3.2363(7)	-3.2337	-0.0026(7)
A_2	-124.6165(5)	-124.6073	-0.0092(5)
B_2	0.7800(21)	0.7800	0.0
C_2	-0.00030(9)	-0.00032	0.00002(9)

$$C_2 = -\frac{1}{50} \nu_2 \left(\frac{1}{2}; \frac{3}{2} \right) + \frac{1}{50} \nu_2 \left(\frac{3}{2}; \frac{5}{2} \right) - \frac{1}{175} \nu_2 \left(\frac{5}{2}; \frac{7}{2} \right). \quad (40)$$

The above listed parameters (36)–(40) were evaluated using both experimental and theoretical transition frequencies $\nu_J(F; F+1)$, and their numerical values are presented in Table II. The experimental values of ν_J (in MHz) were taken from Ref. [14], assuming their uncertainties are statistical and uncorrelated. The resulting experimental values for A_1 and B_1 are the same as in Ref. [14], while for A_2 and B_2 are twice as accurate due to inclusion of the C_2 coefficient. On the other hand, theoretical values were found by diagonalization of the effective Hamiltonian of Eq. (12). Most importantly, the parameter b was fixed by matching theoretical and experimental B_2 , which holds for

$$b = 1.4525(40) \text{ MHz}, \quad (41)$$

where the uncertainty originates from that of the experimental one in B_2 .

The differences between experimental and theoretical values, shown in Table II, are not discrepancies but result from neglected higher order relativistic and QED effects, which are beyond those proportional to the Fermi contact interaction. For the A_2 coefficient, these effects are of relative order 0.07×10^{-3} , while for A_1 they are 0.33×10^{-3} . They are larger for A_1 due to hyperfine mixing of 2^3P_1 with the nearby 2^1P_1 state, which we have not taken into account in our calculation. Moreover, the relative difference for the B_1 coefficient is $\delta B_1/B_1 = 0.80 \times 10^{-3}$, which most probably is also due to this mixing. For this reason, we have chosen B_2 and not B_1 for the determination of the b coefficient.

The centroids of the calculated hyperfine energy levels are shown in Table III in the column “fs levels” for all J values of the 2^3P state. This table contains also experimental [14] and calculated fine-structure transition frequencies as well as their difference. This difference is much larger than the estimate

TABLE III. Theoretical fine structure (fs) energy levels of the 2^3P state of ^9Be , and the comparison of experimental [14] and theoretical fs transition frequencies (in GHz).

J	fs levels	$\nu^{\text{exp}}(J; J+1)$	$\nu^{\text{the}}(J; J+1)$	Diff.
2	33.890			
1	-37.489	71.86(24)	71.379	0.48(24)
0	-56.987	19.41(19)	19.498	-0.09(19)

TABLE IV. Comparison of the electric quadrupole moment (in barns) of the ^9Be nucleus obtained using atomic structure calculations.

Source	$Q_N(^9\text{Be})$
Blachman and Lurio [14]	0.049(3)
Ray <i>et al.</i> [24]	0.052 5(3)
Sinanoğlu and Beck [25]	0.054 94
Beck and Nicolaides [26]	0.055 45
Sundholm and Olsen [27]	0.052 88(38)
Jönsson and Fisher [28]	0.052 56
Nemouchi <i>et al.</i> [9]	0.052 77
Beloy <i>et al.</i> [15]	0.053(3)
This work	0.053 50(14)

of relativistic corrections, which is $\mathcal{O}((Z\alpha)^2)$. Therefore, we presume that these differences are due to inaccurate experimental results, because these fine splittings have not been directly measured but determined from the magnetic field dependence of the hfs splittings [14].

V. DETERMINATION OF $Q_N(^9\text{Be})$

Having calculated the value of $b/Q_N = 27.148\,87(3)$ MHz/barn (see Table I), and fixed b by matching theoretical and experimental values of B_2 , we obtain the electric nuclear quadrupole moment Q_N of ^9Be :

$$Q_N = 0.053\,50(14) \text{ barn.} \quad (42)$$

The uncertainty in Q_N comes from the experimental error in the ν_2 hyperfine frequencies. In this context, the uncertainties from the neglected higher order relativistic and QED effects in the b parameter, as well as the numerical uncertainties, are much smaller and are not shown.

Among the literature results listed in Table IV and depicted in Fig. 1, only the Q_N reported by Beloy *et al.* [15], thanks to its large uncertainty, agrees with our quadrupole moment. The lack of agreement with the other previous studies indicates that it is very difficult to estimate theoretical uncertainties using well-established, atomic structure methods such as multi configuration Hartree-Fock (MCHF) or CI + MBPT.

VI. CONCLUSIONS

We have determined the electric quadrupole moment $Q_N(^9\text{Be})$ with significantly higher accuracy and in disagreement with previous values. The improvement achieved in this work has several independent sources. The first one is the recalculation of the A_2 , B_2 , and C_2 coefficients from experimental hyperfine splitting. More precisely, including

FIG. 1. Comparison of the electric quadrupole moment (in barns) of the ^9Be nucleus obtained using atomic structure calculations.

C_2 in the effective Hamiltonian $H_{\text{hfs,eff}}$ in Eq. (13) enabled us to decrease the experimental uncertainties by a factor of 2. The second improvement is due to the inclusion of the hyperfine mixing by exact diagonalization of the effective fine- and hyperfine-structure Hamiltonian of Eq. (12). The third improvement comes from the very accurate calculation of the expectation values of the fine- and hyperfine-structure operators using explicitly correlated functions, allowing for the complete electron correlations. The fourth one is due to the exact inclusion of the finite nuclear mass in the fine and hyperfine interaction in Eqs. (2) and (5). Finally, the fifth source of improvement is due to accounting for the relativistic, radiative, and nuclear structure corrections by appropriate rescaling of the a_1 parameter.

Regarding our theoretical uncertainties, they come exclusively from neglected higher order relativistic and QED contributions, in particular those due to hyperfine mixing of the 2^3P_1 and 2^1P_1 states. The calculation of the complete $\mathcal{O}(\alpha^2)$ correction is possible but technically difficult. It has been performed for Be^+ in Ref. [29] due to the availability of the exponentially correlated basis functions there. Nevertheless, if a complete $\mathcal{O}(\alpha^2)$ correction is known, one can use both B_1 and B_2 parameters to obtain an even more accurate nuclear quadrupole moment of ^9Be .

ACKNOWLEDGMENTS

This research was supported by National Science Center (Poland) Grant No. 2014/15/B/ST4/05022 as well as by a computing grant from Poznań Supercomputing and Networking Center and by PL-Grid Infrastructure.

- [1] A. A. Filin, V. Baru, E. Epelbaum, H. Krebs, D. Möller, and P. Reinert, Extraction of the Neutron Charge Radius from a Precision Calculation of the Deuteron Structure Radius, *Phys. Rev. Lett.* **124**, 082501 (2020).
- [2] M. Puchalski, J. Komasa, and K. Pachucki, Hyperfine structure of the first rotational level in H_2 , D_2 , and HD Molecules and the

Deuteron Quadrupole Moment, *Phys. Rev. Lett.* **125**, 253001 (2020).

- [3] H.-G. Voelk and D. Fick, The electric moments of ^7Li from Coulomb scattering, *Nucl. Phys. A* **530**, 475 (1991).
- [4] D. Vinciguerra and T. Stovall, Electron scattering from p -shell nuclei, *Nucl. Phys. A* **132**, 410 (1969).

- [5] M. Bernheim, R. Riskalla, T. Stovall, and D. Vinciguerra, Electron scattering form factor of ^9Be , *Phys. Lett. B* **30**, 412 (1969).
- [6] M. Urban and A. J. Sadlej, The nuclear quadrupole moment of Li: Refined calculations of electric field gradients in LiH, LiF, and LiCl, *Chem. Phys. Lett.* **173**, 157 (1990).
- [7] G. H. Diercksen and A. J. Sadlej, A possible determination of the nuclear quadrupole moment of ^9Be from molecular calculations of electric properties of BeH, *Chem. Phys. Lett.* **155**, 127 (1989).
- [8] A. C. Borin, F. B. Machado, and F. R. Ornellas, Nuclear motion dependence of the electric field gradient at the ^9Be nucleus in BeH^+ , *Chem. Phys. Lett.* **196**, 417 (1992).
- [9] M. Nemouchi, P. Jönsson, J. Pinard, and M. Godefroid, Theoretical evaluation of the $^7,^9\text{Be}^- 2s2p^2\ ^4P_{1/2,3/2,5/2}$ hyperfine structure parameters and Be $2s2p\ ^3P^o$ electron affinity, *J. Phys. B* **36**, 2189 (2003).
- [10] E. Tiesinga, P. J. Mohr, D. B. Newell, and B. N. Taylor, CODATA recommended values of the fundamental physical constants: 2018, *Rev. Mod. Phys.* [to be published (2021)].
- [11] J. Krauth, K. Schuhmann, M. Ahmed *et al.*, Measuring the α -particle charge radius with muonic helium-4 ions, *Nature (London)* **589**, 527 (2021).
- [12] K. Pachucki, V. Patkóš, and V. A. Yerokhin, Three-photon-exchange nuclear structure correction in hydrogenic systems, *Phys. Rev. A* **97**, 062511 (2018).
- [13] J. L. Friar and G. L. Payne, Deuteron dipole polarizabilities and sum rules, *Phys. Rev. C* **72**, 014004 (2005).
- [14] A. G. Blachman and A. Lurio, Hyperfine structure of the metastable $(1s^22s2p)\ ^3P$ states of $^4\text{Be}^9$ and the nuclear electric quadrupole moment, *Phys. Rev.* **153**, 164 (1967).
- [15] K. Beloy, A. Derevianko, and W. R. Johnson, Hyperfine structure of the metastable 3P_2 state of alkaline-earth-metal atoms as an accurate probe of nuclear magnetic octupole moments, *Phys. Rev. A* **77**, 012512 (2008).
- [16] A. Hibbert, Developments in atomic structure calculations, *Rep. Prog. Phys.* **38**, 1217 (1975).
- [17] M. I. Eides, H. Grotch, and V. A. Shelyuto, Theory of light hydrogenlike atoms, *Phys. Rep.* **342**, 63 (2001).
- [18] M. Puchalski and K. Pachucki, Fine and hyperfine splitting of the $2P$ state in Li and Be^+ , *Phys. Rev. A* **79**, 032510 (2009).
- [19] A. Kedziorowski, M. Stanke, and L. Adamowicz, Atomic fine-structure calculations performed with a finite-nuclear-mass approach and with all-electron explicitly correlated Gaussian functions, *Chem. Phys. Lett.* **751**, 137476 (2020).
- [20] M. Wang, G. Audi, F. G. Kondev, W. Huang, S. Naimi, and X. Xu, The AME2016 atomic mass evaluation (II): Tables, graphs and references, *Chin. Phys. C* **41**, 030003 (2017).
- [21] N. J. Stone, Nuclear magnetic dipole and electric quadrupole moments: Their measurement and tabulation as accessible data, *J. Phys. Chem. Ref. Data* **44**, 031215 (2015).
- [22] C. Schwartz, Theory of hyperfine structure, *Phys. Rev.* **97**, 380 (1955).
- [23] V. Jaccarino, J. G. King, R. A. Satten, and H. H. Stroke, Hyperfine structure of I^{127} . Nuclear magnetic octupole moment, *Phys. Rev.* **94**, 1798 (1954).
- [24] S. N. Ray, T. Lee, and T. P. Das, Study of the nuclear quadrupole interaction in the excited (2^3P) state of the beryllium atom by many-body perturbation theory, *Phys. Rev. A* **8**, 1748 (1973).
- [25] O. Sinanoğlu and D. R. Beck, HFS constants of Be I $1s^22s2p\ ^3P_0$, Be I $1s^22s^2p^2\ ^4P$, and Be I $1s^22s^2p^2\ ^2D$ obtained from the non-closed shell many-electron theory for excited states, *Chem. Phys. Lett.* **20**, 221 (1973).
- [26] D. R. Beck and C. A. Nicolaides, Fine and hyperfine structure of the two lowest bound states of Be^- and their first two ionization thresholds, *Int. J. Quantum Chem.* **26**, 467 (1984).
- [27] D. Sundholm and J. Olsen, Large MCHF calculations on the hyperfine structure of $\text{Be}(^3P^o)$: The nuclear quadrupole moment of ^9Be , *Chem. Phys. Lett.* **177**, 91 (1991).
- [28] P. Jönsson and C. F. Fischer, Large-scale multiconfiguration Hartree-Fock calculations of hyperfine-interaction constants for low-lying states in beryllium, boron, and carbon, *Phys. Rev. A* **48**, 4113 (1993).
- [29] M. Puchalski and K. Pachucki, Ground-state hyperfine splitting in the Be^+ ion, *Phys. Rev. A* **89**, 032510 (2014).