Comment on “Natural BNS damping of the fast ion instability”

G. V. Stupakov

Stanford Linear Accelerator Center, Stanford University, P.O. Box 4349, Stanford, California 94309

(Received 14 June 1999; published 11 January 2000)

We show that the effect of BNS damping for the fast ion instability found in the recent paper “Natural BNS damping of the fast ion instability” [Phys. Rev. ST Accel. Beams 2, 044403 (1999)] occurs only when initial perturbations in the beam are amplified by many orders of magnitude, and, most likely, the linear approximation breaks down. We also demonstrate that the asymptotic solution found by Raubenheimer and Zimmermann gives a good approximation for the linear regime of the instability contrary to the statement of Pestrikov.

PACS numbers: 29.27.-a, 41.85.-p

In a recent paper [1], Pestrikov revised the previous approach to the derivation of the fast ion instability (FII). Based on the same system of equations as in Refs. [2,3], but using a more sophisticated analysis, he finds that, due to the variation of the betatron frequency in the beam, after an initial growth, the amplitude of the beam oscillation saturates and goes down. He concludes that, asymptotically, the instability is eliminated due to the BNS damping effect. Comparing his solution with the solution of Ref. [2], he also claims that the latter gives a poor representation for both small and large values of time.

The purpose of this paper is to show that although the analysis of Ref. [1] is mathematically correct, the effect of instability saturation occurs only on such time scale when, in most practical cases, the instability goes to the nonlinear regime and the linear approximation breaks down. We will also show that the asymptotic solution found in Refs. [2,3] gives a good approximation for the initial development of the instability, contrary to the statement of Ref. [1].

The analysis of FII in all three papers [1–3] is based on the following equation for the beam centroid offset (we use notation of Ref. [3]):

$$\begin{aligned} \frac{\partial^2}{\partial s^2} y(s, z) + \frac{\omega_\beta^2}{c^2} y(s, z) \\ = -\kappa \int_0^z z' \frac{\partial y(s, z')}{\partial z'} \cos \omega_i(z - z') dz'. \end{aligned} \quad (1)$$

Here s is the distance along the beam orbit, $z = ct - s$ is the coordinate measured along the beam from the head of the bunch train, the factor κ describes the interaction strength of the electrons with the ions and is a small parameter, ω_i is the ion frequency, and ω_β is the betatron frequency. Pestrikov uses a different notation, in which $k = \omega_\beta/c$, $C = c\kappa/2\omega_\beta$, and $\Omega = \omega_i/c$.

First, we remind the reader that the theory developed in Refs. [2,3] assumes that the interaction between the beam and the ions is small,

$$c^2 \kappa l \ll \omega_i^2, \omega_\beta^2, \quad (2)$$

where l denotes the length of the bunch train, so that the instability develops on a time scale much larger than both the betatron period and the period of ion oscillations. In such a situation, the most unstable solution of Eq. (1) can be represented as a wave propagating in the beam with a slowly varying amplitude and phase,

$$y(s, z) = \text{Re} A(s, z) e^{-i\omega_\beta s/c + i\omega_i z/c}, \quad (3)$$

where $A(s, z)$ is the complex amplitude. It is also assumed that A is a “slow” function of its variables,

$$\left| \frac{\partial \ln A}{\partial s} \right| \ll \frac{\omega_\beta}{c}, \quad \left| \frac{\partial \ln A}{\partial z} \right| \ll \frac{\omega_i}{c} \quad (4)$$

(these two conditions were explicitly formulated in Ref. [3]).

Using an averaging method, it was found in [2,3] that, for initial condition $A(s=0, z) = 1$, the solution of Eq. (1) is $A(s, z) = I_0(z\sqrt{\kappa\omega_i s/2\omega_\beta})$, where I_0 is the modified Bessel function of zeroth order. Asymptotically, for $s \gg \omega_\beta/\kappa\omega_i l^2$,

$$A(s, z) \approx \left(2\pi z \sqrt{\frac{\kappa\omega_i s}{2\omega_\beta}} \right)^{-1/2} \exp\left(z \sqrt{\frac{\kappa\omega_i s}{2\omega_\beta}} \right), \quad (5)$$

FIG. 1. (Color) Numerical solution of Eq. (1), $A(s)$, for $z = l$ (curve 1) and $z = l/2$ (curve 2). Dashed curves show the respective analytical solution $A(s, z) = I_0(z\sqrt{\kappa\omega_i s/2\omega_\beta})$. The inset shows the magnified part of curve 1 with its asymptotic counterpart.

TABLE I. Maximum amplification for the asymptotic solution.

	PEP-II HER	NLC PL	Accelerator		HERA e^-	CESR
l (m)	2000	38	NLC ML	ALS	6048	670
$\omega_i/2\pi$ (MHz)	4	104	200	25	0.8	0.6
$A_{\max}$	5.8×10^{21}	1.8×10^{10}	5.4×10^{20}	1.7×10^{13}	5.9×10^{12}	3.5

which indicates an instability with a characteristic time (in the tail of the beam) $\tau \approx 2\omega_\beta/\kappa\omega_i l^2 c$.

The solution (5) is valid only if s is not very large. The applicability condition for this solution can be obtained from the second inequality in (4):

$$s \ll s_{\max} \equiv \frac{2\omega_\beta\omega_i}{\kappa c^2}, \quad (6)$$

or, in notation of Ref. [1], $s \ll \Omega/C$. This result explains why the saturation effect found in Ref. [1] is missing in Eq. (5). Indeed, according to [1], the instability saturates at a time when $sC/\Omega \approx 2$, or $s \approx 2s_{\max}$, which lies beyond the applicability limit of the asymptotic theory developed in Refs. [2,3].

There is also a statement in Ref. [1] that even for $s \ll s_{\max}$ Eq. (5) gives a poor representation of the solution. To verify that statement, we wrote a computer code that integrates Eq. (1) numerically. The validity of the numerical integration obtained from this code is not restricted by the applicability condition [Eq. (6)] of the quasiexponential solution. Using this code, we found numerical solutions of Eq. (1) for several different values of parameters κ , ω_β , and ω_i , with initial condition $A(s=0, z) = 1$. They all showed a good agreement with Eq. (5) when condition (6) is satisfied. An example of numerical solution for $lc^2\kappa/\omega_\beta^2 = 0.05$, $l\omega_i/c = 20$ is shown in Fig. 1. This solution demonstrates the saturation effect found in [1], which, however, occurs only after the initial amplitude grows by many orders of magnitude. For these parameters, $s_{\max}\omega_\beta/c = 400$, and one can expect a good agreement with Eq. (5) for $s\omega_\beta/c < 200$, as, indeed, seen in the inset.

Since the asymptotic solution (5) is valid only for a limited time, a practically important question arises, if this time interval is large enough to describe the growth of the instability. To answer this question, we estimated the maximum growth allowed by the asymptotic solution, putting $s = \alpha s_{\max}$ into Eq. (5) where α is a small factor. This gives the maximum amplification factor in the asymptotic regime as

$$|A|_{\max} = \frac{1}{(2\pi\alpha z\omega_i/c)^{1/2}} e^{\alpha z\omega_i/c}. \quad (7)$$

We calculated $|A|_{\max}$ assuming $\alpha = 0.1$ for the FII in different accelerators using the values for the bunch train l and the ion frequency ω_i from Table I of Ref. [2]. The result is shown in Table I.

As seen from this table, for most accelerators the amplification in the linear regime that is represented by Eq. (5) lies between 10 and 20 orders of magnitude. The BNS damping effect would occur when $\alpha \approx 1$, resulting in an even larger amplification factor. It is clear that, well before the saturation occurs, the amplitude of the oscillations would become comparable to the size of the transverse beam, the linear theory would break down, and the instability would proceed into nonlinear regime.

In conclusion, we found that the solution found by Raubenheimer and Zimmermann in [2], for most practical cases, adequately describes the growth of the oscillations in the beam until the nonlinear effects become important. Although the saturation effect described in Ref. [1] is an interesting mathematical feature of the solution, it occurs after an initial perturbation is amplified by many orders of magnitude, and, most likely, the instability will proceed into nonlinear regime. By direct numerical integration of Eq. (1), we also showed a good agreement between the numerical solution and the asymptotic representation found in [2,3].

ACKNOWLEDGMENT

This work was supported by the U.S. Department of Energy, Office of Basic Energy Sciences, under Contract No. DE-AC03-76SF00515.

-
- [1] D. V. Pestrikov, Phys. Rev. ST Accel. Beams **2**, 044403 (1999).
 - [2] T. O. Raubenheimer and F. Zimmermann, Phys. Rev. E **52**, 5487 (1995).
 - [3] G. V. Stupakov, T. O. Raubenheimer, and F. Zimmermann, Phys. Rev. E **52**, 5499 (1995).