rf driven stable spin-flipping motion of a stored polarized beam

Ya. S. Derbenev and V. A. Anferov*

Physics Department, University of Michigan, Ann Arbor, Michigan 48109-1120

(Received 30 March 2000; published 7 September 2000)

The conventional method of spin flipping a stored polarized beam is based on slowly crossing an rf induced depolarizing resonance. This paper discusses a different approach where the polarization reversal is achieved by trapping the beam polarization into a stable spin-flipping motion on top of rf induced resonance at a half-revolution frequency.

PACS numbers: 41.75.Ak, 29.27.Bd, 29.27.Hj

I. INTRODUCTION

Developing an efficient spin-flipping technique is important for high-energy spin experiments since frequent reversals of the beam polarization can significantly reduce systematic errors in an experiment's spin asymmetry measurements. To spin flip a stored polarized beam, one can slowly ramp the frequency of an rf magnet (either solenoid or dipole) through the rf induced depolarizing resonance. This technique was successfully applied for spin-flipping polarized proton beam stored in the Indiana University cooler ring with and without a Siberian snake [1]. To achieve good spin-flip efficiency, this method requires a slow resonance crossing rate, which also increases the time of each spin flip. Moreover, slow resonance crossing makes the spin motion sensitive to weak synchrotron sideband resonances or higher-order depolarizing resonances that may occur in the vicinity of the rf induced resonance.

In this paper we discuss another possible way of spin flipping the beam polarization by rearranging the stable spin motion in such a way that the polarization direction alternates on every particle turn around the ring without any depolarization.

II. STABLE SPIN MOTION AT AN RF INDUCED RESONANCE

In a circular accelerator or a storage ring with no Siberian snakes, the spin vector of each particle precesses around the vertical magnetic field of the ring's dipole magnets. For a particle moving along the closed orbit, the spin tune ν_s , which is the number of spin precessions during one turn around the ring, is proportional to the beam energy

$$\nu_s = G\gamma, \quad (1)$$

where γ is the Lorentz energy factor and $G = (g - 2)/2$ is the particle's gyromagnetic anomaly. While for protons $G = 1.79285$, it is much smaller for electrons ($G = 0.00116$) and deuterons ($G = -0.1426$).

This vertical spin precession can be perturbed by any horizontal rf magnetic field whenever its frequency is in resonance with the spin motion

$$f_{\text{rf}} = f_{\text{circ}}(k \pm \nu_s), \quad (2)$$

where f_{circ} is the beam circulation frequency and k is an integer. Near the rf induced resonance, the spin precession becomes unstable, which could lead to depolarization of a vertically polarized beam. However, the same rf magnetic field establishes a new stable spin direction in the horizontal plane. To show this, let us consider the spin motion in the presence of an rf spin perturbation $\varepsilon e^{-i\omega\theta}$, where ε is the resonance strength, which is the angle of spin perturbation on top of the resonance per radian of the closed orbit. We also assume that the perturbation frequency $\omega = f_{\text{rf}}/f_{\text{circ}} \pm k$ is close to the resonance condition of Eq. (2). The equation of spin motion can be written for the spinor wave function ψ in the following form [2]:

$$\frac{d\psi}{d\theta} = -\frac{i}{2} \begin{pmatrix} G\gamma & \varepsilon e^{-i\omega\theta} \\ \varepsilon^* e^{i\omega\theta} & -G\gamma \end{pmatrix} \psi, \quad (3)$$

where θ is the azimuthal particle coordinate in the ring. In this notation, the diagonal terms in Eq. (3) represent spin rotation around the vertical axis, while the off diagonal terms correspond to horizontal spin perturbation. Transforming the above equation into the resonance rotating frame, $\psi = e^{-i\omega\theta\sigma_3/2}\xi$, the equation of spin motion becomes

$$\frac{d\xi}{d\theta} = -\frac{i}{2} \begin{pmatrix} G\gamma - \omega & \varepsilon \\ \varepsilon^* & \omega - G\gamma \end{pmatrix} \xi. \quad (4)$$

Note that exactly on top of the rf induced resonance $G\gamma - \omega = 0$, and the spin precesses around the spin perturbing field ε , which rotates with frequency $\omega = f_{\text{rf}}/f_{\text{circ}} \pm k$ in the laboratory frame. Thus, the direction along the rotating spin perturbation field becomes stable for the spin motion when the resonance condition of Eq. (2) is satisfied. Next, we consider how the existence of such a rotating horizontal stable spin direction can be used for spin flipping.

*Present address: Indiana University Cyclotron Facility, Bloomington, IN 47408.

III. SPIN FLIPPING AT SELECTED ENERGIES IN THE ABSENCE OF SIBERIAN SNAKES

As we saw in the previous section, an external rf magnetic field, when at resonance with the spin precession, creates a horizontal stable spin direction which rotates around the ring with angular frequency

$$\omega\theta = (f_{\text{rf}}/f_{\text{circ}} \pm k)\theta = G\gamma\theta. \quad (5)$$

When $G\gamma$ is equal to a half-integer number, the horizontal spin rotates by exactly 180° in one turn around the ring. Thus, the spin would flip its direction after every turn. This opens a possibility to organize the spin motion in such a way that spin would arrive at the experimental section with longitudinal polarization, whose sign alternates after every turn. A practical solution would require installing an rf solenoid at the experimental straight section and matching the injected beam polarization with the longitudinal direction at the experimental section. One can also use radial rf dipole field to create a spin-flipping motion of a horizontal spin. The stable spin direction would be radial near the rf dipole, while in a different straight section it will be rotated towards the longitudinal direction by the angle

$$\phi = G\gamma\theta_{\text{bend}}, \quad (6)$$

where θ_{bend} is the orbit bend between the rf dipole and the point of interest. Note that the spin tune $G\gamma$ should be half-integer for both the rf solenoid and rf dipole induced spin flipping.

There are several effects that could potentially perturb the spin motion driven by the rf magnet. With an energy offset or spread present in the beam, particles will have their spin precession frequency slightly shifted from the rf resonance frequency. An offset from the rf resonance would tilt the stable spin direction out of the horizontal plane by an angle β ,

$$\beta = \tan^{-1}\left(\frac{\Delta\nu_s}{\varepsilon}\right), \quad (7)$$

where $\Delta\nu_s = G\Delta\gamma$ is the spin tune shift due to the energy offset. The tilt in the stable spin direction would in turn reduce the equilibrium horizontal beam polarization P by a factor of

$$\frac{\Delta P}{P} = 1 - \cos^2 \beta \approx \left(\frac{\Delta\nu_s}{\varepsilon}\right)^2. \quad (8)$$

Thus, in order to maintain control over the spin motion ($\Delta P/P < 0.1$), the rf induced resonance should dominate over the energy spread,

$$\varepsilon \geq 3\Delta\nu_s. \quad (9)$$

For example, to overcome a spin tune shift of 0.001 the strength of the rf induced resonance should be about 0.003. The average stable polarization magnitude during the spin-flipping process is plotted against the $\varepsilon/\Delta\nu_s$ ratio in Fig. 1.

FIG. 1. The average stable polarization magnitude during the spin-flipping process is plotted against the $\varepsilon/\Delta\nu_s$ ratio. The solid curve is derived analytically from the spin transformation over two turns around the ring. The dashed curve is stable polarization predicted by Eq. (8).

Similarly, the rf induced resonance should dominate over higher-order horizontal spin perturbations, which would also tend to perturb the stable spin direction. It is also important to note that, in the spin-flipping method described here, the rf solenoid would operate at exactly one-half of the beam circulation frequency. Therefore, one particular part of a continuous beam would always pass the rf magnet when its field is very close to zero and, thus, would not satisfy the stability condition of Eq. (9). To avoid this problem, the stored polarized beam should be bunched and synchronized with the spin-driving rf field. Another solution could be to use a high frequency rf magnet (or a special rf cavity) synchronized with every bunch in the beam. This could be done when there is an odd number of bunches in the ring; the rf magnet would then be operating at the frequency $f_{\text{rf}} = \frac{h}{2}f_{\text{circ}}$, where h is the harmonic number (odd integer) of the main rf cavities.

IV. SPIN FLIPPING IN THE PRESENCE OF SIBERIAN SNAKES

In the presence of a full Siberian snake [3] in the ring, the spin tune becomes half-integer and energy independent. With a proper choice of the betatron tunes, a set of Siberian snakes can overcome all dangerous depolarizing resonances in the ring [4]. Nevertheless, an rf magnetic field in resonance with the half-integer spin tune can depolarize the beam even in the presence of a Siberian snake. These resonances are often called “snake” depolarizing resonances [5]. Similar to the case with no snake in the ring, such an rf induced snake resonance creates a rotating stable spin direction. Moreover, this rotating stable spin direction would be flipped after every turn around the

ring since the spin tune is half-integer in the presence of a Siberian snake.

To create such a stable spin-flipping mode of the spin motion, the rf magnetic field has to be orthogonal to the unperturbed spin. With a single snake in the ring and no spin perturbation present, the stable spin direction is horizontal and coincides with the snake axis in the straight section opposite to the snake location. Therefore, an rf dipole with vertical field operating at $0.5f_{\text{circ}}$ would make the vertical direction stable for the spin motion, while the polarization direction would flip after every turn around the ring.

With an even number of snakes in the ring, the unperturbed stable spin direction is vertical. Ideally, one could use a longitudinal rf field near the interaction region to create the longitudinal spin stable and flipping every turn around the ring. However, solenoids become impractical at high energies since their spin rotation angle varies inversely with the beam momentum. An energy independent spin rotation in a dipole makes dipole magnets more attractive for spin manipulation at high energies. An rf dipole with radial field could create stable spin flipping of the beam that has radial polarization near the rf dipole location, while at the experimental straight section the polarization would be longitudinal.

The stability of the continuous spin-flipping motion could be lost when the spin tune moves out of the induced rf resonance. Such a spin tune shift could be caused by a small error in the snake current or by some high-order spin perturbation. As in the case with no Siberian snake in the ring, the strength of the rf induced resonance determines the tolerable spin tune shift from the half-integer value [as indicated in Eq. (8)]. The effect of the high-order spin perturbation could be reduced by a proper choice of the betatron tunes. Therefore, an adequate strength of the rf magnet (to achieve $\varepsilon \simeq 10^{-3}$) and a snake current precision at the level of 10^{-4} would provide stability of the spin-flipping motion.

In practice, an rf dipole field of 0.1 T could be obtained [6] in the frequency range near 20 kHz, which corresponds to one-half of the circulation frequency in several modern high energy hadron synchrotrons. The spin perturbation strength of a 1 m long rf dipole of this type would be

$$\varepsilon = \frac{Ge}{2\pi mc} \int B dl \sim 0.01 \text{ (for protons)}. \quad (10)$$

Such an rf dipole would certainly have adequate strength to control the spin motion.

V. SPIN MANIPULATION OF A POLARIZED DEUTERON BEAM

The spin-flipping technique discussed above seems especially attractive for polarized deuteron beams. The conventional spin manipulation methods become difficult due to the deuteron's small anomalous magnetic moment. A

full Siberian snake would be practical only at low energies, where solenoidal magnets could provide the required spin rotation (a 200 MeV deuteron beam would require a solenoid with 11 Tm field integral). Similarly, the effect of the rf magnetic field on spin will be rather small, which limits applicability of the conventional spin-flipping technique. In contrast, the method presented here uses the accelerator lattice to flip the spin while the rf field keeps the spin motion stable. This feature becomes an advantage for particles with small anomalous magnetic moment in the energy region where $G\gamma < 10$. In the case of deuteron beam, stable spin-flipping motion could be achieved at the beam kinetic energies corresponding to half-integer $G\gamma$ values,

$$T [\text{GeV}] = 4.7 + 13.156n, \quad (11)$$

where n is an integer. A strong rf field could also create stability for the nonflipping longitudinal polarization direction when $G\gamma$ is an integer. However, the rf field in this case should be strong enough to dominate over the imperfection fields in the accelerator. Note that there are no such first-order spin perturbations near a half-integer $G\gamma$, and the required rf field could be much weaker.

The direct effect of the rf magnetic field on the deuteron's spin is rather small. However, an rf dipole also excites beam oscillations and the spin is mainly driven by the quasis resonant accumulation of the spin kicks from all quadrupoles of the accelerator lattice. To estimate the amplification of the spin perturbation due to the excited beam oscillations, we first calculate the beam oscillation amplitude using the approach discussed in [7]. Considering the beam motion in the phase space coordinates $(x, x'\beta_x)$, it is convenient to transform to the rf rotating frame where the beam kick by the rf dipole is constant. After the rf dipole's kick, the particle's phase space vector would rotate by an angle

$$2\pi\delta = 2\pi\left(\nu_x - \frac{f_{\text{rf}}}{f_{\text{circ}}} + k\right), \quad (12)$$

where ν_x is the horizontal betatron frequency and k is an integer. When the rf dipole is turned on adiabatically, the excited beam oscillations reach a stable mode which is shown in Fig. 2. The beam oscillation amplitude X_0 for this stable mode is given by

$$2\pi\delta X_0 = \beta_0 x' = \beta_0 \frac{1}{2} \int \frac{B_{\text{rf}} d\ell}{B\rho}, \quad (13)$$

where β_0 is the value of the horizontal beta function near the rf dipole. The resulting beam oscillations around the ring can be written as

$$x(s) = X_0 \sqrt{\frac{\beta_x(s)}{\beta_0}} \cos(\nu_{\text{rf}}\theta), \quad (14)$$

where $\nu_{\text{rf}} = k + f_{\text{rf}}/f_{\text{circ}}$ is the rf harmonic closest to the betatron tune (i.e., $\nu_{\text{rf}} \sim \nu_x$). The strength of the induced spin perturbation is given by the spin kick accumulated in the lattice quadrupoles over one turn around the ring,

FIG. 2. Beam oscillations excited by the rf dipole are shown in the phase space rotating with the rf frequency. The amplitude of the excited beam oscillations, X_0 , is determined by the rf kick amplitude and the rf frequency separation from the betatron tune $\delta = \nu_x - \frac{f_{rf}}{f_{circ}} + k$. This stable mode of the excited beam oscillations is achieved when rf dipole is turned on adiabatically.

$$\varepsilon_b = \frac{G\gamma}{2\pi} \oint g(s)x(s) e^{i\nu_s\theta} ds, \quad (15)$$

where $g(s) = \frac{\partial B_y / \partial x}{B\rho}$ is the normalized strength of the quadrupoles around the ring and ν_s is the spin tune. Equation (15) should be compared with the strength of the spin perturbation by the direct effect of the rf magnetic field, which is given by

$$\varepsilon_0 = \frac{G\gamma}{4\pi} \int \frac{B_{RF} d\ell}{B\rho}. \quad (16)$$

In both cases, the effect of transverse magnetic fields is proportional to the anomalous magnetic moment [8]. Substituting Eq. (14) into Eq. (15), one would quickly obtain

$$\varepsilon_b = \varepsilon_0 \frac{\sqrt{\beta_0}}{4\pi\delta} \oint g(s) \sqrt{\beta_x(s)} e^{i(\nu_s \pm \nu_\pi)\theta} ds. \quad (17)$$

When the spin tune is close to the frequency of the induced beam oscillations, $|\nu_s - \nu_{rf}| \ll 1$, one can neglect the exponent in the integral. The remaining integral resembles the definition of the chromaticity function $\xi_x = \frac{-1}{4\pi} \oint g(s) \beta_x(s) \approx -\nu_x$. Therefore, the strength of the induced spin perturbation can be estimated as

$$\varepsilon_b \approx \varepsilon_0 \frac{\nu_x}{2\delta} \frac{\sqrt{\beta_0}}{\langle \sqrt{\beta_x} \rangle}. \quad (18)$$

This estimate corresponds to the maximum amplification of the spin perturbation which occurs in the vicinity of the strong intrinsic depolarizing resonances $\nu_s \approx \nu_x$, where

the spin kicks from all lattice quadrupoles are synchronized with the spin precession. Farther away from these regions, the effect of the induced beam oscillations on spin is reduced by the exponent in Eq. (17); in that case, smaller δ would enhance the excited beam oscillation amplitude as well as the induced spin perturbation.

Finally, we would like to comment that acceleration of polarized deuterons seems possible in modern high energy rings [9]. While full Siberian snakes do not seem practical for high-energy deuterons, the depolarizing resonances for a deuteron beam are 25 times weaker and 25 times farther apart than for protons. Therefore, one could use individual resonance correction techniques developed for the proton beam at the Brookhaven National Laboratory AGS ring. A partial Siberian snake could overcome all imperfection depolarizing resonances [10]. Such a partial snake could be realized using either a solenoid magnet or a set of correction dipoles distributed around the ring to create a controlled closed orbit perturbation. Note that, in an ideal case, reversing the axis of the partial snake could flip the longitudinal polarization of the beam stored near an integer $G\gamma$. However, without additional correction of the natural imperfection resonance this spin-flipping method remains impractical due to inevitable polarization losses. The intrinsic depolarizing resonances could be handled by an rf dipole which induces coherent beam oscillations and makes intrinsic resonances strong enough to spin flip [11]. An rf spin perturbation could also be used as a spin rotator for polarized deuterons. For example, to bring initial vertical beam polarization to the longitudinal direction, a horizontal rf dipole field could be applied for a time period which corresponds to the required $\pi/2$ spin rotation.

VI. SUMMARY

In summary, we found that an external rf magnetic field can be used to create a stable mode of the spin motion, where the polarization direction flips after every turn around the ring. Such stable spin-flipping spin motion can be realized whenever the spin tune is equal to a half-integer value; this is always true in rings equipped with Siberian snakes, while rings without Siberian snakes also can reach half-integer spin tune at certain energies. The applied rf magnetic field should be orthogonal to the unperturbed stable spin direction and should operate at exactly one-half of the circulation frequency. Provided that the rf field is synchronized with the circulating beam and is strong enough to dominate over possible spin tune spread, this spin-flipping motion is stable. Note that the spin flip is driven by the accelerator lattice itself, while the rf magnet provides stability for the spin-flipping motion. This method could be used for spin-flipping polarized beams in storage rings such as the MIT-Bates Storage Ring, Relativistic Heavy Ion Collider, and HERA at DESY. Moreover, as was noted earlier [12], the rf stabilization of the

spin motion against the spin tune spread is an interesting possibility for accelerators with Siberian snakes.

ACKNOWLEDGMENTS

We would like to thank P. Schwandt and Yu. Shatunov for many valuable suggestions improving this paper. This work was supported by a research grant from the U.S. Department of Energy.

Note added.—After submitting the paper for publication, an electron storage ring proposal by Baturin *et al.* [13] was brought to our attention, which also mentions the possibility of organizing the stable longitudinal spin motion on top of an rf induced resonance. We believe that our paper further enhances their pioneering idea by establishing the spin stability criteria and focusing on the spin-flipping applications in rings with and without Siberian snakes.

-
- [1] D.D. Caussyn *et al.*, Phys. Rev. Lett. **73**, 2857 (1994);
B.B. Blinov *et al.*, Phys. Rev. Lett. **81**, 2906 (1998).

- [2] B. W. Montague, Part. Accel. **11**, 219 (1981).
[3] Ya. S. Derbenev and A. M. Kondratenko, Sov. Phys. Dokl. **20**, 562 (1978).
[4] S. Y. Lee and E. D. Courant, Phys. Rev. D **41**, 292 (1990).
[5] S. Y. Lee and S. Tepikian, Phys. Rev. Lett. **56**, 1635 (1986);
R. A. Phelps *et al.*, Phys. Rev. Lett. **78**, 2772 (1997).
[6] B. Parker *et al.*, in *Proceedings of the 1999 Particle Accelerator Conference New York* (IEEE, Piscataway, NJ, 1999), p. 3336; P. Schwandt (private communication).
[7] M. Bai *et al.*, Phys. Rev. E **56**, 6002 (1997).
[8] A. M. Kondratenko, in *Proceedings of the 9th International Symposium on High Energy Spin Physics, Bonn, 1990* (Springer-Verlag, Berlin, 1991), p. 140.
[9] E. D. Courant, in *Proceedings of the Workshop on RHIC Spin Physics, 1998* (BNL Report No. BNL-65615, 1998), p. 275; Spin Note AGS/RHIC/SN 066, 1997.
[10] V. A. Anferov *et al.*, Phys. Rev. A **46**, R7383 (1992);
H. Huang *et al.*, Phys. Rev. Lett. **73**, 2982 (1994).
[11] M. Bai *et al.*, Phys. Rev. Lett. **80**, 4673 (1998).
[12] Ya. S. Derbenev, in *Proceedings of the DESY Workshop on Polarized Protons at High Energies* (DESY Report No. DESY-PROC-1999-03, 1999), p. 225.
[13] P. I. Baturin *et al.*, in *High-Energy Spin Physics*, edited by Kenneth J. Heller, AIP Conf. Proc. No. 187 (AIP, New York, 1989), p. 1028.