



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Resonant nonlinear refraction of 4.3- μm light in CO₂ gas

J. J. Pigeon, D. Tovey, S. Ya. Tochitsky, G. J. Louwrens, I. Ben-Zvi, D. Martyshkin, V. Fedorov, K. Karki, S. Mirov, and C. Joshi

Phys. Rev. A **100**, 011803 — Published 30 July 2019

DOI: [10.1103/PhysRevA.100.011803](https://doi.org/10.1103/PhysRevA.100.011803)

Resonant nonlinear refraction of 4.3- μm light in CO_2 gas

J.J. Pigeon¹, D. Tovey², S. Ya. Tochitsky², G. J. Louwrens², I. Ben-Zvi¹
D. Martyshkin³, V. Fedorov³, K. Karki³, S. Mirov³, and C. Joshi²

¹*Department of Physics and Astronomy, Stony Brook University, Stony Brook, New York, 11794, USA*

²*Department of Electrical Engineering, University of California at Los Angeles, Los Angeles, California 90095, USA*

³*Department of Physics, University of Alabama at Birmingham, Birmingham, Alabama 35294, USA*
jeremy.pigeon@stonybrook.edu

Abstract

We use time- and frequency-resolved measurements of self-focusing and self-defocusing of 4.3 μm radiation in CO_2 gas to study the resonant nonlinear optical response in the vicinity of individual rovibrational lines. Measurements over a range of lines on the 4P and 4R branch of the 000 – 001 transition indicate that the nonlinear response of CO_2 is strongly affected by power broadening, resulting in a sign reversal of the nonlinearity as compared to that of a saturable two-level system.

I. Introduction

The resonant interaction of light with matter is the central pillar of atomic, molecular and optical physics. Moreover, knowledge of the resonant nonlinear optical response of atoms and molecules in the gas phase underpins the progress of a wide range of sub-fields such as nonlinear optics, quantum electronics, and ultrafast science. With the advent of spectrally bright and coherent laser sources, there have been extensive theoretical and experimental studies dedicated to understanding the resonant nonlinear optical interaction of visible and near-IR light with the electronic transitions of atomic gases, usually alkali metal vapors. In doing so, researchers have observed and characterized numerous resonant optical phenomena such as saturable absorption [1], resonant self-focusing [2], self-defocusing [3], electromagnetically induced transparency [4], and the production of slow light [5]. Further sophistication of this effort has even enabled single-photon nonlinear processes thus providing a notable advance in the field of quantum optics [6].

There has been much less effort, however, dedicated to performing such experiments in molecular gases, where the nonlinear optical effects are dominated by rovibrational transitions that are active in the mid- and long-wave IR (LWIR) spectral range. These studies, hindered by a lack of high-power, continuously tunable sources in the LWIR, have been performed using line-tunable CO_2 lasers thereby constraining measurements to coincidental overlap between CO_2 laser lines and the transitions of molecules under investigation. As a result, previous studies on nonlinear self-focusing and self-defocusing have been limited to molecules such as SF_6 , NH_3 , and CDF_3 [7-11] for which the spectra are rather complicated. Both basic and applied research in this field would benefit from extending these studies to smaller molecules such as CO_2 and CO that have fewer modes of vibration. Thus, their relatively simple spectra allow for a systematic study of the

resonant nonlinear optical response in the vicinity of individual rovibrational lines. Furthermore, these molecules exist in substantial quantities in the atmosphere and, as a result, measurements of the nonlinear rovibrational response of such IR-active, minor air constituents are needed to support the burgeoning interest in the LWIR filamentation in air [12] and in the development of a nonlinear refractive index equivalent of the HITRAN database [13].

In this letter, we report on the first study of the resonant rovibrational nonlinearity of CO₂ gas driven by a continuously tunable, Fe:ZnSe laser operating near 4.3 μm. By performing frequency- and time-resolved measurements of self-focusing and self-defocusing of the 4.3 μm beam, we have characterized the dispersion of the resonant Kerr nonlinearity in the vicinity of a single rovibrational line of the CO₂ molecule. We have found that the rovibrational nonlinearity of CO₂ is dominated by the resonant ac-Stark or power broadening effect and that the sign of the nonlinearity is opposite to that predicted by the commonly used two-level system model that describes the resonant nonlinear optics of both atoms and molecules in the gas phase [14]. We determine that the nonlinear refractive index of low-pressure CO₂ is $\sim 10^{-12}$ cm²/W suggesting that the 400 ppm CO₂ concentration can act to increase the nonlinear refractive index of ambient air by four orders of magnitude as compared to the non-resonant values measured in the mid- and LWIR [15-18].

II. Experiment

Continuously tunable mid-IR laser pulses produced by a Fe:ZnSe laser system were used to study the resonant nonlinear optics of CO₂ gas. The room temperature Fe:ZnSe laser was optically pumped by 2.94 μm, 15 mJ laser pulses produced by a Q-switched Er:YAG oscillator to generate 2 mJ, 40 ns laser pulses at 10 Hz, tunable in the range 3.6 – 5.1 μm [19]. Figure 1 depicts a simplified experimental setup. We focused the 4.3 μm beam to a full width at half maximum (FWHM) beam diameter of ~ 600 μm resulting in a peak intensity of ~ 15 MW/cm² at the entrance of the gas-filled cell. After propagating through the 10 cm long cell, the beam was split by a 50/50 beamsplitter to perform simultaneous temporal and spatial measurements. The exit of the cell was imaged and magnified by a factor of 5 using a telescope and measured with a pyroelectric camera. Frequency-resolved measurements of self-focusing and self-defocusing were performed using a spectrometer with the pyrocamera as a readout device. The resolution of the spectrometer is < 1 nm. It should be noted that the FWHM of the Fe:ZnSe laser bandwidth is < 2 nm and that the spacing between rovibrational lines of the 4P branch of the CO₂ molecule (see Fig. 1b) is ~ 4 nm. As a result, the laser spectrum only interacts with an individual line on the long-wavelength side of the CO₂ absorption spectrum (4P branch) where the line spacing is the greatest. The entire experimental set-up was purged with argon to avoid absorption related to CO₂ in the air.




Figure 1: (a) Simplified experimental setup used for measurements of resonant self-focusing and self-defocusing of $4.3 \mu\text{m}$ light in CO_2 gas and (b) absorption spectrum of CO_2 (100 torr, 295 K) obtained from the HITRAN database showing the rovibrational structure of the so-called 4R and 4P branches [13]. The inset is a simplified energy level diagram of CO_2 showing only the $4.3 \mu\text{m}$ $000 - 00n$ (v_3) vibrational asymmetric stretching mode.

III. Results

We have observed self-focusing, self-defocusing, and transverse beam break-up indicative of multiple filamentation after propagating the $4.3 \mu\text{m}$ laser pulses tuned to various spectral locations within the 4P branch of pure CO_2 . Figure 2, below, shows typical two-dimensional beam profiles that demonstrate these physical effects and were measured in ~ 50 torr of CO_2 . Figure 2a shows the initial beam profile, measured when the cell was evacuated. Note that for the self-focusing and self-defocusing cases (Fig. 2b and 2c, respectively), the laser was tuned in the vicinity of a rovibrational line on the long wavelength wing of the 4P branch. The beam profile showing transverse beam breakup (Fig. 2d) was observed for a case when the laser was tuned to the self-focusing side of a rovibrational line near the peak of the 4P branch. Typical absorption was measured to be between 30 – 70% depending on CO_2 pressure and laser wavelength resulting in an absorbed energy of 0.6 – 1.4 mJ.




Figure 2: (a) Initial $4.3 \mu\text{m}$ beam profile and beam profiles for a case when the laser was tuned to the (b) self-focusing and (c) self-defocusing sides of a rovibrational line located on the wing of the $000 - 001$ $4P$ branch of CO_2 . (d) Beam profile showing transverse beam break-up indicative of multiple filamentation for a case when the laser was tuned to the self-focusing side of a rovibrational line near the peak of the $4P$ branch.

To characterize the spatial changes to the beam caused by nonlinear focusing we have measured the evolution of the beam profile with variable pressure. Figure 3, below, is a plot of the FWHM beam diameter as a function of cell pressure for a case when the laser was tuned to the self-focusing and self-defocusing sides of resonance. As can be seen in Fig. 3, the FWHM beam diameter decreased from $\sim 600 \mu\text{m}$ to $200 \mu\text{m}$ for the self-focusing case and increased to $\sim 800 \mu\text{m}$ for the self-defocusing case when the pressure was varied from $0 - 100$ torr.




Figure 3: The evolution of the FWHM beam profile as a function of CO_2 pressure for a case when the laser was tuned to the self-focusing (blue squares) and self-defocusing (orange circles) sides of resonance. The dashed lines are quadratic fits to the data intended to guide the eye.

To determine the sign of the nonlinear refractive index on either side of resonance we performed frequency-resolved measurements of nonlinear focusing. Figure 4 shows the results of these measurements for a case when the laser spectrum was tuned around the 4P(28) (~ 4301 nm) line of the 000 – 001 transition of CO_2 . The pyrocamera images presented in Fig. 4 represent wavelength versus space plots. Here the location of the rovibrational line is denoted with a dashed white line. Note that the location of the line was identified by observing the attenuation of the blue wing of the laser's spectrum and is an estimate to within 1 nm of the location of the resonance at low laser intensity. Figure 4a depicts the unaffected beam detuned from the rovibrational line. As can be seen in Fig. 4b, the laser beam begins to focus as the rovibrational line is approached from the red side of resonance. Figure 4c shows the beam focusing and defocusing on either side of the resonance since the 4P(28) line is in the middle of the laser bandwidth. Finally, Fig. 4d shows that the beam is defocused on the blue side of the rovibrational line. It should be noted that the dumbbell shape visible on Fig. 4d is the cross-section of the donut-shaped beam that is produced by strong self-defocusing. This donut shape has also been observed in early experiments on self-defocusing in Rb vapor [3]. These measurements indicate that the sign of the nonlinearity around individual rovibrational lines of CO_2 is opposite to what has been observed in other atomic and molecular gases and that predicted by the two-level model developed by Javan and Kelley [14]. The potential origin of this sign reversal is discussed in detail below. Finally, it should be noted that the width of the rovibrational lines of CO_2 at 100 torr of pressure are ~ 0.5 Å and is beyond the resolution of the spectrometer used for these measurements.




Figure 4: Frequency-resolved measurements of self-focusing and self-defocusing in the vicinity of a rovibrational line of the 000 – 001 transition of CO_2 . (a) Shows the unaffected beam where the location of the line is marked by the vertical dashed line. (b) Shows the beam self-focusing as the rovibrational line is approached from the red side of resonance. (c) Shows the laser spectrum both focusing and defocusing on either side of resonance. (d) Shows the laser defocusing on the blue side of resonance. It should be noted that the dumbbell shape is the one-dimensional cross-section of the donut-shaped beam that is produced from self-defocusing and that the elongated beam profiles result from imaging the slit of the spectrometer.

By performing frequency-resolved measurements of nonlinear focusing for strong absorption lines, we have observed intensity dependent absorption that is consistent with power broadening. Indeed, we expect that the resonant ac-Stark effect should play a major role in this experiment since the $4.3\ \mu\text{m}$ ($000 - 001$) transitions of CO_2 have a relatively large dipole moment of $\sim 0.3\ \text{D}$ [20]. This effect causes the rovibrational line to broaden and, eventually, split as the applied field strength is increased, the degree of which scales in proportion to the Rabi frequency, $\Omega_R = 2\pi\mu E/h$, where μ is the dipole moment of the transition, E is the electric field strength, and h is Planck's constant. For our experimental conditions, we calculate a maximum frequency of $\sim 15\ \text{GHz}$ that is much larger than the $\sim 300\ \text{MHz}$ width of rovibrational lines of CO_2 at 100 torr pressure. Figure 5a shows a frequency-resolved self-focusing measurement where the power broadening effect manifests as an intensity ring. The radius of the ring results from a relative increase in absorption caused by the broadening of the rovibrational line that is largest on-axis where the intensity is maximum. Figure 5b shows the calculated CO_2 absorption spectrum in the vicinity of the 4P(30) ($\sim 4305\ \text{nm}$) line at 100 torr pressure and 295 K [13] where the dashed line depicts the anticipated degree of splitting based on the calculated Rabi frequency. It should be noted that the distance between rovibrational lines with a high rotational quantum number, J , is $\sim 60\ \text{GHz}$.




Figure 5: (a) Frequency-resolved self-focusing measurement showing apparent power broadening of the 4P(30) line of CO_2 . The field dependence of the Rabi frequency gives rise to a ring-shaped intensity distribution since the maximum shift is on-axis where the laser intensity is maximum. (b) The absorption spectrum of CO_2 in the vicinity of the 4P(30) and 4P(32) line from the HITRAN database with a depiction of the level splitting that may be achieved based on the calculated Rabi frequency.

Measurements of the self-focused or self-defocused temporal pulse profile indicate that the medium has a nonlinear response time $\leq 4\ \text{ns}$ FWHM. Figure 6 summarizes the results of these measurements and shows the spatial beam profile and temporal pulse profile of the initial beam (Fig. 6a and 6d) and the beam for cases when the laser was tuned to the self-focusing (Fig. 6b and 6e) and self-defocusing (Fig. 6c and 6f) sides of resonance. As can be inferred from Fig. 6d, the initial pulse, as measured by a fast HgCdTe detector, was comprised of a $\sim 4\ \text{ns}$ spike followed by a $\sim 30\ \text{ns}$ tail. When the laser was tuned to the self-focusing (Fig. 6b and 6e) side of resonance only the initial 4 ns

long spike was self-focused by the CO₂ cell and subsequently detected by the on-axis detector. When the laser was tuned on the self-defocusing side (Fig. 6c and 6f) of the resonance, the 4 ns long spike was defocused and the on-axis detector detected only the long tail.

Finally, we estimated the nonlinear refractive index of CO₂ near the 4P(28) and 4P(30) lines of the 000 – 001 band of CO₂. By using a Gaussian beam decomposition method [11, 22] we have fit our data using an analytic expression derived to calculate the change in beam radius of a Gaussian beam after it is subjected to a nonlinear phase shift. The value we have calculated for the nonlinear refractive index is $\sim 10^{-12}$ cm²/W for 100 torr of CO₂ corresponding to a critical power for Kerr self-focusing of ~ 10 kW. This value is consistent with our estimation of the critical power for self-focusing obtained by observing the transition from a single to multiple filamentation regime (see Fig. 2d). Using a concentration of 400 ppm of CO₂ in air, we estimate that the critical power for self-focusing of 4.3 μ m radiation in the atmosphere is approximately 10 MW, $\sim 10^4$ times less than the value calculated using measurements of the non-resonant nonlinear refractive index in the mid- and LWIR range, $\sim 10^{-19}$ cm²/W [15-18].




Figure 6: Beam profile measurements of (a) the initial beam, (b) the self-focused, and (c) self-defocused beam that correspond to temporal pulse profile measurements (d), (e), and (f), respectively. Note that pulse consisted of a 4 ns FWHM spike followed by a 30 ns tail and that only the initial spike was subjected to nonlinear focusing. Note that the vertical axes of (d), (e), and (f) are normalized to the peak amplitude of the initial laser pulse. These measurements were performed in the vicinity of the 4P(30) line where the absorption was $\sim 50\%$ for (b), (e) and $\sim 70\%$ for (c), (f).

IV. Discussion

The most interesting finding of this experimental study is the sign reversal of the nonlinearity of CO_2 as compared to that of a saturable two-level system (*i.e.* n_2 is negative at high-frequencies and positive at low-frequencies relative to the rovibrational line center). To the best of the authors' knowledge, this is the first report of a reversal in the dispersion of the Kerr nonlinearity in either atomic or molecular gases. To understand how this nonlinearity can manifest it is useful to review the sign of the nonlinearity of a saturable, two-level system [14]. Figure 7a and 7c show the imaginary and real part of a Lorentzian lineshape function that has decreased in amplitude due to saturation. For this system, the resonant nonlinearity results in self-focusing and self-defocusing on the high- and low-frequency side of resonance, respectively (see Fig. 7c). The CO_2 molecule, however, cannot be approximated as a saturable two-level system due to the existence of higher vibrational levels that are slightly detuned due to small anharmonicity and because of its rotational structure (see Fig 1b and the inset of Fig. 1). These differences make the $4.3 \mu\text{m}$ absorption channel difficult to saturate [22], particularly at the intensities used in this experiment. Indeed, it has been reported that the absorption of similar simple molecules such as SO_2 , OCS , NO_2 , and O_3 will not saturate even at intensities of $10^9 - 10^{11} \text{ W/cm}^2$ since power broadening can cause adjacent rovibrational lines to overlap resulting in absorption by a large number of quasi-resonant rovibrational transitions [23].




Figure 7: Imaginary and real parts of the susceptibility (χ) when modified by (a), (c) saturation and (b), (d) power broadening, respectively. Note that $\omega_L - \omega_0$ is the detuning where ω_L is the laser frequency and ω_0 is the resonant frequency. SF and SD denote regions of self-focusing and self-defocusing, respectively.

Since the 4.3 μm absorption channel is not saturable, other effects must be responsible for the observed optical nonlinearity. An increase in absorption as a function of intensity can certainly lead to the qualitative behavior that is observed in experiment and this may be possible since excitation of population from 000 – 001 opens an absorption channel from 001 - 002. It is unlikely that this factor is responsible for our observations, however, since we observe the same sign of nonlinearity over a wide range of rovibrational lines across both the 4P and 4R branch. Figure 8 shows calculated CO_2 absorption spectra for unexcited and excited CO_2 that correspond to vibrational temperatures (T_3) of 295 K ($T_3 = T$, Fig. 8a) and 2500 K ($T_3 \neq T$, Fig. 8b), respectively. Here we have only shown lines that correspond to the 000 – 001 (blue solid line) and 001 – 002 (red dashed line) transitions. As can be inferred from Fig. 8b, the 001 – 002 transitions only coincide with the 000 – 001 transitions for the 4P branch. It should be noted that vibrational temperatures of 2500 K have been measured for an optically pumped CO_2 laser under similar experimental conditions [22].




Figure 8: Calculated absorption spectra showing lines corresponding to the 000 – 001 and 001 – 002 transitions of the CO_2 molecule for a vibrational temperature of (a) 295 K and (b) 2500 K.

A more likely explanation for our experimental observations is that the nonlinearity is dominated by power broadening, as our calculations of the Rabi frequency and measurements of frequency-resolved self-focusing (see Fig. 5a) indicate that this effect plays a major role in the experiment. To explain the qualitative sign of the nonlinearity we have used a simplified model of the power broadening process where we have approximated the level-splitting by a summation of two Lorentzian lineshape functions that are separated by the Rabi frequency. Figure 7b and 7d show the imaginary and real part of this function, respectively, where the degree of level splitting is comparable to that expected for our experimental conditions. As can be seen from Fig. 7d, this level splitting may cause self-focusing on the low-frequency and self-defocusing on the high-frequency wing of the absorption line, consistent with what has been observed in experiment. Figure 7d also suggests that the nonlinearity changes sign within the bandwidth of the unperturbed resonant line. This fine structure was not observed in experiment, due to

limitations in this simplified model or for experimental reasons such as high absorption or insufficient resolution.

The time-domain data presented in Fig. 6, suggesting that the nonlinear response time is <4 ns, is consistent with the instantaneous response time that is expected from a field effect such as power broadening. A nonlinear response originating from excitation, on the other hand, should respond on a time scale related to the upper state lifetime, ~ 40 μ s for the 100 torr pressures used in our experiment [24, 25]. While this finding suggests that power broadening is the dominant mechanism in this experiment, the resonant nonlinear optics of simple molecules such as CO₂ are complicated by a combination of excitation and power broadening effects. A comparison between high-resolution spectral measurements and calculations using a self-consistent, density matrix approach is necessary to completely elucidate the physics.

V. Conclusions

We have used time- and frequency-resolved measurements of self-focusing and self-defocusing to characterize the resonant rovibrational nonlinear optical response of the CO₂ molecule at 4.3 μ m. In doing so, we have performed measurements of the qualitative dispersion of the resonant nonlinearity in the vicinity of individual rovibrational lines. When subjected to intense 4.3 μ m laser pulses, the CO₂ molecule exhibits a nonlinearity which is opposite in sign to that predicted by the two-level model and that observed in other atomic and molecular gases, thus underscoring the importance of performing such measurements for a number of simple molecules. The data suggest that, owing to the relatively large dipole moment of the asymmetric stretching mode of CO₂, power broadening and level-splitting dominates the nonlinear response. Future work will be dedicated to performing a detailed comparison between high-resolution measurements and theory to fully understand the physics of this resonant rovibrational nonlinearity. These measurements are directly applicable to the development of optically pumped molecular lasers and could prove useful for establishing a nonlinear refractive index equivalent of the HITRAN database, a task that is critical to support research on the propagation of high-power mid- and LWIR beams in the atmosphere.

Acknowledgements

This material is based upon work supported by the Office of Naval Research (ONR) MURI (4-442521-JC-22891) and the Department of Energy (DOE) Office of Science Accelerator Stewardship award DE-SC0018378.

References

- [1] S. L. McCall and E. L. Hahn, "Self-induced transparency by pulsed coherent light", *Physical Review Letters*, **18**, 21 (1967).
- [2] D. Grischkowsky, "Self-focusing of light by potassium vapor", *Physical Review Letters*, **24**, 16 (1970).

- [3] D. Grischkowsky and J. A. Armstrong, "Self-Defocusing of Light by Adiabatic Following in Rubidium Vapor", *Physical Review A*, **6**, 4 (1972).
- [4] K. -J. Boller, A. Imamoğlu, and S. E. Harris, "Observation of Electromagnetically Induced Transparency", *Physical Review Letters*, **66**, 20, (1991)
- [5] A. Kasapi, M. Jain, G. Y. Yin, and S. E. Harris, "Electromagnetically Induced Transparency: Propagation Dynamics", *Physical Review Letters*, **74**, 13 (1995).
- [6] M. D. Lukin and A. Imamoğlu, "Nonlinear Optics and Quantum Entanglement of Ultraslow Single Photons", *Physical Review Letters*, **84**, 7 (2000).
- [7] P. Bernard, P. Galarneau, and S. L. Chin, "Self-focusing of CO₂ laser pulses in low-pressure SF₆", *Optics Letters*, **6**, 3 (1981).
- [8] I. A. Al-Saidi, D. J. Biswas, C. A. Emschary, and R. G. Harrison, "Self focussing of CO₂ laser radiation in NH₃ gas", *Optics Communications*, **52**, 5 (1985).
- [9] B. K. Deka, R. S. Joshi, and M. A. Rob, "Self-Focusing and Defocusing of TEA CO₂ Laser Radiation in NH₃", *Applied Physics B*, **44**, 1-4 (1987).
- [10] Y. Beaudoin, P. Galarneau, A. Normandin, and S. L. Chin, "An Experimental Study of Self-Focusing and Self-Defocusing of a TEA CO₂ Laser Pulse in CDF₃", *Applied Physics B*, **42**, 225-231 (1987).
- [11] C. H. Oh and S. S. Lee, "Measurement of nonlinear refractive index coefficients of NH₃ gas for transversely excited atmospheric CO₂ laser lines", *Journal of Applied Physics* **65**, 1 (1989).
- [12] S. Tochitsky, E. Welch, M. Polyanskiy, I. Pogorelsky, P. Panagiotopoulos, M. Kolesik, E. M. Wright, S. W. Koch, J. V. Moloney, J. Pigeon and C. Joshi, "Megafilament in air formed by a self-guided terawatt long-wavelength infrared laser". *Nature Photonics*, **13**, 41-46 (2019).
- [13] I. E. Gordon, L. S. Rothman, C. Hill, *et. al.*, "The HITRAN2016 Molecular Spectroscopic Database", *Journal of Quantitative Spectroscopy and Radiative Transfer*, **203**, 3 – 69 (2017).
- [14] A. Javan and P. L. Kelley, "Possibility of Self-Focusing Due to Intensity Dependent Anomalous Dispersion", *IEEE Journal of Quantum Electronics*, **QE-2**, 9 (1966).
- [15] S. Zahedpour, J. K. Wahlstrand, and H. M. Milchberg, "Measurement of the nonlinear refractive index of air constituents at mid-infrared wavelengths", *Optics Letters*, **40**, 24 (2015).
- [16] J. J. Pigeon, S. Ya. Tochitsky, E. C. Welch, and C. Joshi "Measurements of the nonlinear refractive index of air, N₂, and O₂ at 10 μm using four-wave mixing", *Optics Letters*, **41**, 17 (2016).
- [17] J. J. Pigeon, S. Ya. Tochitsky, E. C. Welch, and C. Joshi, "Experimental study of the third-order nonlinearity of atomic and molecular gases using 10-μm laser pulses", *Physical Review A*, **97**, 043829 (2018).
- [18] S. Zahedpour, S. W. Hancock, and H. M. Milchberg, "Ultrashort infrared 2.5 – 11 μm pulses: spatiotemporal profiles and absolute nonlinear response of air constituents", *Optics Letters*, **44**, 4 (2019).
- [19] V. Fedorov, D. Martyshkin, K. Karki, S. Mirov, "Gain Switched and Q-Switched Fe:ZnSe Lasers tunable over 3.60-5.15 μm", in *Advanced Solid State Laser Conference*, Boston, MA, November 2018, paper AW3A.6.
- [20] R. K. Brimacombe, and J. Reid, "Influence of the dynamic Stark effect on the small-signal gain of optically pumped 4.3-μm CO₂ lasers", *Journal of Applied Physics*, **58**, 3,

(1985).

[21] D. Weaire, B. S. Wherrett, D. A. B. Miller, and S. D. Smith, "Effect of low-power nonlinear refraction on laser-beam propagation in InSb", *Optics Letters*, **4**, 19 (1979).

[22] R. C. Y. Auyeung, and J. Reid, "High Vibrational Temperatures in Optically-Pumped CO₂", *IEEE Journal of Quantum Electronics*, **24**, 3 (1988).

[23] A. V. Chugunov, M. S. Djidjoev, S. V. Ivanov, and V. Ya. Panchenko, "Nonlinear absorption of strong infrared radiation by ozone", *Optics Letters*, **10**, 12 (1985).

[24] F. Lepoutre, G. Louis, and H. Manceau, "Collisional relaxation in CO₂ between 180 K and 400 K measured by the spectrophone method", *Chemical Physics Letters*, **48**, 3 (1977).

[25] G. Inoue, and S. Tsuchiya, "Vibrational relaxation of CO₂(00⁰1) in CO₂, He, Ne, and Ar in the temperature range from 300~140K", *Journal of the Physical Society of Japan*, **38**, 3 (1975).