

This is the accepted manuscript made available via CHORUS. The article has been published as:

Near-threshold scaling of resonant inelastic collisions at ultralow temperatures

Rebekah Hermsmeier, Adrien Devolder, Paul Brumer, and Timur V. Tscherbul

Phys. Rev. A **105**, L011302 — Published 14 January 2022

DOI: [10.1103/PhysRevA.105.L011302](https://doi.org/10.1103/PhysRevA.105.L011302)

Near-threshold scaling of resonant inelastic collisions at ultralow temperatures

Rebekah Hermsmeier¹, Adrien Devolder², Paul Brumer², and Timur V. Tscherbul¹

¹*Department of Physics, University of Nevada, Reno, NV, 89557, USA*

²*Chemical Physics Theory Group, Department of Chemistry,
and Center for Quantum Information and Quantum Control,
University of Toronto, Toronto, Ontario, M5S 3H6, Canada*

(Dated: December 3, 2021)

We show that the cross sections for a broad range of resonant *inelastic* processes accompanied by excitation exchange (such as spin-exchange, Förster resonant, or angular momentum exchange) exhibit a near-threshold scaling $E^{\Delta m_{12}}$, where E is the collision energy, $\Delta m_{12} = m'_1 + m'_2 - m_1 - m_2$, and m_i and m'_i are the initial and final angular momentum projections of the colliding species ($i = 1, 2$). In particular, the inelastic cross sections for $\Delta m_{12} = 0$ transitions display an unconventional E^0 scaling similar to that of elastic cross sections, and their rates vanish as $T^{\Delta m_{12}+1/2}$. For collisions dominated by even partial waves (such as those of identical bosons in the same internal state) the scaling is modified to $\sigma_{\text{inel}} \propto E^{\Delta m_{12}+1}$ if Δm_{12} is odd. We present accurate quantum scattering calculations that illustrate these modified threshold laws for resonant spin exchange in ultracold Rb + Rb and O₂ + O₂ collisions. Our results illustrate that the well-known k^{-1} threshold scaling of inelastic cross sections only applies to exothermic, rather than resonant, inelastic processes.

I. INTRODUCTION

The unique controllability of ultracold atomic and molecular collisions [1–10] gives rise to numerous applications of ultracold quantum gases in quantum information science and on precision tests of foundations of chemical and statistical physics [5, 11]. Further, much progress has been achieved over the last few years in our ability to observe the reactants and products of molecular collisions and chemical reactions in single, well-controlled quantum states [6–9, 11–13].

Ultracold atoms and molecules are typically prepared and trapped in quantum states with a fixed value of the angular momentum projection m on a space-fixed quantization axis (typically fixed by an external electromagnetic field). Subsequent angular momentum projection-changing collisions play a key role in ultracold chemistry. If such collisions are exothermic, they convert the intramolecular energy into translational energy of the relative motion, leading to undesirable trap loss and heating. In contrast, resonant excitation exchange (EE) collisions conserve the internal energy but lead to a coherent transfer of excitations (such as spin polarization) from one particle to another. Examples of resonant EE processes include spin-exchange in atomic, molecular, and ionic collisions [14–25], Förster resonances in collisions of Rydberg atoms [26–31], atom-dimer exchange chemical reactions [20, 32], excitation exchange between identical atoms or molecules [33], charge transfer in cold ion-atom collisions [34–36], and rotational angular momentum projection-changing collisions [10]. [In this Letter, we resolve a long-standing controversy regarding the threshold behavior of EE cross sections.](#)

Spin-exchange collisions, in particular, play an important role in a wide array of research fields, ranging from spin-exchange optical pumping [14–17], cold chemistry [20, 21, 32], precision magnetometry [18], and astrophysics [37–41] to quantum many-body physics [22–25] and quantum information processing, where they are

used to generate entangled states [42, 43] and to operate quantum logic gates [44–48]. Recent experiments observed spin-exchange collisions of ultracold alkali-metal atoms [49, 50], Förster resonant energy transfer in Rydberg atom collisions [28–30], and electric field-induced resonances in collisions of rotationally excited KRb molecules [7]. We have recently found that ultracold EE collisions are amenable to extensive coherent control [10] due to the suppression of m_{12} -changing collisions at ultralow temperatures.

However, despite the ever-increasing interest in ultracold EE collisions, there remains a considerable uncertainty about the scaling of their integral cross sections (ICS) near collision thresholds [51, 52]. It is commonly believed that the inelastic ICSs scale as k^{-1} with the incident collision wavevector k in the limit $k \rightarrow 0$ [53, 54]. While some computational studies did observe the k^{-1} scaling [16, 21, 39] others reported that the ICS for atomic EE collisions approach a constant value [35, 38, 40, 41], an unexpected observation that has remained unexplained. Several authors attributed the k^0 scaling to a deficiency of the degenerate internal state (DIS) (or elastic) approximation [55–57] [widely used to model EE processes](#) [32, 35, 38, 40, 41, 55–69], [which](#) neglects the internal structure of colliding atoms [19, 38, 39]. The exact reason for this deficiency has remained unclear. More recently, the constant near-threshold scaling of spin-exchange cross sections was claimed to be incorrect [39], further deepening the existing controversy surrounding the threshold behavior of EE collisions.

Krems and Dalgarno presented a derivation of the threshold laws for m -changing collisions of an open-shell atom (or rotating molecule) with a spherically symmetric particle [70] in the limit $k \rightarrow 0$. [They showed that the elastic ICSs exhibit a \$k^0\$ scaling, whereas the inelastic ICSs for \$m\$ -changing collisions vanish as \$k^{2\Delta m}\$ when \$\Delta m\$ is even and as \$k^{2\(\Delta m+1\)}\$ when \$\Delta m\$ is odd](#) [70]. However, their analysis cannot be applied to describe

EE processes where both species exchange angular momenta. As such, a comprehensive understanding of the threshold behavior of EE processes is still lacking, limiting our ability to control a wide range of ultracold EE collisions that are key to implementing quantum logic gates [44–48] and to generating entanglement [42, 43] and strong correlations [22–25] in quantum many-body systems.

As noted above, here we resolve the long-standing controversy regarding the threshold behavior of EE cross sections. By deriving the threshold laws for ultracold EE collisions of two species *both of which possess internal angular momenta*, we show that the ICS for these processes display a collision energy scaling $\sigma_{\text{EE}} \propto E^{|\Delta m_{12}|}$. Here, $\Delta m_{12} = m'_1 + m'_2 - m_1 - m_2$, and m_i and m'_i are the initial and final angular momentum projections of the colliding species ($i = 1, 2$). In the important particular case $\Delta m_{12} = 0$, the *s*-wave *inelastic* ICS approaches a constant value and the corresponding rate vanishes as $T^{1/2}$ as typically expected of elastic collisions [54]. These results provide the first conclusive explanation for the unconventional k^0 near-threshold scaling of the resonant EE processes obtained in the widely used DIS approximation [32, 35, 38, 40, 41, 55–57] and for the heretofore unexplained change in the threshold scaling of spin exchange cross sections from k^{-1} to k^0 caused by this approximation [38].

To illustrate our findings, we present numerically exact quantum scattering calculations of ultracold $\text{O}_2 + \text{O}_2$ and $\text{Rb} + \text{Rb}$ collisions, which confirm our analytic results and provide a blueprint for observing the unconventional threshold scaling of inelastic cross sections in ultracold atomic and molecular collision experiments. (For clarification, note that inelastic refers to any collision where the nature or internal state of a colliding partner changes [71, 72]).

Theory. Consider a binary collision of two atoms and/or molecules, each possessing *total* internal angular momenta $\mathbf{j}_1$ and $\mathbf{j}_2$. The nature of the operators $\mathbf{j}_i$ is set by the details of the internal structure of the colliding species, which is immaterial for the discussion below. We focus on the case of weak external fields, where j_i remain good quantum numbers, and consider as numerical examples, collisions of two open-shell $^{17}\text{O}_2(^3\Sigma)$ molecules ($\mathbf{j} = \mathbf{S} + \mathbf{N}$, where $\mathbf{S}$ is the electron spin and $\mathbf{N}$ is the rotational angular momentum) and of two ^{87}Rb atoms ($\mathbf{j} = \mathbf{S} + \mathbf{I}$, where $\mathbf{I}$ is the atomic nuclear spin).

The starting point for our discussion is the expression for the integral cross section for two particles initially colliding in well-defined angular momentum eigenstates $|j_i m_i\rangle$ (below, the subscripts j_1 and j_2 will be omitted for brevity unless stated otherwise)

$$\sigma_{m_1 m_2 \rightarrow m'_1 m'_2} = \frac{\pi}{k^2} \sum_{l, m_l} \sum_{l', m'_l} |T_{m_1 m_2, l m_l \rightarrow m'_1 m'_2, l' m'_l}|^2 \quad (1)$$

where $|l m_l\rangle$ and $|l' m'_l\rangle$ are the eigenstates of the orbital angular momentum $\hat{L}^2$ of the collision complex and its

projection $\hat{L}_z$ on the space-fixed quantization axis. We are interested in the threshold scaling of the ICS with the collision energy E , which is determined by that of the transition T -matrix elements $T_{m_1 m_2, l m_l \rightarrow m'_1 m'_2, l' m'_l}$. To make the threshold scaling more explicit, we rewrite the T -matrix in the total angular momentum representation [73, 74]

$$\begin{aligned} T_{j_1 m_1 j_2 m_2, l m_l \rightarrow j'_1 m'_1 j'_2 m'_2, l' m'_l} &= \sum_{J, M} \sum_{j_{12}, m_{12}} \sum_{j'_{12}, m'_{12}} (2J+1) \\ &\sqrt{(2j_{12}+1)(2j'_{12}+1)} (-1)^{l'+l} T_{j_1 j_2 j_{12} l \rightarrow j'_1 j'_2 j'_{12} l'}^J \\ &(-1)^{j_{12}+j'_{12}+m_{12}+m'_{12}} \begin{pmatrix} j_1 & j_2 & j_{12} \\ m_1 & m_2 & -m_{12} \end{pmatrix} \begin{pmatrix} j_{12} & l & J \\ m_{12} & m_l & -M \end{pmatrix} \\ &(-1)^{j_1+j'_1-j_2-j'_2} \begin{pmatrix} j'_1 & j'_2 & j'_{12} \\ m'_1 & m'_2 & -m'_{12} \end{pmatrix} \begin{pmatrix} j'_{12} & l' & J \\ m'_{12} & m'_l & -M \end{pmatrix} \quad (2) \end{aligned}$$

where $j_{12} = |\mathbf{j}_{12}| = |\mathbf{j}_1 + \mathbf{j}_2|$ is the total internal angular momentum, and $J = |\mathbf{J}| = |\mathbf{j}_1 + \mathbf{j}_2 + \mathbf{l}|$ is the total angular momentum of the collision partners, which is conserved in the absence of external fields.

The advantages of using Eq. (2) are two-fold. First, the threshold behavior of the T -matrix elements $T_{j_1 j_2 j_{12} l \rightarrow j'_1 j'_2 j'_{12} l'}^J$ only depends on l and l' through the Wigner threshold law [51, 52]

$$T_{\gamma l, \gamma' l'} \propto k^{l+1/2} (k')^{l'+1/2}, \quad (3)$$

where k and k' are the incident and final wavevectors [51, 52, 75] and γ stands for all quantum numbers other than l , i.e the internal quantum numbers j_1, j_2, j_{12} or quantum numbers describing other simultaneous resonant processes. Note that Eq. (3) assumes the absence of near-threshold resonance, bound, and virtual states. A different k scaling would result if such states were present [52, 75, 76]. In addition, we assume that the interactions between the collision partners are short-ranged and neglect the modification of the threshold laws by long-range interactions [52]. In the case of resonant scattering in the absence of external fields considered below, the initial and final states are degenerate ($k = k'$) and the near-threshold dependence of T -matrix elements takes the form

$$T_{\gamma l, \gamma' l'} \propto k^{l+l'+1}. \quad (4)$$

In the limit of zero collision energy ($k \rightarrow 0$) it follows from Eq. (4) that the T -matrix elements with the lowest l and l' provide the dominant contributions to the sum in Eq. (2) and hence to the ICS (1).

Second, the $3j$ -symbols ($:::$) in Eq. (2) make explicit the rotational symmetry restrictions on the possible values of l, m_l, l' , and m'_l . The $3-j$ symbols must satisfy the selection rule $m_{12} + m_l = M = m'_{12} + m'_l$, $m_1 + m_2 = m_{12}$, and $m'_1 + m'_2 = m'_{12}$, which implies conservation of the total angular momentum projection, $M = m_1 + m_2 + m_l = m'_1 + m'_2 + m'_l$. The values of m_l

and m'_l are then restricted by the change of the internal projection $\Delta m_{12} = m_{12} - m'_{12} = m'_l - m_l$. To illustrate the restrictions on l and l' , consider the s -wave scattering ($l = m_l = 0$) of two distinguishable particles that changes $m_{12} = m_1 + m_2$ by 1. As stated above, since M is conserved, we have $\Delta m_{12} = m_{12} - m'_{12} = m'_l - m_l = \pm 1$ and thus $m'_l = \pm 1$. In the absence of additional symmetry restrictions on the lowest possible values of l or l' in Eq. (1), the dominant partial wave contributions are $l = 0$ and $l' = |\Delta m_{12}| = |\Delta m_{12}|$. Substituting these values into Eq. (4) we find $T_{\gamma l, \gamma' l'} \propto k^{|\Delta m_{12}|+1}$. If the parity of molecular (atomic) states does not change, the conservation of the total inversion parity p implies that l and l' must have the same parity. This is, for example, the case of spin-exchange transitions, which only change the projections of the internal angular momenta ($j_1 = j'_1$ and $j_2 = j'_2$). For ultracold collisions $l = 0$, and thus l' must be even. The relation $l' = |\Delta m_{12}|$ still holds for even $|\Delta m_{12}|$ while for odd values of $|\Delta m_{12}|$, it must be replaced by $l' = |\Delta m_{12}| + 1$.

For collisions of identical bosons (fermions) in the same internal state, the permutation symmetry must also be taken into account. Only even (odd) partial waves will be present in Eq. (1) [16, 77]. For ultracold collisions of identical bosons $l = 0$, and all permissible values of l' are even as the case of j_i -conserving spin-exchange transitions. For ultracold collisions of identical fermions, $l = 1$ and all permissible values of l' are odd. The relation $l' = |\Delta m_{12}|$ holds for odd $|\Delta m_{12}|$ while for even values of $|\Delta m_{12}|$, it must be replaced by $l' = |\Delta m_{12}| - 1$. Again, we find the same near-threshold scaling $T_{\gamma l, \gamma' l'} \propto k^{|\Delta m_{12}|+1}$ for even $|\Delta m_{12}|$ and $T_{\gamma l, \gamma' l'} \propto k^{|\Delta m_{12}|+2}$ for odd $|\Delta m_{12}|$. Note that $|\Delta m_{12}| = 0$ is a special case: the lowest value of l' is 1 and $T_{\gamma l, \gamma' l'} \propto k^3$. Thus, ultracold inelastic collisions of identical fermions with $|\Delta m_{12}| = 0, 1$ and 2 all follow a near-threshold scaling identical to that of p -wave elastic scattering, i.e., $T_{\gamma l, \gamma' l'} \propto k^3$.

Summarizing the above discussion, the threshold behavior of the T -matrix elements in the absence of l -restricting symmetries is given by $T_{\gamma l, \gamma' l'} \propto k^{|\Delta m_{12}|+1}$. By taking the absolute magnitude squared of the T -matrix elements and dividing by $k^2 = 2\mu E$, where μ is the reduced mass of the collision complex [see Eq. (1)], we obtain the threshold scaling of the ICS for collision-induced EE processes in the general case

$$\sigma_{j_1 m_1 j_2 m_2 \rightarrow j'_1 m'_1 j'_2 m'_2} \simeq E^{|\Delta m_{12}|}. \quad (5)$$

A modified threshold scaling applies in the case of even partial wave scattering (spin-exchange transitions with $j_1 = j'_1$ and $j_2 = j'_2$ or those between identical bosons in the same internal states)

$$\begin{aligned} \sigma_{j_1 m_1 j_2 m_2 \rightarrow j'_1 m'_1 j'_2 m'_2} &\simeq E^{|\Delta m_{12}|} \quad (|\Delta m_{12}| \text{ even}), \\ &\simeq E^{|\Delta m_{12}|+1} \quad (|\Delta m_{12}| \text{ odd}). \end{aligned} \quad (6)$$

Significantly, unlike in the case of collisions with a structureless target [70], m_i can be nonzero for *both* of the

colliding species, and thus $|\Delta m_{12}| = 0$ can correspond to inelastic as well as to elastic scattering. Therefore, remarkably, these expressions show that the threshold scaling of the ICS for an inelastic EE process that conserves m_{12} (such as flip-flop spin-exchange collisions) approach a constant value in the limit of ultralow collision energies. This behavior stands in contrast with the expected $E^{-1/2}$ scaling of the inelastic ICSs in the limit $E \rightarrow 0$ [4, 54], and was reported without explanation in several previous quantum scattering calculations of near-resonant charge exchange [35] and spin-exchange in cold H + H collisions [38]. Note that m_{12} in Eqs. (5) and (6) is the projection of the *two-particle* angular momentum $\mathbf{j}_{12}$ in the incident collision channel. In contrast, in the work of Krems and Dalgarno [70], m refers to a *single-particle* angular momentum projection. The inclusion of the internal structure of the second collision partner is an essentially new aspect of this work, which leads to a qualitatively different behavior compared to the case of structureless atom-molecule collisions considered before [70]. In particular, $\Delta m = 0$ implies that a collision is elastic [70], whereas $\Delta m_{12} = 0$ does not.

The physical origin of the unconventional Wigner threshold scaling can be traced back to the threshold behavior of the T -matrix elements, which depends *only on the initial and final values of collision wavevectors k and k' and on l and l'* (see above and Eq. (3) [51]). As a result, as long as a collision process remains resonant ($k = k'$), the threshold dependence will be governed by Δm_{12} , regardless of whether the process is elastic or inelastic. Thus, as shown above, both the elastic and inelastic collisions that conserve k and m_{12} have the same constant near-threshold scaling, usually characteristic of the elastic ICS. This explains why the DIS approximation, which assumes $k = k'$ for all the incident and final channels [55–57] gives a constant threshold scaling.

When the internal structure is included, both exothermic (change of j_i) and resonant (only change of projections) processes must be considered. For the exothermic processes, the discussion in the above paragraphs can be straightforwardly extended. Assuming that k' is outside of the threshold regime, the dependence of the T -matrix on k' is negligibly weak, and $T_{\gamma l, \gamma' l'} \propto k^{l+1/2}$. As a result, the relaxation ICSs assume their conventional k^{-1} scaling with relaxation rates approaching constant values in the $T \rightarrow 0$ limit. Then, the theory and rigorous CC calculations give a k^{-1} scaling for the exothermic processes, while for the resonant processes (see below for the CC calculations), they give the scaling predicted above. If one sums these two contributions, only the k^{-1} scaling is observed [38], solving the above-mentioned controversy. Our work also explains why the averaging of the resonant processes over the initial and final values of m_1 and m_2 gives a k^0 scaling [39]. In the limit $E \rightarrow 0$, the m_{12} -changing ICSs vanish, so only the m_{12} -conserving contributions, which scale as k^0 , remain.

Numerical examples: Spin exchange in ultracold $O_2 + O_2$ and $Rb + Rb$ collisions. To verify the unconventional Wigner threshold scaling of inelastic EE




FIG. 1. (a) State-to-state inelastic ICSs for ultracold $\text{O}_2 + \text{O}_2$ collisions plotted as a function of collision energy. The initial and final values of m_i for the O_2 molecules are indicated in the legend as $m_1 m_2$ to $m'_1 m'_2$. The fits as E^0 and E^2 are represented by dashed lines. (b) Same as panel (a) but for the temperature dependence of state-to-state inelastic collision rates $K_{m_1 m_2 \rightarrow m'_1 m'_2}$. The fits as $T^{1/2}$ and $T^{5/2}$ are represented by dashed lines.

processes derived above, we carried out numerically exact quantum scattering calculations of spin exchange in ultracold $\text{O}_2 + \text{O}_2$ and $\text{Rb} + \text{Rb}$ collisions.

We first consider ultracold collisions of identical $\text{O}_2(X^3\Sigma^-)$ molecules initially in their ground rovibrational states $|N_i = 0, j_i = 1, m_i\rangle$, where $\mathbf{j}_i = \mathbf{N}_i + \mathbf{S}_i$ ($i = 1, 2$) and m_i is the eigenvalue of j_{iz} . Since all $j_i > 1$ states are energetically unavailable at ultralow collision energies, the initial and final two-molecule states can be denoted $|m_1 m_2\rangle$ with $m_i = -1, 0, 1$ (here, we neglect the molecular hyperfine structure for simplicity).

Figure 1(a) shows the ICS for spin exchange in $^{17}\text{O}_2 + ^{17}\text{O}_2$ collisions calculated as a function of collision energy as described in the Supplemental Material [78]. The ICS for the $|\Delta m_{12}|$ conserving $|-1, +1\rangle \rightarrow |00\rangle$ transition is seen to approach a constant value in the limit $E \rightarrow 0$ as predicted by Eq. (6). The corresponding inelastic rate tends to zero as $T^{1/2}$ as illustrated in Fig. 1(b). Note that for $j = 1$, the only transition with $|\Delta m_{12}| = 0$ is between the states $|0, 0\rangle$ and $|-1, +1\rangle$.

In Fig. 1(a) we plot the ICS for $|\Delta m_{12}|$ changing transitions for two identical bosons in the same internal state ($|0, 0\rangle \rightarrow |+1, +1\rangle$ and $|0, 0\rangle \rightarrow |0, +1\rangle$), and for two identical bosons in different internal states ($|-1, +1\rangle \rightarrow |0, +1\rangle$). As follows from Eq. (6), these




FIG. 2. (a) State-to-state inelastic ICSs for ultracold s -wave $^{87}\text{Rb} + ^{87}\text{Rb}$ collisions plotted vs. collision energy at a magnetic field of 0.01 G. The initial and final hyperfine states of the Rb atoms $F_i m_{F_i}$ are indicated in the legend. The fits as E^0 are represented by dashed lines. (b) Same as panel (a) but for the state-to-state inelastic collision rates $K_{m_{F_1} m_{F_2} \rightarrow m'_{F_1} m'_{F_2}}$ as a function of temperature. The fits as $T^{1/2}$ are represented by dashed lines.

three transitions must have the same E^2 scaling, as is indeed observed. The corresponding inelastic rate tends to zero as $T^{5/2}$ as illustrated by the fit in Fig. 1(b).

We next consider ultracold spin-exchange collisions of ^{87}Rb atoms, which have been the subject of much experimental study (for a review, see [2]). The two-atom threshold states $|j_1 m_1\rangle |j_2 m_2\rangle = |F_1 m_{F_1}\rangle |F_2 m_{F_2}\rangle$, where $|j_i m_{F_i}\rangle = |F_i m_{F_i}\rangle$ are the atomic hyperfine states and $\mathbf{j}_i = \mathbf{S}_i + \mathbf{I}_i$ are the total atomic angular momenta, which are vector sums of the electron and nuclear spins of the i -th atom ($i = 1, 2$). The quantum scattering problem for $^{87}\text{Rb} + ^{87}\text{Rb}$ is solved using the standard CC approach as described in, e.g., Ref. [83] by integrating the CC equations on a grid of R values from 2 to $400a_0$ with a step size of $5 \times 10^{-3}a_0$. The interaction potentials for the singlet and triplet states of Rb_2 with long-range scaling $\propto \frac{C_6}{R^6}$ give good agreement with the experimentally measured positions of Feshbach resonances for ^{87}Rb - ^{85}Rb [83].

The s -wave ICSs for $^{87}\text{Rb} + ^{87}\text{Rb}$ spin-exchange collisions are plotted as a function of collision energy in Fig. 2(a) for three representative hyperfine transitions $|10\rangle|10\rangle \rightarrow |1, -1\rangle|1, +1\rangle$, $|20\rangle|20\rangle \rightarrow |2, -1\rangle|2, +1\rangle$, and $|20\rangle|20\rangle \rightarrow |2, -2\rangle|2, +2\rangle$. As all of these transitions have $\Delta m_{12} = 0$, Eq. (6) establishes that their ICSs

should scale as E^0 in the limit of zero collision energy, and their rates should decrease as $T^{1/2}$ as $T \rightarrow 0$. The predicted trends are clearly observable in Fig. 2. Interestingly, the cross sections for the spin-exchange transition $|20\rangle|20\rangle \rightarrow |2-2\rangle|2-2\rangle$ is strongly suppressed compared to $|20\rangle|20\rangle \rightarrow |2-1\rangle|21\rangle$. This is caused by the vanishing matrix element of the spin-dependent Rb-Rb interaction potential $\langle 20|\langle 20|\hat{V}(R)|2-2\rangle|22\rangle$ (in contrast, $\langle 20|\langle 20|\hat{V}(R)|2-1\rangle|2+1\rangle \neq 0$), making the spin-exchange transition $|20\rangle|20\rangle \rightarrow |2-2\rangle|2+2\rangle$ forbidden in first order [78]. The minimum near 100 μK in the collision dependence of the $|20\rangle|20\rangle \rightarrow |2-2\rangle|22\rangle$ ICS shown in Fig. 2(a) is due to the Ramsauer-Townsend effect [84, 85], which occurs due to the scattering phase shift approaching π (caused by a near-threshold resonance) shifting the onset of the threshold regime to lower collision energies.

In conclusion, we have shown that the near-threshold scaling of the ICS for resonant inelastic EE processes (such as spin exchange) is given by $\sigma_{\text{inel}} \propto E^{\Delta m_{12}}$, which only depends on the difference between the combined angular momentum projections Δm_{12} in the incident and final collision channels. For $\Delta m_{12} = 0$ the scaling of the inelastic ICS is the same as that of the elastic ICS, i.e., $\sigma_{\text{inel}} \propto E^0$. This work resolves the uncertainty con-

cerning the threshold behavior of EE cross sections when the internal angular momentum projection of both collision partners can change. More generally, it suggests that the k^{-1} scaling of inelastic transitions is limited to exothermic processes, and must be changed to a Δm_{12} dependent scaling for resonant processes.

Our results demonstrate a universal $T^{\Delta m_{12}+1/2}$ suppression of a wide class of resonant EE processes at ultralow temperatures. This suppression and the k^0 scaling of some inelastic transitions could be observed experimentally for, e.g., spin-exchange atom-atom collisions in optical tweezers [49, 86], atom-ion collisions in an optical lattice setup, which allows for high collision energy resolution [87], as well as in cold and ultracold collisions of Yb atoms [88], Ti atoms [89], Rydberg atoms [28, 29] and polar molecules [7, 11]. Our work also reveals a subtle deficiency of the DIS approximation [55–57], which reproduces the correct threshold scaling of EE cross sections only when the initial and final thresholds (including the internal structure of the collision partners) are exactly degenerate.

We are grateful to Bernard Zygelman for bringing Ref. [38] to our attention. This work was supported by the U.S. Air Force Office for Scientific Research (AFOSR) under Contract No. FA9550-19-1-0312 and partially by the NSF (Grant No. PHY-1912668).

-
- [1] R. V. Krems, *Molecules in Electromagnetic Fields* (Wiley-VCH, 2019).
 - [2] C. Chin, R. Grimm, P. Julienne, and E. Tiesinga, Feshbach resonances in ultracold gases, *Rev. Mod. Phys.* **82**, 1225 (2010).
 - [3] L. D. Carr, D. DeMille, R. V. Krems, and J. Ye, Cold and ultracold molecules: science, technology and applications, *New J. Phys.* **11**, 055049 (2009).
 - [4] N. Balakrishnan, Perspective: Ultracold molecules and the dawn of cold controlled chemistry, *J. Chem. Phys.* **145**, 150901 (2016).
 - [5] J. L. Bohn, A. M. Rey, and J. Ye, Cold molecules: Progress in quantum engineering of chemistry and quantum matter, *Science* **357**, 1002 (2017).
 - [6] J. Wolf, M. Deiß, A. Krüchow, E. Tiemann, B. P. Ruzic, Y. Wang, J. D’Incao, P. S. Julienne, and J. H. Denschlag, State-to-state chemistry for three-body recombination in an ultracold rubidium gas, *Science* **358**, 921 (2017).
 - [7] K. Matsuda, L. De Marco, J.-R. Li, W. G. Tobias, G. Valtolina, G. Quémener, and J. Ye, Resonant collisional shielding of reactive molecules using electric fields, *Science* **370**, 1324 (2020).
 - [8] Z. Z. Yan, J. W. Park, Y. Ni, H. Loh, S. Will, T. Karman, and M. Zwierlein, Resonant dipolar collisions of ultracold molecules induced by microwave dressing, *Phys. Rev. Lett.* **125**, 063401 (2020).
 - [9] Y. Liu, M.-G. Hu, M. A. Nichols, D. D. Grimes, T. Karman, H. Guo, and K.-K. Ni, Photo-excitation of long-lived transient intermediates in ultracold reactions, *Nat. Phys.* **16**, 1132 (2020).
 - [10] A. Devolder, P. Brumer, and T. V. Tscherbul, Complete quantum coherent control of ultracold molecular collisions, *Phys. Rev. Lett.* **126**, 153403 (2021).
 - [11] Y. Liu, M.-G. Hu, M. A. Nichols, D. Yang, D. Xie, H. Guo, and K.-K. Ni, Precision test of statistical dynamics with state-to-state ultracold chemistry, *arXiv:2012.15842* (2020).
 - [12] Y. Segev, M. Pitzer, M. Karpov, N. Akerman, J. Narevicius, and E. Narevicius, Collisions between cold molecules in a superconducting magnetic trap, *Nature* **572**, 189 (2019).
 - [13] L. W. Cheuk, L. Anderegg, Y. Bao, S. Burchesky, S. S. Yu, W. Ketterle, K.-K. Ni, and J. M. Doyle, Observation of collisions between two ultracold ground-state CaF molecules, *Phys. Rev. Lett.* **125**, 043401 (2020).
 - [14] T. G. Walker and W. Happer, Spin-exchange optical pumping of noble-gas nuclei, *Rev. Mod. Phys.* **69**, 629 (1997).
 - [15] T. G. Walker, Fundamentals of spin-exchange optical pumping, *J. Phys. C* **294**, 012001 (2011).
 - [16] T. V. Tscherbul, P. Zhang, H. R. Sadeghpour, and A. Dalgarno, Collision-induced spin exchange of alkali-metal atoms with ^3He : An ab initio study, *Phys. Rev. A* **79**, 062707 (2009).
 - [17] T. V. Tscherbul, P. Zhang, H. R. Sadeghpour, and A. Dalgarno, Anisotropic hyperfine interactions limit the efficiency of spin-exchange optical pumping of ^3He nuclei, *Phys. Rev. Lett.* **107**, 023204 (2011).
 - [18] I. M. Savukov and M. V. Romalis, Effects of spin-exchange collisions in a high-density alkali-metal vapor in low magnetic fields, *Phys. Rev. A* **71**, 023405 (2005).
 - [19] T. Sikorsky, M. Morita, Z. Meir, A. A. Buchachenko, R. Ben-shlomi, N. Akerman, E. Narevicius, T. V.

- Tscherbul, and R. Ozeri, Phase locking between different partial waves in atom-ion spin-exchange collisions, *Phys. Rev. Lett.* **121**, 173402 (2018).
- [20] J. Rui, H. Yang, L. Liu, D.-C. Zhang, Y.-X. Liu, J. Nan, Y.-A. Chen, B. Zhao, and J.-W. Pan, Controlled state-to-state atom-exchange reaction in an ultracold atom-dimer mixture, *Nat. Phys.* **13**, 699 (2017).
- [21] Y.-X. Liu, J. Nan, D.-C. Zhang, L. Liu, H. Yang, J. Rui, B. Zhao, and J.-W. Pan, Observation of a threshold behavior in an ultracold endothermic atom-exchange process involving Feshbach molecules, *Phys. Rev. A* **100**, 032706 (2019).
- [22] Y. Kawaguchi and M. Ueda, Spinor Bose-Einstein condensates, *Phys. Rep.* **520**, 253 (2012).
- [23] D. M. Stamper-Kurn and M. Ueda, Spinor Bose gases: Symmetries, magnetism, and quantum dynamics, *Rev. Mod. Phys.* **85**, 1191 (2013).
- [24] Y. Nishida, SU(3) orbital Kondo effect with ultracold atoms, *Phys. Rev. Lett.* **111**, 135301 (2013).
- [25] J. Bauer, C. Salomon, and E. Demler, Realizing a Kondo-correlated state with ultracold atoms, *Phys. Rev. Lett.* **111**, 215304 (2013).
- [26] K. A. Safinya, J. F. Delpech, F. Gounand, W. Sandner, and T. F. Gallagher, Resonant Rydberg-Atom-Rydberg-Atom collisions, *Phys. Rev. Lett.* **47**, 405 (1981).
- [27] R. A. D. S. Zanon, K. M. F. Magalhaes, A. L. de Oliveira, and L. G. Marcassa, Time-resolved study of energy-transfer collisions in a sample of cold rubidium atoms, *Phys. Rev. A* **65**, 023405 (2002).
- [28] I. I. Ryabtsev, D. B. Tretyakov, I. I. Beterov, and V. M. Entin, Observation of the Stark-tuned Förster resonance between two Rydberg atoms, *Phys. Rev. Lett.* **104**, 073003 (2010).
- [29] J. Nipper, J. B. Balewski, A. Krupp, B. Butscher, R. Low, and T. Pfau, Highly resolved measurements of Stark-tuned forster resonances between Rydberg atoms, *Phys. Rev. Lett.* **108**, 113001 (2012).
- [30] S. Ravets, H. Labuhn, D. Barredo, L. Béguin, T. Lahaye, and A. Browaeys, Coherent dipole-dipole coupling between two single Rydberg atoms at an electrically-tuned Förster resonance, *Nat. Phys.* **10**, 914 (2014).
- [31] A. L. Win, W. D. Williams, T. J. Carroll, and C. I. Sukenik, Catalysis of Stark-tuned interactions between ultracold Rydberg atoms, *Phys. Rev. Lett.* **108**, 032703 (2012).
- [32] T. Sikorsky, Z. Meir, R. Ben-shlomi, N. Akerman, and R. Ozeri, Spin-controlled atom-ion chemistry, *Nat. Commun.* **9**, 920 (2018).
- [33] M. Bouledroua, A. Dalgarno, and R. Côté, Diffusion and excitation transfer of excited alkali atoms, *Phys. Rev. A* **65**, 012701 (2001).
- [34] R. Côté and A. Dalgarno, Ultracold atom-ion collisions, *Phys. Rev. A* **62**, 012709 (2000).
- [35] E. Bodo, P. Zhang, and A. Dalgarno, Ultra-cold ion-atom collisions: near resonant charge exchange, *New J. Phys.* **10**, 033024 (2008).
- [36] P. Zhang, A. Dalgarno, and R. Côté, Scattering of Yb and Yb⁺, *Phys. Rev. A* **80**, 030703 (2009).
- [37] B. Zygelman, Hyperfine level-changing collisions of hydrogen atoms and tomography of the dark age Universe, *Astrophys. J.* **622**, 1356 (2005).
- [38] B. Zygelman, Electronic spin-flipping collisions of hydrogen atoms, *Phys. Rev. A* **81**, 032506 (2010).
- [39] M. Li and B. Gao, Proton-hydrogen collisions at low temperatures, *Phys. Rev. A* **91**, 032702 (2015).
- [40] A. E. Glassgold, P. S. Krstić, and D. R. Schultz, H⁺ + H scattering and ambipolar diffusion heating, *Astrophys. J.* **621**, 808 (2005).
- [41] S. R. Furlanetto and M. R. Furlanetto, Spin exchange rates in proton-hydrogen collisions, *Mon. Not. R. Astron. Soc.* **379**, 130 (2007).
- [42] L.-M. Duan, J. I. Cirac, and P. Zoller, Quantum entanglement in spinor Bose-Einstein condensates, *Phys. Rev. A* **65**, 033619 (2002).
- [43] M.-S. Chang, Q. Qin, W. Zhang, L. You, and M. S. Chapman, Coherent spinor dynamics in a spin-1 Bose condensate, *Nat. Phys.* **1**, 111 (2005).
- [44] D. Jaksch, H.-J. Briegel, J. I. Cirac, C. W. Gardiner, and P. Zoller, Entanglement of atoms via cold controlled collisions, *Phys. Rev. Lett.* **82**, 1975 (1999).
- [45] T. Calarco, E. A. Hinds, D. Jaksch, J. Schmiedmayer, J. I. Cirac, and P. Zoller, Quantum gates with neutral atoms: Controlling collisional interactions in time-dependent traps, *Phys. Rev. A* **61**, 022304 (2000).
- [46] D. Hayes, P. S. Julienne, and I. H. Deutsch, Quantum logic via the exchange blockade in ultracold collisions, *Phys. Rev. Lett.* **98**, 070501 (2007).
- [47] J. H. M. Jensen, J. J. Sørensen, K. Mølmer, and J. F. Sherson, Time-optimal control of collisional $\sqrt{\text{swap}}$ gates in ultracold atomic systems, *Phys. Rev. A* **100**, 052314 (2019).
- [48] K.-K. Ni, T. Rosenband, and D. D. Grimes, Dipolar exchange quantum logic gate with polar molecules, *Chem. Sci.* **9**, 6830 (2018).
- [49] F. Schmidt, D. Mayer, Q. Bouton, D. Adam, T. Lausch, J. Nettersheim, E. Tiemann, and A. Widera, Tailored single-atom collisions at ultralow energies, *Phys. Rev. Lett.* **122**, 013401 (2019).
- [50] F. Fang, J. A. Isaacs, A. Smull, K. Horn, L. D. Robledo-De Basabe, Y. Wang, C. H. Greene, and D. M. Stamper-Kurn, Collisional spin transfer in an atomic heteronuclear spinor Bose gas, *Phys. Rev. Research* **2**, 032054 (2020).
- [51] E. P. Wigner, On the behavior of cross sections near thresholds, *Phys. Rev.* **73**, 1002 (1948).
- [52] H. R. Sadeghpour, J. L. Bohn, M. Cavagnero, B. D. Esry, I. I. Fabrikani, J. H. Macek, and A. R. P. Rau, Collisions near threshold in atomic and molecular physics, *J. Phys. B* **33**, R39 (2000).
- [53] N. Balakrishnan, V. Kharchenko, R. C. Forrey, and A. Dalgarno, Complex scattering lengths in multi-channel atom-molecule collisions, *Chem. Phys. Lett.* **280**, 5 (1997).
- [54] R. V. Krems, Molecules near absolute zero and external field control of atomic and molecular dynamics, *Int. Rev. Phys. Chem.* **24**, 99 (2005).
- [55] A. Dalgarno and D. R. Bates, Spin-change cross-sections, *Proc. R. Soc. London. Ser. A* **262**, 132 (1961).
- [56] A. Dalgarno, M. R. H. Rudge, and D. R. Bates, Spin-change cross-sections for collisions between alkali atoms, *Proc. R. Soc. London. Ser. A* **286**, 519 (1965).
- [57] A. J. Moerdijk, B. J. Verhaar, and T. M. Nagtegaal, Collisions of dressed ground-state atoms, *Phys. Rev. A* **53**, 4343 (1996).
- [58] R. Côté and A. Dalgarno, Elastic scattering of two Na atoms, *Phys. Rev. A* **50**, 4827 (1994).
- [59] A. J. Moerdijk and B. J. Verhaar, Prospects for Bose-Einstein condensation in atomic ⁷Li and ²³Na, *Phys.*

- Rev. Lett. **73**, 518 (1994).
- [60] E. R. I. Abraham, W. I. McAlexander, J. M. Gerton, R. G. Hulet, R. Côté, and A. Dalgarno, Triplet s-wave resonance in ^6Li collisions and scattering lengths of ^6Li and ^7Li , Phys. Rev. A **55**, R3299 (1997).
 - [61] S. J. J. M. F. Kokkelmans, H. M. J. M. Boesten, and B. J. Verhaar, Role of collisions in creation of overlapping bose condensates, Phys. Rev. A **55**, R1589 (1997).
 - [62] R. Côté, A. Dalgarno, H. Wang, and W. C. Stwalley, Potassium scattering lengths and prospects for Bose-Einstein condensation and sympathetic cooling, Phys. Rev. A **57**, R4118(R) (1998).
 - [63] E. Timmermans and R. Côté, Superfluidity in sympathetic cooling with atomic Bose-Einstein condensates, Phys. Rev. Lett. **80**, 3419 (1998).
 - [64] M. J. Jamieson, A. Dalgarno, B. Zygelman, P. S. Krtic, and D. R. Schultz, Collisions of ground-state hydrogen atoms, Phys. Rev. A **61**, 014701 (1999).
 - [65] O. Makarov, R. Côté, H. Michels, and W. W. Smith, Radiative charge-transfer lifetime of the excited state of $(\text{NaCa})^+$, Phys. Rev. A **67**, 042705 (2003).
 - [66] D. H. Santamore and E. Timmermans, Pseudospin and spin-spin interactions in ultracold alkali atoms, New J. Phys. **13**, 023043 (2011).
 - [67] B. Zygelman, A. Dalgarno, M. J. Jamieson, and P. C. Stancil, Multichannel study of spin-exchange and hyperfine-induced frequency shift and line broadening in cold collisions of hydrogen atoms, Phys. Rev. A **67**, 042715 (2003).
 - [68] A. Pandey, M. Niranjana, N. Johi, A. Rangwala, R. Vexiau, and O. Dulieu, Interaction potentials and ultracold scattering cross sections for the $^7\text{Li}^+ + ^7\text{Li}$ ion-atom system, Phys. Rev. A **101**, 052702 (2020).
 - [69] A. Mohammadi, A. Krukow, A. Mahdian, M. Deiss, J. Pérez-Ríos, H. da Silva Jr., M. Raoult, O. Dulieu, and J. H. Denschlag, Life and death of a cold BaRb^+ molecule inside an ultracold cloud of Rb atoms, Phys. Rev. Res. **3**, 013196 (2021).
 - [70] R. V. Krems and A. Dalgarno, Threshold laws for collisional reorientation of electronic angular momentum, Phys. Rev. A **67**, 050704 (2003).
 - [71] C. Cohen-Tannoudji, B. Diu, and F. Laloë, *Quantum Mechanics* (Wiley, New York, 1977) p. 951.
 - [72] J. Weiner, V. S. Bagnato, S. Zilio, and P. S. Julienne, Experiments and theory in cold and ultracold collisions, Rev. Mod. Phys. **71**, 1 (1999).
 - [73] A. M. Arthurs and A. Dalgarno, The theory of scattering by a rigid rotator, Proc. R. Soc. London. Ser. A. **256**, 540 (1960).
 - [74] M. D. Rowe and A. J. McCaffery, Transfer of state multipoles in excited $\text{A } ^1\Sigma_u^+ \text{ } ^7\text{Li}_2$ following rotationally inelastic collisions with He: Experiment and theory, Chem. Phys. **43**, 35 (1979).
 - [75] M. H. Ross and G. L. Shaw, Multichannel effective range theory, Ann. Phys. **13**, 147 (1961).
 - [76] I. Simbotin, S. Ghosal, and R. Côté, Threshold resonance effects in reactive processes, Phys. Rev. A **89**, 040701 (2014).
 - [77] S. Green, Rotational excitation in $\text{H}_2\text{-H}_2$ collisions: Close-coupling calculations, J. Chem. Phys. **62**, 2271 (1975).
 - [78] See Supplemental Material at [URL], which includes Refs. [79–82] for details of quantum scattering calculations and convergence tests.
 - [79] T. V. Tscherbul, Y. V. Suleimanov, V. Aquilanti, and R. V. Krems, Magnetic field modification of ultracold molecule-molecule collisions, New J. Phys. **11**, 055021 (2009).
 - [80] V. Aquilanti, D. Ascenzi, M. Bartolomei, D. Cappelletti, S. Cavalli, M. d. C. Vitores, and F. Pirani, Quantum interference scattering of aligned molecules: Bonding in O_4 and role of spin coupling, Phys. Rev. Lett. **82**, 69 (1999).
 - [81] V. Aquilanti, D. Ascenzi, M. Bartolomei, D. Cappelletti, S. Cavalli, M. d. C. Vitores, and F. Pirani, Molecular beam scattering of aligned oxygen molecules. The nature of the bond in the $\text{O}_2\text{-O}_2$ dimer, J. Am. Chem. Soc. **121**, 10794 (1999).
 - [82] R. N. Zare, *Angular Momentum* (Wiley, 1988).
 - [83] Z. Li, S. Singh, T. V. Tscherbul, and K. W. Madison, Feshbach resonances in ultracold $^{85}\text{Rb}\text{-}^{87}\text{Rb}$ and $^6\text{Li}\text{-}^{87}\text{Rb}$ mixtures, Phys. Rev. A **78**, 022710 (2008).
 - [84] S. Aubin, S. Myrskog, M. H. T. Extavour, L. J. LeBlanc, D. McKay, A. Stummer, and J. H. Thywissen, Rapid sympathetic cooling to Fermi degeneracy on a chip, Nat. Phys. **2**, 384 (2006).
 - [85] A. M. Thomas, *Ultra-cold Collisions and Evaporative Cooling of Caesium in a Magnetic Trap*, Ph.D. thesis, Oxford University (2004).
 - [86] P. Sompet, S. S. Szigeti, E. Schwartz, A. S. Bradley, and M. F. Andersen, Thermally robust spin correlations between two ^{85}Rb atoms in an optical microtrap, Nat. Commun. **10**, 1889 (2019).
 - [87] R. Ben-shlomi, M. Pinkas, Z. Meir, T. Sikorsky, O. Katz, N. Akerman, and R. Ozeri, High-energy-resolution measurements of an ultracold-atom-ion collisional cross section, Phys. Rev. A **103**, 032805 (2021).
 - [88] S. Uetake, R. Murakami, J. M. Doyle, and Y. Takahashi, Spin-dependent collision of ultracold metastable atoms, Phys. Rev. A **86**, 032712 (2012).
 - [89] M.-J. Lu, V. Singh, and J. D. Weinstein, Inelastic titanium-titanium collisions, Phys. Rev. A **79**, 050702 (2009).