

This is the accepted manuscript made available via CHORUS. The article has been published as:

Entangling qubit registers via many-body states of ultracold atoms

R. G. Melko, C. M. Herdman, D. Iouchtchenko, P.-N. Roy, and A. Del Maestro

Phys. Rev. A **93**, 042336 — Published 25 April 2016

DOI: [10.1103/PhysRevA.93.042336](https://doi.org/10.1103/PhysRevA.93.042336)

Entangling qubit registers via many-body states of ultracold atoms

R. G. Melko,^{1,2} C. M. Herdman,^{1,3,4} D. Iouchtchenko,⁴ P.-N. Roy,⁴ and A. Del Maestro^{5,*}

¹*Department of Physics and Astronomy, University of Waterloo, Ontario, N2L 3G1, Canada*

²*Perimeter Institute for Theoretical Physics, Waterloo, Ontario N2L 2Y5, Canada*

³*Institute for Quantum Computing, University of Waterloo, Ontario, N2L 3G1, Canada*

⁴*Department of Chemistry, University of Waterloo, Ontario, N2L 3G1, Canada*

⁵*Department of Physics, University of Vermont, Burlington, VT 05405, USA*

Inspired by the experimental measurement of the Rényi entanglement entropy in a lattice of ultracold atoms by Islam *et al.* [Nature **528**, 77 (2015)], we propose a method to entangle two spatially-separated qubits using the quantum many-body state as a resource. Through local operations accessible in an experiment, entanglement is transferred to a qubit register from atoms at the ends of a one-dimensional chain. We compute the *operational entanglement*, which bounds the entanglement physically transferable from the many-body resource to the register, and discuss a protocol for its experimental measurement. Finally, we explore measures for the amount of entanglement available in the register after transfer, suitable for use in quantum information applications.

I. INTRODUCTION

Islam *et al.*¹ have performed a measurement of the Rényi entanglement entropy in a one-dimensional optical lattice of ⁸⁷Rb atoms by exploiting a many-body analogue of the Hong-Ou-Mandel² photon interference effect. After interfering two proximate copies of an L -site lattice using the atomic control of a quantum gas microscope³, a measurement of the parity of the site resolved particle occupation number provides access to the state overlap of the two copies. If the initial copies are identical, this gives the purity of the state⁴. Hence, if a globally pure state is partitioned into spatial subregions, the many-body interference/parity measurement protocol localized to a subregion yields the Rényi entropy, a measure of entanglement between subregions⁵. This provides an experimental probe of a remarkable feature of quantum mechanics with no classical analogue: information may be encoded in a composite system in such a way that it is inaccessible from independent measurements of its component parts.

The advantage of measuring the Rényi entropy as in Ref. [1] is that it encodes the entanglement between subsystems in a scalar quantity that can be accessed through the expectation values of local operators⁴. This is in contrast to other entanglement measures calculated directly from the full density matrix, which is generally inaccessible in experiments without using full state tomography⁶. In particular, there is currently no scalable scheme for its reconstruction for N interacting itinerant particles. This fact makes the two-copy Rényi entropy, $S_2(A) = -\log(\text{Tr} \rho_A^2)$, particularly well-suited for exploration in a quantum many-body system bipartitioned into a spatial region A and its complement \bar{A} .

S_2 has proved fruitful for the general characterization of many-body phases and quantum phase transitions, *e.g.* through the exploration of its scaling with subsystem size⁷. Additionally, given that entanglement is a physical resource that can be used for quantum information processing^{8,9}, it is natural to ask whether this many-




Figure 1. (Color online) A schematic setup whereby entanglement can be transferred from a quantum many-body state $|\Psi\rangle$ to a quantum register composed of two spatially separated qubits (Bloch spheres).

body entanglement can be harnessed for these tasks^{10–13}. One route to exploit entanglement between spatial regions of a many-body state for quantum information processing is to transfer entanglement of the many-body system to an external register of localized qubits using local operations; in this way the many-body state acts as an entanglement reservoir for the quantum register. To quantify the entanglement that is usable, one must take into account physical restrictions that limit the amount of entanglement that may be transferred to the register. For itinerant particles, a super selection rule (SSR) due to particle number conservation provides one key limitation¹⁴. Further restrictions are imposed if one wants to entangle spatially separated qubits with only local operations on the many-body system¹⁵.

In this paper we propose the general scheme shown in Fig. 1 and present an experimental protocol, using the basic capabilities of Islam *et al.*¹, to transfer some of the entanglement in a many-body state of ultracold atoms to two spatially-separated qubits composing an external quantum register. We present this protocol within the context of the Islam experiment, but the general concept of entangling qubits using many-body states is relevant to many other systems^{10–13}. We emphasize the importance of the *operational entanglement* as a bound on the

transferable entanglement, and discuss its measurement in the many-body state. The demonstration of this transfer would be proof of principle confirmation that a quantum register can be entangled in current experimental apparatuses for ultracold atoms.

II. ENTANGLEMENT IN THE BOSE-HUBBARD MODEL

The ^{87}Rb atoms of the Islam experiment are confined to move in a deep one-dimensional optical lattice. In their weakly interacting regime, the low energy dynamics of the atoms are accurately governed by the lattice Bose-Hubbard Hamiltonian with N particles on L sites:

$$H = -J \sum_{i=1}^{L-1} (b_i^\dagger b_{i+1} + \text{h.c.}) + \frac{U}{2} \sum_{i=1}^L n_i (n_i - 1), \quad (1)$$

where $b_i^\dagger (b_i)$ creates (annihilates) a boson, and $n_i = b_i^\dagger b_i$ counts the number of atoms on site i . J sets the rate of tunneling between sites while U parametrizes the strength of the on-site repulsion between atoms. In an experiment, the interaction strength between ^{87}Rb atoms is fixed by their s -wave scattering length, while J can be tuned by manipulating the height of the optical lattice. In the thermodynamic limit at unit filling ($N = L$), Eq. (1) exhibits two distinct phases: a Mott insulator for $U/J \gg 1$ and a superfluid for $U/J \ll 1$, both of which are observed experimentally. A quantum phase transition separates these two phases at $(U/J)_c \approx 3.3$ ^{16–19}.

The spatially delocalized nature of particles in the superfluid phase suggests that it should be significantly more entangled under a spatial bipartition than a Mott insulator with localized particles. This is manifest as an increase in S_2 accompanying the onset of delocalization at $U/J \sim \text{O}(1)$ observed in the experiment for $N = 4$ atoms¹. The same experimental capabilities that allow for the measurement of the entanglement in an optical lattice can also be used to transfer entanglement to spatially separated qubits, that can be employed as a quantum register for information processing tasks via logic gates. This entanglement transfer procedure is limited by the SSR that forbids the creation of a coherent superposition of states with different local particle number¹⁴. Thus entanglement that arises *solely* due to particle fluctuations between subregions is not physically transferable to a register without a global phase reference²⁰.

A. Operational entanglement

To address this issue, Wiseman and Vaccaro¹⁴ introduced the concept of *operational entanglement* – the amount of entanglement that can be extracted from a resource (many-body state) and transferred to a quantum register in the presence of a SSR. Conceptually it is

the weighted sum of the spatial entanglement when projecting onto states of fixed local particle number. For the two copy Rényi entropy it is defined as:

$$S_2^{\text{op}}(A) = \sum_n P_n S_2(A_n), \quad (2)$$

where $S_2(A_n)$ is the Rényi entropy evaluated for the reduced density matrix

$$\rho_{A_n} = \frac{1}{P_n} \hat{P}_n \rho_A \hat{P}_n \quad (3)$$

projected by \hat{P}_n onto states of fixed local particle number n in subsystem A . The summation is over all possible local particle number states in the subregion with $n = 0, \dots, N$, each having probability $P_n = \langle \Psi | \hat{P}_n | \Psi \rangle$. This projection is a local operation that can only decrease entanglement¹⁵ so $S_2^{\text{op}} \leq S_2$.

Thus it is S_2^{op} , not S_2 which bounds the amount of entanglement that can be generated in the register using local operations and classical communication (LOCC). A measurement of S_2^{op} is possible with a simple modification of the experimental interference/parity measurement procedure^{1,4}. This requires that a projection onto states of definite subsystem particle number n be made after interference, which can be achieved in the experimental system by measuring the total particle number in each of the B subsystems, n_B after the beam splitter operation is performed. Given that each copy is initialized to a state with N particles, P_n is the probability of having $n_B = N - n$ in a single copy; $S_2(A_n)$ is computed from the parity measurements of instances $n_B = N - n$ in both copies, with which must be binned accordingly to compute S_2^{op} . Therefore, by collecting these n -resolved statistics of the Rényi entropy, Eq. (2) can be used to experimentally measure the operational entanglement.

In order to explore which parameter regime maximizes operational entanglement, we calculate S_2^{op} in the Bose-Hubbard model. Experiments on ^{87}Rb in the near future should be possible with $4 < N \lesssim 10$, and we study the ground states of systems with sizes of this order via exact numerical diagonalization of Eq. (1). In Fig. 2, we compare the two-copy Rényi entropy for a symmetric spatial bipartition to the operational entanglement for a range of U/J and N relevant for experiments. Unlike the entropy under a spatial bipartition, which is maximum deep in the superfluid phase (or the particle entanglement, which is maximum deep in the Mott phase²¹), S_2^{op} displays a peak at an intermediate value of the interaction. While for these system sizes, the peak is not positioned directly at the thermodynamic-limit critical point $(U/J)_c \approx 3.3$, it appears to approach this value as L is increased.²² This suggests that the appropriate experimental parameters for maximizing the transfer of many-body entanglement to a system of quantum registers will be those that tune the system to near the superfluid-Mott transition.

As seen in Fig. 2, S_2^{op} is necessarily smaller than S_2 , as it does not include entanglement generated by par-




Figure 2. (Color online) The spatial second Rényi entropy S_2 and operational entanglement S_2^{op} for symmetric bipartitions $\ell = L/2$ of the Bose-Hubbard model. Curves increase in saturation for $L = N = 6, 8, 10, 12$. The dashed vertical line indicates the location of the thermodynamic phase transition.

ticle fluctuations between subsystems that is not physically accessible due to the SSR. Additionally, S_2^{op} is reduced as interactions in Eq. (1) are strictly onsite and occur at fixed subsystem occupation through 2nd order processes. Thus, the behavior of the physically accessible entanglement differs from S_2 both qualitatively and quantitatively.

III. EXTRACTING MANY-BODY ENTANGLEMENT

Given that the operational entanglement indicates that some of the entanglement between spatial subregions of the many-body ground state may be transferred to an external quantum register using LOCC, we now describe an experimental procedure to do so. This allows the many-body state to act as an entanglement resource for quantum information protocols. We concentrate on the minimal $L = N = 6$ Bose-Hubbard system where entanglement may be transferred to two spatially separated qubits. Each qubit is comprised of one atom occupying one of two neighboring lattice sites adjacent to the Bose-Hubbard chain; the two locations of the atom provide the logical states. Thus, the physical system we describe consists of 10 total lattice sites, which must be doubled as shown in Fig. 3 if a two-copy Rényi measurement is to be made on the final entanglement between the qubits.

The starting point is the isolation of a 6×4 array of atoms that can be prepared deep in the Mott phase. This array includes the many-body entanglement resource, which will be partitioned into three spatial subregions with two sites each (labeled A, B, C), two qubit registers, Q_A and Q_B , and a copy that will be employed to




Figure 3. (Color online) Upper left: an array of 20 optical lattice sites forming the two copies necessary to measure the second Rényi entropy. Remaining panels: the protocol (described in the text) to transfer entanglement from a many-body state in $A \cup B \cup C$ to spatially separated qubits Q_A and Q_B . Solid lines correspond to a large tunnel barrier, double arrows represent the application of a SWAP operation and single arrows indicate performing many-body interference.

read out the amount of entanglement generated between Q_A and Q_B . To manipulate and measure entanglement in the system, we define a pairwise hopping unitary operator:

$$U_{ij}(\phi) \equiv \exp \left[i\phi \left(b_i^\dagger b_j + \text{h.c.} \right) \right]. \quad (4)$$

This is a trivial generalization of the beam-splitter operation reported in Ref. [1] (where $\phi = \pi/4$) and $\phi = \pi/2$ corresponds to a SWAP gate between i and j within the $n_{i,j} = 0, 1$ subspace. Additionally this physical operation can be used to perform single qubit rotations when applied within a single qubit. As U_{ij} will not generally preserve particle number within the resource and qubits (and thus not remain in the logical subspace of the qubits), subsystem resolved particle occupation number measurements must be used to post-select states that have exactly one particle in each of A and B .

Transfer of many-body entanglement to the register and its subsequent measurement can be accomplished via the three-step procedure depicted in Fig. 3.

- 1 The optical lattice within the array is manipulated such that large barriers (as indicated by solid lines) isolate the many-body resource. Each qubit must be constructed with exactly one particle between its two sites, with the barrier between them remaining high throughout the experiment. The many-body resource can be prepared identically to Ref. [1] with the lattice strength tuned near the critical value $(U/J)_c$ to maximize the operational entanglement as discussed above.
- 2 A SWAP operation (double arrow) is performed be-

tween $A \Leftrightarrow Q_A$ by applying the unitary hopping operator

$$U_{1,1'}(\pi/2)U_{2,2'}(\pi/2), \quad (5)$$

where sites 1, 2 are in region A , while $1', 2'$ label adjacent sites in Q_A . This is repeated for $B \Leftrightarrow Q_B$ and the identical procedure is performed in the copy. Thus entanglement is transferred from the many-body resource to the spatially separated qubits.

- ③ To read out this entanglement, a beam-splitter operation (single arrow) is performed between the two copies of Q_A and Q_B , followed by a subsystem resolved particle number measurement where instances with one atom in each qubit are post-selected, which is discussed below.

The above procedure will transfer many-body entanglement to a quantum register. As only A and B are swapped with the register, its density matrix ρ_{Q_A, Q_B} will generically be in a mixed state, even if the initial many-body state (ρ_{ABC}) was pure. Consequently, the mutual information

$$I_2(AB) = S_2(A) + S_2(B) - S_2(AB) \quad (6)$$

will have contributions from both classical correlations and quantum entanglement. $I_2(AB)$ is measurable in current experiments combined with post-selection to conserve particle number in $Q_{A/B}$.

A. Post-selection in the qubit subspace

Because the protocol discussed above requires a tripartite partitioning of each copy of the many-body state, post-selection of experimental instances that are in the qubit subspace of $Q_{A/B}$ after the SWAP operation is non-trivial. In measuring $S_2(AB)$, post selecting to $n_B = N - 2$ is insufficient to solely select instances in the qubit subspace. We define the number of particles in each qubit to be $n_{Q_{A/B}}$ and $n_{\bar{Q}_{A/B}}$. Since the beam splitter operation conserves the sum of the number of particles in each qubit and its copy, instances within the qubit subspace always result in $n_{Q_A} + n_{\bar{Q}_A} = 2$ and $n_{Q_B} + n_{\bar{Q}_B} = 2$. If the data is additionally post-selected such that $n_C = n_{\bar{C}} = N - 2$, the only non-qubit instances arise from those of the form:

$$|02\rangle_{Q_A}|00\rangle_{Q_B}|00\rangle_{\bar{Q}_A}|02\rangle_{\bar{Q}_B} \quad (7)$$

and 8 related permutations, before the beam splitter operation. As such states have a zero expectation value of parity after the beam splitter (or equivalently the expectation value of the SWAP operator is zero for such states), it is only necessary to determine the normalization of parity measurement, i.e. the probability of instances in the qubit subspace. For a given initial state,

the qubit probability P_{QB} can be independently measured from separate experiments where no beam-splitter operation is performed and only the site-resolved number of particles is measured. Then the beam-splitter parity measurements are performed and post-selected for states with $n_{Q_A} + n_{\bar{Q}_A} = 2$ and $n_{Q_B} + n_{\bar{Q}_B} = 2$ and $n_C = n_{\bar{C}} = N - 2$; we define the result of this measurement to be Σ_{AB} , and the probability of such instances to be \tilde{P}_{AB} . The desired value can be then computed as

$$S_2(Q_A Q_B) = -\log \left(\frac{\tilde{P}_{AB}}{P_{QB}} \Sigma_{AB} \right). \quad (8)$$

B. Quantifying entanglement between qubits

To quantify only the desired generation of quantum entanglement between the qubits, we compute various measures of mixed state entanglement for the reduced density matrix ρ_{AB} of the many-body ground state. Unlike for pure states, where the von Neumann entropy is the unique and appropriate entanglement measure, for mixed states, there are a variety of entanglement measures with different physical meanings. For example, the entanglement of formation E_F , roughly defined as the amount of entanglement required to form the mixed state, can be directly computed for any two qubit density matrix²³. The logarithmic negativity E_N depends on the sum of the negative eigenvalues of the density matrix after a partial transpose, and thus is readily computable for any density matrix²⁴. It provides an upper bound to the amount of entanglement that can be extracted from the mixed state using LOCC.

In Fig. 4 we have plotted $I_2(AB)$, E_F , and E_N of ρ_{AB} for the ground state of Eq. (1) in the 6-site geometry of Fig. 3, projected onto states with a single particle occupying A and B . We find that all these measures peak near the quantum phase transition^{25–27}. The peaks coincide with the parameter region of maximal operational entanglement desired for optimal transfer between resource and register. $E_N > 0$ is a necessary and sufficient condition for a two qubit state to be inseparable²⁸ such that it can be distilled to form a maximally entangled state²⁹. This implies that near the critical point the many-body resource has entanglement that can be extracted and distilled. Although there is no general relationship between I_2 and the entanglement measures E_F and E_N , in this case we can compute the relationship exactly for the Bose-Hubbard model. Thus, measurement of I_2 in an experimental regime where the Bose-Hubbard parameters are known will provide an estimate of the entanglement that can be generated between the qubits through the relationship calculated in Fig. 4.




Figure 4. (Color online) The logarithmic negativity E_N , entanglement of formation E_F , and two copy Rényi mutual information $I_2(AB)$ of the spatially separated qubits obtained from the $L = N = 6$ Bose-Hubbard ground state. The inset shows the probability of projecting onto a state with a single particle in each of A and B . The dashed vertical line indicates the location of the thermodynamic phase transition.

IV. DISCUSSION

In conclusion, we have introduced an experimental procedure for the transfer of entanglement from a many-body resource to spatially separated qubits forming a register suitable for quantum information processing. Conservation of particle number limits the amount of entanglement transferable from the resource, as quantified by the operational entanglement. The precise control of the current generation of quantum emulator experiments enables the faithful creation of lattice Bose-Hubbard models using ultracold atoms. This allows us to quantify the operational entanglement using exact calculations, and we find that the transferable entanglement is maximized

near the quantum phase transition between the Mott insulator and superfluid phases. This is in contrast to the naive expectation that transfer should occur in the superfluid phase, where experiments have confirmed that the two-copy Rényi entanglement is largest¹.

We have introduced a measurement protocol to experimentally probe the entanglement transferred by this procedure that employs a variation of a many-body interference technique^{1,4}. It is explicitly described for the transfer of entanglement from a 6-site resource to a register composed of two 2-site qubits – 20 lattice sites in total. It can be easily scaled to arbitrary size as experimental technology progresses. Our Bose-Hubbard calculations quantify the relationship between a mutual information accessible by this protocol and well-known measures for entanglement in mixed states.

The ability to engineer a wealth of variations of the Bose-Hubbard model will open up exciting prospects for extensions and optimizations of our results, through inhomogeneous parameters, topologies, and dimensionality. The experimental implementation of our protocol will demonstrate the potential of using many-body states of ultracold atoms as an entanglement resource for quantum information processing.

V. ACKNOWLEDGEMENTS

This work would not have been possible without discussions with R. Islam and A. Kaufmann. We thank J. Carrasquilla for his insights into the 1D Bose-Hubbard model and A. Brodutch for discussions about entanglement in mixed states. This research was supported by NSERC of Canada, the Canada Research Chair Program, the Perimeter Institute for Theoretical Physics (PI) and the National Science Foundation under Grant No. NSF PHY11-25915. Research at PI is supported by the Government of Canada through Industry Canada and by the Province of Ontario through the Ministry of Economic Development & Innovation.

* Adrian.DelMaestro@uvm.edu

¹ Rajibul Islam, Ruichao Ma, Philipp M Preiss, M Eric Tai, Alexander Lukin, Matthew Rispoli, and Markus Greiner, “Measuring entanglement entropy in a quantum many-body system,” *Nature* **528**, 77–83 (2015).

² C K Hong, Z Y Ou, and L Mandel, “Measurement of subpicosecond time intervals between two photons by interference,” *Phys. Rev. Lett.* **59**, 2044–2046 (1987).

³ W S Bakr, A Peng, M E Tai, R Ma, J Simon, J I Gillen, S Fölling, L Pollet, and M Greiner, “Probing the Superfluid-to-Mott Insulator Transition at the Single-Atom Level,” *Science* **329**, 547–550 (2010).

⁴ A J Daley, H Pichler, J Schachenmayer, and P Zoller, “Measuring Entanglement Growth in Quench Dynamics of Bosons in an Optical Lattice,” *Phys. Rev. Lett.* **109**,

020505 (2012).

⁵ Ryszard Horodecki, Michał Horodecki, and Karol Horodecki, “Quantum entanglement,” *Rev. Mod. Phys.* **81**, 865–942 (2009).

⁶ Daniel F V James, Paul G Kwiat, William J Munro, and Andrew G White, “Measurement of qubits,” *Phys. Rev. A* **64**, 052312 (2001).

⁷ Luigi Amico, Andreas Osterloh, and Vlatko Vedral, “Entanglement in many-body systems,” *Rev. Mod. Phys.* **80**, 517–576 (2008).

⁸ Charles H Bennett, Gilles Brassard, Claude Crépeau, Richard Jozsa, Asher Peres, and William K Wootters, “Teleporting an unknown quantum state via dual classical and Einstein-Podolsky-Rosen channels,” *Phys. Rev. Lett.* **70**, 1895–1899 (1993).

- ⁹ Guifré Vidal, “Efficient Classical Simulation of Slightly Entangled Quantum Computations,” *Phys. Rev. Lett.* **91**, 147902 (2003), 0301063.
- ¹⁰ Leonardo Banchi, Abolfazl Bayat, Paola Verrucchi, and Sougato Bose, “Nonperturbative Entangling Gates between Distant Qubits Using Uniform Cold Atom Chains,” *Phys. Rev. Lett.* **106**, 140501 (2011).
- ¹¹ N. Y. Yao, L. Jiang, A. V. Gorshkov, Z.-X. Gong, A. Zhai, L.-M. Duan, and M. D. Lukin, “Robust Quantum State Transfer in Random Unpolarized Spin Chains,” *Phys. Rev. Lett.* **106**, 040505 (2011).
- ¹² Salvatore M. Giampaolo and Fabrizio Illuminati, “Long-distance entanglement in many-body atomic and optical systems,” *New J. Phys.* **12**, 025019 (2010).
- ¹³ L Campos Venuti, S M Giampaolo, F Illuminati, and P Zanardi, “Long-distance entanglement and quantum teleportation in XX spin chains,” *Phys. Rev. A* **76**, 052328 (2007).
- ¹⁴ H M Wiseman and John A Vaccaro, “Entanglement of Indistinguishable Particles Shared between Two Parties,” *Phys. Rev. Lett.* **91**, 097902 (2003).
- ¹⁵ Michał Horodecki, Paweł Horodecki, and Ryszard Horodecki, “Limits for Entanglement Measures,” *Phys. Rev. Lett.* **84**, 2014–2017 (2000).
- ¹⁶ M A Cazalilla, R Citro, T Giamarchi, E Orignac, and M Rigol, “One dimensional bosons: From condensed matter systems to ultracold gases,” *Rev. Mod. Phys.* **83**, 1405–1466 (2011).
- ¹⁷ Juan Carrasquilla, Salvatore R Manmana, and Marcos Rigol, “Scaling of the gap, fidelity susceptibility, and Bloch oscillations across the superfluid-to-Mott-insulator transition in the one-dimensional Bose-Hubbard model,” *Phys. Rev. A* **87**, 043606 (2013).
- ¹⁸ G Boéris, L Gori, M D Hoogerland, A Kumar, E Lucioni, L Tanzi, M Inguscio, T Giamarchi, C D’Errico, G Carleo, G Modugno, and L Sanchez-Palencia, “Mott Transition for Strongly-Interacting 1D Bosons in a Shallow Periodic Potential,” (2015), 1509.04742.
- ¹⁹ Grigory E Astrakharchik, Konstantin V Krutitsky, Maciej Lewenstein, and Ferran Mazzanti, “One-dimensional Bose gas in optical lattices of arbitrary strength,” (2015), 1509.01424.
- ²⁰ Yakir Aharonov and Leonard Susskind, “Charge Superselection Rule,” *Phys. Rev.* **155**, 1428–1431 (1967).
- ²¹ C M Herdman, Stephen Inglis, P N Roy, R G Melko, and A Del Maestro, “Path-integral Monte Carlo method for Rényi entanglement entropies,” *Phys. Rev. E* **90**, 013308 (2014).
- ²² Note, however, that the choice of bipartition is arbitrary, and an exact scaling theory is unknown.
- ²³ William K Wootters, “Entanglement of Formation of an Arbitrary State of Two Qubits,” *Phys. Rev. Lett.* **80**, 2245–2248 (1998).
- ²⁴ G. Vidal and R. F. Werner, “Computable measure of entanglement,” *Phys. Rev. A* **65**, 032314 (2002), 0102117.
- ²⁵ A Osterloh, Luigi Amico, G Falci, and Rosario Fazio, “Scaling of entanglement close to a quantum phase transition,” *Nature* **416**, 608 (2002).
- ²⁶ Tobias Osborne and Michael Nielsen, “Entanglement in a simple quantum phase transition,” *Phys. Rev. A* **66**, 032110 (2002).
- ²⁷ Irénée Frérot and Tommaso Roscilde, “Entanglement entropy across the superfluid-insulator transition : a signature of bosonic criticality,” (2015), 1512.00805.
- ²⁸ Michał Horodecki, Paweł Horodecki, and Ryszard Horodecki, “Separability of mixed states: necessary and sufficient conditions,” *Phys. Lett. A* **223**, 1 (1996).
- ²⁹ Michał Horodecki, Paweł Horodecki, and Ryszard Horodecki, “Inseparable Two Spin-1/2 Density Matrices Can Be Distilled to a Singlet Form,” *Phys. Rev. Lett.* **78**, 574 (1997).