

This is the accepted manuscript made available via CHORUS. The article has been published as:

## Density-matrix correlations in the relaxation theory of electron broadening

T. A. Gomez, T. Nagayama, D. P. Kilcrease, M. H. Montgomery, and D. E. Winget

Phys. Rev. A **98**, 012505 — Published 16 July 2018

DOI: [10.1103/PhysRevA.98.012505](https://doi.org/10.1103/PhysRevA.98.012505)

# Density-Matrix Correlations in the Relaxation Theory of Electron Broadening

T. A. Gomez,<sup>1,\*</sup> T. Nagayama,<sup>1</sup> D. P. Kilcrease,<sup>2</sup> M. H. Montgomery,<sup>3</sup> and D. E. Winget<sup>3</sup>

<sup>1</sup>*Sandia National Laboratories*

<sup>2</sup>*Los Alamos National Laboratory*

<sup>3</sup>*University of Texas at Austin*

(Dated: June 28, 2018)

Spectral lines of radiating atoms are broadened by perturbations due to the surrounding plasma environment. In line-broadening calculations, the statistical average of the perturbation is weighted by the density matrix, which—in thermal equilibrium—contains correlations between the radiator and plasma. These correlations, however, have been neglected by all line-broadening theories except for the kinetic theory. The relaxation theory of line broadening is a mathematically exact derivation containing only one physical approximation: neglect of density-matrix correlations. We revisit this derivation and improve it by including the correlations. The line-broadening operator derived with the updated relaxation theory differs from that derived from the kinetic theory, though both derivations are considered to be exact. The kinetic theory derivation predicts that density-matrix correlations result a strong static shift of spectral lines. Our derivation, on the other hand, predicts that the correlations are a frequency-dependent effect that affects the line wings, and there is not shift of the line due to correlations. In addition, we predict that changes in the line shape due to correlations are only noticeable at extremely high densities. To distinguish the more appropriate model, we compare the shifts calculated with the relaxation and kinetic theory with data. The comparison shows support for the relaxation theory and casts into doubt the accuracy of the derivation of the kinetic theory of line broadening.

## I. INTRODUCTION

The broadening of spectral lines is important for many applications, including diagnostics of laboratory and astrophysical plasmas [1, 2] and modeling opacities [3–6]. Atomic lines are broadened because the radiating atom is perturbed by the charged particles (such as electrons and ions) in the surrounding plasma.

There are three primary methods for calculating electron broadening: the impact theory [7, 8], the relaxation theory [9], and the kinetic theory [10]; the different methods all give slightly different functional forms for the electron-broadening operator. We will not discuss the impact theory further since it contains a number of simplifying assumptions. Since both the relaxation and kinetic theories are more mathematically rigorous, we will focus on the differences between them. There are two fundamental differences between the kinetic and relaxation theories: how the broadening is related to the electron-collision amplitude, and the treatment of the density matrix. Here, we will focus on the differences in the density matrix treatment.<sup>1</sup>

The electron-broadening operator is closely related to the thermally averaged collision amplitude, which is weighted by the density matrix [25]. The time-dependence of the density matrix is governed by the time

evolution of the total atom plus plasma-electron system.

$$\rho(t) = e^{-iHt} \rho_0 e^{iHt}, \quad (1)$$

where  $\rho_0 \equiv \rho(t=0)$  is the density matrix at a given initial time [9, 25], and  $H$  is the Hamiltonian of the total system, consisting of radiator and plasma Hamiltonians ( $H_0^a$  and  $H_0^p$ , respectively), and the interaction ( $V^{rp}$ ) between the radiator and plasma,

$$H = H_0^r + H_0^p + V^{rp}, \quad (2)$$

where the superscripts  $r$  and  $p$  denote radiator of interest and perturbing plasma, respectively. The expected form of  $\rho_0$  is that it is a thermal distribution of the total atom-plasma system,

$$\begin{aligned} \rho_0 &= e^{-\beta H} / \text{Tr}\{e^{-\beta H}\} \\ &= e^{-\beta(H_0^r + H_0^p + V^{rp})} / \text{Tr}\{e^{-\beta(H_0^r + H_0^p + V^{rp})}\}, \end{aligned} \quad (3)$$

where  $\beta = 1/k_B T$ . The presence of  $V^{rp}$  in the density matrix creates correlations between the radiator and the plasma. The kinetic theory is the only derivation that uses this form. The relaxation theory neglect  $V^{rp}$  in  $\rho_0$  and the density matrix becomes the populations of the individual systems,

$$\rho_{0,relax} \approx \rho_{r,u} \rho_{p,u}, \quad (4)$$

thus ignoring the correlations in  $\rho_0$  between the atom and plasma; this is a common approximation when calculating the time evolution of the density matrix [7, 11–14].

This is not to say that the omitting correlations in  $\rho_0$  neglects correlations entirely. According to Eq (1), correlations will build up in  $\rho(t)$  as the plasma and radiator evolve in time and several collisions have taken place

\* thogome@sandia.gov

<sup>1</sup> The density matrix is an important tool in calculating kinetics of quantum systems. Methods for solving for the time evolution of the density matrix vary, which include—but not limited to—projection-operator techniques [11–15], solving BBGKY hierarchy of differential equations [16–19], and Green’s function methods [20–24].

[7, 9, 11–15]. In other words, the relaxation theory [9] only neglects correlations at the initial time,  $t = 0$ , and the correlations for  $t \gg 0$  will be dominated by the collisions. In this picture, the correlations in  $\rho_0$  will only be important for early time evolution of  $\rho(t)$ , and for the application of spectral line broadening, this should only be important for the wings of the line (where  $\Delta\omega$  is large). Both Baranger [7] and Fano [9] have predicted that the effect of the correlations in  $\rho_0$  should only be important for plasma conditions where the coupling is high.

The kinetic-theory-based formalism that accounts for initial density-matrix correlations predicts a large frequency-independent shift of spectral lines [26]. Frequency-independent shifts imply that the correlations in  $\rho(t)$  are always influenced by  $\rho_0$  regardless of how much time has passed and how many collisions have occurred. This seems to contradict the above prediction of Fano [9] and Baranger [7] that the correlations in  $\rho_0$  affect only the initial time evolution of  $\rho(t)$ .

In this paper, we investigate the prediction of the kinetic theory in two ways. First, we check it theoretically by including density-matrix correlations in another line-shape theory. Second, we check predictions of both the relaxation and kinetic theories against measured line shifts. In order to accomplish our first goal, we revisit the relaxation theory of line broadening, which is mathematically exact and the only physical approximation is the Eq (4). We do this with a straight-forward step involving a specific definition of the identity operator. The corrections to the line shape in the formalism presented here differ from what is predicted in the kinetic theory of line broadening [10]. In the derivation here, the initial correlations have a frequency-dependent effect on the line shape, with the strongest changes in the wings, and the smallest in the core; the overall change in the line shape due to initial correlations is a minor asymmetry. We then compare calculations using the relaxation and kinetic theory formulae with experiment [27], and show that the kinetic theory formulae over-predicts the line shifts by several hundred percent. The differences in derivation and the comparison with experiment cast the kinetic theory results [10, 26, 28–30] into question.

The paper is organized as follows: we first show the reader Fano’s derivation of the relaxation theory; second, we update the relaxation theory to include a more general density matrix; third, we perform a second-order evaluation of the line shape operator and identify the new parts due to correlations; we then provide a means of evaluating these new correlation terms; lastly, we compare our results with the kinetic theory and experiment.

## II. FANO’S RELAXATION THEORY

The line shape is given by the real part of the Fourier transform of the dipole-dipole autocorrelation function

[15, 31],

$$I(\omega) = \frac{1}{\pi} \Re \int_0^\infty dt e^{i\omega t} \text{Tr} \{ \vec{D} \cdot \vec{D}(t) \rho_0 \}, \quad (5)$$

where the trace is performed over all plasma and radiator states; the time evolution of the dipole moment is given in the Heisenberg picture,

$$\vec{D}(t) = e^{iHt} \vec{D} e^{-iHt},$$

where  $H$  is the Hamiltonian of total radiator-plasma system (given in Eq 2); and  $\rho_0$  is the density matrix. In the Heisenberg picture  $\rho_0$  does not evolve in time, a cyclic permutation of the operators can include  $\rho_0$  in the time evolution,

$$\begin{aligned} I(\omega) &= \frac{1}{\pi} \Re \int_0^\infty dt e^{i\omega t} \text{Tr} \{ D \cdot [e^{-iHt} \rho_0 D e^{iHt}] \} \\ &= \Re \int_0^\infty dt e^{i\omega t} \text{Tr} \{ D \cdot e^{-iLt} (\rho_0 D) \} \end{aligned} \quad (6)$$

where the only difference between this and Eq (1) is that the density matrix is multiplied by  $\vec{D}$ . In Eq (6), we have used the Liouville notation as a shorthand,

$$L(\rho_0 \vec{D}) = [H, (\rho_0 \vec{D})] = H(\rho_0 \vec{D}) - (\rho_0 \vec{D})H;$$

see Fano [9] or Zwanzig [15] for more details on Liouville formalism. The line shape formalism can be written as a function of frequency by moving the time integral inside the trace and performing the integral,

$$I(\omega) = \frac{-1}{\pi} \Im \text{Tr} \left\{ D \frac{1}{\omega - L} (\rho_0 D) \right\}. \quad (7)$$

The total system can be separated into the radiator and the perturbing plasma coordinates, where we can split the trace accordingly,

$$I(\omega) = \frac{-1}{\pi} \Im \text{Tr}_r \left[ D \text{Tr}_p \left\{ \frac{1}{\omega - L} \rho_0 \right\} D \right]. \quad (8)$$

The Liouville operator can be separated into the unperturbed and interaction Liouvilles, analogous to Eq (2),

$$L = L_0^r + L_0^p + L_I. \quad (9)$$

where

$$L_0^r \rho_0 = [H_0^r, \rho_0] \quad (9a)$$

$$L_0^p \rho_0 = [H_0^p, \rho_0] \quad (9b)$$

$$L_I \rho_0 = [V^{r,p}, \rho_0] \quad (9c)$$

For simplicity in the derivation, we will contain all non-interacting Liouvilles into one operator,  $L_0 = L_0^r + L_0^p$ . In order to evaluate the effect of the perturbers on the radiator’s time evolution, it is convenient to isolate  $L_I$ . Fano rewrote the quantity inside the curly bracket of Eq (8) as follows:

$$\frac{1}{\omega - L_0 - L_I} \rho_0 = \frac{1}{\omega - L_0} \left[ 1 + T(\omega) \frac{1}{\omega - L_0} \right] \rho_0, \quad (10)$$

where

$$T(\omega) = L_I + L_I(\omega - L_0)^{-1}T(\omega) \quad (11a)$$

$$= \frac{1}{1 - L_I(\omega - L_0)^{-1}}L_I. \quad (11b)$$

Here,  $T(\omega)$  is the Liouville-generalized scattering “T”-matrix (or collision amplitude) and contains the frequency-dependent interaction between the radiator and the perturbors. This is a particularly convenient form since the interaction between the radiator and plasma is isolated in  $T(\omega)$  and is now linearized, so that a density matrix expansion can be performed, as Fano did in the original publication. The density expansion permits everything to be written in terms of two- or three-particle interactions.

Up to this point, Fano has not introduced any approximation, and thus the expression is exact; Eq (10) will be our starting point when we include a more general density matrix.

Fano introduced the approximation that the density matrix could be approximated as

$$\rho_0 \approx \rho_{r,u}\rho_{p,u}, \quad (12)$$

$$\rho_{r,u} = e^{-\beta H_0^r}/Tr\{e^{-\beta H_0^r}\} \quad (12a)$$

$$\rho_{p,u} = e^{-\beta H_0^p}/Tr\{e^{-\beta H_0^p}\} \quad (12b)$$

where  $\rho_{r,u}$  is the density matrix of the isolated radiator, and  $\rho_{p,u}$  is the density matrix of the perturbors; the additional subscript,  $u$  denotes uncorrelated density matrices, which is to distinguish these from the more general definition,

$$\rho_r = Tr_p\{\rho_0\}, \quad (13)$$

which contains the correlations. This assumes that correlations between the radiator and plasma are negligible. Equation (12) has the property that it commutes with  $(\omega - L_0)^{-1}$ . The perturber density matrix was moved to the other side of  $(\omega - L_0)^{-1}$  so it is adjacent to the “T”-matrix,

$$\frac{1}{\omega - L}\rho_{r,u}\rho_{p,u} = \frac{1}{\omega - L_0}\left[\rho_{p,u} + T(\omega)\rho_{p,u}\frac{1}{\omega - L_0}\right]\rho_{r,u}.$$

Then the trace in Eq (8) can be re-written as follows:

$$Tr_p\left\{\frac{1}{\omega - L}\rho_{r,u}\rho_p\right\} = \frac{1}{\omega - L_0^r}\left[1 + \langle T(\omega) \rangle_p \frac{1}{\omega - L_0^r}\right]\rho_{r,u}, \quad (14)$$

where

$$\langle T(\omega) \rangle_p = Tr_p\{T(\omega)\rho_{p,u}\}. \quad (15)$$

The average  $T(\omega)$  contains all the effects of the interaction of the radiator with the plasma.

It is desirable to connect Eq (14) to more commonly-used expression for the line-shape function [7, 31, 32]:

$$I(\omega) = \frac{-1}{\pi} \Im Tr_r \left[ D \frac{1}{\omega - L_0^r - \mathcal{H}(\omega)} \rho_{r,u} D \right], \quad (16)$$

where  $\mathcal{H}(\omega)$  is the line-broadening operator and governs the time evolution of the radiator due to the average perturbation from the plasma. In other words, we need to find a functional form for  $\mathcal{H}(\omega)$  that satisfies

$$\frac{1}{\omega - L_0^r - \mathcal{H}(\omega)} = \frac{1}{\omega - L_0^r} \left[ 1 + \langle T(\omega) \rangle_p \frac{1}{\omega - L_0^r} \right],$$

where the expression for  $\mathcal{H}(\omega)$  that satisfies this equation is

$$\mathcal{H}(\omega) = \frac{1}{1 + \langle T(\omega) \rangle_p (\omega - L_0^r)^{-1}} \langle T(\omega) \rangle_p. \quad (17)$$

This is Fano’s final expression for the electron-broadening operator, and is valid for conditions when density matrix correlations are not important.

### III. INCLUDING OFF-DIAGONAL DENSITY MATRIX TERMS

We now want to derive a line-broadening function that contains a density matrix that includes correlations; our starting point is Eq (10). We need to manipulate Eq (10) so that when we perform the trace, it becomes a form similar to Eq (14),

$$Tr_p \left\{ \frac{1}{\omega - L_0} \left[ 1 + T(\omega) \frac{1}{\omega - L_0} \right] \rho_0 \right\} = \frac{1}{\omega - L_0^r} \left[ 1 + \langle \mathfrak{T}(\omega) \rangle_p \frac{1}{\omega - L_0^r} \right] \rho_r, \quad (18)$$

where the unknown function  $\mathfrak{T}(\omega)$  contains all of the radiator-perturber interactions (including the density matrix) and would replace  $T(\omega)$  in evaluating the electron-broadening operator in Eq (17); here,  $\rho_r$  is defined according to Eq (13).

When the trace is performed on the first term on the left-hand side of Eq (18), it is similar in form to Eq (14),

$$Tr_p\{(\omega - L_0)^{-1}\rho_0\} = (\omega - L_0^r)^{-1}\rho_r.$$

To put the second term (on the left-hand side of Eq 18) in the desired form, we multiply on the right by the unit operator

$$I = \rho_r^{-1}(\omega - L_0^r) \frac{1}{\omega - L_0^r} \rho_r, \quad (19)$$

so that

$$\frac{1}{\omega - L_0} T(\omega) \frac{1}{\omega - L_0} \rho_0 = \quad (20)$$

$$= \frac{1}{\omega - L_0} T(\omega) \frac{1}{\omega - L_0} \rho_0 \rho_r^{-1} (\omega - L_0^r) \frac{1}{\omega - L_0^r} \rho_r$$

$$= \frac{1}{\omega - L_0} \mathfrak{T}(\omega) \frac{1}{\omega - L_0^r} \rho_r$$

$$\mathfrak{T}(\omega) = T(\omega) \frac{1}{\omega - L_0} \rho_0 \rho_r^{-1} (\omega - L_0^r). \quad (21)$$

When we perform the trace over the plasma coordinates, this becomes the right-hand side of Eq (18). Now the electron-broadening operator is defined as

$$\mathcal{H}(\omega) = \frac{1}{1 + \langle \mathfrak{T}(\omega) \rangle (\omega - L_0^r)^{-1}} \langle \mathfrak{T}(\omega) \rangle \quad (22)$$

$$\mathfrak{T}(\omega) = L_I \frac{1}{1 - (\omega - L_0)^{-1} L_I} \frac{1}{\omega - L_0} \rho_0 \rho_r^{-1} (\omega - L_0^r). \quad (23)$$

Use of Eq (19) is required because  $\rho_0$  no longer commutes with  $L_0$ . One property of Eq (23) is that if  $\rho_0$  is defined as in Fano [9], then  $\mathfrak{T}(\omega)$  reduces back to  $T(\omega)$ . We note that we have made no approximations about the structure of  $\rho_0$ , therefore this formula can be applied to a system with arbitrary initial conditions.

#### IV. SECOND-ORDER DENSITY MATRIX CORRELATION CORRECTION

Calculations of the full N-body line-broadening problem are extremely difficult and would require a large amount of computing time; computer simulations can do this numerically [33–36], but these line-broadening simulations are currently limited to classical particles. Some simplifying approximations for semi-analytic evaluation include Taylor expanding the electron-broadening operator (and  $\mathfrak{T}(\omega)$ ) to second order and collecting all terms that are second-order in  $L_I$ . In addition, the line-broadening problem is often reduced to a binary-collision result [7, 8], where the interactions between plasma particles are approximated with a screening factor [37–39]; these approximations are used by every semi-analytic calculation. The second-order binary-collision result for the electron broadening is [9, 40]

$$\mathcal{H}(\omega) \approx \mathcal{H}^{(1)} + \mathcal{H}^{(2)}(\omega) - \mathcal{H}^{(1)}(\omega) \frac{1}{\omega - L_0^r} \mathcal{H}^{(1)}(\omega) \quad (24)$$

$$\mathcal{H}^{(1)}(\omega) = \text{Tr}_p \{ L_I G(\omega) \rho_0 \} \rho_r^{-1} (\omega - L_0^r) \quad (24a)$$

$$\mathcal{H}^{(2)}(\omega) = \text{Tr}_p \{ L_I G(\omega) L_I G(\omega) \rho_0 \} \rho_r^{-1} (\omega - L_0^r) \quad (24b)$$

where  $N$  is the number of particles in the volume considered in the plasma, and  $G(\omega)$  is a Green's function, defined to be

$$G(\omega) = \frac{1}{\omega - L_0}.$$

In order to properly evaluate the density matrix, we need to define the Hamiltonian as explicit sums over the  $N$  plasma particles,

$$H_0^p = \sum_i^N H_0^i + \sum_i^N \sum_{j < i} V^{ij}$$

$$V^{rp} = \sum_i^N V^{r,i}$$

and insert this into Eq (2) to get a total Hamiltonian,

$$H = H_0^r + \sum_i^N H_0^i + \sum_i^N \sum_{j < i} V^{ij} + \sum_i^N V^{ri}. \quad (25)$$

We can write the density matrix in a slightly simplified form,

$$\rho_0 \propto \exp \left\{ -\beta \left( H_0^r + \sum_{i=1}^N H_0^i + \sum_{i=1}^N V^{ri} \right) \right\}, \quad (26)$$

by assuming that we can account for the  $V^{ij}$  terms by screening  $V^{ri}$  [10, 28, 41]. Trying to solve for this density matrix explicitly is impractical, and a simpler method is required.

One of the goals of this section is to clearly show how the density matrix correlations affect the line shape. We therefore use an integral expansion [7, 15, 26, 29, 42] of Eq (26),

$$e^{-\beta(H_0^r + \sum_i H_0^i + \sum_i V^{ri})} = e^{-\beta H_0^r} e^{-\beta \sum_i H_0^i} - e^{-\beta H_0^r} e^{-\beta \sum_i H_0^i} \int_0^\beta d\tau e^{\tau(H_0^r + \sum_j H_0^j)} \sum_k V^{rk} e^{-\tau(H_0^r + \sum_{j'} H_0^{j'} + \sum_{k'} V^{rk'})}, \quad (27)$$

so that the first term is the same uncorrelated density matrix that Fano [9] used. We define the uncorrelated and

correlated density matrices as

$$\rho_0 = [\hat{\rho}_{r,u}\hat{\rho}_{p,u} + \hat{\rho}_c]/Tr\{\hat{\rho}_{r,u}\hat{\rho}_{p,u} + \hat{\rho}_c\} \quad (28)$$

$$\hat{\rho}_{r,u} = e^{-\beta H_0^r} \quad (28a)$$

$$\hat{\rho}_{p,u} = e^{-\beta \sum_i H_0^i} \quad (28b)$$

$$\hat{\rho}_c = -e^{-\beta H_0^r} e^{-\beta \sum_i H_0^i} \int_0^\beta d\tau e^{\tau(H_0^r + \sum_j H_0^j)} \sum_k V^{rk} e^{-\tau(H_0^r + \sum_{j'} H_0^{j'} + \sum_{k'} V^{rk'})}. \quad (28c)$$

where the density matrix with hats denote unnormalized density matrices.

Since our goal is to calculate  $\mathcal{H}(\omega)$  to second order in the interaction potential, we do not consider any density-matrix-correlations for terms in Eq (24) that are already second-order in  $L_I$ ; for these we will simply replace  $\rho_0$  with  $\rho_{r,u}\rho_{p,u}$  and they will become the original Fano [9] results. Therefore, the only second-order contribution from correlations will come from  $\mathcal{H}^{(1)}(\omega)$ . Using Eq (28) to create “uncorrelated” and “correlated” terms,  $\mathcal{H}^{(1)}(\omega)$  becomes

$$\begin{aligned} \mathcal{H}^{(1)}(\omega) &= Tr_p \{L_I(\omega - L_0)^{-1} \rho_0\} \rho_r^{-1}(\omega - L_0^r) \\ &= [Tr_p \{L_I \hat{\rho}_{p,u} \hat{\rho}_{r,u}\}(\omega - L_0^r)^{-1} \rho_r^{-1}(\omega - L_0^r) + Tr_p \{L_I(\omega - L_0)^{-1} \hat{\rho}_c\} \rho_r^{-1}(\omega - L_0^r)] / Tr\{\hat{\rho}_{r,u} \hat{\rho}_{p,u} + \hat{\rho}_c\}, \end{aligned} \quad (29)$$

where commutation of  $[L_0, \rho_{r,u}\rho_{p,u}] = 0$  has been used for the first term on the right-hand side; no further simplifications can be made to this equation without approximation. We desire to connect this first term to the original derivation of Fano, we therefore will approximate—though not entirely accurate—the inverse radiator density matrix as uncorrelated [10, 26]

$$\rho_r^{-1} \approx \rho_{r,u}^{-1}.$$

This simplification allows us to take advantage of commutation relationships and remove the frequency-dependence of this term; we will use this approximation for the rest of the paper. The uncorrelated line-broadening term approximately becomes the static shift as originally derived by Fano,

$$Tr_p \{L_I \hat{\rho}_{p,u} \hat{\rho}_{r,u}(\omega - L_0^r)^{-1}\} \rho_r^{-1}(\omega - L_0^r) / Tr\{\hat{\rho}_{r,u} \hat{\rho}_{p,u} + \hat{\rho}_c\} \approx Tr_p \{L_I \hat{\rho}_p\} / Tr\{\hat{\rho}_{r,u} \hat{\rho}_{p,u} + \hat{\rho}_c\}, \quad (30)$$

The second term then becomes the lowest-order correction for the density matrix correlations,

$$Tr_p \{L_I(\omega - L_0)^{-1} \hat{\rho}_c\} \rho_r^{-1}(\omega - L_0^r) / Tr\{\hat{\rho}_{r,u} \hat{\rho}_{p,u} + \hat{\rho}_c\}. \quad (31)$$

## V. EVALUATION OF THE CORRELATION CORRECTION

To properly evaluate the correction, we first need to find the proper normalization factor for  $\rho_0$ , which requires evaluating the trace of the total density matrix [9, 25]. The largest contribution to  $Tr\{\rho_0\}$  is  $\hat{\rho}_{r,u}\hat{\rho}_{p,u}$  (see appendix A), we therefore approximate the trace as

$$Tr_{rp} \{e^{-\beta H}\} \approx \left[ \sum_r e^{-\beta E_r} \right] \left( \frac{\mathcal{V}}{\lambda_D^3} \right)^N; \quad (32)$$

this will be accurate for the neutral hydrogen cases we explore here. The lowest-order correlated broadening term (Eq 31) becomes

$$\begin{aligned} Tr\{L_I(\omega - L_0)^{-1} \rho_c\} \rho_r^{-1}(\omega - L_0^r) &\approx \\ -Tr \left\{ \sum_i \sum_k L_I^i(\omega - L_0)^{-1} \left( \frac{\lambda_D^3}{\mathcal{V}} \right)^N e^{-\beta \sum_j H_0^j} e^{-\beta H_0^r} \int_0^\beta d\tau e^{\tau(H_0^r + H_0^k)} V^{rk} e^{-\tau(H_0^r + H_0^k)} \right\} e^{\beta H_0^r}(\omega - L_0^r). \end{aligned} \quad (33)$$

The resulting interaction will have single-perturber ( $i = k$ ) and two-perturber interactions ( $i \neq k$ ), with the former of order  $N$ , and the latter of order  $N^2$ . In addition,  $\exp(-\beta \sum_j H_0^j)$  will give a factor of  $\mathcal{V}/\lambda_D^3$  for every term where  $j \neq k$  or  $j \neq i$ . Using  $u$  and  $k$  to denote radiator and free-perturber states, the lowest-order density-matrix-correlation

correction to the line broadening is (upper-state broadening only)

$$\begin{aligned}
L_I(\omega - L_0)^{-1} \rho_c \rho_r^{-1} (\omega - L_0^r) \approx \\
-n_e^2 \lambda_D^6 \int_0^\infty d^3 \vec{k}_1 e^{-\beta \frac{1}{2} k_1^2} \langle u \vec{k}_1 | V^{r1} | u'' \vec{k}_1 \rangle (\omega - E_{u''} + E_l)^{-1} \int_0^\infty d^3 \vec{k}_2 e^{-\beta \frac{1}{2} k_2^2} \langle u'' \vec{k}_2 | V^{r2} | u' \vec{k}_2 \rangle \frac{1 - e^{-\beta(E_{u''} - E_u)}}{E_{u''} - E_{u'}} (\omega - E_{u'} + E_l) \\
- n_e \lambda_D^3 \int_0^\infty d^3 \vec{k}_1 \int_0^\infty d^3 \vec{k}_1' \langle u \vec{k}_1 | V^{r1} | u'' \vec{k}_1' \rangle \left( \omega - E_{u''} + E_l - \frac{1}{2} k_1'^2 + \frac{1}{2} k_1^2 \right)^{-1} \langle u'' \vec{k}_1' | V^{r1} | u' \vec{k}_1 \rangle e^{-\beta \frac{1}{2} k_1^2} \times \\
\frac{1 - e^{-\beta(E_{u''} - E_{u'} + \frac{1}{2} k_1'^2 - \frac{1}{2} k_1^2)}}{E_{u''} - E_{u'} + \frac{1}{2} k_1'^2 - \frac{1}{2} k_1^2} (\omega - E_{u'} + E_l), \quad (34)
\end{aligned}$$

where the superscripts on  $V$  denote on which particle the operator is acting. This formula can be reduced to radial integrals by performing a partial wave expansion and using Wigner 3j and 6j symbols [29, 43–46], which is done in detail in appendix B.

The inverse,  $(\omega - L_0)^{-1}$ , can be solved in the same way as in previous calculations where it is separated into its real and imaginary parts [9, 43],

$$(\omega - L_0)^{-1} = \frac{PV}{\omega - L_0} - i\pi\delta(\omega - L_0)$$

where  $PV$  is the principal value, and  $\delta$  is the Dirac delta function. There are multiple ways of evaluating these terms, such as the method in O'Brien and Hooper [43] and we will not discuss them further. Unfortunately, there is no simplification for the real part in Eq (34), but there is considerable simplification for the imaginary part. The imaginary part of Eq (34) can be evaluated in the same way as [29, 43, 45], but with a correction factor (showing only upper-state broadening),

$$\begin{aligned}
\Im Tr_p \{ L_1(\omega - L_0)^{-1} \rho_c \rho_r^{-1} (\omega - L_0^r) \} \approx \\
- \Im Tr_{k_1} e^{-\beta \frac{1}{2} k_1^2} \langle u k_1 | V(\omega - L_0)^{-1} V | u' k_1 \rangle \times \\
\left[ 1 - e^{-\beta(\omega - E_{u'} + E_l)} \right]. \quad (35)
\end{aligned}$$

This term does nothing to the center of the line, since the correction factor is zero when  $\Delta\omega_{u'l} = \omega - E_{u'} + E_l = 0$ . If  $\Delta\omega_{u'l}$  becomes appreciable compared to  $\beta$ , the blue part of the wings of the line will become depressed due to the reduction in the broadening, and the red wing on the other hand will become enhanced due to the increase of the broadening. In addition, the real part will also be zero at the line center due to the presence of  $(\omega - L_0)$ , but as the detuning becomes large, both the red and the blue sides of the profile exhibit a redshift, meaning that the depression of the blue wing, and the enhancement of the red wing will become stronger. The overall effect therefore, is to skew the wings of the profile toward the red, rather than being symmetric about the line center.

The effects of the density matrix are going to be most pronounced when the temperatures are small and the line shapes are wide. For typical laboratory conditions, such as those found in the Wiese *et al.* [27] experiment, the




FIG. 1. Comparison of line profile with and without inclusion of the correlated density-matrix correction (the second term in Eq (29)). The example here is hte Ly $\beta$  transition of neutral hydrogen at  $T_e = 1\text{eV}$  and  $n_e = 3.2 \times 10^{18} \text{e/cm}^3$ . The effect of the correlations is to slightly skew the wings of the profile.

changes are not noticeable. Therefore, to illustrate how the density matrix correlations change the line shape, we chose to compare line shapes at extremely high density. Figure 1 shows the line shape of Ly $\beta$  neutral hydrogen with and without the density matrix correlations (the details of the electron broadening are in [47] and repeated in appendix B). As discussed above, the correlations skew the wings so that the red wing is enhanced, and the blue wing is depressed. We can therefore conclude that except for highly-coupled plasmas, the correlations in the initial

density matrix are not important.

## VI. COMPARISON WITH THE KINETIC THEORY

First and foremost, we want to say that our derivation does not match the results of Hussey *et al.* [10]. The easiest way to see this is to write both results in the Lippmann-Schwinger [48] expansion. Hussey's electron broadening operator (Eq 4.3 in [10]) is

$$\mathcal{H}_{kin}(\omega) = Tr_p\{\hat{\mathcal{H}}_{kin}(\omega)\rho_0\}\rho_r^{-1} \quad (36)$$

where

$$\hat{\mathcal{H}}_{kin}(\omega)\rho_0 = L_I\rho_0 + \hat{\mathcal{H}}_{kin}(\omega)\rho_0 G_{kin}(\omega)\mathcal{L}_I \quad (37)$$

$$G_{kin}(\omega) = \frac{1}{\omega - L_0^r - L_0^p - C_{r,p}(\omega)} \quad (38)$$

where  $\mathcal{L}_I$  is a screened interaction,  $C_{r,b}(\omega)$  contains mean-field operators. The generalized relaxation theory results, which were derived in appendix C, are

$$\mathcal{H}_{relax}(\omega) = Tr_p\left\{\hat{\mathcal{H}}_{relax}(\omega)\rho_0\right\}\rho_r^{-1}(\omega - L_0^r) \quad (39)$$

where

$$\begin{aligned} \hat{\mathcal{H}}(\omega)\rho &= L_I(\omega - L_0^r - L_0^p)^{-1}\rho_0 \\ &+ L_I(\omega - L_0^r - L_0^p)^{-1}[1 - P]\hat{\mathcal{H}}(\omega)\rho_0, \end{aligned} \quad (40)$$

where  $P$  is a projection operator [11],

$$PA\rho_0 = \rho_0\rho_r^{-1}Tr_p\{A\rho_0\};$$

this form is fundamentally different than the kinetic theory in two principle ways: how the density matrix affects the broadening, and the presence of the projection operator.

The presence of the projection operator is unique to the relaxation theory of line-broadening<sup>2</sup>. Hussey *et al.* [10] shows that (after using  $C_{rp}(\omega)$  to screen  $L_I$  in frequency-dependent terms of  $\mathcal{H}(\omega)$ ) that his derivation is formally identical with the unified theory of Smith *et al.* [38]. We point out that even though the final form of the unified theory does not contain a projection operator, it is formally present in the derivation and is thrown out—a point of criticism by Lee [49]. Therefore, because a projection operator is formally present in both the relaxation and unified theories of line broadening, the lack of one in the kinetic theory raises concerns. However, this is not the proper place to discuss or justify the presence of the

projection operator in  $\mathcal{H}(\omega)$ ; this will be further examined in a later publication. The rest of this section is devoted to comparing how the density-matrix correlations change the line broadening according to each prediction (kinetic or relaxation theory).

Both our work and the Hussey *et al.* [10] derivation established that the lowest-order corrections due to density-matrix correlations are in the first term of  $\mathcal{H}(\omega)$ , which we denote as  $\mathcal{H}^{(1)}(\omega)$ . Splitting the density matrix into its uncorrelated and correlated terms,  $\rho_0 = \rho_p\rho_r + \rho_c$ , the  $\mathcal{H}^{(1)}(\omega)$  for the relaxation and kinetic theories are

$$\begin{aligned} \mathcal{H}_{kin}^{(1)}(\omega) &= Tr_b\{L_I\rho_{p,u}\} + \\ &Tr_p\{L_I\rho_c\}\rho_{r,u}^{-1}, \end{aligned} \quad (41)$$

$$\begin{aligned} \mathcal{H}_{rel}^{(1)}(\omega) &= Tr_p\{L_I\rho_{p,u}\} + \\ &Tr_p\{L_I(\omega - L_0)^{-1}\rho_c\}\rho_{r,u}^{-1}(\omega - L_0^r). \end{aligned} \quad (42)$$

Both theories have the same uncorrelated shift term. But the correlated-density-matrix correction is different: kinetic theory is static (i.e. frequency independent), relaxation theory is frequency-dependent.

Hussey *et al.* [10] predicts a static shift (meaning the whole line is shifted by this amount) that is second-order in  $V$  (since both  $L_I$  and  $\rho_c$  contain a factor of  $V$ ), which makes this term dipole allowed. For near-neutral one-electron atoms, dipole-allowed interactions are typically the strongest interactions, which means that this shift term can be quite large, and similar in magnitude to the electron broadening. Interestingly, the first implementation of the Hussey *et al.* [10] results do not include the static shift [28], though it is implemented in later calculations [26, 29, 30].

In our derivation, the density-matrix correlations are a frequency-dependent effect. The far wings of the line are shifted by the amount predicted by [10] and [26], which is achieved by evaluating Eq (42) at the limit where  $\omega \rightarrow \infty$ . The line core, on the other hand, does not have this shift; the presence of  $(\omega - L_0^r)$  in Eq (42) results in zero shift at the line core, and a slight asymmetry of the spectral lines. This asymmetry will enhance the red wing of the profile and depress the blue wing of the profile (as shown in the previous section).

There is experimental evidence that questions the accuracy of the kinetic-theory shift (Eq 41). The experiment by Wiese *et al.* [27] measured high-density hydrogen at temperatures of  $\sim 1$  eV and electron densities between  $10^{16} - 10^{17} \text{ e/cm}^3$ ; the scatter of the measured shifts for  $\text{H}\beta$  are less than  $0.1\text{\AA}$ . We computed  $\mathcal{H}(\omega)$  according to the relaxation and kinetic theory formulae for  $\text{H}\beta$ , the  $n = 4 \rightarrow n = 2$  transition; we use an expanded basis<sup>3</sup> to ensure the accuracy of our calculation. The shifts are compared in figure 2. At the highest-density conditions

<sup>2</sup> Projection operators have never been included to calculate a line shape; they have always been ignored.

<sup>3</sup> We include the  $n = 5$  states in addition to the  $n = 2$  and  $n = 4$  states; see Stambulchik *et al.* [50], Gomez *et al.* [36], or Kilcrease *et al.* [51] for a discussion of basis set accuracy.






FIG. 2. Predictions of shifts for the Relaxation theory and the Kinetic theory compared to the measured shifts by Wiese *et al.* [27]. The shifts from the relaxation theory, while not perfect, agrees fairly well with the measured values. The shifts from the kinetic theory, on the other hand, is several times larger than the measured value.

( $n_e = 10^{17} \text{ e/cm}^3$ ), the shift from Eq (41) over predicts the measured value by 675%, while the relaxation theory under-predicts by 35%.

## VII. CONCLUSION

Most line-broadening calculations assume that the density matrix which describes the radiator-plasma system is uncorrelated, meaning off-diagonal terms are neglected. Only the kinetic theory [10] of spectral line broadening has included these off-diagonal terms of the density matrix in its derivation. We have updated the mathematically rigorous relaxation theory of Fano [9] to include the off-diagonal terms of the density matrix. However, the line-broadening formulae we derive are different from that of Hussey's [10] kinetic theory. The largest difference between our work and the kinetic theory is the appearance of a line shift in the latter. Comparison with experiment shows that the kinetic theory formalism over-predicts the shifts. This work questions the accuracy of the kinetic theory line-broadening derivation of [10], and closer inspection of the derivation is needed.

## ACKNOWLEDGMENTS

We thank Dr. Carlos Iglesias, Dr. Stephanie Hansen, Dr. Andrew Baczewski, and Dr. Daniel Jensen for stimulating discussions. We also thank Marc Schaeuble for help with the manuscript.

T.A.G. acknowledges support from the National Science Foundation Graduate Research Fellowship under grant DGE-1110007. T.A.G. was also supported by the

U.S. Department of Energy, Office of Science Early Career Research Program, Office of Fusion Energy Sciences under FWP-14-017426. M.H.M., and D.E.W. acknowledge support from the United States Department of Energy under grant DE-SC0010623, the Wootton Center for Astrophysical Plasma Properties under the United States Department of Energy under grant DE-FOA-0001634, and the National Science Foundation grant NSF-AST 1707419. Sandia National Laboratories is a multimission laboratory managed and operated by National Technology and Engineering Solutions of Sandia, LLC., a wholly owned subsidiary of Honeywell International, Inc., for the U.S. Department of Energy's National Nuclear Security Administration under contract DE-NA-0003525. The views expressed in the article do not necessarily represent the views of the U.S. Department of Energy or the United States Government.

## Appendix A: Normalizing the Correlated Density Matrix

The density matrix, by definition, needs to be normalized to unity [25],

$$\text{Tr}\{\rho_0\} = 1. \quad (\text{A1})$$

Thus, whatever form the density matrix takes, needs to be divided by the trace. In the discussion above, we have separated the density matrix into its correlated and uncorrelated parts:

$$\rho_0 = \rho_{r,u}\rho_{p,u} + \rho_c$$

and the trace is evaluated as the sum of both the uncorrelated and correlated parts of the density matrix. The trace of the uncorrelated density matrix has been previously evaluated by Smith and Hooper [40] and O'Brien and Hooper [43],

$$\text{Tr}_r\{\rho_{r,u}\} = \sum_i e^{-\beta E_i^r} \quad (\text{A2})$$

$$\text{Tr}_p\{\rho_{p,u}\} = \left(\frac{\mathcal{V}}{\lambda_D^3}\right)^N, \quad (\text{A3})$$

where  $\mathcal{V}$  is the volume,  $\lambda_D$  is the thermal deBroglie wavelength, and  $N$  is the number of particles inside of the volume. We use this section to evaluate the trace of the density matrix that includes correlations.

In the expansion of the density matrix given in Eq (27), we obtain a first-order correction,

$$\text{Tr}\{\rho_c\} \approx \text{Tr}\left\{e^{-\beta E_i^r} e^{-\beta \sum_l H_0^l} \times \int_0^\beta d\tau e^{\tau(H_0^r + \sum_j H_0^j)} \sum_i V^{ri} e^{-\tau(H_0^r - \sum_k H_0^k)}\right\} \quad (\text{A4})$$

Because of the trace, and the properties of the single-particle operators in the matrix exponentials, the exponentials in the integral will vanish, so that the trace reduces to

$$\text{Tr}\{\rho_c\} \approx \left(\frac{\mathcal{V}}{\lambda_D^3}\right)^{N-1} \text{Tr}\left\{e^{-\beta E_i^r} e^{-\beta H_0^1} N V^{r1} \beta\right\}. \quad (\text{A5})$$

---

We can factor out volume and de Broglie wavelength terms out of  $\text{Tr}\{\rho_c\}$  to make it proportional to  $\text{Tr}\{\rho_{r,u}\rho_{p,u}\}$ ,

$$\begin{aligned} \text{Tr}\{\rho_0\} &= \text{Tr}\{\rho_{r,u}\rho_{p,u}\} + \text{Tr}\{\rho_c\} \\ &\approx \sum_i e^{-\beta E_i^r} \left(\frac{\mathcal{V}}{\lambda_D^3}\right)^N - \sum_i e^{-\beta E_i^r} \left(\frac{\mathcal{V}}{\lambda_D^3}\right)^N \lambda_D^3 \frac{N}{V} \text{Tr}\left\{e^{-\beta E_i^r} e^{-\beta H_0^1} V^{ri} \beta\right\} \left[\sum_i e^{-\beta E_i^r}\right]^{-1}, \end{aligned} \quad (\text{A6})$$

so that we can write the trace of the total density matrix as

$$\text{Tr}\{\rho_0\} \approx \text{Tr}\{\rho_{r,u}\rho_{p,u}\} \left[1 - \lambda_D^3 \frac{N}{V} \text{Tr}\left\{e^{-\beta E_i^r} e^{-\beta H_0^1} V^{r1} \beta\right\} \left(\sum_i e^{-\beta E_i^r}\right)^{-1}\right]. \quad (\text{A7})$$

In most cases, the density is small compared to unity, meaning that the term in square brackets can almost always be approximated as unity,

$$\begin{aligned} \text{Tr}\{\rho\} &= \text{Tr}\{\rho_{r,u}\rho_{p,u}\} + \text{Tr}\{\rho_c\} \\ &\approx \text{Tr}\{\rho_{r,u}\rho_{p,u}\}, \end{aligned} \quad (\text{A8})$$

which is what we use as the normalization in section V.

## Appendix B: Line-Broadening Details

This is an outline of the electron-broadening treatment that we use for calculation of the profiles in figure 1 and the shifts in figure 2. This was a code developed in Gomez [47], and we repeat the details here for the reader's convenience. Our electron-broadening calculation treats both the atomic and plasma electrons quantum mechanically. This was modeled after the quantum-mechanical treatments of Junkel *et al.* [29] and Woltz and Hooper [45], but differ in that we include exchange effects in our calculation. Since calculations of electron exchange in line broadening are not documented for a second-order line broadening theory (in the same way that direct-only calculations are documented in [29, 45]), we show the work

---

here. The point here is not to open a discussion about exchange; this will be given strict attention in a later paper. Rather, this is to inform the reader of our electron-broadening calculation.

For our electron-broadening treatment, we evaluate  $\mathcal{H}(\omega)$  to second-order in  $L_I$ , repeated from appendix C,

$$\begin{aligned} \mathcal{H}(\omega) &= \langle L_I G(\omega) \rho_0 \rangle \rho_r^{-1}(\omega - L_0^r) \\ &\quad + \langle L_I G(\omega) L_I G(\omega) \rho_0 \rangle \rho_r^{-1}(\omega - L_0^r) \\ &\quad - \langle L_I G(\omega) \rho_0 \rangle \rho_r^{-1} \langle L_I G(\omega) \rho_0 \rangle \rho_r^{-1}(\omega - L_0^r) \end{aligned}$$

In the main text, we reduce this to a binary-collision approximation and we keep everything second order in the interaction, including terms with the density matrix correlations. The resulting approximation for  $\mathcal{H}(\omega)$  is

$$\begin{aligned} \mathcal{H}(\omega) &= N \langle L_I \rho_p \rangle + N \langle L_I G(\omega) \rho_c \rangle \rho_r^{-1}(\omega - L_0^r) \\ &\quad + N \langle L_I G(\omega) L_I G(\omega) \rho_p \rangle; \end{aligned} \quad (\text{B1})$$

we are not currently evaluating the third term. The reason for this is that  $N \langle L_I \rho_p \rangle$  for neutral hydrogen is so small that the third term is negligible compared to  $N \langle L_I G(\omega) L_I G(\omega) \rho_p \rangle$ .

To calculate line shapes with the above, we use the same full-Coulomb treatment as Junkel *et al.* [29], where the individual terms are (showing only the upper-state broadening term)

---


$$\begin{aligned} \mathcal{H}_{u,u'}(\omega) &= n_e \lambda_D^3 \int_0^\infty e^{-\beta \frac{1}{2} k_1^2} \langle u \vec{k}_1 | V | u' \vec{k}_1 \rangle d^3 \vec{k}_1 + n_e \lambda_D^3 \int_0^\infty e^{-\beta \frac{1}{2} k_1^2} \int_0^\infty \langle u \vec{k}_1 | V | u'' \vec{k}_2 \rangle G(\omega) \langle u'' \vec{k}_2 | \rho_c | u' \vec{k}_1 \rangle d^3 \vec{k}_1 \\ &\quad + n_e \lambda_D^3 \int e^{-\beta \frac{1}{2} k_1^2} \int \langle u \vec{k}_1 | V | u'' \vec{k}_2 \rangle G(\omega) \langle u'' \vec{k}_2 | V | u' \vec{k}_1 \rangle d^3 \vec{k}_1, \end{aligned} \quad (\text{B2})$$

where  $u$  denotes a set of atomic states and  $\vec{k}$  denote a set of free-electron states. Here, we have neglected the two-electron term in Eq (34), and all  $k$  subscripts denote the different states of only one electron (in the main text,

the different subscripts were used to distinguish between electrons, but that is unnecessary here and the distinction between states becomes clearer).

As in Junkel *et al.* [29] and Woltz and Hooper [45], our interaction potential,  $V$  is a Coulomb interaction, and contain nuclear potential terms,

$$V(r_1, r_2) = \left[ \frac{1}{|\vec{r}_1 - \vec{r}_2|} - \frac{1}{r_1} \right], \quad \frac{1}{|\vec{r}_1 - \vec{r}_2|} = \sum_k \sum_q \frac{r_{<}^k}{r_{>}^{k+1}} \frac{4\pi}{2k+1} Y_{kq}^*(\hat{r}_1) Y_{kq}(\hat{r}_2), \quad (\text{B3})$$

where the electron-electron repulsion is Taylor expanded for easy evaluation, and  $r_{<}$  and  $r_{>}$  are the minimum and maximum, respectively, of  $r_1$  and  $r_2$ . To more simply evaluate the Coulomb matrix elements, we perform a partial-wave expansion of the free-electron states, and assume that the atomic wavefunction can also be separable in radial and angular components [29, 44, 45, 52]:

$$\langle r | \vec{k}_1 \rangle = \frac{2}{\pi} \sum_{l_1 m_1} i^l \frac{1}{k_1 r} F_{l_1}(k_1, r) Y_{l_1 m_1}^*(\hat{k}_1) Y_{l_1 m_1}(\hat{r}), \quad \langle r | u \rangle = \frac{1}{r} R_u Y_{lm}(\hat{r}), \quad (\text{B4})$$

where the radial wavefunctions  $F_l(k, r)$  are Coulomb waves (or plane waves for the case of neutral atoms). To evaluate the Coulomb matrix elements, it becomes convenient to couple the angular momentum together to a total angular momentum,  $L$ , with magnetic quantum number  $M$ . Coulomb matrix elements are independent of  $M$ , and depend on  $m_1$  only through a Clebsch-Gordan coupling Coefficient [44]. We can evaluate the  $M$  and  $m_1$  averaged interaction via a 3j-symbol sum rule,

$$\langle u k_1 l | \mathcal{H}(\omega) | u' k_1 l \rangle = \sum_L \langle u l_1 k_1 L | \mathcal{H}(\omega) | u' l k_1 L \rangle (2L+1) \sum_{m_1, M} \begin{pmatrix} l_u & l_1 & L \\ m_u & m_1 & -M \end{pmatrix}^2 = \sum_L \langle u l_1 k_1 L | \mathcal{H}(\omega) | u' l k_1 L \rangle \frac{(2L+1)}{2l_u+1}$$

After inserting this into Eq (B2), coupling the angular momentum of the atom and plasma electron, then performing the sum over the magnetic quantum numbers, the electron-broadening operator becomes,

$$\mathcal{H}_{u,u'}(\omega) = \sum_{L,S} \frac{(2L+1)(2S+1)}{(2l_u+1)(2s+1)} n_e \frac{\lambda_D^3}{2} \left\{ \int_0^\infty e^{-\beta \frac{1}{2} k_1^2} A_{u,u'}(k_1) dk_1 + \int_0^\infty e^{-\beta \frac{1}{2} k_1^2} \int_0^\infty dk_2 dk_1 B_{u,u'',u'}(k_1, k_2) \times \right. \\ \left. \left[ \frac{1 - e^{-\beta(E_{u''} - E_u + \frac{1}{2} k_2^2 - \frac{1}{2} k_1^2)}}{E_{u''} - E_u + \frac{1}{2} k_2^2 - \frac{1}{2} k_1^2} (\omega - E_{u'} + E_l) + 1 \right] \left[ \frac{PV}{\omega - E_{u''} + E_l - \frac{1}{2} k_2^2 + \frac{1}{2} k_1^2} - i\pi \delta \left( \omega - E_{u''} + E_l - \frac{1}{2} k_2^2 + \frac{1}{2} k_1^2 \right) \right] \right\}$$

where the additional factor of 1/2 is from the density matrix now including plasma-electron spin, and  $s$  is the spin of the atomic electron. In addition, we have used the following for shorthand [46]

$$A_{u,u'}(k_1) = \frac{2}{\pi} \delta_{l_u l_{u'}} \sum_{l_1} \left[ \sum_k \langle u l_1 k_1 L S | D^k | u' l_1 k_1 L S \rangle + \langle u l_1 k_1 L S | E^k | u' l_1 k_1 L S \rangle \right] \quad (\text{B5})$$

$$B_{u,u'',u'}(k_1, k_2) = \frac{4}{\pi^2} \delta_{l_u, l_{u''}} \frac{1}{2} \sum_{l_1, l_2} \left[ \sum_k \langle u l_1 k_1 L S | D^k | u'' l_2 k_2 L S \rangle + \langle u l_1 k_1 L S | E^k | u'' l_2 k_2 L S \rangle \right] \times \\ \left[ \sum_k \langle u'' l_2 k_2 L S | D^k | u' l_1 k_1 L S \rangle + \langle u'' l_2 k_2 L S | E^k | u' l_1 k_1 L S \rangle \right] \quad (\text{B6})$$

$$\langle u l_1 k_1 L S | D^k | u'' l_2 k_2 L S \rangle = (-1)^{L+l_u+l_{u''}} \begin{pmatrix} l_u & k & l_{u''} \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l_1 & k' & l_2 \\ 0 & 0 & 0 \end{pmatrix} \begin{Bmatrix} l_u & l_1 & L \\ l_2 & l_{u''} & k \end{Bmatrix} \times \\ \int_0^\infty \int_0^\infty dr_2 dr_1 F_l(k_1, r_2) R_u(r_1) \left[ \frac{r_{>}^k}{r_{<}^{k+1}} - \frac{\delta_{k,0}}{r_1} \right] R_{u'}(r_1) F_l(k_1, r_2) \\ \langle u l_1 k_1 L S | E^k | u'' l_2 k_2 L S \rangle = (-1)^{S+l_u+l_{u''}} \begin{pmatrix} l_u & k & l_2 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l_1 & k' & l_{u''} \\ 0 & 0 & 0 \end{pmatrix} \begin{Bmatrix} l_u & l_1 & L \\ l_{u''} & l_2 & k \end{Bmatrix} \times \\ \int_0^\infty \int_0^\infty dr_2 dr_1 F_l(k_1, r_2) R_u(r_1) \left[ \frac{r_{>}^k}{r_{<}^{k+1}} - \frac{\delta_{k,0}}{r_1} \right] F_l(k_1, r_1) R_{u'}(r_2)$$

where we have labeled  $D$  for direct interaction and  $E$  for exchange interaction, and we have used the 3j and 6j symbols [44] to perform the angular integrals. The factor of 1/2 in  $B$  is due to the anti-symmetrization operator and the structure of the scattering “T”-matrix for hydrgeon-like systems [53–56].




FIG. 3. Comparison of line profile calculated with the electron-broadening model here vs the *Xenomorph* code [36]. For this comparison, the ion dynamics have been turned off in *Xenomorph* [36]. The two calculations use the same ion microfields so that the only difference between the codes is the electron-broadening.

We want to present some level of validation of the electron-broadening model presented here. This code has compared well with other validated codes in Alexiou *et al.* [57]. In addition, we compare this calculation with Gomez *et al.* [36] (which has compared well with the Wiese *et al.* [27] experiment), which we show in figure 3 for typical laboratory conditions. To do a proper comparison, the two methods have the same static-ion broadening so that the only difference between this work and *Xenomorph* [36] is the electron broadening. And, as shown, in figure 3, the two calculations give near identical results.

### Appendix C: Derivation of the Lippmann-Schwinger Recursion Relationship for the Electron-Broadening Operator

For purposes of comparison with the kinetic theory, and to link this work back to the Lippmann-Schwinger [48] “T”-matrix form, we want to write the electron-broadening operator in a recursion-relationship form. The electron-broadening operator,  $\mathcal{H}(\omega)$  is given as a function of  $\mathfrak{T}(\omega)$ , and both can be Taylor expanded,

$$\begin{aligned}\mathcal{H}(\omega) &= \frac{1}{1 + \langle \mathfrak{T}(\omega) \rangle (\omega - L_0^r)^{-1}} \langle \mathfrak{T}(\omega) \rangle \\ &= \langle \mathfrak{T}(\omega) \rangle \sum_{k=0}^{\infty} [-(\omega - L_0^r)^{-1} \langle \mathfrak{T}(\omega) \rangle]^k \\ \mathfrak{T}(\omega) &= L_I \frac{1}{1 - (\omega - L_0)^{-1} L_I} \frac{1}{\omega - L_0} \rho_0 \rho_r^{-1} (\omega - L_0^r). \\ &= L_I \sum_{k=0}^{\infty} [(\omega - L_0)^{-1} L_I]^k \frac{1}{\omega - L_0} \rho_0 \rho_r^{-1} (\omega - L_0^r).\end{aligned}$$

If we insert the Taylor-expanded definition for  $\mathfrak{T}(\omega)$  into the Taylor-expanded definition for  $\mathcal{H}(\omega)$  and collect all of the terms of the same  $L_I$ , the resulting equation for line-broadening is (out to third order),

$$\begin{aligned}\mathcal{H}(\omega) &= \langle L_I G(\omega) \rho_0 \rangle \rho_r^{-1} (\omega - L_0^r) \\ &+ \langle L_I G(\omega) L_I G(\omega) \rho_0 \rangle \rho_r^{-1} (\omega - L_0^r) - \langle L_I G(\omega) \rho_0 \rangle \rho_r^{-1} \langle L_I G(\omega) \rho_0 \rangle \rho_r^{-1} (\omega - L_0^r) \\ &+ \langle L_I G(\omega) L_I G(\omega) L_I G(\omega) \rho_0 \rangle \rho_r^{-1} (\omega - L_0^r) - \langle L_I G(\omega) \rho_0 \rangle \rho_r^{-1} \langle L_I G(\omega) L_I G(\omega) \rho_0 \rangle \rho_r^{-1} (\omega - L_0^r) \\ &- \langle L_I G(\omega) L_I G(\omega) \rho_0 \rangle \rho_r^{-1} \langle L_I G(\omega) \rho_0 \rangle \rho_r^{-1} (\omega - L_0^r) + \langle L_I G(\omega) \rho_0 \rangle \rho_r^{-1} \langle L_I G(\omega) \rho_0 \rangle \rho_r^{-1} \langle L_I G(\omega) \rho_0 \rangle \rho_r^{-1} (\omega - L_0^r),\end{aligned}\tag{C1}$$

where  $G(\omega) = (\omega - L_0)^{-1}$ . There is a common factor of  $\rho_r^{-1} (\omega - L_0^r)$  on the right-hand side of all the terms; this

can be factored out, and we can seek a function  $\hat{\mathcal{H}}(\omega)$  where

$$\mathcal{H}(\omega) = \text{Tr}_b\{\hat{\mathcal{H}}(\omega)\}\rho_r^{-1}(\omega - L_0^r) \quad (\text{C2})$$

Based on Eq (C1), we can write

$$\hat{\mathcal{H}}(\omega) = L_I(\omega - L_0)^{-1}\rho_0 + L_I(\omega - L_0)^{-1}[1 - P]\hat{\mathcal{H}}(\omega) \quad (\text{C3})$$

where  $P$  is a projection operator that performs a trace

over the perturber coordinates,

$$PA\rho_0 = \rho_0\rho_r^{-1}\text{Tr}_p\{A\rho_0\}. \quad (\text{C4})$$

This projection operator has slightly different properties than the one presented in Fano [9] or Smith and Hooper [40]. Because of the lack of correlations, the projection operators there commuted with  $(\omega - L_0)^{-1}$ —ours does not.

- 
- [1] R. E. Falcon, G. A. Rochau, J. E. Bailey, T. A. Gomez, M. H. Montgomery, D. E. Winget, and T. Nagayama, *Astrophys. J.* **806**, 214 (2015), arXiv:1505.03801 [astro-ph.SR].
  - [2] T. Nagayama, J. E. Bailey, R. C. Mancini, C. A. Iglesias, S. B. Hansen, C. Blancard, H. K. Chung, J. Colgan, P. Cosse, G. Faussurier, R. Florido, C. J. Fontes, F. Gilleron, I. E. Golovkin, D. P. Kilcrease, G. Loisel, J. J. MacFarlane, J.-C. Pain, G. A. Rochau, M. E. Sherrill, and R. W. Lee, *High Energy Density Physics* **20**, 17 (2016).
  - [3] M. J. Seaton, *Journal of Physics B Atomic Molecular Physics* **23**, 3255 (1990).
  - [4] C. A. Iglesias, *High Energy Density Physics* **18**, 14 (2016).
  - [5] I. Hubeny and D. Mihalas, *Theory of Stellar Atmospheres*, by I. Hubeny and D. Mihalas. Princeton, NJ: Princeton University Press, 2014 (2014).
  - [6] P.-E. Tremblay and P. Bergeron, *Astrophys. J.* **696**, 1755 (2009), arXiv:0902.4182 [astro-ph.SR].
  - [7] M. Baranger, *Physical Review* **112**, 855 (1958).
  - [8] A. C. Kolb and H. Griem, *Physical Review* **111**, 514 (1958).
  - [9] U. Fano, *Physical Review* **131**, 259 (1963).
  - [10] T. Hussey, J. W. Dufty, and C. F. Hooper, *Phys. Rev. A* **12**, 1084 (1975).
  - [11] R. Zwanzig, *J. Chem. Phys.* **33**, 1338 (1960).
  - [12] K. Burnett, J. Cooper, R. J. Ballagh, and E. W. Smith, *Phys. Rev. A* **22**, 2005 (1980).
  - [13] V. L. Jacobs, J. Cooper, and S. L. Haan, *Phys. Rev. A* **50**, 3005 (1994).
  - [14] V. L. Jacobs, *Phys. Rev. A* **90**, 033841 (2014).
  - [15] R. Zwanzig, *Nonequilibrium Statistical Mechanics* (OUP USA, 2001).
  - [16] N. N. Bogoliubov, *Journal of Experimental and Theoretical Physics* **16**, 691 (1946).
  - [17] J. G. Kirkwood, *The Journal of Chemical Physics* **14**, 180 (1946).
  - [18] J. G. Kirkwood, *The Journal of Chemical Physics* **15**, 72 (1947).
  - [19] M. Born and H. S. Green, *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences* **188**, 10 (1946).
  - [20] D. W. Ross, *Annals of Physics* **36**, 458 (1966).
  - [21] L. Klein, *J. Quant. Spec. Radiat. Transf.* **9**, 199 (1969).
  - [22] B. Bezzerides, *Physical Review* **159**, 3 (1967).
  - [23] R. Lee, *Journal of Physics A Mathematical General* **5**, 950 (1972).
  - [24] R. W. Lee, *Journal of Physics B Atomic Molecular Physics* **5**, 1271 (1972).
  - [25] U. Fano, *Reviews of Modern Physics* **29**, 74 (1957).
  - [26] D. B. Boercker and C. A. Iglesias, *Phys. Rev. A* **30**, 2771 (1984).
  - [27] W. L. Wiese, D. E. Kelleher, and D. R. Paquette, *Phys. Rev. A* **6**, 1132 (1972).
  - [28] T. W. Hussey, J. W. Dufty, and C. F. Hooper, Jr., *Phys. Rev. A* **16**, 1248 (1977).
  - [29] G. C. Junkel, M. A. Gunderson, C. F. Hooper, Jr., and D. A. Haynes, Jr., *Phys. Rev. E* **62**, 5584 (2000).
  - [30] M. A. Gunderson, G. C. Junkel-Vives, and C. F. Hooper, *J. Quant. Spec. Radiat. Transf.* **71**, 373 (2001).
  - [31] P. W. Anderson, *Physical Review* **76**, 647 (1949).
  - [32] H. R. Griem, *Spectral line broadening by plasmas* (New York, Academic Press, Inc. (Pure and Applied Physics. Volume 39), 1974. 421 p., 1974).
  - [33] R. Stamm, E. W. Smith, and B. Talin, *Phys. Rev. A* **30**, 2039 (1984).
  - [34] M. A. Gigoso and V. Cardeñoso, *Journal of Physics B Atomic Molecular Physics* **29**, 4795 (1996).
  - [35] E. Stambulchik and Y. Maron, *J. Quant. Spec. Radiat. Transf.* **99**, 730 (2006).
  - [36] T. A. Gomez, T. Nagayama, D. P. Kilcrease, M. H. Montgomery, and D. E. Winget, *Phys. Rev. A* **94**, 022501 (2016).
  - [37] P. Kepple and H. R. Griem, *Physical Review* **173**, 317 (1968).
  - [38] E. W. Smith, J. Cooper, and C. R. Vidal, *Physical Review* **185**, 140 (1969).
  - [39] C. R. Vidal, J. Cooper, and E. W. Smith, *ApJS* **25**, 37 (1973).
  - [40] E. W. Smith and C. F. Hooper, *Physical Review* **157**, 126 (1967).
  - [41] J. W. Dufty and D. B. Boercker, *J. Quant. Spec. Radiat. Transf.* **16**, 1065 (1976).
  - [42] M. Baranger, *Physical Review* **111**, 481 (1958).
  - [43] J. T. O'Brien and C. F. Hooper, Jr., *J. Quant. Spec. Radiat. Transf.* **14**, 479 (1974).
  - [44] R. D. Cowan, *The Theory of Atomic Structure and Spectra* (Los Alamos Series in Basic and Applied Sciences, Berkeley: University of California Press, 1981, 1981).
  - [45] L. A. Woltz and C. F. Hooper, Jr., *Phys. Rev. A* **30**, 468 (1984).
  - [46] H. Nguyen, M. Koenig, D. Benredjem, M. Caby, and G. Coulaud, *Phys. Rev. A* **33**, 1279 (1986).
  - [47] T. A. Gomez, *Improving Calculations of the Interaction Between Atoms and Plasma Particles and its Effect on Spectral Line Shapes*, Ph.D. thesis, University of Texas at Austin (2017).

- [48] B. A. Lippmann and J. Schwinger, Physical Review **79**, 469 (1950).
- [49] R. Lee, Journal of Physics B Atomic Molecular Physics **4**, 1640 (1971).
- [50] E. Stambulchik, D. V. Fisher, Y. Maron, H. R. Griem, and S. Alexiou, High Energy Density Physics **3**, 272 (2007).
- [51] D. P. Kilcrease, R. C. Mancini, and C. F. Hooper, Jr., Phys. Rev. E **48**, 3901 (1993).
- [52] H. A. Bethe and E. E. Salpeter, *Quantum Mechanics of One- and Two-Electron Atoms*, New York: Academic Press, 1957 (New York: Academic Press, 1957).
- [53] D. H. Madison, I. Bray, and I. McCarthy, Physical Review Letters **64**, 2265 (1990).
- [54] D. H. Madison, I. Bray, and I. E. McCarthy, Journal of Physics B Atomic Molecular Physics **24**, 3861 (1991).
- [55] D. H. Madison, R. P. McEachran, and M. Lehmann, Journal of Physics B Atomic Molecular Physics **27**, 1807 (1994).
- [56] M. L. Goldberger and K. M. Watson, *Collision theory*, by Goldberger, Marvin L.; Watson, Kenneth M. New York : Wiley, c1964. *Structure of matter series* (1964).
- [57] S. Alexiou, T. Stambulchik, E. Gomez, and M. Koubiti, “The Fourth Workshop on Lineshape Code Comparison: Line Merging,” (2018), in press.