



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Screening of an oscillating external electric field in atoms

V. A. Dzuba, J. C. Berengut, J. S. M. Ginges, and V. V. Flambaum

Phys. Rev. A **98**, 043411 — Published 5 October 2018

DOI: [10.1103/PhysRevA.98.043411](https://doi.org/10.1103/PhysRevA.98.043411)

Screening of oscillating external electric field in atoms

V. A. Dzuba,¹ J. C. Berengut,¹ J. S. M. Ginges,² and V. V. Flambaum¹

¹*School of Physics, University of New South Wales, Sydney, NSW 2052, Australia*

²*School of Mathematics and Physics, The University of Queensland, Brisbane QLD 4072, Australia*

We study the screening of a homogeneous oscillating external electric field E_0 in noble-gas atoms using atomic many-body calculations. At zero frequency of the oscillations ($\omega = 0$) the screened field $E(r)$ vanishes at the nucleus, $E(0) = 0$. However, the profile of the field $E(r)$ is complicated, with the magnitude of the field exceeding the external field E_0 at certain points. For $\omega > 0$ the field $E(r, \omega)$ strongly depends on ω and at some points may exceed the external field E_0 many times. The field at the nucleus is not totally screened and grows with ω faster than ω^2 . It can even be enhanced when ω comes close to resonance with a frequency of an atomic transition. This field interacts with CP-violating nuclear electric dipole moments creating new opportunities for studying them. The screening of the external field by atomic electrons may strongly suppress (or enhance near an atomic resonance) the low energy nuclear electric dipole transitions.

PACS numbers: 31.15.A-,11.30.Er

I. INTRODUCTION

Nuclear forces that violate the conservation of combined charge-conjugation and parity (CP) produce CP-violating nuclear moments that in turn may produce observable effects in atoms and molecules. The study of these effects provides a powerful probe of new physics beyond the Standard Model; see, e.g., reviews [1–5]. According to the Schiff theorem [6], the lowest-order CP-violating moment, the nuclear electric dipole moment (EDM), is unobservable in neutral atoms. Indeed, a neutral atom (and its nucleus) is not accelerated in a homogeneous static external electric field. Considering the nucleus to be point-like, the external electric field is completely screened at the nucleus by atomic electrons, and the nuclear EDM has nothing to interact with.

This screening is a big obstacle in the study of CP-violating nuclear forces. One has to go to higher-order moments or include some small corrections. For example, the screening is not complete if finite nuclear size is taken into account [6]. Indeed, while the total force on the nucleus is zero, the electric field does not have to vanish at each point across the nucleus. A convenient way to consider this effect is by introducing the so-called nuclear Schiff moment which induces atomic and molecular EDMs [7–10]. It can be roughly described as what is left from the nuclear EDM when the screening of the external electric field by electrons is taken into account. Refs. [7, 8] considered the effect of the proton EDM. The Schiff moments produced by CP-violating nuclear forces were introduced and calculated for the first time in [9, 10]. Among other possibilities is the atomic EDM generated by the nuclear magnetic quadrupole moment [9].

The electric field is not totally screened in ions. However, the strong constant electric field would remove ions from the trap. Another possibility is to use an oscillating electric field. It was stated in Ref. [11] that an oscillating electric field is not totally screened in atoms and may even lead to an enhancement of the field at the nucleus when the frequency of the field oscillations is close

to the frequency of an atomic transition. In a recent work [12], a formula was derived which states that the screened oscillating field at the nucleus is proportional to $\omega^2 \alpha_{zz}(\omega)$, where ω is the frequency of the oscillating field and $\alpha_{zz}(\omega)$ is the dynamic polarizability of the atom at this frequency. At sufficiently large frequencies the screening is significantly reduced. The field at the nucleus may even be enhanced in the resonance situation when the frequency of the external field is close to the frequency of an atomic transition.

In this paper we study the effect of screening of the external oscillating electric field in the noble-gas atoms numerically using the relativistic time-dependent Hartree-Fock method which is also known as the random-phase approximation. We demonstrate that the numerical calculations agree practically exactly with the formula for the screened electric field at the nucleus from Ref. [12]. Thus we have checked that the problem of finding the screened oscillating field in atoms is reduced to the calculations (or measurements) of the atomic dynamic polarizabilities. This in turn may lead to new ways of studying nuclear EDMs. Another application is the calculation of the effect of the electron screening on the probabilities of the nuclear electric dipole transitions.

In this paper we have also calculated the screened electric field inside an atom at all distances. The screened field oscillates and can actually exceed the external field.

II. CALCULATIONS

It has been shown in Ref. [12] that an external oscillating electric field is screened at the atomic nucleus with the value

$$E = \frac{E_0}{Z} [Z_i - \tilde{\omega}^2 \tilde{\alpha}_{zz}], \quad (1)$$

where E_0 is the amplitude of an external field directed along the z axis, Z is the nuclear charge, Z_i is the ionization degree, $\tilde{\omega} = \frac{\omega}{e^2/\hbar a_b}$ is the oscillation frequency ω in

atomic units, $\tilde{\alpha}_{zz} = \frac{\alpha_{zz}(\omega)}{a_b^3}$ is the dynamic polarizability of the atom $\alpha_{zz}(\omega)$ in atomic units, and a_b is the Bohr radius. For $Z_i = 0$ and $\omega = 0$, the field at the nucleus is totally screened, $E = 0$, in agreement with the Schiff theorem.

It is known that the Schiff theorem is fulfilled exactly in the random-phase approximation (RPA) [11]. RPA can be considered as a self-consistent Hartree-Fock approximation in a weak external field so that only terms linear in the external field are kept. It is also known that the RPA approximation gives very accurate values for the atomic polarizabilities for noble-gas atoms (see, e.g., Ref. [13] and below). This means that the use of the RPA method for noble-gas atoms is a good starting point for studying the screening of the external electric field in atoms. Note that for the closed-shell atoms the α_{zz} polarizability in Eq. (1) is just the scalar polarizability α_0 .

A. Random-phase approximation

We start from the Hartree-Fock equations for the single-electron orbital ψ_a (atomic units, $\hbar = m_e = |e| = 1$):

$$\begin{aligned} (\hat{H}_0 - \epsilon_a)\psi_a &= 0, \\ \hat{H}_0 &= c\boldsymbol{\alpha} \cdot \mathbf{p} + (\beta - 1)c^2 + V_{\text{nuc}} + \hat{V}. \end{aligned} \quad (2)$$

\hat{H}_0 is the relativistic Hartree-Fock Hamiltonian, $V_{\text{nuc}} \approx -Z/r$ is the finite-size nuclear potential, $\psi_a(\mathbf{r})$ is a four-component Dirac spinor,

$$\psi_a(\mathbf{r}) = \frac{1}{r} \begin{pmatrix} f(r)\Omega_{\kappa m} \\ ig(r)\Omega_{-\kappa m} \end{pmatrix}$$

and \hat{V} is the self-consistent electronic potential

$$\begin{aligned} \hat{V}\psi_a(\mathbf{r}) &= \sum_b \int d^3r' \frac{\psi_b^\dagger(\mathbf{r}')\psi_b(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} \psi_a(\mathbf{r}) \\ &\quad - \sum_b \int d^3r' \frac{\psi_b^\dagger(\mathbf{r}')\psi_a(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} \psi_b(\mathbf{r}) \end{aligned} \quad (3)$$

where the index b enumerates the electrons in the core. An applied weak periodic field

$$\hat{F} = \hat{f}^{-i\omega t} + \hat{f}^\dagger e^{i\omega t}, \quad (4)$$

modifies the atomic orbitals, adding to them small oscillating corrections

$$\tilde{\psi}_b = \psi_b + \chi_b e^{-i\omega t} + \eta_b e^{i\omega t}, \quad (5)$$

which can be found by solving the RPA equations

$$\begin{aligned} (\hat{H}_0 - \epsilon_b - \omega)\chi_b &= -(\hat{f} + \delta\hat{V})\psi_b, \\ (\hat{H}_0 - \epsilon_b + \omega)\eta_b &= -(\hat{f}^\dagger + \delta\hat{V}^\dagger)\psi_b, \end{aligned} \quad (6)$$

where $\delta\hat{V}$ is the correction to the self-consistent Hartree-Fock potential due to the external field. We consider the case where \hat{f} is the electric dipole operator (in length form $\hat{f} = z$). Equations (6) are solved self-consistently for all states in the core.

Detailed equations for (6) can be found in [14, 15]. Briefly, we expand the χ_b and η_b in partial waves (χ_β and η_β) with fixed angular momentum j_β and parity $\pi = (-1)^{l_\beta + K}$ for electric 2^K -pole excitations where K is the rank of \hat{f} ($K = 1$ for the electric dipole operator) and $|j_b - K| \leq j_\beta \leq j_b + K$. The reduced matrix elements required are

$$\begin{aligned} &\langle \chi_\alpha | \delta\hat{V} | \psi_a \rangle \\ &= \sum_{b\beta} \frac{\langle \kappa_\alpha | C^K | \kappa_a \rangle \langle \kappa_\beta | C^K | \kappa_b \rangle}{2K + 1} \end{aligned} \quad (7a)$$

$$\begin{aligned} &(R^K(\chi_\alpha \psi_b, \psi_a \chi_\beta) + R^K(\chi_\alpha \psi_b, \psi_a \eta_\beta)) \\ &+ \sum_{b\beta k} (-1)^{K+k} \langle \kappa_\beta | C^k | \kappa_a \rangle \langle \kappa_b | C^k | \kappa_\alpha \rangle \\ &\quad \left\{ \begin{matrix} j_a & j_\beta & k \\ j_b & j_\alpha & K \end{matrix} \right\} R^k(\chi_\alpha \psi_a, \psi_b \eta_\beta) \end{aligned} \quad (7b)$$

$$\begin{aligned} &+ \sum_{b\beta k} (-1)^{K+k} \langle \kappa_b | C^k | \kappa_a \rangle \langle \kappa_\alpha | C^k | \kappa_\beta \rangle \\ &\quad \left\{ \begin{matrix} j_a & j_b & k \\ j_\beta & j_\alpha & K \end{matrix} \right\} R^k(\chi_\alpha \psi_a, \chi_\beta \psi_b) \end{aligned} \quad (7c)$$

where b runs over core states and the β enumerate the partial waves of their respective corrections. The reduced spherical tensor matrix elements are defined by

$$\begin{aligned} \langle \kappa_a | C^k | \kappa_b \rangle &= (-1)^{j_a + 1/2} \sqrt{(2j_a + 1)(2j_b + 1)} \\ &\quad \begin{pmatrix} j_a & j_b & k \\ -1/2 & 1/2 & 0 \end{pmatrix} \xi(l_a + l_b + k) \end{aligned}$$

with $\xi(n) = [(-1)^n + 1]/2$, while the radial Slater integrals are defined

$$\begin{aligned} R^k(\psi_a \psi_b, \psi_c \psi_d) &= \int dr (f_a(r)f_c(r) + g_a(r)g_c(r)) Y_{\psi_b \psi_d}^k(r) \\ Y_{\psi_b \psi_d}^k(r) &= \int dr' \frac{r_{<}^k}{r_{>}^{k+1}} (f_b(r')f_d(r') + g_b(r')g_d(r')), \end{aligned}$$

where $r_{<} = \min(r, r')$ and $r_{>} = \max(r, r')$.

The conjugate equations $\langle \eta_\alpha | \delta\hat{V}^\dagger | \psi_a \rangle$ are similar to (7) but with $\chi_\alpha \rightarrow \eta_\alpha$ and $\chi_\beta \leftrightarrow \eta_\beta$ exchanged. In this work we are interested in the electric dipole polarizability, hence $K = 1$.

B. Scalar polarizability

The dynamic scalar polarizability of a closed-shell atom in the RPA method is given by

$$\alpha_0(\omega) = -\frac{1}{3} \sum_{b\beta} \left(\langle \psi_b | \hat{f} | \chi_\beta \rangle + \langle \psi_b | \hat{f} | \eta_\beta \rangle \right) \quad (8)$$

TABLE I. Static dipole polarizabilities of the noble-gas atoms calculated using the RPA method, experimental values for the polarizabilities, and the experimental positions of the first excitation which gives the dominant contribution to the polarizability.

Atom	α_0 (a.u.)		$\hbar\omega$ (cm ⁻¹) [22]
	RPA	Expt.	
He	1.322	1.383759 (13) [16]	169087
Ne	2.380	2.66110 (3) [17]	134042
Ar	10.77	11.083 (2) [18]	93750
Kr	16.47	16.740 [19]	80917
Xe	26.97	27.292 [20]	68045
Rn	35.00		55989

where b runs over core states. Note that one can use summation over a complete set of the single-electron basis states $|n\rangle$ to calculate the corrections χ_b and η_b

$$\chi_b = \sum_n \frac{\langle n | \hat{f} + \delta\hat{V} | b \rangle}{\epsilon_b - \epsilon_n + \omega} |n\rangle, \quad (9)$$

$$\eta_b = \sum_n \frac{\langle n | \hat{f}^\dagger + \delta\hat{V}^\dagger | b \rangle}{\epsilon_b - \epsilon_n - \omega} |n\rangle. \quad (10)$$

This would lead to a more commonly used expression for the dynamic polarizability of a closed-shell atom,

$$\alpha_\omega = -\frac{2}{3} \sum_{bn} \frac{(\epsilon_b - \epsilon_n) \langle b | \hat{f} | n \rangle \langle n | \hat{f} + \delta\hat{V} | b \rangle}{(\epsilon_b - \epsilon_n)^2 - \omega^2}. \quad (11)$$

Summation in (11) goes over occupied single-electron states b and vacant states n . We do not use expressions (9), (10) and (11) in the calculations of the present work. However, having these expressions is useful for a discussion of the polarizability behaviour near a resonance ($\omega \approx \epsilon_n - \epsilon_b$).

The induced electric potential inside the atom can be extracted from the direct term of (7a) as

$$\delta V(r) = \frac{1}{3} \sum_{b\beta} \langle \kappa_\beta | C^1 | \kappa_b \rangle \left(Y_{\psi_b \chi_\beta}^1(r) + Y_{\psi_b \eta_\beta}^1(r) \right), \quad (12)$$

and the total screened electric field inside the atom is given by

$$E(r) = E_0 + \varepsilon(r) = E_0 \left(1 + \frac{d}{dr} \delta V(r) \right). \quad (13)$$

Note that the derivatives of $Y_{\psi_b \psi_d}^1(r)$ can be expressed

$$\begin{aligned} \frac{d}{dr} Y_{\psi_b \psi_d}^1(r) &= -\frac{2}{r^3} \int_0^r r' (f_b(r') f_d(r') + g_b(r') g_d(r')) dr' \\ &+ \int_r^\infty (f_b(r') f_d(r') + g_b(r') g_d(r')) / r'^2 dr'. \end{aligned}$$

Static dipole polarizabilities of the noble-gas atoms calculated using the RPA equations (6) and (8) at $\omega = 0$ are




FIG. 1. Dynamic polarizabilities of noble-gas atoms calculated in the RPA approximation. Dots at $\omega = 0$ show experimental values for static polarizabilities.

presented in Table I and compared to the most accurate experimental values. The difference is only a few per cent and tends to be better for the heavier atoms. There is no experimental value for Rn, however our calculated value 35.00 a.u. agrees within 5.5% with the value 33.18 a.u. obtained in the more sophisticated coupled-cluster calculations of Ref. [21]. Table I also presents experimental energies of the first excitation from the ground state which gives the dominant contribution to the polarizability at small frequency. These energies decrease in value monotonically from He to Rn. This explains the larger polarizabilities for the heavier atoms and their faster increase with ω (see Fig. 1). Fig. 1 shows dynamic polarizabilities of the noble-gas atoms calculated in the RPA method using Eqs. (6) and (8).

Fig. 2 shows the screened electric field on the nuclei as a function of the frequency of the field oscillations. Note the excellent agreement between the two methods of calculation, using (1) or (13). For $\omega < 0.24$ a.u. the largest electric field on the nucleus is in the He atom, for $\omega > 0.24$ a.u. the largest field is in Xe and Rn. For most of the noble-gas atoms (with an exception of Ne) the screening is less than a tenth for $\omega > 0.3$ a.u. ($\lambda < 152$ nm). Note that the field can even be enhanced [12] when its frequency comes close to a resonance with an atomic transition. The numerical method used in this paper does not allow us to come close to a resonance. Therefore, we leave this for a future study.

In contrast to the formula (1) which gives the screened electric field at one point, $r = 0$, the formula (13) gives the screened electric field at any distance from the nucleus. Fig. 3 shows the screened electric field in Xe at two




FIG. 2. Screened electric field at the nuclei of He, Ne, Kr and Xe as a function of the frequency of the field oscillations. Solid line shows the result of the RPA calculations using formula (13), dots come from formula (1) with the calculated dynamic polarizability as in Fig. 1. The graphs for Ar and Rn are not shown because they are similar to those for Kr and Xe, respectively.




FIG. 3. Screened external electric field in Xe as a function of the distance (r is in atomic units). Dotted line corresponds to $\omega = 0$, solid line is for $\omega = 0.3$ a.u.

values of the frequency, $\omega = 0$ and $\omega = 0.3$ a.u. The field $E = E_0$ at large distances and its screened value at short distances is equal to what is given by formula (1). However, inside the atom the behaviour is very complicated, reflecting the shell structure of the atom and oscillations of the wave functions of external electrons. Note the




FIG. 4. Calculated screened external electric field at $r = 0$ in Xe as a function of the iteration number.

strong enhancement of the peaks at $\omega > 0$. This complicated behaviour is a collective effect caused by the fine tuning of electron orbitals affected by the external field and the change in other orbitals. The collectiveness of the effect is illustrated in Fig. 4. It shows the electric field at the nucleus of the Xe atom at $\omega = 0$ as a function of the iteration number. The iterations are used to solve the RPA equations (6) starting from $\chi = 0$ and $\eta = 0$. Each iteration corresponds to the next order of perturbation theory in the residual Coulomb interaction. It takes about twenty iterations to get the correct field at the nucleus. This illustrates that the effect is not perturbative and has a collective nature. Similar pictures for Tl^+ were presented in our earlier work [11]. Note that the figure captions were misplaced in that work.

For atoms other than the noble-gas atoms the correlations between external electrons and between external electrons and the core electrons play an important role (see, e.g., [13, 23]). These correlations are not included in the RPA calculations. This means that neither formula (8) for the dynamic polarizability nor formula (13) for the screened electric field are likely to give accurate results. However, the formula (1) was obtained without any assumptions about electron structure and should work well for any atom. This reduces the problem of screening to the problem of the dynamic polarizability of an atom which can be found from calculations or measurements.

For atoms with total angular momentum $J > 1$ in the ground state the scalar polarizability $\alpha_0(\omega)$ should be replaced by $\alpha_{zz}(\omega)$ (see Eq. (1)) which may have vector and tensor contributions. Calculation of the polarizabilities can be performed to very high accuracy for atoms with few valence electrons above closed shells (see, e.g.,

[13, 24]). However, even for atoms with more complicated electron structure, e.g. for atoms with an open f -shell, the polarizabilities can still be calculated with reasonably good accuracy [25, 26].

ACKNOWLEDGMENTS

This work was funded in part by the Australian Research Council through a Discovery Project DP150101405 and a Future Fellowship (J.G.) FT170100452 and by the National Science Foundation under grant No. NSF PHY11-25915. V.F. is grateful to Kavli Institute for Theoretical Physics at Santa Barbara for hospitality.

-
- [1] J. S. M. Ginges and V. V. Flambaum, *Phys. Rep.* **397**, 63 (2004).
- [2] M. Pospelov and A. Ritz, *Ann. Phys.* **318**, 119 (2005).
- [3] J. Engel, M. J. Ramsey-Musolf, and U. van Kolck, *Prog. Part. Nucl. Phys.* **71**, 21 (2013).
- [4] M.S. Safronova, D. Budker, D. DeMille, Derek F. Jackson Kimball, A. Derevianko, and Charles W. Clark, arXiv:1710.01833 (2017).
- [5] T. E. Chupp, P. Fierlinger, M. J. Ramsey-Musolf, and J. T. Singh, arXiv:1710.02504 (2017).
- [6] L. I. Schiff, *Phys. Rev.* **132**, 2194 (1963).
- [7] P. G. H. Sandars, *Phys. Rev. Lett.* **19**, 1396 (1967).
- [8] E. A. Hinds, and P. G. H. Sandars, *Phys. Rev. A* **21**, 471 (1980).
- [9] O. P. Sushkov, V. V. Flambaum, and I.B. Khriplovich, *Zh. Exp. Teor. Fiz.* **87**, 1521 (1984) [*Sov. JETP* **60**, 873 (1984)].
- [10] V.V. Flambaum, I.B. Khriplovich, and O.P. Sushkov, *Nucl. Phys. A* **449**, 750 (1986).
- [11] V. A. Dzuba, V. V. Flambaum, P. G. Silvestrov, and O. P. Sushkov, *Phys. Lett. A* **118**, 177 (1986).
- [12] V. V. Flambaum, arXiv:1804.08898 (2018).
- [13] J. Mitroy, M. S. Safronova, and C. W. Clark, *J. Phys. B: At. Mol. Phys.* **43**, 202001 (2010).
- [14] W. R. Johnson, C. D. Lin, K. T. Cheng, and C. M. Lee, *Physica Scripta* **21**, 409 (1980).
- [15] V. A. Dzuba, V. V. Flambaum, and O. P. Sushkov, *J. Phys. B: At. Mol. Phys.* **17**, 1953 (1984).
- [16] J. W. Schmidt, R. M. Gaiosio, E. F. May, and M. R. Moldover, *Phys. Rev. Lett.* **98**, 254504 (2007).
- [17] C. Gaiser and B. Fellmuth, *Europhys. Lett.* **90**, 63002 (2010).
- [18] A. C. Newell and R. C. Baird, *J. Appl. Phys.* **36**, 3751 (1965).
- [19] P. W. Langhoff and M. Karplus, *J. Op. Soc. Am.* **59**, 863 (1969).
- [20] A. Dalgarno and A. E. Kingston, *Proc. R. Soc. A* **259**, 424 (1960).
- [21] T. Nakajima and K. Hirao, *Chem. Lett.* **30**, 706 (2001).
- [22] A. Kramida, Yu. Ralchenko, J. Reader, and NIST ASD Team (2018). NIST Atomic Spectra Database (ver. 5.5.6), [Online]. Available: <https://physics.nist.gov/asd> [2018, April 27]. National Institute of Standards and Technology, Gaithersburg, MD.
- [23] V. A. Dzuba, V. V. Flambaum, and M. G. Kozlov, *Phys. Rev. A* **54**, 3948 (1996).
- [24] V. A. Dzuba and A. Derevianko, *J. Phys. B* **43**, 074011 (2010).
- [25] V. A. Dzuba, V. V. Flambaum, and B. L. Lev, *Phys. Rev. A* **83**, 032502 (2011).
- [26] V. A. Dzuba, A. Kozlov, and V. V. Flambaum, *Phys. Rev. A* **89**, 042507 (2014).