



# CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

## Emergent mode and bound states in single-component one-dimensional lattice fermionic systems

Yuchi He, Binbin Tian, David Pekker, and Roger S. K. Mong

Phys. Rev. B **100**, 201101 — Published 7 November 2019

DOI: [10.1103/PhysRevB.100.201101](https://doi.org/10.1103/PhysRevB.100.201101)

# Emergent mode and bound states in single-component one-dimensional lattice fermionic systems

Yuchi He,<sup>1,2,\*</sup> Binbin Tian,<sup>2,3,\*</sup> David Pekker,<sup>2,3</sup> and Roger S. K. Mong<sup>2,3</sup>

<sup>1</sup>*Department of Physics, Carnegie Mellon University, Pittsburgh, PA, 15213, USA*

<sup>2</sup>*Pittsburgh Quantum Institute, Pittsburgh, PA, 15260, USA*

<sup>3</sup>*Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh, PA, 15260, USA*

We study the formation of bound states in a one-dimensional, single-component Fermi chain with attractive interactions. The phase diagram, computed from DMRG (density matrix renormalization group), shows not only a superfluid of paired fermions (pair phase) and a liquid of three-fermion bound states (trion phase), but also a phase with two gapless modes. We show that the latter phase is described by a 2-component Tomonaga-Luttinger liquid (TLL) theory, consisting of one charged and one neutral mode. We argue based on our numerical data, that the single, pair, and trion phases are descendants of the 2-component TLL theory. We speculate on the nature of the phase transitions amongst these phases.

Tomonaga-Luttinger liquid (TLL) theory captures the physics of many 1-D quantum systems such as spin chains, spin ladders, nanotubes [1], nanowires [2], and cold atoms confined to 1-D tubes [3–7]. In higher dimensional systems, TLL is a tool that is often used, e.g. in edge theory [8] and coupled-wire constructions [9–11].

Recently, there has been significant interest in the study of 1D systems that cannot be described by the standard TLL theory [12–18]. In describing 1D interacting fermions, TLL theory naturally arises through bosonization that maps fermionic modes to bosonic modes. Nearby phases (i.e., descendants) such as charge density order appear as instabilities of the parent TLL theory [19–26]. This approach breaks down at the weak to strong pairing transition, i.e., the transition to the  $p$ -wave paired liquid. As recently pointed out in Ref. [27, 28], the  $p$ -wave pairing phase cannot be described as a descendant phase of a single-mode TLL; instead the transition is described by an emergent mode theory, with the weak and strong pairing phases being descendants of this theory. Which raises the question: what other phases, beyond  $p$ -wave pairing, can appear in one-component interacting fermions and how are these phases connected to some emergent-mode description?

In this paper, we investigate the formation of multi-fermion bound states in 1D single-component systems. We perform DMRG numerics on a lattice model with finite-range interactions, and find liquids of singles, pairs, trions, etc. in addition to an extended phase with two gapless modes (2M phase). We unify these findings by constructing an effective theory with an emergent mode that characterizes the 2M phase, the descendants of this theory describe the liquid phases of single fermions as well as multi-fermion bound states (i.e. bound states of 2, 3, 4, ... fermions). Our construction (Eq. (4)) is not equivalent to the band-bending construction in Ref. [28] (see supplement [29]) but is similar to Ref. [27] (see discussion at the end of the paper).

*Microscopic Model* – We study the lattice Hamiltonian

$$H = \sum_i \left[ -\frac{1}{2} (c_i^\dagger c_{i+1} + c_{i+1}^\dagger c_i) + \sum_{m=1}^3 V_m n_i n_{i+m} \right], \quad (1)$$

where  $c_i$  and  $c_i^\dagger$  are the fermion annihilation and creation operators at lattice site  $i$ ,  $n_i = c_i^\dagger c_i$  is the number operator, and  $V_m$  defines the shape of the fermion-fermion interaction potential. We choose short-ranged attractive interactions ( $V_1 < 0$  and  $V_2 < 0$ ) to promote the formation of pairs and trions, but with  $V_3 > 0$  to prevent phase-separation [27]. To decrease the parameter space we restrict our attention to the subspace  $V_1 = V_2$ . We expect that extending the range of attractive interactions will result in more liquid phases of multi-fermion bound states. For example, we demonstrate that extending the attractive interactions to three sites results in a quaternion liquid phase [29].

We use iDMRG [30–32] to study the ground state properties of the Hamiltonian (1) with focus on the 1/5 filling. The accuracy of iDMRG is controlled by the bond dimension  $\chi$ , the result becomes exact as  $\chi \rightarrow \infty$  [29]. To identify the various phases, we use two types of diagnostics: central charge  $c$  and various two-point correlators.

We obtain  $c$  as follows. We study the bipartite entanglement entropy  $S$ , i.e., the von Neumann entropy of DMRG ground state traced over either half the system. Both  $S$  and the correlation length  $\xi$  are infinite for the true ground state, but are cut off by finite  $\chi$ . The manner in which these two variables diverge gives the central charge:  $S = \frac{c}{6} \log(\xi) + \text{const}$  [33, 34].

We also compute the single, pair, and trion two-point correlators

$$G_1(r) = \langle c_i^\dagger c_{i+r} \rangle, \quad (2a)$$

$$G_2(r) = \langle (c_i c_{i+1})^\dagger c_{i+r} c_{i+r+1} \rangle, \quad (2b)$$

$$G_3(r) = \langle (c_i c_{i+1} c_{i+2})^\dagger c_{i+r} c_{i+r+1} c_{i+r+2} \rangle. \quad (2c)$$




FIG. 1. Central charge as a function of interactions in the lattice model (1) computed at filling fraction  $1/5$ . Four phases labeled are identified with further analysis of correlation functions. We have checked that all the reported phases exist for  $V_1 = 2V_2$  and we expect the qualitative features of the phase diagram to hold for generic values of  $V_1 \neq V_2$  in the vicinity of  $V_1 = V_2$ .

In the single phase all correlators decay algebraically; in the pair phase only  $G_2$  decays algebraically while  $G_1$  and  $G_3$  decay exponentially; in the trion phase  $G_3$  is algebraic while  $G_1$  and  $G_2$  are exponential. This behavior implies that there is a gap to adding a single fermion into the pair/trion phase but no gap to adding two/three fermions.

Figure 1 shows  $c$  as a function of the interaction parameters  $V_1 = V_2$  and  $V_3$ . The blue regions denote the single-mode phases with  $c = 1$ , we identify these as single, pair, and trion phases based on their two-point correlators [Eq. (2)]. While we observe a direct transition between the pair and single phases [27, 28], we do not find a direct transition between the pair and trion phases; instead we find an intermediate phase with  $c \approx 2$  which we call the 2M (2-mode) phase. The 2M phase neighbors all other phases and indicates a parent theory with an emergent mode, which enables a unified description of the multi-fermion bound-state phases and their transitions.

Our strategy for the remainder of the paper is as follows. First, we introduce a field theory to describe the 2M phase, writing down its operators and free Hamiltonian. Next, we introduce the possible interaction terms, and examine the descendant phases that result. Finally, we show that these prediction are consistent with our numerics, justifying our theoretical model.

*Theory of the emergent mode* – Motivated by Refs. [27, 28] and our data, we introduce a theory with two modes. In this theory, the charge-1 operators in the lowest har-

monic are [35]:

$$\begin{aligned}\psi_{0,\eta}^{(\pm)} &= e^{\pm i\theta_1} e^{i\theta_0 + i\eta(\phi_0 + k_F x)}, \\ \psi_{1,\eta}^{(\pm)} &= e^{i\theta_0} e^{\pm i\theta_1 \pm i\eta(\phi_1 + k' x)},\end{aligned}\quad (3)$$

where  $\eta = +1/-1$  denotes a right/left mover;  $\theta_\mu$  is dual field of the compact bosonic field  $\phi_\mu$  and satisfies  $[\partial_x \theta_\mu(x), \phi_\nu(x')] = i\pi \delta_{\mu\nu} \delta(x - x')$ . The charge is carried by the  $\theta_0$  mode, while  $\theta_1$  is neutral, as a result  $k_F$  is fixed by the density of electrons while  $k'$  is a free parameter.

The set of local physical operators can be generated via products of operators from Eq. (3), i.e.,  $(\psi_{0,1}^+)^l (\psi_{0,-1}^+)^m (\psi_{1,1}^-)^n \dots$ . (Note that the generators Eq. (3) are over-complete.) As a result, primary operators of charge  $q$  take the form:

$$c(x)^q \sim \sum_{q_1, r_0, r_1} e^{i(q\theta_0 + q_1\theta_1 + r_0(\phi_0 + k_F x) + r_1(\phi_1 + k' x))}, \quad (4)$$

$$\text{where } q_1 \equiv r_0 + r_1 \equiv q \pmod{2}.$$

Due to the restrictions on the coefficients  $q_1$ ,  $r_0$ , and  $r_1$  of physical operators, we cannot simply treat this theory as a product of decoupled  $\theta_0/\phi_0$  and  $\theta_1/\phi_1$  theories.

The theory must obey charge conservation, and be invariant under both parity ( $\phi_{0,1} \rightarrow -\phi_{0,1}$  and  $x \rightarrow -x$ ) and time-reversal ( $\theta_{0,1} \rightarrow -\theta_{0,1}$ ,  $i \rightarrow -i$ , and  $t \rightarrow -t$ ). The kinetic part of the Hamiltonian takes the form:

$$\mathcal{H}_{\text{KE}} = \sum_{\mu,\nu} [A_{\mu\nu}(\partial_x \theta_\mu)(\partial_x \theta_\nu) + B_{\mu\nu}(\partial_x \phi_\mu)(\partial_x \phi_\nu)]. \quad (5)$$

$\mathcal{H}_{\text{KE}}$  describes a 2-mode TLL, which we later demonstrate to be consistent with the 2M phase found in the numerics.

*Single-mode phases as descendants of the 2M theory* –

The single-mode phases (single, pair, trion, ...) are constructed by introducing locking terms, shown in Table I, to the Hamiltonian (5). For a term to appear, it must be of the form of Eq. (4) with  $q = 0$ , and also respect parity and time-reversal. At large interaction strength, some of these terms may ‘lock’ [26]; taking an expectation value and reducing the theory to a 1-component TLL.

Our analysis for the locking terms follows [36]. For an interaction term to lock it should have no oscillation (i.e.,  $x$  dependence), which places constraints on the Fermi momenta. For each locking term of the form  $\cos \Lambda$ , we find linear combinations of the  $\theta$ s and  $\phi$ s that commute with  $\Lambda$ . Among this set we find a conjugate pair which we denote as  $\theta_+$  and  $\phi_+$ . The set of gapless operators are then generated by  $e^{i\Lambda}$ ,  $e^{i\theta_+}$ , and  $e^{i\phi_+}$ , and must be a subset of Eq. (4) [37]. We show that the minimal (unit) charges for these operators are indeed  $q_{\min} = 1, 2, 3$  for the single, pair, trion phases respectively, from the given locking terms. We extend our analysis to arbitrary  $q_{\min}$  in the supplement [29].

Locking term	$\cos(2\theta_1)$	$\cos(2k'x + 2\phi_1)$	$\cos[(3k' - k_F)x + 3\phi_1 - \phi_0]$	...
Resulting phase	single	pair	trion	...
Single correlator $G_1(r)$	$\sum_n \frac{\sin[(2n+1)k_F r ]}{ r ^{(1/K+(2n+1)^2K)/2}}$			
Pair correlator $G_2(r)$	$\sum_n \frac{\cos[(2n)k_F r ]}{ r ^{2/K+2n^2K}}$	$\sum_n \frac{\cos[(2n)\frac{k_F}{2} r ]}{ r ^{(1/K+(2n)^2K)/2}}$		
Trion correlator $G_3(r)$	$\sum_n \frac{\sin[(2n+1)k_F r ]}{ r ^{(9/K+(2n+1)^2K)/2}}$		$\sum_n \frac{\sin[(2n+1)\frac{k_F}{3} r ]}{ r ^{(1/K+(2n+1)^2K)/2}}$	

TABLE I. Locking terms and correlators of single-mode phases. The first line lists interaction terms and the second line shows the corresponding phases when interaction terms get locked. The remaining rows show the algebraic decay form of correlators  $G_{1,2,3}$ ; the coefficient of each term is neglected for simplicity. Figure 2 shows the numeric data verifying the predicted dependence.

We first analyze the locking term  $\cos(2\theta_1)$  which induces the single phase. The gapless mode is described by the dual fields  $\theta_+ = \theta_0$  and  $\phi_+ = \phi_0$ . Thus the gapless operators take the form  $c(x) \sim \sum e^{ia\theta_1} e^{i\theta_0} e^{i(2n+1)(\phi_0+k_F x)}$  where  $n$  is an integer and  $a$  an odd integer. (The dual field  $\phi_1$  is disordered and cannot appear here.) As  $e^{i\theta_1}$  is a constant,  $c(x)$  reduces to the standard bosonization form of a fermion mode [26, 38].

Next, we show that the locking term  $\cos(2\phi_1 + 2k'x)$  induces the pair phase. Notably, for this term to lock we must enforce  $k' = 0$ . As  $\theta_1$  is disordered, it cannot appear in a gapless operator, i.e.,  $q_1 = 0$ . From the parity relation (4), we see that  $q$  must be an even integer and thus the single and trion correlators decay exponentially. Letting  $\theta_+ = 2\theta_0$  and  $\phi_+ = \phi_0/2$ , we recover the standard bosonization expansion of a boson mode [26, 38] for the pair operator:  $c(x)^2 \sim b(x) \approx \sum e^{i\theta_+} e^{i(2n)(\phi_+ + k_B x)}$  with  $n \in \mathbb{Z}$  and  $k_B = k_F/2$ . We interpret this descendant theory as a TLL of fermion pairs, with the density of pairs being half of the density of elementary fermions.

Finally, we address the locking term  $\Lambda = 3(\phi_1 + k'x) - (\phi_0 + k_F x)$  which yields the trion phase while fixing  $k' = k_F/3$ . As  $\Lambda$  commutes with  $\theta_+ = 3\theta_0 + \theta_1$  and  $\phi_+ = \phi_1$ , the gapless operators take the form  $c(x)^q \sim \sum e^{i(q/3)\theta_+} e^{ia(\phi_+ + k'x) + ibL}$ . Mapping the expression to Eq. (4), we get  $q_1 = q/3$ ,  $r_0 = -b$ , and  $r_1 = a + 3b$ ; we determine the consistency conditions  $q/3, a, b \in \mathbb{Z}$  and  $a \equiv q \pmod{2}$ . Hence for any gapless operator,  $q$  must be a multiple of 3, which implies exponential decay of  $G_1$  and  $G_2$ . The trion operator expansion reduces to:  $c(x)^3 \sim \sum e^{i\theta_+} e^{i(2n+1)(\phi_+ + k'x)}$ , where  $k' = k_F/3$  is the Fermi wavevector of the trions and  $n$  is an integer.

Within the low energy theory for each of the three single TLL mode phases,  $c(x)^{q_{\min}}$  admits a standard bosonization expansion in terms of  $\theta_+$  and  $\phi_+$ . The effective Hamiltonian is thus

$$\mathcal{H}_+ = \frac{v_+}{2\pi} [K(\partial_x \theta_+)^2 + \frac{1}{K}(\partial_x \phi_+)^2], \quad (6)$$

where  $K$  is the Luttinger parameter.

*Fourier spectra of the correlators* – The long-distance




FIG. 2. Verification of predicted decay exponents. Left: leading decay exponent ( $\eta_2$ ) of pair correlator in pair phase at  $V_1 = V_2 = -0.8$ ,  $V_3 = 1.4$ . Right: leading decay exponent ( $\eta_3$ ) of trion correlator in trion phase at  $V_1 = V_2 = -1$ ,  $V_3 = 1.4$ . The two lines are prediction from TLL theory (cf. Tab. I),  $\eta_2 = \frac{1}{2K}$  and  $\eta_3 = \frac{1}{2}(K + \frac{1}{K})$ . The values of Luttinger parameter  $K$  are extracted from the neutral sector [29]. In order to cover larger range of  $K$ , we use DMRG data from fillings (left to right)  $\frac{1}{5}, \frac{1}{6}, \dots, \frac{1}{10}$ .

behavior of the correlation functions of gapless operators can be written as a sum of algebraically decaying terms of the form

$$\frac{\cos(k_{\text{osc}}|r| + \varphi)}{|r|^\eta}. \quad (7)$$

Our theory puts a restriction on the allowed values of  $k_{\text{osc}}$  in the 2M phase and the single-mode phases. Table I summarizes the long distance behavior of the correlation functions within the single-mode phases; observe that the (leading) decay exponents  $\eta$  of all harmonics  $k_{\text{osc}}$  depend only on the Luttinger parameter  $K$ ; this is verified in Fig. 2.

To connect the effective theory to our microscopic model, we compare the  $k_{\text{osc}}$  in correlation functions obtained from field theory and DMRG. We perform Fourier transforms on the correlation functions  $G_{1,2,3}(r)$  and take the  $n^{\text{th}}$  derivative, such that terms of the form Eq. (7) with  $\eta < n + 1$  will show a divergent peak at  $k_{\text{osc}}$ . We then match the set of predicted oscillation wavevectors to peaks in the Fourier transforms. Figure 3 presents the correlation functions along cuts at constant  $V_3$ . Panels (a-c) show a cut through the trion, 2M and pair phases;




FIG. 3. Spectra  $G_1$ ,  $G_2$  and  $G_3$  (from top to bottom) as a function of wavevector and interaction strength ( $V_1 = V_2$ ), showing agreement of peak locations between DMRG and theory. The data is taken at cuts shown in Fig. 1. Plots (a–c) taken at  $V_3 = 1.56$  show the trion, 2M, and pair phases; plots (d–f) taken at  $V_3 = 1.3$  show the trion and single phases (with a possible 2M phase in between). Darker (Blue) colors represent larger values of amplitudes. The peak in the data of  $G_1$ , which continuously varying between 0 and  $k_F/3$  in the 2M/single phase is identified as  $k'$ . The lines added to the color plot are theoretic predictions with the determined parameter  $k'$ . The solid lines denote several long distance  $k_{\text{osc}}$  associated with algebraic-decay; the dotted lines denote several exponential-decay “peaks”, which are possibly visible if the decay-length-scale is large. The parameters for the plots are explained in the supplement [29].

while panels (d–f) cut from trion to single phase (with a possible intervening 2M phase).

In the 2M phase, both modes are gapless and the allowed  $k_{\text{osc}}$ 's are given by the oscillatory part of  $c(x)^q$  in Eq. (4):

$$k_{\text{osc}} = r_0 k_F + r_1 k'. \quad (8)$$

For  $G_1$  and  $G_3$ ,  $r_0 + r_1$  is odd, hence the first several  $k_{\text{osc}}$  are  $k'$ ,  $k_F$ ,  $k_F \pm 2k'$ , and  $2k_F \pm k'$ . For  $G_2$ ,  $r_0 + r_1$  is even and so  $k_{\text{osc}} = 0$ ,  $k_F \pm k'$ ,  $2k'$ ,  $2k_F$ , etc. These

wavevectors are fitted to numerical data and are marked by the dotted lines in Fig. 3. As  $k_F$  is fixed,  $k'$  is the only fitting parameter at each point of phase space. In the 2M region of panels (a–c), we observe unambiguous peaks at the predicted wavevectors. (In the numerics, we have not resolved peaks at some of the predicted  $k_{\text{osc}}$ , as these peaks are too weak or have large exponent  $\eta$ .)

A key feature of the DMRG data in the 2M phase is that  $k'$  varies continuously between the two limiting values:  $k' = k_F/3$  on the trion side and  $k' = 0$  on the pair side. The variation of this wavevector is a clear sign of a neutral emergent mode and confirms our effective two-mode TLL [39].

In the single-mode phases, some of the peaks found for the 2M phase persist while others are no longer divergent as modes become gapped out. The trion phase is characterized by the absence of singular behavior in panels (a,b,d,e) as  $G_{1,2}$  decay exponentially. We observe that  $G_3$  decays algebraically with peaks in Fig. 3(c,f) at odd multiples of  $k' = k_F/3$ . In the pair phase,  $k' = 0$  and only  $G_2$  shows divergent peaks at multiples of  $k_F$ , as predicted. [The features at 0,  $k_F$  in panel (a) are not divergent and broadened out due to  $G_1$  being gapped. Deep in the pair phase, they become invisible.] Finally, for the single phase all three correlators are algebraically decaying [Fig. 3(d–f)] with peaks at multiples of  $k_F$ . Remarkably, we also observe exponentially decaying features at the moving  $k'$  which are remnants of the 2M parent theory.

*Phase transitions* – There are five potential phase transitions in our phase diagram. The locking mechanisms give hints about the possible phase transitions, which we discuss in relation to our data.

- Single-pair transition. The transition is controlled by the competition between the terms  $\cos(2\theta_1)$  and  $\cos(2\phi_1)$ , and results in a quantum Ising transition [27, 28, 40–42]. In the supplement [29], we provide the definitive evidence that the single-pair transition is Ising via finite- $\chi$  scaling.
- 2M-single transition. This transition is driven by the term  $\cos(2\theta_1)$ , and is likely a Berezinskii-Kosterlitz-Thouless (BKT) transition.
- 2M-pair/trion transition. Both 2M-to-pair and 2M-to-trion transitions are accompanied by  $k'$  reaching a commensurate value. This suggests a commensurate-incommensurate transition
- Single-trion transition. We are unable to determine if there is a direct transition between the trion and single phase, or whether there is an intervening 2M phase which extends down as  $V_3$  is decreased. In both cases, our numerical analysis suggests a (at least one) first-order transition (see the supplement [29]).

*Discussion* – In summary we find conclusive evidence for an emergent mode in a one-dimensional attractive

fermion chain. This emergent mode results in the formation of a stable 2M phase with two Fermi surfaces. We argue that the multi-fermion bound state liquids are not descendants of the single-mode TLL phase but are rather descendants of this 2M phase. Here the 2M parent theory is written as a mixture of charged/neutral modes. Curiously, we can also rewrite the theory in terms of a mixture of charge-1/charge-2 modes [27], or more generally charge- $n$ /charge- $(n + 1)$  modes.

The two ingredients required to realizing the proposed phenomenology are (1) confining the fermions to one-dimension and (2) controlling the form of the interaction potential between the fermions. In the setting of solid state systems the two ingredients could be realized in nanowires made of superconducting semiconductors [43–48]. In ultracold atoms confinement could be provided by either optical lattices [3, 6, 49], or atom chips [5] and tunable long-range interaction by the use of dipolar interactions [50, 51], or Rydberg state-mediated interactions [52].

The 1D systems studied here can also be used to construct higher dimensional topological phases via the coupled-wire construction [9, 27, 53, 54]. TLL enriched by emergent mode(s) may give a pathway to a wide range of new phases in condensed matter.

We acknowledge enlightening discussions with J. Levy. This work was supported by the Charles E. Kaufman foundation and NSF PIRE-1743717.

---

\* These two authors contributed equally

- [1] M. Bockrath, D. H. Cobden, J. Lu, A. G. Rinzler, R. E. Smalley, L. Balents, and P. L. McEuen, *Nature* **397**, 598 (1999).
- [2] V. V. Deshpande, M. Bockrath, L. I. Glazman, and A. Yacoby, *Nature* **464**, 209 (2010).
- [3] T. Kinoshita, T. Wenger, and D. S. Weiss, *Science* **305**, 1125 (2004).
- [4] S. Capponi, G. Roux, P. Lecheminant, P. Azaria, E. Boulat, and S. White, *Physical Review A* **77**, 013624 (2008).
- [5] M. Gring, M. Kuhnert, T. Langen, T. Kitagawa, B. Rauer, M. Schreitl, I. Mazets, D. A. Smith, E. Demler, and J. Schmiedmayer, *Science* **337**, 1318 (2012).
- [6] T. L. Yang, P. Grišins, Y. T. Chang, Z. H. Zhao, C. Y. Shih, T. Giamarchi, and R. G. Hulet, *Phys. Rev. Lett.* **121**, 103001 (2018).
- [7] G. Salomon, J. Koepsell, J. Vijayan, T. Hilker, J. Nespolo, L. Pollet, I. Bloch, and C. Gross, *Nature* **565** (2018), 10.1038/s41586-018-0778-7.
- [8] X.-G. Wen, *Advances in Physics* **44**, 405 (1995).
- [9] C. L. Kane, R. Mukhopadhyay, and T. C. Lubensky, *Phys. Rev. Lett.* **88**, 036401 (2002).
- [10] M. Cazalilla, A. Ho, and T. Giamarchi, *New Journal of Physics* **8**, 158 (2006).
- [11] D. F. Mross, J. Alicea, and O. I. Motrunich, *Phys. Rev. Lett.* **117**, 016802 (2016).
- [12] A. Imambekov and L. I. Glazman, *Science* **323**, 228 (2009).
- [13] A. Imambekov, T. L. Schmidt, and L. I. Glazman, *Reviews of Modern Physics* **84**, 1253 (2012).
- [14] M. Mattioli, M. Dalmonte, W. Lechner, and G. Pupillo, *Physical review letters* **111**, 165302 (2013).
- [15] M. Dalmonte, W. Lechner, Z. Cai, M. Mattioli, A. Läuchli, and G. Pupillo, *Physical Review B* **92**, 045106 (2015).
- [16] S. Rossotti, M. Teruzzi, D. Pini, D. E. Galli, and G. Bertaina, *Phys. Rev. Lett.* **119**, 215301 (2017).
- [17] T. Scaffidi, D. E. Parker, and R. Vasseur, *Phys. Rev. X* **7**, 041048 (2017).
- [18] H.-C. Jiang, Z.-X. Li, A. Seidel, and D.-H. Lee, *Science Bulletin* **63**, 753 (2018).
- [19] A. Luther and V. J. Emery, *Phys. Rev. Lett.* **33**, 589 (1974).
- [20] T. Giamarchi and H. Schulz, *Journal de Physique* **49**, 819 (1988).
- [21] I. Affleck, *Nuclear Physics B* **305**, 582 (1988).
- [22] K. Penc and F. Mila, *Physical Review B* **49**, 9670 (1994).
- [23] P. Lecheminant, E. Boulat, and P. Azaria, *Phys. Rev. Lett.* **95**, 240402 (2005).
- [24] E. Burovski, G. Orso, and T. Jolicoeur, *Phys. Rev. Lett.* **103**, 215301 (2009).
- [25] G. Orso, E. Burovski, and T. Jolicoeur, *Phys. Rev. Lett.* **104**, 065301 (2010).
- [26] G. Roux, E. Burovski, and T. Jolicoeur, *Physical Review A* **83**, 053618 (2011).
- [27] C. L. Kane, A. Stern, and B. I. Halperin, *Phys. Rev. X* **7**, 031009 (2017).
- [28] J. Ruhman and E. Altman, *Phys. Rev. B* **96**, 085133 (2017).
- [29] See Supplemental Material at for numerical methods, additional numerical data, and discussions of the 2M theory.
- [30] S. R. White, *Phys. Rev. Lett.* **69**, 2863 (1992).
- [31] I. P. McCulloch, arXiv preprint arXiv:0804.2509 (2008).
- [32] J. A. Kjäll, M. P. Zaletel, R. S. K. Mong, J. H. Bardarson, and F. Pollmann, *Phys. Rev. B* **87**, 235106 (2013).
- [33] P. Calabrese and J. Cardy, *Journal of Statistical Mechanics: Theory and Experiment* **2004**, P06002 (2004).
- [34] F. Pollmann, S. Mukerjee, A. M. Turner, and J. E. Moore, *Physical review letters* **102**, 255701 (2009).
- [35] We omit Klein factors for notational simplicity.
- [36] A. O. Gogolin, A. A. Nersesyan, and A. M. Tsvelik, *Bosonization and strongly correlated systems* (Cambridge university press, 2004).
- [37] Although the operators  $e^{ib\Lambda}$  are not gapless, they are long-ranged correlated (ordered).
- [38] F. Haldane, *Physical Review Letters* **47**, 1840 (1981).
- [39] We discuss the incompatibility of this result and Ref. [28] in the supplement [29].
- [40] P. Lecheminant, A. O. Gogolin, and A. A. Nersesyan, *Nuclear Physics B* **639**, 502 (2002).
- [41] M. Sitte, A. Rosch, J. Meyer, K. Matveev, and M. Garst, *Physical review letters* **102**, 176404 (2009).
- [42] O. Alberton, J. Ruhman, E. Berg, and E. Altman, *Physical Review B* **95**, 075132 (2017).
- [43] V. Mourik, K. Zuo, S. M. Frolov, S. Plissard, E. P. Bakkers, and L. P. Kouwenhoven, *Science* **336**, 1003 (2012).
- [44] J. Chen and P. Yu, *Sci. Adv.* **3**, e1701476 (2017).
- [45] M. Deng, S. Vaitiekėnas, E. B. Hansen, J. Danon, M. Lei-

- juse, K. Flensberg, J. Nyg ard, P. Krogstrup, and C. M. Marcus, *Science* **354**, 1557 (2016).
- [46] A. Annadi, G. Cheng, H. Lee, J.-W. Lee, S. Lu, A. Tylan-Tyler, M. Briggeman, M. Tomczyk, M. Huang, D. Pekker, C.-B. Eom, P. Irvin, and J. Levy, *Nano Letters* **18**, 4473 (2018).
- [47] G. Cheng, M. Tomczyk, S. Lu, J. P. Veazey, M. Huang, P. Irvin, S. Ryu, H. Lee, C.-B. Eom, C. S. Hellberg, *et al.*, *Nature* **521**, 196 (2015).
- [48] G. Cheng, M. Tomczyk, A. B. Tacla, H. Lee, S. Lu, J. P. Veazey, M. Huang, P. Irvin, S. Ryu, C.-B. Eom, *et al.*, *Physical Review X* **6**, 041042 (2016).
- [49] I. Bloch, *Rev. Mod. Phys.* **80**, 885 (2008).
- [50] M. A. Baranov, M. Dalmonte, G. Pupillo, and P. Zoller, *Chemical Reviews* **112**, 5012 (2012).
- [51] C. Gross and I. Bloch, *Science* **357**, 995 (2017).
- [52] J. Zeiher, J.-y. Choi, A. Rubio-Abadal, T. Pohl, R. van Bijnen, I. Bloch, and C. Gross, *Physical Review X* **7**, 041063 (2017).
- [53] J. C. Y. Teo and C. L. Kane, *Phys. Rev. B* **89**, 085101 (2014).
- [54] R. S. Mong, D. J. Clarke, J. Alicea, N. H. Lindner, P. Fendley, C. Nayak, Y. Oreg, A. Stern, E. Berg, K. Shtengel, *et al.*, *Physical Review X* **4**, 011036 (2014).