



This is the accepted manuscript made available via CHORUS. The article has been published as:

Microscopic theory of field-tuned topological transitions in the Kitaev honeycomb model

Jagannath Das, Aman Kumar, and Vikram Tripathi

Phys. Rev. B **111**, L041111 — Published 21 January 2025

DOI: [10.1103/PhysRevB.111.L041111](https://doi.org/10.1103/PhysRevB.111.L041111)

Microscopic theory of field tuned topological transitions in the Kitaev honeycomb model

Jagannath Das,^{1,*} Aman Kumar,^{2,3,†} and Vikram Tripathi^{1,‡}

¹*Department of Theoretical Physics, Tata Institute of Fundamental Research,
Homi Bhabha Road, Colaba, Mumbai 400005, India.*

²*National High Magnetic Field Laboratory, Tallahassee, Florida 32310, USA*

³*Department of Physics, Florida State University, Tallahassee, Florida 32306, USA*

We microscopically construct an abelian mutual Chern-Simons lattice gauge theory for magnetic field-tuned topological transitions in the Kitaev model and obtain a complete characterization of the phases, including their quasiparticles. At low fields for both ferro and antiferromagnetic (FM(AFM)) Kitaev interactions, we demonstrate nonabelian Ising topological order (ITO), and explicitly construct the Majorana anyon as an intrinsic excitation – a twist defect in our *abelian* gauge theory. For the AFM case, an abelian chiral phase appears at intermediate fields with trivial topological order and fermionic bulk excitations. Remarkably, both the ITO phase and the intermediate phase have the same chiral central charge $c = 1/2$, implying no change in the quantized thermal Hall response across the transition. For the FM case, there is a direct transition from ITO to a partially polarized nontopological phase. Our study completes the proof of Kitaev’s original proposal of the low-field ITO with $c = 1/2$ going beyond his mean-field arguments by including the crucial effect of the gauge fluctuations, and provides a resolution of the debate surrounding the intermediate field phase in the AFM case.

Study of topological orders and their stability against perturbations is a central topic of current interest in condensed matter physics [1–13]. It was originally proposed by Kitaev [12], using a mapping to a mean-field chiral p -wave superconductor (p SC) model, that the Zeeman-perturbed isotropic Kitaev model has nonabelian Ising topological order (ITO) because (a) Majorana zero modes associated with *extrinsic* twist defects (π -flux particles) of the p SC obeyed Ising braiding rules, and (b) the chiral central charge of the mean field model was $c = 1/2$. However the effect of the gauge sector must be included to demonstrate the nonabelian Majorana anyon as an *intrinsic* bulk excitation, and also obtain the correct values of c , topological entanglement entropy γ , and ground state degeneracy (GSD). *For mean field p SC models, it is long known that three or four different ground states on the torus are possible [14]; however, such ground states are unique and do not simultaneously belong to a degenerate ground state sector. Specifically, the three-fold degenerate ITO state in this case is known to require nonabelian gauge fluctuations [14] while in Kitaev’s fermionization scheme, the gauge fields that appear naturally are abelian Z_2 . In the low field regime, recent perturbative treatments [15, 16] emphasize the important role of vison (gauge field) fluctuation effects, and numerical studies suggest a low-field ITO phase [15–19] but the understanding of quasiparticles, necessary for complete characterization of the topological phases, is lacking. In the AFM case there is an ongoing debate regarding the nature of the intermediate field phase [17, 19–29]. The high field phase is topologically trivial and partially polarized. For the FM Kitaev model, there is a direct field-tuned transition from the KSL to the partially polarized phase.*

Experimentally, there is great interest in the possibility

of ITO revival in Kitaev materials near field-suppressed magnetic order [30, 31]. Such questions have also inspired phenomenological studies of topological phases and excitations in relevant matter Chern-Simons field theories [32, 33]. However before attempting a microscopic treatment of field-tuning ITO in Kitaev materials, it is important to first understand the behavior in the Kitaev limit.

We present here a microscopic derivation and analysis of an effective topological gauge theory for the isotropic Kitaev model subjected to a (001) Zeeman and (Haldane mass) three-spin perturbations. We use a recent Chern-Simons spin fermionization scheme [34] valid for arbitrary 2D lattices that avoids enlargement of the local Hilbert space, with the fermions attached in specific ways to two abelian Z_2 gauge fields respectively on the lattice and dual lattice. Our analysis gives the first complete characterization of low-field ITO as well as intermediate field phases in the AFM Kitaev model starting from a microscopic theory.

Our main results are as follows (see Table I). At low fields, in both FM and AFM cases, the fusion rules, 3-fold GSD on the torus, $\gamma = \ln 2$ and $c = 1/2$ substantiate the phase has ITO, proving Kitaev’s original proposal. For the FM case, ITO is lost beyond a critical field h_c and the transition is to a topologically trivial partially polarized state characterized by $\gamma = c = 0$. For the AFM case, the system transitions at h_{c1} (appreciably larger than h_c of the corresponding FM case) to a chiral abelian phase with fermion quasiparticles. This phase which persists at intermediate fields $h_{c1} < h < h_{c2}$ has trivial topological order i.e. $\gamma = 0$, no GSD, and $c = 1/2$, *and belongs to the class of a $p + ip$ superconductor where fermion number parity is not gauged.* Significantly, *none* of these facts about the intermediate phase can be captured by parton mean field theory, and neglecting gauge fluctuations leads to erroneous results such as sign reversal of half-quantized thermal Hall response [23] at h_{c1} . Beyond h_{c2} , the partially polarized phase appears like the FM case.

We begin with the following $SO(3)$ Majorana fermion-

* jagannath.das@theory.tifr.res.in

† akumar@magnet.fsu.edu

‡ vttripathi@theory.tifr.res.in

← Isotropic AFM Kitaev model		← Isotropic FM Kitaev model →	
Degeneracy	Three fold GSD	No GSD	Three fold GSD
Entanglement Entropy (γ)	$\ln 2$	0	$\ln 2$
Chiral central charge in the edge theory (c)	1/2	1/2	1/2
Phase	ITO {1, ψ , σ }	An abelian phase with trivial topological order and fermionic excitations	ITO {1, ψ , σ }
field (h) → 0		h_{c1} h_{c2} 0	0 h_c → field (h)

TABLE I. The table gives a complete characterization of the different field-tuned topological phases of both the FM and AFM Kitaev model.

ization of the spin-1/2 ($\mathbf{S} = \boldsymbol{\sigma}/2$) [35],

$$\sigma_i^x = -i\eta_i^y \eta_i^z, \quad \sigma_i^y = -i\eta_i^z \eta_i^x, \quad \sigma_i^z = -i\eta_i^x \eta_i^y. \quad (1)$$

The singlet operators $\gamma_i = -i\eta_i^x \eta_i^y \eta_i^z$, anticommute with all fermions except with the η_i fermions at the same site i , and act as Jordan-Wigner (JW) strings [36, 37]. Field theoretically, the 2D JW transformation [38–41] is implemented by Chern-Simons (CS) flux attachment [42–45], but on the lattice, consistency is achieved only if local one to one face-vertex correspondence is satisfied [46]. However enlarging the gauge fields by including the dual lattice allows construction of consistent mutual CS theories on arbitrary 2D lattices [47–50] with Lagrangian (see Supplemental Material (SM) [51] for details)

$$\mathcal{L}_{\text{CS}} = \frac{k}{4\pi} [\xi_{f^*e^*}^* a_{e^*}^* A_v \delta_{f^*v} + D_{v^*e^*}^* a_{v^*}^* A_e \delta_{ee^*} - \partial_0 a_{e^*}^* A_e \delta_{ee^*}] - \frac{k}{4\pi} [\xi_{fe} A_e a_{v^*}^* \delta_{fv^*} + D_{ve} A_v a_{e^*}^* \delta_{ee^*} - \partial_0 A_e a_{e^*}^* \delta_{ee^*}], \quad (2)$$

where ξ_{fe} and D_{ve} are respectively the lattice analogs of curl and gradient [46] operations and “*” everywhere labels the dual lattice. The δ -functions are defined as follows: $\delta_{e,e^*} = 1$ if e and e^* are links dual to each other, and zero otherwise, and similarly for δ_{fv^*} etc. Hereafter we make the gauge choice that all the temporal components of the electromagnetic fields are zero, i.e. $A_v = a_{v^*}^* = 0$. We fermionize the spin operators, $\sigma_i^\alpha = \gamma_i \eta_i^\alpha$, using the JW string, $\gamma_i = e^{i \int_C (B_e + A_e)}$, where $B_e = \frac{k}{2}(a_{e1^*}^* + a_{e2^*}^* + a_{e3^*}^* + a_{e4^*}^*)$ is a specific combination of dual lattice gauge fields [34] enclosing the link e , chosen to satisfy the spin commutation relations¹. $\mathcal{C}$ denotes a path that starts from some reference point and ends at lattice site i . We choose both A_e and $a_{e^*}^*$ as Z_2 , so gauge invariance constraints models we study to those conserving fermion and vortex number parity. From the property $\sigma_i^2 = 1$, we additionally require

¹ The normalization in B_e is $\frac{k}{2}$ which differs from Ref. [34] by a factor of two, and corrects it.

$e^{i \int_e 2B_e} = 1$. For Z_2 gauge fields, 2-body spin interactions on any link $e = \langle ij \rangle$ have a simple form,

$$\sigma_i^\alpha \sigma_j^\beta = \eta_i^\alpha e^{i(A_e + B_e)} \eta_j^\beta. \quad (3)$$

Comparing with the hermitian conjugate of Eq. 3 and using the anticommutation property of the η -fermions, we obtain the constraint $e^{iB_e} = e^{i \frac{(2n+1)\pi}{2}}$ with n an integer. The conditions $e^{2iB_e} = -1$ and $e^{i \int_e 2B_e} = 1$ ensure the correct spin statistics [51]. As a corollary of Eq. 3, for every link $e = \langle ij \rangle$, $\langle \gamma_i \gamma_j \rangle = e^{i(A_e + B_e)}$. We choose the level $k = 1$ which describes a trivial state [34].

We now proceed to implement Chern-Simons fermionization of the isotropic Kitaev model,

$$\mathcal{H}_K = -K \sum_{\langle pq \rangle \in \vartheta - \text{links}} \sigma_p^\vartheta \sigma_q^\vartheta. \quad (4)$$

Here p, q are the vertices associated with the corresponding link ϑ , with $\vartheta = x, y$, or z . Using the $SO(3)$ representation of Eq. 1 we write $\sigma_i^x \sigma_j^x = i(-i\eta_i^y \eta_j^y) \eta_i^z \eta_j^z = ie^{iA_e^x} \eta_i^z \eta_j^z$ and $\sigma_i^y \sigma_j^y = i(-i\eta_i^x \eta_j^x) \eta_i^z \eta_j^z = ie^{iA_e^y} \eta_i^z \eta_j^z$, where $e^{iA_e^x} = -i\eta_i^y \eta_j^y$ for the x -link and $e^{iA_e^y} = -i\eta_i^x \eta_j^x$ for the y -link are chosen such that commutation relations of these η bilinears are preserved. We perform a Hubbard-Stratonovich decoupling of the four-fermion interaction on the z -bond following Ref. [23], $-\sigma_i^z \sigma_j^z = -(\gamma_i \eta_i^z)(\gamma_j \eta_j^z) = -m_i \gamma_j \eta_j^z - m_j \gamma_i \eta_i^z + m_i m_j - i\delta \eta_i^z \eta_j^z - \gamma_i i \gamma_j \Delta + \delta \Delta$, with the mean fields $m_i = \langle \gamma_i \eta_i^z \rangle$, $\delta = \langle i \gamma_i \gamma_j \rangle$ and $\Delta = \langle \eta_i^z \eta_j^z \rangle$. We now introduce the following time-reversal symmetry breaking terms that preserve fermion number parity: $\mathcal{H}_Z = -h \sum_i \sigma_i^z$, is a Zeeman (001) perturbation and $\mathcal{H}_\kappa = -\kappa \sum_{j,k,l} \sigma_j^x \sigma_k^y \sigma_l^z$, corresponds to the scalar spin chirality (Haldane mass term). The (001) orientation has garnered interest recently because of enhanced field-robustness of the KSL [22, 23]. For small deviations of the field direction from (001), the Haldane mass term – necessary for chiral phases – is also generated. The model now has two parameters h and κ ; however, $\mathcal{H}_\kappa$ commutes with the vison (flux) operators and therefore will not cause the vison fluctuations necessary to bring about the topological transitions; also see SM [51]. The mean field Hamiltonian in momentum space is [23] (see SM [51] for details)

$$\mathcal{H}_K = \sum_{\vec{k}} \Psi_{\vec{k}}^\dagger \mathcal{M}_K \Psi_{\vec{k}} + NK_z \delta \Delta + NK_z m^2, \quad (5)$$

with $\Psi_{\vec{k}}^T = [i\gamma_{\vec{k}A} \ i\gamma_{\vec{k}B} \ \eta_{\vec{k}A}^z \ \eta_{\vec{k}B}^z]$ and

$$\mathcal{M}_K = \begin{bmatrix} 0 & \frac{iK\Delta}{2} & -\frac{i(Km+h)}{2} & 0 \\ -\frac{iK\Delta}{2} & 0 & 0 & -\frac{i(Km+h)}{2} \\ \frac{i(Km+h)}{2} & 0 & \frac{W}{2} & -\frac{iKf}{2} \\ 0 & \frac{i(Km+h)}{2} & \frac{iKf^*}{2} & -\frac{W}{2} \end{bmatrix}, \quad (6)$$

where $W^2 = \kappa^2 \{ \delta \sin(0.5k_x + 0.5\sqrt{3}k_y) + \delta \sin(0.5k_x - 0.5\sqrt{3}k_y) - \sin(k_x) \}^2$ and $f = e^{-ik_1} + e^{-ik_2} + \delta$, and $m_{iA} = m_{iB} = m$. N is the number of unit cells, A and B are sublattice indices, and $k_1 = \vec{k} \cdot \hat{n}_1$, $k_2 = \vec{k} \cdot \hat{n}_2$ where




FIG. 1. The configurations that give CS terms. The green lines represent the $\hat{T}$ matrices and red lines are associated with Green function elements.

$\hat{n}_1$ and $\hat{n}_2$ are the unit vector along the x and y -links respectively. The mean fields are calculated self-consistently using Eqs. 5 and 6 [23]. The low-lying bands are chiral, with the Dirac points located at $\vec{k}_0 = (k_0, 0)$, where

$$\cos \frac{k_0}{2} = -\frac{1}{2\Delta} \{ \Delta\delta + (Km + h)^2 \} = -\frac{\xi}{2}. \quad (7)$$

An advantage of this mean field scheme over earlier paragon mean field studies, e.g. [27, 28], is the avoidance of Hilbert space enlargement that can introduce extra topological phases. Most of the numerical studies [17, 20, 25] broadly show only two field-tuned transitions like we find here. See SM [51] for a validation of our mean field choice by benchmarking with extreme limit results.

To obtain the effective gauge action, we first integrate out the massive fermions yielding

$$S_e = -\frac{1}{2} \text{Tr} \ln \{ G_0^{-1} + \hat{T} \} + S_{CS} + NK_z(\delta\Delta + m^2). \quad (8)$$

Here $G_0^{-1} = (-i\omega_n + \mathcal{M}_K)$ is the inverse Green function corresponding to the mean-field Hamiltonian (i.e. without the gauge fluctuations). The matrix $\hat{T}$ contains the gauge fluctuation parts of both Kitaev and the scalar spin chirality terms [51].

Expanding the logarithm term in Eq. (8) we get first non-trivial contributions coming from the second order for the links terms. For the $x(y)$ -bonds we get Maxwell-type electric field terms, $\mathcal{L}_{x(y)} = E_c^{-1} (\dot{A}_e^{x(y)})^2$, with $E_c \approx \frac{W}{2} \frac{[\Delta^2 + (m + \frac{h}{K})^2]^2}{(m + \frac{h}{K})^2}$, and for the z -bonds, $\mathcal{L}_z = \delta^2 E_c^{-1} (\dot{A}_e^z + \dot{B}_e^z)^2$. As a high field benchmarking we check that $E_c \sim h^3/K^2$ grows rapidly with h , while the order parameter δ vanishes resulting in an effectively one-dimensional character of the low-excitations in the x - y backbone reported earlier [22, 34]. Chern-Simons terms arise from processes shown in Fig. 1. For example the configuration in Fig. 1(a) gives [51]:

$$\Delta S_{CS} = \frac{i}{4\pi} \text{sign}(\xi) \text{sign}(W') \sum_{\langle\langle ee' \rangle\rangle} \int d\tau A_e^x \frac{\partial A_{e'}^y}{\partial \tau}, \quad (9)$$

with $e = j_A \rightarrow l_B$ and $e' = i_A \rightarrow j_B$. It is easy to obtain similar CS contributions from the diagrams in Fig. 1b

and 1c. Note that the two link fields in our CS action are separated by precisely one link and do not share a vertex. In our formalism, the z -links are associated with the gauge fields $A_e + B_e$. Consequently, diagrams containing one z -link (e.g. Fig. 1b, 1c) also generate a correction to the mutual CS interaction:

$$\Delta S_{MCS} = \frac{i}{4\pi} \text{sign}(\xi) \text{sign}(W') \sum_{\langle\langle ee' \rangle\rangle} \int d\tau A_e^y \frac{\partial B_{e'}^z}{\partial \tau}. \quad (10)$$

In our model, the sign of effective gap W' does not change unless κ changes sign. However, $\text{sign}(\xi)$ is controlled by the strength of the Zeeman field [51]. In the AFM case, ξ changes sign at h_{c1} . Surprisingly, the sign change of the CS level (chirality) comes not from band inversion but reversal of a velocity component [23].

In the remaining part of the paper, we study the nature of the topological phases in different field regimes. For this purpose we go over to the continuum limit. The lattice gauge field A_e is a line integral of the vector potential $\int_e \vec{A} \cdot d\vec{l}$. The lattice sum passes over to the continuum limit as $\sum_{\text{links}} = \frac{1}{V} \int d^2x$, where $V = 3\sqrt{3}a^2/2$ is the unit cell volume. The continuum limit CS theory has two abelian Z_2 gauge fields and effective (field-dependent) level matrix,

$$K = \begin{bmatrix} \text{sign}(\xi) \text{sign}(W') & 1 + \text{sign}(\xi) \text{sign}(W') \\ 1 + \text{sign}(\xi) \text{sign}(W') & 0 \end{bmatrix}. \quad (11)$$

At low fields ($0 \leq h < h_{c1}$), $\text{sign}(\xi) \text{sign}(W') = 1$, so $K_{\text{low}} = \begin{bmatrix} 1 & 2 \\ 2 & 0 \end{bmatrix}$, which is different from a Z_2 spin liquid for which $K_{Z_2} = \begin{bmatrix} 0 & 2 \\ 2 & 0 \end{bmatrix}$. In the AFM case for $h_{c1} \leq h < h_{c2}$,

we have $\text{sign}(\xi) \text{sign}(W') = -1$ and $K_{\text{inter}} = \begin{bmatrix} -1 & 0 \\ 0 & 0 \end{bmatrix}$. At even higher fields $h > h_{c2}$ corresponding to the partially polarized phase, $\text{sign}(\xi) \text{sign}(W') = 0$ and we revert to the bare trivial K -matrix. For the FM case, single phase transition occurs at some h_c , from K_{low} to the trivial one. Having obtained the level matrices in the different field regimes, we now describe the nature of these phases.

Consider first Z_2 topological order with level K_{Z_2} . The (abelian) quasiparticles (e, m, ϵ) have the general form $X = e^i \int_C l_X^\alpha A_{\alpha\mu} dx^\mu$ where $l_X^\alpha \in Z$, $A_{1\mu} = A_\mu$ and $A_{2\mu} = a_\mu^*$. The quasiparticle charges are $l_e = (1 \ 0)^T$, $l_m = (0 \ 1)^T$, and $l_\epsilon = (1 \ 1)^T$. The fusion rules and braiding statistics follow from the canonical commutation relations $[A_\mu(x), a_\nu^*(x')] = 2\pi i \epsilon_{\mu\nu} K_{Z_2}^{-1} \delta^2(x - x') = \pi i \epsilon_{\mu\nu} \delta^2(x - x')$. The Wilson loop $e^i \oint_C \vec{A} \cdot d\vec{l}$ gives +1 when it encloses the trivial quasiparticle **1** or the e quasiparticle, and -1 otherwise. Similarly the Wilson loop $e^i \oint_C \vec{a}^* \cdot d\vec{l}$ is +1 when it encloses **1** or the m quasiparticle, and -1 otherwise. The ϵ particle corresponds to both the eigenvalues being -1. Any abelian phase and its topological properties can be completely described by the abelian K matrix. But that is not true in general for any non-abelian phase like Ising topological order (ITO) which we will describe now. For $K = K_{\text{low}}$, Z_2 order is clearly lost since the above fusion rules are no longer satisfied.

For an anyon model which is symmetric under some permutations of their topological charges (e.g. $e \leftrightarrow m$ for Z_2), one can describe a non-abelian phase by introducing twist defects [52–55] such as dislocations. This is related to a local symmetry operation $G = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ on K_{Z_2} that amounts to interchanging the two species of gauge fields locally. For Z_2 topological order, it is known that such twist defects (σ) are nonabelian ITO anyons [52, 56].

Now we show that the structure of K_{low} naturally admits a twist defect that satisfies the ITO fusion rules – specifically, the diagonal entry in K_{low} introduces a single twist that breaks e and m exchange symmetry. We show that the Wilson loop $\Gamma = e^{i \oint_{c_2} \vec{a}^* \cdot d\vec{l}}$, where the closed loop c_2 winds twice (say anticlockwise) with only one crossing plays the role of detector of a twist defect σ . This is analogous to the double winding string detectors of twist defects $\sigma_{\pm}$ in Z_2 spin liquids [52] – the main difference in the Z_2 case is the time reversal symmetry allows two distinct twist defects of opposite chirality, and respective eigenvalues $+i$ or $-i$ for the corresponding double winding detector. In our case, only one of these eigenvalues is realized because all defects have the same chirality. Mathematically, the imaginary eigenvalue for the double winding loop comes from the non-trivial commutation relation $[a_{\mu}^*(x), a_{\nu}^*(x')] = 2\pi i \epsilon_{\mu\nu} K_{\text{low}}^{-1} \delta^2(x - x') = -\frac{\pi}{2} i \epsilon_{\mu\nu} \delta^2(x - x')$ in our model. On the other hand, because of the presence of the diagonal term, e and m charges cannot be defined consistently. We now obtain the fusion rules of these twist defects. In the absence of other excitations, for two closed strings that wind twice, say Γ_1 and Γ_2 , we have $\Gamma_1 \Gamma_2 = -1$ in Fig. 2 (a), where the minus sign comes from the presence of two nontrivial crossings, and the remaining dotted curve counts the fermion number parity within. The ψ fermionic quasiparticle can be realised with the string, $e^{i2 \int \vec{a}^* \cdot d\vec{l}}$, and introducing a ψ into any double winding loop of the type Γ simply changes the fermion number parity. The fermion number parity can be ± 1 i.e., modulo 2, the charge within is 1 or ψ . In Fig. 2 (b), introduction of a ψ (yellow line) into Γ clearly does not change the sign, consistent with the fusion rule $\sigma \times \psi = \sigma$.

Now we define the Majorana fermion by an open string $C_j = e^{i \int \vec{a}^* \cdot d\vec{l}}$ that loops around a twist defect at j and all Majorana strings thus defined are to have the same end points. Two such strings (say, C_j and C_{j+1}) with same end points, when braided, results in the change in the orientation of their intersection, yielding the braiding relations $C_j \rightarrow C_{j+1}$, $C_{j+1} \rightarrow -C_j$, consistent with Refs. [52, 57]. The total charge upon fusion is $-iC_j C_{j+1}$, taking values either $+1$ for $\mathbf{1}$ or -1 for ψ respectively. The anyon sector $\{\mathbf{1}, \psi, \sigma\}$ gives ITO.

In anyon-CFT correspondence [58–61], it is well known that these twist fields in $1+1$ dimensions are either holomorphic or antiholomorphic, which is the case for our model. From the edge theory one can also calculate the conformal weight of these fields and the braiding statistics of ITO. The GSD on the torus for this ITO phase is three due to three superselection sectors discussed above. The




FIG. 2. Illustration of fusion rules for twist defects. The blue and yellow line represent the string $e^{i \int \vec{a}^* \cdot d\vec{l}}$ and $e^{i2 \int \vec{a}^* \cdot d\vec{l}}$ respectively. In (a) fusion of two closed strings that wind twice are shown. The dotted line measures the fermion number parity within. The fusion between the quasiparticles ψ and σ is shown in (b).

topological entanglement entropy (γ) can be expressed as $\gamma = \ln D$ where D is the total quantum dimension. In case of the ITO, $D^2 = \sum_{\alpha} d_{\alpha}^2$ with $d = 1$ for vacuum and ψ and $d = \sqrt{2}$ for quasi-particle σ , hence $\gamma = \ln 2$ in ITO as expected. On the other hand, the field regime $h_{c1} \leq h < h_{c2}$ corresponds to a chiral abelian phase with trivial topological order and fermionic bulk excitations, so $D = 1$, $\gamma = 0$. Note that in the ITO phase, strings composed of the gauge field A_{μ} do not have any nontrivial commutation relation with the σ or ψ quasiparticles of ITO. On the other hand, in the intermediate phase of AFM Kitaev model, they are the sole gauge fields, and give the parity and time reversal symmetry breaking CS term in the theory. The gauge sector gives nontrivial contribution to the central charge in this regime, and is associated with the nonzero vison Chern number. It is understood [57, 60] that the $p + ip$ superconductor supports intrinsic non-abelian Ising anyons upon gauging the fermion number parity. In our theory, the fermion number parity is associated with the a^* dynamical gauge field and A is associated with the vortex number parity. The intermediate field phase is deconfined through A dynamical gauge field which is associated with vortex number parity.

Chiral central charge: From the bulk-edge correspondence [62–64], the effective edge theories of K_{low} imply the following Kac-Moody algebra of chiral bosons (ϕ), $[\phi_I(x), \partial_y \phi_J(y)] = \pm 2\pi i K_{\text{low}, I, J}^{-1} \delta(x - y)$. The fact that number of positive and negative eigenvalues are equal, results in same number of left-movers and right-movers on the edge, corresponds to zero net chirality ($c = 0$). Thus the central charge in the ITO phase, coming solely from bare chiral Majoranas is $\frac{1}{2}$ [64]. On the other hand, the intermediate phase of AFM Kitaev model, has an effective abelian CS theory of level $k = -1$ and leads to $c = -1$ from the edge theory [64], while for the bare chiral Majoranas, the Chern number (-1) is flipped w.r.t. the low field phase [23]. The resultant chiral central charge in this phase taking into account both the Majorana and gauge sectors is $c = c_{\text{Majorana}} - c_{\text{gauge}} = \frac{1}{2}$.

Discussion: In summary, we presented a microscopic derivation of abelian lattice Chern-Simons gauge theories for the FM and AFM Kitaev models subjected to time re-

versal symmetry breaking Zeeman (001) and scalar spin chirality perturbations. We obtained a comprehensive understanding of the topological phases and emergent excitations in different field regimes. In the low field phase, we proved the long proposed ITO by constructing the intrinsic nonabelian σ anyons as twist defects with a specific chirality. In the AFM case, the intermediate field phase associated with trivial topological order was found to be chiral with $c = 1/2$, (same as the ITO phase, with no sign change) and the same half-quantized thermal Hall response as the ITO phase (see SM [51] and Ref. [65] for details). Experimentally, the transition to the intermediate phase could be detected as a sharp drop in the plaquette flux value from unity [51] but lacking the 1D excitations of the polarized high field phase. **However extrinsic Majorana anyons could still be induced in such a phase by freezing spins locally [66, 67].** Our technique provides a way for understanding other open questions, e.g. the possibility of field-revived ITO in the simultaneous presence of competing ITO degrading perturbations such as Zeeman and exchange interactions.

Our CS fermionization approach is generally applicable to 2D spin-1/2 systems with fermion number parity symmetry, and could be useful in understanding the effects of

perturbations on the stability and phase transitions in different spin liquid systems. From a practical perspective, our approach could be useful for realizing fault tolerant nonabelian qubits in an abelian gauge theory setting.

ACKNOWLEDGMENTS

JD and VT acknowledge support of the Department of Atomic Energy, Government of India, under Project Identification No. RTI 4002, and Department of Theoretical Physics, TIFR, for computational resources. AK thanks the National High Magnetic Field Laboratory, which is supported by National Science Foundation Cooperative Agreement No. DMR-2128556* and the State of Florida. AK was supported through a Dirac postdoctoral fellowship at NHMFL. JD and VT thank specially Subir Sachdev for useful comments on an earlier version of this work, and Shiraz Minwalla, Ashvin Vishwanath, Yuan-Ming Lu, and Nandini Trivedi for fruitful discussions.

Statement of author contributions: JD and VT conceived the problem and performed the theoretical analysis. AK performed the numerical calculations. JD and VT wrote the paper with inputs from AK.

-
- [1] P. W. Anderson, The resonating valence bond state in La_2CuO_4 and superconductivity, *Science* **235**, 1196 (1987).
 - [2] G. Baskaran, Z. Zou, and P. W. Anderson, The resonating valence bond state and high- T_c superconductivity—a mean field theory, *Solid state commun.* **88**, 853 (1993).
 - [3] V. Kalmeyer and R. Laughlin, Equivalence of the resonating-valence-bond and fractional quantum Hall states, *Phys. Rev. Lett.* **59**, 2095 (1987).
 - [4] S. A. Kivelson, D. S. Rokhsar, and J. P. Sethna, Topology of the resonating valence-bond state: Solitons and high- T_c superconductivity, *Phys. Rev. B* **35**, 8865 (1987).
 - [5] I. Affleck and J. B. Marston, Large- n limit of the heisenberg-hubbard model: Implications for high- T_c superconductors, *Phys. Rev. B* **37**, 3774 (1988).
 - [6] N. Read and S. Sachdev, Large- n expansion for frustrated quantum antiferromagnets, *Phys. Rev. Lett.* **66**, 1773 (1991).
 - [7] X.-G. Wen, Vacuum degeneracy of chiral spin states in compactified space, *Phys. Rev. B* **40**, 7387 (1989).
 - [8] F. J. Wegner, Duality in generalized Ising models and phase transitions without local order parameters, *J. of Math. Phys.* **12**, 2259 (1971).
 - [9] T. Senthil and M. P. Fisher, Z_2 gauge theory of electron fractionalization in strongly correlated systems, *Phys. Rev. B* **62**, 7850 (2000).
 - [10] J. B. Kogut, An introduction to lattice gauge theory and spin systems, *Rev. Mod. Phys.* **51**, 659 (1979).
 - [11] R. Moessner and S. Sondhi, Resonating valence bond liquid physics on the triangular lattice, *Phys. Rev. Lett.* **86**, 1881 (2001).
 - [12] A. Kitaev, Anyons in an exactly solved model and beyond, *Ann. Phys. (N. Y.)* **321**, 2 (2006).
 - [13] L. Savary and L. Balents, Quantum spin liquids: A review, *Rep. on Prog. in Phys.* **80**, 016502 (2016).
 - [14] N. Read and D. Green, Paired states of fermions in two dimensions with breaking of parity and time-reversal symmetries and the fractional quantum Hall effect, *Phys. Rev. B* **61**, 10267 (2000).
 - [15] A. P. Joy and A. Rosch, Dynamics of visons and thermal hall effect in perturbed Kitaev models, *Phys. Rev. X* **12**, 041004 (2022).
 - [16] C. Chen and I. S. Villadiejo, Nature of visons in the perturbed ferromagnetic and antiferromagnetic Kitaev honeycomb models, *Phys. Rev. B* **107**, 045114 (2023).
 - [17] M. Gohlke, R. Moessner, and F. Pollmann, Dynamical and topological properties of the Kitaev model in a [111] magnetic field, *Phys. Rev. B* **98**, 014418 (2018).
 - [18] A. Kumar and V. Tripathi, Thermal Hall conductivity near field-suppressed magnetic order in a Kitaev-Heisenberg model, *Phys. Rev. B* **107**, L220406 (2023).
 - [19] C. Hickey and S. Trebst, Emergence of a field-driven $U(1)$ spin liquid in the Kitaev honeycomb model, *Nat. Commun.* **10**, 530 (2019).
 - [20] Z. Zhu, I. Kimchi, D. Sheng, and L. Fu, Robust non-abelian spin liquid and a possible intermediate phase in the antiferromagnetic Kitaev model with magnetic field, *Phys. Rev. B* **97**, 241110 (2018).
 - [21] Y. Teng, Y. Zhang, R. Samajdar, M. S. Scheurer, and S. Sachdev, Unquantized thermal Hall effect in quantum spin liquids with spinon fermi surfaces, *Phys. Rev. Res.* **2**, 033283 (2020).
 - [22] S. Feng, A. Agarwala, and N. Trivedi, Dimensional reduction of Kitaev spin liquid at quantum criticality, *Phys. Rev. Res.* **6**, 013298 (2024).
 - [23] J. Nasu, Y. Kato, Y. Kamiya, and Y. Motome, Successive Majorana topological transitions driven by a magnetic field

- in the Kitaev model, *Phys. Rev. B* **98**, 060416 (2018).
- [24] S. Liang, M.-H. Jiang, W. Chen, J.-X. Li, and Q.-H. Wang, Intermediate gapless phase and topological phase transition of the Kitaev model in a uniform magnetic field, *Phys. Rev. B* **98**, 054433 (2018).
- [25] N. D. Patel and N. Trivedi, Magnetic field-induced intermediate quantum spin liquid with a spinon fermi surface, *PNAS* **116**, 12199 (2019).
- [26] C. Hickey, M. Gohlke, C. Berke, and S. Trebst, Generic field-driven phenomena in Kitaev spin liquids: Canted magnetism and proximate spin liquid physics, *Phys. Rev. B* **103**, 064417 (2021).
- [27] S.-S. Zhang, G. B. Halász, and C. D. Batista, Theory of the Kitaev model in a [111] magnetic field, *Nat. Commun.* **13**, 399 (2022).
- [28] M.-H. Jiang, S. Liang, W. Chen, Y. Qi, J.-X. Li, and Q.-H. Wang, Tuning topological orders by a conical magnetic field in the Kitaev model, *Phys. Rev. Lett.* **125**, 177203 (2020).
- [29] J. Das, S. Kundu, A. Kumar, and V. Tripathi, Field tuning Kitaev systems for spin fractionalization and topological order, *J. Phys. Condens. Matter* **36**, 443001 (2024).
- [30] T. Yokoi, S. Ma, Y. Kasahara, S. Kasahara, T. Shibauchi, N. Kurita, H. Tanaka, J. Nasu, Y. Motome, C. Hickey, *et al.*, Half-integer quantized anomalous thermal Hall effect in the Kitaev material candidate α -RuCl₃, *Science* **373**, 568 (2021).
- [31] J. Bruin, R. Claus, Y. Matsumoto, N. Kurita, H. Tanaka, and H. Takagi, Robustness of the thermal Hall effect close to half-quantization in α -RuCl₃, *Nat. Phys.* **18**, 401 (2022).
- [32] C. Nayak, S. H. Simon, A. Stern, M. Freedman, and S. D. Sarma, Non-abelian anyons and topological quantum computation, *Rev. of Mod. Phys.* **80**, 1083 (2008).
- [33] L. Zou and Y.-C. He, Field-induced QCD 3-Chern-Simons quantum criticalities in Kitaev materials, *Phys. Rev. Res.* **2**, 013072 (2020).
- [34] J. Das, A. Kumar, A. Maity, and V. Tripathi, Jordan-Wigner fermionization of quantum spin systems on arbitrary two-dimensional lattices: A mutual Chern-Simons approach, *Phys. Rev. B* **108**, 085110 (2023).
- [35] A. Tsvelik, New fermionic description of quantum spin liquid state, *Phys. Rev. Lett.* **69**, 2142 (1992).
- [36] J. Fu, Properties and application of SO(3) Majorana representation of spin: Equivalence with Jordan-Wigner transformation and exact Z_2 gauge theories for spin models, *Phys. Rev. B* **98**, 214432 (2018).
- [37] J. Fu, J. Knolle, and N. B. Perkins, Three types of representation of spin in terms of majorana fermions and an alternative solution of the Kitaev honeycomb model, *Phys. Rev. B* **97**, 115142 (2018).
- [38] T. Sedrakyan, R. Moessner, and A. Kamenev, Helical spin liquid in a triangular xxz magnet from chern-simons theory, *Phys. Rev. B* **102**, 024430 (2020).
- [39] T. A. Sedrakyan, V. M. Galitski, and A. Kamenev, Topological spin ordering via Chern-Simons superconductivity, *Phys. Rev. B* **95**, 094511 (2017).
- [40] A. Lopez, A. Rojo, and E. Fradkin, Chern-simons theory of the anisotropic quantum heisenberg antiferromagnet on a square lattice, *Phys. Rev. B* **49**, 15139 (1994).
- [41] K. Kumar, K. Sun, and E. Fradkin, Chern-simons theory of magnetization plateaus of the spin-1/2 quantum xxz heisenberg model on the kagome lattice, *Phys. Rev. B* **90**, 174409 (2014).
- [42] E. Fradkin, Jordan-Wigner transformation for quantum-spin systems in two dimensions and fractional statistics, *Phys. Rev. Lett.* **63**, 322 (1989).
- [43] Y. Wang, Ground state of the two-dimensional antiferromagnetic Heisenberg model studied using an extended Wigner-Jordan transformation, *Phys. Rev. B* **43**, 3786 (1991).
- [44] M. Azzouz, Interchain-coupling effect on the one-dimensional spin-1/2 antiferromagnetic Heisenberg model, *Phys. Rev. B* **48**, 6136 (1993).
- [45] O. Derzhko, Jordan-Wigner fermionization for spin-1/2 systems in two dimensions: A brief review, *arXiv preprint cond-mat/0101188* (2001).
- [46] K. Sun, K. Kumar, and E. Fradkin, Discretized Abelian Chern-Simons gauge theory on arbitrary graphs, *Phys. Rev. B* **92**, 115148 (2015).
- [47] T. Banks and B. Zhang, Lattice bf theory, dumbbells, and composite fermions, *Nucl. Phys. B* **981**, 115877 (2022).
- [48] R. Kantor and L. Susskind, A lattice model of fractional statistics, *Int. J. Mod. Phys. B* **5**, 2701 (1991).
- [49] S. Sen, S. Sen, J. C. Sexton, and D. H. Adams, Geometric discretization scheme applied to the Abelian Chern-Simons theory, *Phys. Rev. E* **61**, 3174 (2000).
- [50] Y.-A. Chen, A. Kapustin, and D. Radičević, Exact bosonization in two spatial dimensions and a new class of lattice gauge theories, *Ann. of Phys.* **393**, 234 (2018).
- [51] See Supplemental Material for (a) details of lattice mutual Chern-Simons action, (b) Chern-Simons fermionization of spins on arbitrary lattices using two abelian gauge fields and the mean-field results, (c) details of derivation of the effective gauge theory and renormalization of Chern-Simons level, and (d) benchmarking and numerical evidence.,.
- [52] H. Bombin, Topological order with a twist: Ising anyons from an Abelian model, *Phys. Rev. Lett.* **105**, 030403 (2010).
- [53] J. C. Teo, T. L. Hughes, and E. Fradkin, Theory of twist liquids: gauging an anyonic symmetry, *Ann. of Phys.* **360**, 349 (2015).
- [54] Y.-Z. You, C.-M. Jian, and X.-G. Wen, Synthetic non-Abelian statistics by Abelian anyon condensation, *Phys. Rev. B* **87**, 045106 (2013).
- [55] H. Bombin and M. Martin-Delgado, Family of non-Abelian Kitaev models on a lattice: Topological condensation and confinement, *Phys. Rev. B* **78**, 115421 (2008).
- [56] H. Zheng, A. Dua, and L. Jiang, Demonstrating non-Abelian statistics of majorana fermions using twist defects, *Phys. Rev. B* **92**, 245139 (2015).
- [57] D. A. Ivanov, Non-Abelian statistics of half-quantum vortices in p -wave superconductors, *Phys. Rev. Lett.* **86**, 268 (2001).
- [58] N. Read and E. Rezayi, Beyond paired quantum Hall states: Parafermions and incompressible states in the first excited Landau level, *Phys. Rev. B* **59**, 8084 (1999).
- [59] N. Read and G. Moore, Fractional quantum Hall effect and nonabelian statistics, *Prog. of Theor. Phys. Supp.* **107**, 157 (1992).
- [60] G. Moore and N. Read, Nonabelions in the fractional quantum Hall effect, *Nucl. Phys. B* **360**, 362 (1991).
- [61] X. Chen, A. Roy, J. C. Teo, and S. Ryu, From orbifolding conformal field theories to gauging topological phases, *Phys. Rev. B* **96**, 115447 (2017).
- [62] X.-G. Wen, Topological orders and edge excitations in fractional quantum Hall states, *Adv. in Phys.* **44**, 405 (1995).
- [63] E. Witten, Quantum field theory and the Jones polynomial, *Commun. in Math. Phys.* **121**, 351 (1989).
- [64] P. Francesco, P. Mathieu, and D. Sénéchal, *Conformal Field*

- Theory* (Springer Science & Business Media, 2012).
- [65] A. Kapustin and L. Spodyneiko, Thermal Hall conductance and a relative topological invariant of gapped two-dimensional systems, [Phys. Rev. B **101**, 045137 \(2020\)](#).
 - [66] K. Dhochak, R. Shankar, and V. Tripathi, Magnetic impurities in the honeycomb Kitaev model, [Phys. Rev. Lett. **105**, 117201 \(2010\)](#).
 - [67] S. D. Das, K. Dhochak, and V. Tripathi, Kondo route to spin inhomogeneities in the honeycomb Kitaev model, [Phys. Rev. B **94**, 024411 \(2016\)](#).