

This is the accepted manuscript made available via CHORUS. The article has been published as:

Numerical search for diabolical points in the energy spectrum of the single-molecule magnet $\text{Fe}_{\{8\}}$

Feifei Li and Anupam Garg

Phys. Rev. B **83**, 132401 — Published 4 April 2011

DOI: [10.1103/PhysRevB.83.132401](https://doi.org/10.1103/PhysRevB.83.132401)

Lost and found: The missing diabolical points in the Fe_8 molecular magnet

Feifei Li and Anupam Garg*

Department of Physics and Astronomy, Northwestern University, Evanston, Illinois 60208

Certain diabolical points in the tunneling spectrum of the single-molecule magnet Fe_8 were previously believed to have been eliminated as a result of a weak fourth-order anisotropy. As shown by Bruno, this is not so, and the points are only displaced in the magnetic field space along the medium anisotropy direction. The previously missing points are numerically located by following the lines of the Berry curvature. The importance of an experimental search for these rediscovered points is discussed.

PACS numbers: 75.50.Xx, 03.65.Vf, 03.65.Db

The purpose of this note is to report on a numerical search of certain diabolical points (DP's) in the energy spectrum of the single-molecule magnet (also known as a molecular magnet or nanomagnet) Fe_8 that were earlier believed to be missing, but are in fact not so [1]. Several other DP's *have* been seen experimentally in Fe_8 [2], and their observation provides the best evidence of spin orientation tunneling between deep levels in all single-molecule magnets studied to date. Observation of even some of the missing DP's would strengthen our understanding of Fe_8 substantially.

For a system whose Hamiltonian depends on some parameter, a DP is a point in parameter space where two (or more) energy levels are degenerate [3]. In Fe_8 , the parameter is the static applied magnetic field, and the locations of the DP's so far observed (as well as many other experimental measurements) are well described by the anisotropy Hamiltonian,

$$\mathcal{H} = k_1 J_x^2 + k_2 J_y^2 - C(J_+^4 + J_-^4) - g\mu_B \mathbf{J} \cdot \mathbf{H}. \quad (1)$$

Here $\mathbf{J} = (J_x, J_y, J_z)$ is the spin, \mathbf{H} is an external magnetic field, g is a g-factor, μ_B is the Bohr magneton, and k_1, k_2 , and C are anisotropy coefficients. Experimentally, $J = 10$, $k_1 \simeq 0.338$ K, $k_2 \simeq 0.246$ K, $C \simeq 29 \mu\text{K}$, and $g \simeq 2$. The DP's can be understood as arising when tunneling between two states with (nearly) oppositely oriented magnetic moment is quenched because of destructive interference between instantons (spin tunneling trajectories) [4, 5].

The model Hamiltonian (1) was first analyzed in 1993 [4] with $C = 0$, and it was found that for ground state tunneling it had 10 DP's along the positive H_x axis, corresponding to $J = 10$. In reality only 4 DP's are seen [2], which was explained in Ref. [6] as follows. When $C \neq 0$, we get two new (but noninterfering) instantons, which are discontinuous at the end points. One of these instantons has the least action when H_x exceeds a certain value H^* , and since this instanton has no interfering partner, there are no more DP's for $H_x > H^*$. For the values of the anisotropy coefficients quoted above, H^* lies just beyond the location of the fourth DP, which explains why the last six DP's are not seen in experiment or by

direct numerical diagonalization of Eq. (1). We can further check this picture by decreasing C so as to make H^* larger. It is seen that as this happens, one successively sees six and then eight, and finally all ten DP's, all in accord with direct numerical diagonalization results [6].

However, as shown by Bruno [1], the above picture, though correct, is incomplete. For any energy level, the sum of the Chern numbers for all DP's involving that level is a topological invariant as parameters like k_1, k_2 , or C are varied. Since DP's in any system are generically simple, we expect this to be so in Fe_8 also, and the Chern number for any one DP should be ± 1 whether $C = 0$ or $C \neq 0$. Hence the six missing DP's must be present elsewhere in magnetic field space. A similar conclusion applies to the DP's associated with tunneling between other pairs of levels [7]. For tunneling between the ground states, the DP's merely move off the x axis into the xy plane. For the higher energy levels, they move off the xz plane into the full three dimensional \mathbf{H} space. This point can also be understood by noting that for a system with purely four-fold symmetry ($k_2 = k_1, C \neq 0$), the ground state DP's lie on the $\pm \hat{x} \pm \hat{y}$ axes, while for the excited states they lie in the planes formed by these axes and the \hat{z} axis [8]. When both two-fold and four-fold anisotropies are present ($k_2 \neq k_1, C \neq 0$), it is then not surprising that the location of some of the DP's is also intermediate [1].

Observation of these rediscovered DP's would be interesting in itself, and also provide an important test of the validity of the model (1) vs. other models [10] that add extra 6th and 8th order anisotropies because the location of the DP's is very sensitive to the higher order anisotropies. With this motivation, we have undertaken a search for the DP's for the ground state and some of the excited states. We stress that the key insight that these points should exist in the first place is due to Bruno, and our contribution is only to find their specific locations. Nevertheless, finding them is not without challenge as we discuss next.

A direct search for the DP's by numerical minimization of the energy differences fails because the energy surface is like a golf course with rolling hills on which




FIG. 1: Diabolical points of Fe_8 corresponding to tunneling from the Zeeman level $m = -10$ to $m = 10$ (layer 1), to $m = 9$ (layer 2), and to $m = 8$ (layer 3), all projected onto the xy plane. $H_c = 2k_1 J/g\mu_B$.

the DPs are the holes. Because the holes are so localized, unless one starts close to one of them by luck, any numerical algorithm will in general simply head for the valleys of the course and miss the holes entirely. Because \mathcal{H} is not real for general \mathbf{H} , we also cannot use the method of Ref. [8], which is to corral the DPs by using the Herzberg and Longuet-Higgins theorem [9] to find and successively bisect a sign-reversing circuit. We therefore proceed as follows. Let us denote the eigenstates and eigenvalues of Eq. (1) for fixed \mathbf{H} by $|n(\mathbf{H})\rangle$ and $E_n(\mathbf{H})$, $n = 1, 2, \dots, 21$, and order them so that $E_n \geq E_{n-1}$ for every \mathbf{H} . Except at degeneracies (the DPs), the Berry curvature for the n th level is defined by [11]

$$\mathbf{B}_n = -\text{Im} \sum_{n' \neq n} \frac{\langle n | \nabla_{\mathbf{H}} \mathcal{H} | n' \rangle \times \langle n' | \nabla_{\mathbf{H}} \mathcal{H} | n \rangle}{(E_{n'} - E_n)^2}. \quad (2)$$

Further, the Chern number associated with a degeneracy is given by

$$Q_n = -\frac{1}{2\pi} \oint_S \mathbf{B}_n \cdot d^2\mathbf{s}, \quad (3)$$

where the integral is independent of the choice of the surface S , as long as it encloses the degeneracy, since $\nabla_{\mathbf{H}} \cdot \mathbf{B}_n = 0$ away from the degeneracy. The Chern number is always an integer, and for a simple double degeneracy, it equals ± 1 . In other words, near a DP, \mathbf{B}_n has the form of a monopole field with flux equal to $\pm 2\pi$. Hence, to find the DPs, we numerically evaluate \mathbf{B}_n for an initial \mathbf{H} , and follow the lines of \mathbf{B}_n in the direction of increasing strength until we hit a monopole. Since the number of DPs where levels n and $n+1$ are degenerate is topologically fixed and known, all the DPs can be found by taking sufficiently many initial values of \mathbf{H} . The DPs for successive pairs of levels occur in layers, with H_z essentially constant in a layer. With $\mathbf{h} \equiv g\mu_B \mathbf{H}/2k_1 J$, the first three layers are at $h_z = 0$ (exactly), 0.0427 , and 0.0854 . It should be noted, however, that in a given layer,

TABLE I: Locations of selected diabolical points

(m, m')	Layer	(h_x, h_y, h_z)
$(-10, 10)$	1	$(0.0404, 0, 0)$
		$(0.1207, 0, 0)$
		$(0.2011, 0, 0)$
		$(0.2565, 0, 0)$
		$(0.3139, \pm 0.0642, 0)$
		$(0.3875, \pm 0.1334, 0)$
		$(0.4656, \pm 0.2043, 0)$
$(-10, 9)$	2	$(0, 0, 0.0426)$
		$(0.0787, 0, 0.0426)$
		$(0.1568, 0, 0.0427)$
		$(0.2276, 0, 0.0427)$
		$(0.2893, \pm 0.0447, 0.0427)$
		$(0.3594, \pm 0.1109, 0.0427)$
		$(0.4340, \pm 0.1788, 0.0427)$
$(-10, 8)$	3	$(0.0385, 0, 0.0853)$
		$(0.1150, 0, 0.0853)$
		$(0.1892, 0, 0.0853)$
		$(0.2650, \pm 0.0268, 0.0854)$
		$(0.3326, \pm 0.0900, 0.0854)$
		$(0.4038, \pm 0.1548, 0.0854)$
$(-9, 9)$	1	$(0.0384, 0, 0)$
		$(0.1147, 0, 0)$
		$(0.1889, 0, 0)$
		$(0.2651, \pm 0.0261, 0)$
		$(0.3325, \pm 0.0891, 0)$
		$(0.4034, \pm 0.1535, 0)$
$(-8, 8)$	1	$(0.0365, 0, 0)$
		$(0.1089, 0, 0)$
		$(0.1805, 0, 0)$
		$(0.2339, 0, 0)$
		$(0.2831, \pm 0.0511, 0)$
		$(0.3468, \pm 0.1092, 0)$

one can have DP's corresponding to tunneling between levels with different pairs of Zeeman quantum numbers. For example, layer 1 contains points corresponding to tunneling between $m = -10 \leftrightarrow 10$, $-9 \leftrightarrow 9$, $-8 \leftrightarrow 8$, etc. Likewise, layer 2 contains points corresponding to tunneling between $m = -10 \leftrightarrow 9$, $-9 \leftrightarrow 8$, $-8 \leftrightarrow 7$, etc. In Table I, we show the DP's for $(-10 \leftrightarrow 10)$ ($-10 \leftrightarrow 9$), $(-10 \leftrightarrow 8)$, $(-9 \leftrightarrow 9)$, and $(-8 \leftrightarrow 8)$ tunneling. For the first three of these pairs of levels, the projections of the DP's onto the xy plane are shown in Fig. 1. We note that it is just these three pairs of levels for which tunnel splittings were reported in Ref. [2]. However, except for the ground pair of levels (layer 1), not even all the DP's on the x axis are found there.

In the rest of this note, we discuss the form of the Berry curvature near a DP in more detail. For simplicity, we

will divide \mathcal{H} by $k_1 J^2$. Since $\nabla_{\mathbf{h}} \mathcal{H}$ and E_n are both divided by this factor, it follows from Eq. (2) that \mathbf{B}_n is unchanged. With this preliminary remark, let us suppose that $E_n = E_{n-1} \equiv E_{n,n-1}$ at $\mathbf{h} = \mathbf{h}_{n,n-1}$, and denote

$$\mathbf{r} = \mathbf{h} - \mathbf{h}_{n,n-1}. \quad (4)$$

Further, let us make a particular choice of the two degenerate states at $\mathbf{r} = 0$, and denote them by $|a\rangle$ and $|b\rangle$, with $\langle b|a\rangle = 0$. (Any orthogonal linear combination of $|a\rangle$ and $|b\rangle$ would also work.) It suffices to truncate the Hamiltonian to this two dimensional subspace since the sum in Eq. (2) is dominated by degenerate states. Hence, at $\mathbf{r} = 0$, we have

$$\mathcal{H} = E_{n,n-1} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}. \quad (5)$$

For small enough \mathbf{r} , we can take $|a\rangle$ and $|b\rangle$ to be unchanged, so

$$\nabla_{\mathbf{h}} \mathcal{H} = -\frac{2}{J} \begin{pmatrix} \mathbf{J}_{aa} & \mathbf{J}_{ab} \\ \mathbf{J}_{ba} & \mathbf{J}_{bb} \end{pmatrix}, \quad (6)$$

where $\mathbf{J}_{aa} = \langle a|\mathbf{J}|a\rangle$ etc. Next, let us define $\mathbf{J}_{aa} + \mathbf{J}_{bb} = J\mathbf{t}$, $\mathbf{J}_{aa} - \mathbf{J}_{bb} = J\mathbf{u}$, $\mathbf{J}_{ab} = J(\mathbf{v} + i\mathbf{w})/2$, where \mathbf{t} , \mathbf{u} , \mathbf{v} , and \mathbf{w} are real vectors. In terms of these vectors, we have

$$\mathcal{H} = -\frac{\mathbf{t} \cdot \mathbf{r}}{J} - \begin{pmatrix} \mathbf{u} \cdot \mathbf{r} & (\mathbf{v} + i\mathbf{w}) \cdot \mathbf{r} \\ (\mathbf{v} - i\mathbf{w}) \cdot \mathbf{r} & -\mathbf{u} \cdot \mathbf{r} \end{pmatrix}, \quad (7)$$

where we have ignored the constant $E_{n,n-1}$. Similarly ignoring the overall shift $-\mathbf{t} \cdot \mathbf{r}/J$, the eigenvalues of this matrix are $\pm\epsilon(\mathbf{r})$, with

$$\epsilon(\mathbf{r}) = [(\mathbf{u} \cdot \mathbf{r})^2 + (\mathbf{v} \cdot \mathbf{r})^2 + (\mathbf{w} \cdot \mathbf{r})^2]^{1/2}. \quad (8)$$

To write the eigenvectors compactly, we define

$$\cos \theta(\mathbf{r}) = \mathbf{u} \cdot \mathbf{r} / \epsilon(\mathbf{r}), \quad (9)$$

$$\sin \theta(\mathbf{r}) e^{i\varphi(\mathbf{r})} = (\mathbf{v} + i\mathbf{w}) \cdot \mathbf{r} / \epsilon(\mathbf{r}). \quad (10)$$

The eigenvectors are then

$$|-\rangle = \begin{pmatrix} \cos \frac{1}{2}\theta \\ \sin \frac{1}{2}\theta e^{-i\varphi} \end{pmatrix}, \quad |+\rangle = \begin{pmatrix} \sin \frac{1}{2}\theta \\ -\cos \frac{1}{2}\theta e^{-i\varphi} \end{pmatrix}. \quad (11)$$

Further abbreviating $c = \cos \frac{1}{2}\theta$ and $s = \sin \frac{1}{2}\theta$, and $\mathbf{g} = \langle +|\nabla_{\mathbf{h}} \mathcal{H}|-\rangle$, we have

$$\mathbf{g} = -2cs\mathbf{u} - s^2 e^{-i\varphi}(\mathbf{v} + i\mathbf{w}) + c^2 e^{i\varphi}(\mathbf{v} - i\mathbf{w}). \quad (12)$$

It then follows that

$$\mathbf{g} \times \mathbf{g}^* = 2i[\cos \theta(\mathbf{v} \times \mathbf{w}) + \sin \theta \cos \varphi(\mathbf{w} \times \mathbf{u}) + \sin \theta \sin \varphi(\mathbf{u} \times \mathbf{v})], \quad (13)$$

so that for the level labeled +,

$$\begin{aligned} \mathbf{B}_+ &= \frac{-1}{2\epsilon^3(\mathbf{r})} [(\mathbf{u} \cdot \mathbf{r})(\mathbf{v} \times \mathbf{w}) + (\mathbf{v} \cdot \mathbf{r})(\mathbf{w} \times \mathbf{u}) \\ &\quad + (\mathbf{w} \cdot \mathbf{r})(\mathbf{u} \times \mathbf{v})] \\ &= \frac{-1}{2\epsilon^3(\mathbf{r})} [\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w})] \mathbf{r}. \end{aligned} \quad (14)$$

This is clearly of monopole form with appropriately scaled and sheared axes. It is not difficult to show that $\mathbf{B}_- = -\mathbf{B}_+$ and that $\nabla \cdot \mathbf{B}_+ = 0$.

To find the Chern number, we must evaluate the integral

$$Q_{\pm} = -\frac{1}{2\pi} \oint_S \mathbf{B}_{\pm} \cdot d^2\mathbf{s} \quad (15)$$

for a suitable surface S . Let us take S to be the surface of the parallepiped with vertices at $(\pm\mathbf{u}^*, \pm\mathbf{v}^*, \pm\mathbf{w}^*)$, where \mathbf{u}^* , \mathbf{v}^* , and \mathbf{w}^* are the reciprocal vectors

$$\mathbf{u}^* = \frac{\mathbf{v} \times \mathbf{w}}{\Omega}, \quad \mathbf{v}^* = \frac{\mathbf{w} \times \mathbf{u}}{\Omega}, \quad \mathbf{w}^* = \frac{\mathbf{u} \times \mathbf{v}}{\Omega}; \quad (16)$$

$$\Omega = \mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}). \quad (17)$$

Then $\mathbf{u} \cdot \mathbf{u}^* = 1$, $\mathbf{u} \cdot \mathbf{v}^* = \mathbf{u} \cdot \mathbf{w}^* = 0$, etc. Let us consider the integral over the face of the parallepiped that has edges along \mathbf{v}^* and \mathbf{w}^* , and thus has a normal parallel to $+\mathbf{u}$ (the others have normals along $-\mathbf{u}$, $\pm\mathbf{v}$, and $\pm\mathbf{w}$). On this face, we may parametrize \mathbf{r} as

$$\mathbf{r} = \mathbf{u}^* + \alpha\mathbf{v}^* + \beta\mathbf{w}^*, \quad (18)$$

where $-1 \leq \alpha \leq 1$, $-1 \leq \beta \leq 1$. Then, since the area of this face is $4|\mathbf{v}^* \times \mathbf{w}^*| = 4|\mathbf{u}|/\Omega$,

$$d^2\mathbf{s} = \frac{\mathbf{u}}{\Omega} d\alpha d\beta. \quad (19)$$

Finally, $\mathbf{u} \cdot \mathbf{r} = 1$, $\mathbf{v} \cdot \mathbf{r} = \alpha$, and $\mathbf{w} \cdot \mathbf{r} = \beta$. Thus, the contribution from this face to Q_{\pm} is given by

$$\pm \frac{1}{4\pi} \int_{-1}^1 \int_{-1}^1 \frac{d\alpha d\beta}{(1 + \alpha^2 + \beta^2)^{3/2}} = \pm \frac{1}{6}. \quad (20)$$

(The integral is elementary, and can be performed by standard trigonometric substitutions.) The contributions from the other faces are identical, so

$$Q_{\pm} = \pm 1. \quad (21)$$

Thus the Chern number associated with the lower energy level is always -1 irrespective of the details, and this is why the sum of the numbers for a given energy level cannot be altered by varying the anisotropy parameters.

The reader cannot have failed to notice the regularity of the DP pattern for the present problem. This is in sharp contrast to the generic situation where no pattern is expected apart from a general scaling of the density

with energy [3]. The reason for this regularity is unclear to us.

This work is supported by the NSF via grant no. PHY-0854896.

* e-mail address: agarg@northwestern.edu

- [1] P. Bruno, Phys. Rev. Lett. **96**, 117208 (2006).
- [2] W. Wernsdorfer and R. Sessoli, Science **284**, 133 (1999).
- [3] M. V. Berry and M. Wilkinson, Proc. Roy. Soc. London A **392**, 15 (1984).
- [4] A. Garg, Europhys. Lett. **22**, 205 (1993).
- [5] D. Loss, D. P. DiVincenzo, and G. Grinstein, Phys. Rev. Lett. **69**, 3232 (1992); J. von Delft and C. L. Henley, *ibid* **69**, 3236 (1992).
- [6] Ersin Keçecioglu and A. Garg, Phys. Rev. Lett. **88**, 237205 (2002); Phys. Rev. B **67**, 054406 (2003).
- [7] J. Villain and A. Fort, Eur. Phys. J. B **17**, 69 (2000). Ersin Keçecioglu and A. Garg, Phys. Rev. B **63**, 064422 (2001);
- [8] Chang-Soo Park and A. Garg, Phys. Rev. B **65**, 064411 (2002).
- [9] G. Herzberg and H. C. Longuet-Higgins, Discuss. Faraday Soc. **35**, 77 (1963).
- [10] M. N. Leuenberger and D. Loss, Phys. Rev. B **61**, 12200 (2000). The relative merits of this proposal vs. Eq. (1) are discussed in Ersin Keçecioglu and A. Garg, Phys. Rev. B **76**, 134405 (2007).
- [11] M. V. Berry, Proc. Roy. Soc. London A **392**, 45 (1984).