

This is the accepted manuscript made available via CHORUS. The article has been published as:

Phase of phonon-induced resistance oscillations in a high-mobility two-dimensional electron gas

A. T. Hatke, M. A. Zudov, L. N. Pfeiffer, and K. W. West

Phys. Rev. B **84**, 121301 — Published 2 September 2011

DOI: [10.1103/PhysRevB.84.121301](https://doi.org/10.1103/PhysRevB.84.121301)

The phase of phonon-induced resistance oscillations in a high-mobility 2D electron gas

A. T. Hatke,¹ M. A. Zudov,^{1,*} L. N. Pfeiffer,² and K. W. West²

¹*School of Physics and Astronomy, University of Minnesota, Minneapolis, Minnesota 55455, USA*

²*Princeton University, Department of Electrical Engineering, Princeton, NJ 08544, USA*

(Received 11 July 2011)

We report on experimental studies of magnetoresistance oscillations which originate from the resonant interaction of two-dimensional electrons with thermal transverse acoustic phonons in very high-mobility GaAs/AlGaAs quantum wells. We find that the oscillation maxima consistently occur when a frequency of a phonon with twice the Fermi momentum *exceeds* an integer multiple of the cyclotron frequency. This observation is in contrast to all previous experiments associating resistance maxima with magneto-phonon resonance and its harmonics. Our experimentally obtained resonant condition is in excellent quantitative agreement with recent theoretical proposals.

PACS numbers: 73.43.Qt, 73.63.Hs, 73.21.-b, 73.40.-c

Resonant interaction of a two-dimensional electron gas (2DEG) with thermal *optical* phonons was predicted¹ and observed² in magnetotransport measurements in GaAs/AlGaAs heterostutures a long time ago. Experimentally, this interaction gives rise to a rather weak enhancement of the longitudinal resistivity occurring whenever the phonon frequency ω_o equals an integer multiple of the cyclotron frequency, $\omega_c = eB/m^*$, where B is the magnetic field and m^* is the effective electron mass. In GaAs,³ $\omega_o \sim 10^{13} \text{ s}^{-1}$ and observation of the resulting magnetoresistance oscillations requires both high temperatures, $T \gtrsim 10^2 \text{ K}$, and strong magnetic fields, $B \gtrsim 10^2 \text{ kG}$.

About a decade ago another class of magnetoresistance oscillations was discovered⁴ in a high-mobility ($\mu_e \sim 10^6 \text{ cm}^2/\text{Vs}$) 2DEG at much lower temperatures, $T \sim 1 - 10 \text{ K}$, and magnetic fields, $B \sim 1 - 10 \text{ kG}$. These so called phonon-induced resistance oscillations (PIRO) are currently understood in terms of resonant interaction of 2D electrons with thermal *acoustic* phonons which was first observed in experiments on phonon drag in silicon inversion layers.⁵

Resonant electron-acoustic phonon scattering in 2DEG is made possible by virtue of a special “selection rule” which results in electron *backscattering* off of an acoustic phonon.⁴ In this scenario, even though phonons of many different energies are present, *only* the most energetic phonons electrons can scatter off can be considered to describe the effect. Such phonons carry momentum $\simeq 2k_F$ (k_F is the Fermi wavenumber) and, as a result, their frequency ω_s is determined entirely by the parameters of the 2DEG, i.e. the sound velocity v_s and the electron density n_e ,

$$\omega_s = 2k_F v_s. \quad (1)$$

Due to absorption and emission of $2k_F$ phonons an electron undertakes *indirect* transitions between Landau levels and the longitudinal resistivity acquires a periodic in $1/B$ oscillatory correction. Similar to magnetoresistance oscillations originating from optical phonons,² this correction oscillates with the ratio of the $2k_F$ phonon fre-

quency ω_s to the cyclotron frequency,

$$\epsilon = \frac{\omega_s}{\omega_c}, \quad (2)$$

and can be described by

$$\delta\rho \simeq A(\epsilon) \cos(2\pi\epsilon + \varphi), \quad \epsilon = \frac{\omega_s}{\omega_c}. \quad (3)$$

In contrast to other low-field transport phenomena, such as microwave-induced⁶⁻³⁴ and Hall field-induced³⁵⁻⁴¹ resistance oscillations, zero resistance⁴²⁻⁵⁴ and zero differential resistance⁵⁵⁻⁵⁷ states, PIRO have not received much attention until recently⁵⁸⁻⁶⁶ and several unsolved issues remain.

One such issue is the value of the phase φ appearing in Eq. (3). All of the existing experiments^{4,58-62,67,68} associated PIRO maxima and minima with integer and half-integer ϵ values, respectively, suggesting that the value of the phase is close to zero, $\varphi \approx 0$. On the other hand, recent theories⁶³⁻⁶⁵ predict that the phase is not necessarily zero and is determined by the specifics of the phonon spectrum, the width of the quantum well, and the temperature, which controls excitation of different phonon modes. However, to date there exist no experimental study which examines these theoretical predictions.

In this Rapid Communication we systematically investigate the period and the phase of phonon-induced resistance oscillations in several very high mobility ($\mu_e > 10^7 \text{ cm}^2/\text{Vs}$) 2DEGs. Extremely low level of disorder in our samples allows reliable measurement of up to eight oscillations at relatively low temperatures. The latter condition is rather crucial to ensure that contribution from more energetic longitudinal phonon modes is sufficiently weak and can be neglected. Analysis of the oscillation waveform reveals that in all of our samples the dominant mode has a velocity $v_s \approx 3.4 \text{ km/s}$ suggesting a dominant contribution from the transverse branch of the phonon spectrum. More importantly, in contrast to previous experimental studies which assumed $\varphi = 0$, we find that all of the observed oscillations are best described by a nonzero phase, $\varphi \approx -0.24\pi$, in all of the samples studied. We discuss our findings in terms of recent theoretical proposals^{64,65} and find excellent quantitative agreement.




FIG. 1: (color online) Magnetoresistivity $\rho(B)$ measured in sample A at $T = 3.5$ K (a), in sample B at $T = 3.0$ K (b), and in sample C at $T = 4.2$ K (c). Integers mark the oscillation order n . Vertical lines are drawn to illustrate higher magnetic field of the first PIRO peak in sample A (right line) than in samples B and C (left line).

All of our samples (A, B, and C) are lithographically defined Hall bars fabricated from very high mobility symmetrically doped GaAs/AlGaAs quantum well structures grown by molecular beam epitaxy. The electron density n_e (in units of 10^{11} cm^{-2}), the disorder-limited mobility μ_e (in units of $10^7 \text{ cm}^2/\text{Vs}$), and the quantum well width (in nm), were 3.7, 1.0, and 30 (sample A), 3.0, 1.2, and 30 (sample B), and 2.9, 2.4, and 28 (sample C), respectively.

Magnetoresistivity $\rho(B)$ was measured in a ^3He cryostat using low frequency lock-in technique.

In Fig. 1 we present the magnetoresistivity $\rho(B)$ measured in sample A at $T = 3.5$ K (a), in sample B at $T = 3.0$ K (b), and in sample C at $T = 4.2$ K (c). All samples reveal several orders (up to $n = 8$ in sample A) of pronounced PIRO which are marked by integers. Oscillations become progressively stronger with the magnetic field reflecting $A \propto \lambda^2$ scaling of the oscillation amplitude. Here, $\lambda = \exp(-\pi/\omega_c\tau_q)$ is the Dingle factor and τ_q is the quantum lifetime.^{60,64,65}

Close examination of Fig. 1 reveals that PIRO occur at somewhat higher magnetic fields in sample A (cf. right vertical line) compared to samples B and C (cf. left vertical line). This difference is well accounted for by




FIG. 2: (color online) Oscillation order n (solid circles) for the maxima and $n + 1/2$ (open circles) for the minima (left axis) and the oscillatory part of the resistivity $\delta\rho$ (right axis) vs $1/B$ measured in sample A (a), sample B (b), and sample C (c). Lines are linear fits to the oscillation order j . Insets illustrate that the fits extrapolate to non-zero value at $1/B = 0$ for all samples.

$\omega_s \propto k_F \propto \sqrt{n_e}$ scaling of the phonon frequency, see Eq. (1). Finally, Fig. 1 demonstrates that PIRO appear regardless of the sign and the magnitude of the magnetoresistance effect. Indeed, in samples A, B, and C, oscillations are superimposed on a slowly varying background which is decreasing, increasing, and roughly constant, respectively.

To examine the period of the oscillations we extract the oscillatory part of the magnetoresistivity $\delta\rho$ by subtracting the background from the data presented in Fig. 1. The result of such extraction is presented in Fig. 2 (right axis) clearly showing periodicity in $1/B$ for all three samples, in agreement with Eq. (3). Following earlier experimental studies,^{4,58–62,67} we associate the maxima and the minima of $\delta\rho$ with integer n and half-integer $n + 1/2$ values of the oscillation order, respectively. These values for the maxima (solid circles) and the minima (open circles) are plotted in Fig. 2 (left axis) as a function of $1/B$. For all samples studied we observe universal linear dependence on $1/B$ for both the maxima and the minima. From the slope of the linear fits (solid lines) we obtain the value of the sound velocity, $v_s \approx 3.4 \text{ km/s}$ for all three samples, again confirming $\omega_s \propto \sqrt{n_e}$ scaling.




FIG. 3: (color online) Normalized PIRO amplitude $A\epsilon^{1/2}$ (left axis) and the oscillatory part of the resistivity $\delta\rho$ (right axis) vs $\epsilon = \omega_s/\omega_c$ measured in sample A (a), sample B (b), and sample C (c). Solid lines are fits to the data. Dotted (dashed) vertical lines are drawn at $\epsilon^+ = n + 1/8$ ($\epsilon^- = n + 5/8$).

This value is close to the velocity of the bulk transverse acoustic mode, $v_T = 3.35$ km/s.

We further notice that none of the fits pass through zero at $1/B = 0$. As shown in the insets of Fig. 2, all of the fits reveal a *negative* intercept at $-\delta \approx -0.12$ suggesting that the positions of the PIRO maxima⁽⁺⁾ and minima⁽⁻⁾ in all of our samples can be described by

$$\epsilon^+ = n + \delta, \quad \epsilon^- = n + \frac{1}{2} + \delta, \quad \delta \approx 0.12. \quad (4)$$

From this value of the intercept δ we obtain $\varphi \approx -0.24\pi$ appearing in Eq. (3).

To verify the validity of Eq. (4) for individual maxima and minima, we convert $1/B$ to ϵ and present the oscillatory part of the resistivity $\delta\rho$ as a function of $\epsilon = \omega_s/\omega_c$ in Fig. 3 (right axis). Plotted in this way, the data confirm that the period is equal to unity and that *all* maxima (minima), in all of our samples, are *consistently* shifted from integer (half-integers) ϵ to higher values. Except for a few deviations, the value of the shift is roughly the same for all extrema as demonstrated by vertical dotted (dashed) lines drawn at $\epsilon^+ = n + 1/8$ ($\epsilon^- = n + 5/8$) for the maxima (minima). This observation further confirms that PIRO in all of our samples under the conditions of

the experiment can be well described considering a single phonon mode.

We now discuss our experimental findings in terms of recent theoretical proposals.^{64,65} The first study,⁶⁴ considering *anisotropic* phonon dispersion, identified important contributions from three phonon modes – piezoelectric transverse mode polarized along [001] (out-of-plane) direction and two deformational longitudinal modes polarized along [100] and [110] (in-plane) directions. Analytical expressions for partial contributions from these modes, obtained for the case of narrow quantum wells, $k_F b \lesssim 1$ and in the limit of $\epsilon \gg 1$, revealed the phase φ of $-\pi/2$, $\pi/4$, and $-\pi/4$, respectively.⁷⁰ However, in all of our samples $k_F b \gtrsim 4$, and numerical calculations,⁶⁴ performed for sample A used in the present study, revealed a considerably *reduced* phase of the transverse contribution for *all* of the experimentally detected oscillation orders (see, Fig. 3 (b) in Ref. 64). While in this calculation the phase does approach $\varphi = -\pi/2$, it happens only at very high oscillation orders, $\epsilon^{1/2} \sim k_F b$, not accessible in our experiment.

In a more recent study⁶⁵ it was argued that for the case of wide quantum wells, $k_F b \gg 1$, only phonons with very small out-of-plane momentum, $q_\perp \sim b^{-1} \ll k_F$, interact with electrons, provided that ϵ is not too large, $\epsilon^{1/2} \lesssim k_F b$.⁷¹ As a result, the electron-phonon scattering form-factor can be approximated by a delta-function and in the limit of high temperature, $k_B T \gg \hbar\omega_c$, $\hbar\omega_s$,⁷² the oscillatory part of the resistivity is given by

$$\begin{aligned} \frac{\delta\rho}{\rho} &= \frac{g^2 k_B T \tau}{\hbar} \lambda^2 [J_0(2\pi\epsilon) - J_2(2\pi\epsilon)] \\ &\simeq \frac{2g^2 k_B T \tau}{\pi \hbar \sqrt{\epsilon}} \lambda^2 \cos(2\pi\epsilon - \pi/4). \end{aligned} \quad (5)$$

Here, g is the dimensionless electron-phonon coupling constant,⁷³ and τ is the transport scattering time. The second line of Eq. (5) is justified for $2\pi\epsilon \gg 1$, i.e. it works well already at $\epsilon = 1$. Direct comparison with Eq. (3) clearly reveals $\varphi = -\pi/4$, which is in excellent quantitative agreement with our experimental result, $\varphi \simeq -0.24\pi$.

Finally, we analyze the oscillation amplitude A on ϵ in terms of Eq. (5) which predicts $A \propto \epsilon^{-1/2} \lambda^2 = \epsilon^{-1/2} e^{-\alpha\epsilon}$, where $\alpha = 2\pi/\tau q \omega_s$. In Fig. 3 (left axis) we present reduced oscillation amplitude $A\epsilon^{1/2}$ as a function of ϵ and observe roughly exponential dependence, which further confirms the dominant contribution of a single phonon mode in our experiments. The deviations from this dependence are likely due to subleading contributions from other modes, e.g. interface phonons, originally proposed to explain PIRO^{4,69} and/or weakly excited longitudinal modes^{64,65} which have subleading contributions under the conditions of our experiment.

In summary, we have investigated the period and the phase of phonon-induced resistance oscillations in a very-high mobility 2DEG. Due to a very low level of disorder in our samples we were able to reliably measure up to eight

oscillations at relatively low temperatures which ensured dominant contribution of the transverse acoustic phonon mode. The extracted value of the sound velocity is in good agreement with the known value of $v_T = 3.35$ km/s. Furthermore, in all of the samples studied we found that the oscillations exhibit a nonzero phase, $\varphi \approx -0.24\pi$. This value is in excellent agreement both with the analytical expression⁶⁵, obtained for the experimentally relevant case of wide quantum well and not too large ϵ , and with numerical calculations.⁶⁴

We thank I. Dmitriev and O. Raichev for discussions. The work at Minnesota was supported by NSF Grant No. DMR-0548014. The work at Princeton was partially funded by the Gordon and Betty Moore Foundation as well as the NSF MRSEC Program through the Princeton Center for Complex Materials (DMR-0819860).

-
- * Corresponding author: zudov@physics.umn.edu
- ¹ V. L. Gurevich and Y. A. Firsov, *Sov. Phys. JETP* **13**, 137 (1961).
 - ² D. C. Tsui, T. Englert, A. Y. Cho, and A. C. Gossard, *Phys. Rev. Lett.* **44**, 341 (1980).
 - ³ J. L. T. Waugh and G. Dolling, *Phys. Rev.* **132**, 2410 (1963).
 - ⁴ M. A. Zudov, I. V. Ponomarev, A. L. Efros, R. R. Du, J. A. Simmons, et al., *Phys. Rev. Lett.* **86**, 3614 (2001).
 - ⁵ N. V. Zavaritskii and Z. D. Kvon, *JETP Lett.* **38**, 97 (1983).
 - ⁶ V. I. Ryzhii, *Sov. Phys. Solid State* **11**, 2078 (1970).
 - ⁷ V. I. Ryzhii, R. A. Suris, and B. S. Shchamkhalova, *Sov. Phys. Semicond.* **20**, 1299 (1986).
 - ⁸ M. A. Zudov, R. R. Du, J. A. Simmons, and J. L. Reno, *Phys. Rev. B* **64**, 201311(R) (2001).
 - ⁹ P. D. Ye, L. W. Engel, D. C. Tsui, J. A. Simmons, J. R. Wendt, et al., *Appl. Phys. Lett.* **79**, 2193 (2001).
 - ¹⁰ A. C. Durst, S. Sachdev, N. Read, and S. M. Girvin, *Phys. Rev. Lett.* **91**, 086803 (2003).
 - ¹¹ X. L. Lei and S. Y. Liu, *Phys. Rev. Lett.* **91**, 226805 (2003).
 - ¹² I. A. Dmitriev, A. D. Mirlin, and D. G. Polyakov, *Phys. Rev. Lett.* **91**, 226802 (2003).
 - ¹³ R. L. Willett, L. N. Pfeiffer, and K. W. West, *Phys. Rev. Lett.* **93**, 026804 (2004).
 - ¹⁴ R. G. Mani, J. H. Smet, K. von Klitzing, V. Narayana-murti, W. B. Johnson, et al., *Phys. Rev. Lett.* **92**, 146801 (2004).
 - ¹⁵ I. A. Dmitriev, M. G. Vavilov, I. L. Aleiner, A. D. Mirlin, and D. G. Polyakov, *Phys. Rev. B* **71**, 115316 (2005).
 - ¹⁶ S. I. Dorozhkin, J. H. Smet, V. Umansky, and K. von Klitzing, *Phys. Rev. B* **71**, 201306(R) (2005).
 - ¹⁷ S. A. Studenikin, M. Potemski, A. Sachrajda, M. Hilke, L. N. Pfeiffer, et al., *Phys. Rev. B* **71**, 245313 (2005).
 - ¹⁸ X. L. Lei and S. Y. Liu, *Phys. Rev. B* **72**, 075345 (2005).
 - ¹⁹ C. L. Yang, R. R. Du, L. N. Pfeiffer, and K. W. West, *Phys. Rev. B* **74**, 045315 (2006).
 - ²⁰ X. L. Lei, *Phys. Rev. B* **73**, 235322 (2006).
 - ²¹ W. Zhang, M. A. Zudov, L. N. Pfeiffer, and K. W. West, *Phys. Rev. Lett.* **98**, 106804 (2007).
 - ²² I. A. Dmitriev, A. D. Mirlin, and D. G. Polyakov, *Phys. Rev. Lett.* **99**, 206805 (2007).
 - ²³ S. A. Studenikin, A. S. Sachrajda, J. A. Gupta, Z. R. Wasilewski, O. M. Fedorych, et al., *Phys. Rev. B* **76**, 165321 (2007).
 - ²⁴ A. T. Hatke, H.-S. Chiang, M. A. Zudov, L. N. Pfeiffer, and K. W. West, *Phys. Rev. B* **77**, 201304(R) (2008).
 - ²⁵ A. T. Hatke, H.-S. Chiang, M. A. Zudov, L. N. Pfeiffer, and K. W. West, *Phys. Rev. Lett.* **101**, 246811 (2008).
 - ²⁶ M. Khodas and M. G. Vavilov, *Phys. Rev. B* **78**, 245319 (2008).
 - ²⁷ I. A. Dmitriev, M. Khodas, A. D. Mirlin, D. G. Polyakov, and M. G. Vavilov, *Phys. Rev. B* **80**, 165327 (2009).
 - ²⁸ X. L. Lei, *Phys. Rev. B* **79**, 115308 (2009).
 - ²⁹ A. T. Hatke, M. A. Zudov, L. N. Pfeiffer, and K. W. West, *Phys. Rev. Lett.* **102**, 066804 (2009).
 - ³⁰ O. M. Fedorych, M. Potemski, S. A. Studenikin, J. A. Gupta, Z. R. Wasilewski, et al., *Phys. Rev. B* **81**, 239903 (2010).
 - ³¹ M. Khodas, H. S. Chiang, A. T. Hatke, M. A. Zudov, M. G. Vavilov, et al., *Phys. Rev. Lett.* **104**, 206801 (2010).
 - ³² Y. Dai, R. R. Du, L. N. Pfeiffer, and K. W. West, *Phys. Rev. Lett.* **105**, 246802 (2010).
 - ³³ A. T. Hatke, M. A. Zudov, L. N. Pfeiffer, and K. W. West, *Phys. Rev. B* **83**, 121301(R) (2011).
 - ³⁴ A. T. Hatke, M. A. Zudov, L. N. Pfeiffer, and K. W. West, *Phys. Rev. B* **83**, 201301(R) (2011).
 - ³⁵ C. L. Yang, J. Zhang, R. R. Du, J. A. Simmons, and J. L. Reno, *Phys. Rev. Lett.* **89**, 076801 (2002).
 - ³⁶ A. A. Bykov, J. Zhang, S. Vitkalov, A. K. Kalagin, and A. K. Bakarov, *Phys. Rev. B* **72**, 245307 (2005).
 - ³⁷ W. Zhang, H.-S. Chiang, M. A. Zudov, L. N. Pfeiffer, and K. W. West, *Phys. Rev. B* **75**, 041304(R) (2007).
 - ³⁸ X. L. Lei, *Appl. Phys. Lett.* **90**, 132119 (2007).
 - ³⁹ M. G. Vavilov, I. L. Aleiner, and L. I. Glazman, *Phys. Rev. B* **76**, 115331 (2007).
 - ⁴⁰ A. T. Hatke, M. A. Zudov, L. N. Pfeiffer, and K. W. West, *Phys. Rev. B* **79**, 161308(R) (2009).
 - ⁴¹ A. T. Hatke, M. A. Zudov, L. N. Pfeiffer, and K. W. West, *Phys. Rev. B* **83**, 081301(R) (2011).
 - ⁴² R. G. Mani, J. H. Smet, K. von Klitzing, V. Narayana-murti, W. B. Johnson, et al., *Nature (London)* **420**, 646 (2002).
 - ⁴³ M. A. Zudov, R. R. Du, L. N. Pfeiffer, and K. W. West, *Phys. Rev. Lett.* **90**, 046807 (2003).
 - ⁴⁴ C. L. Yang, M. A. Zudov, T. A. Knuuttila, R. R. Du, L. N. Pfeiffer, et al., *Phys. Rev. Lett.* **91**, 096803 (2003).
 - ⁴⁵ A. V. Andreev, I. L. Aleiner, and A. J. Millis, *Phys. Rev. Lett.* **91**, 056803 (2003).
 - ⁴⁶ M. A. Zudov, *Phys. Rev. B* **69**, 041304(R) (2004).
 - ⁴⁷ J. H. Smet, B. Gorshunov, C. Jiang, L. Pfeiffer, K. West, et al., *Phys. Rev. Lett.* **95**, 116804 (2005).
 - ⁴⁸ A. Auerbach, I. Finkler, B. I. Halperin, and A. Yacoby, *Phys. Rev. Lett.* **94**, 196801 (2005).
 - ⁴⁹ M. A. Zudov, R. R. Du, L. N. Pfeiffer, and K. W. West, *Phys. Rev. B* **73**, 041303(R) (2006).
 - ⁵⁰ M. A. Zudov, R. R. Du, L. N. Pfeiffer, and K. W. West, *Phys. Rev. Lett.* **96**, 236804 (2006).
 - ⁵¹ A. A. Bykov, A. K. Bakarov, D. R. Islamov, and A. I. Toropov, *JETP Lett.* **84**, 391 (2006).
 - ⁵² S. Wiedmann, G. M. Gusev, O. E. Raichev, A. K. Bakarov, and J. C. Portal, *Phys. Rev. Lett.* **105**, 026804 (2010).
 - ⁵³ I. V. Andreev, V. M. Muravev, I. V. Kukushkin, S. Schmult, and W. Dietsche, *Phys. Rev. B* **83**, 121308(R) (2011).
 - ⁵⁴ S. I. Dorozhkin, L. Pfeiffer, K. West, K. von Klitzing, and J. H. Smet, *Nature Physics* **7**, 336 (2011).
 - ⁵⁵ A. A. Bykov, J.-Q. Zhang, S. Vitkalov, A. K. Kalagin, and A. K. Bakarov, *Phys. Rev. Lett.* **99**, 116801 (2007).
 - ⁵⁶ A. T. Hatke, H.-S. Chiang, M. A. Zudov, L. N. Pfeiffer, and K. W. West, *Phys. Rev. B* **82**, 041304(R) (2010).
 - ⁵⁷ G. M. Gusev, S. Wiedmann, O. E. Raichev, A. K. Bakarov, and J. C. Portal, *Phys. Rev. B* **83**, 041306 (2011).
 - ⁵⁸ W. Zhang, M. A. Zudov, L. N. Pfeiffer, and K. W. West, *Phys. Rev. Lett.* **100**, 036805 (2008).
 - ⁵⁹ M. A. Zudov, H.-S. Chiang, A. T. Hatke, W. Zhang, L. N. Pfeiffer, et al., *Int. J. of Mod. Phys. B* **23**, 2684 (2009).
 - ⁶⁰ A. T. Hatke, M. A. Zudov, L. N. Pfeiffer, and K. W. West, *Phys. Rev. Lett.* **102**, 086808 (2009).
 - ⁶¹ A. A. Bykov and A. V. Goran, *JETP Lett.* **90**, 578 (2009).
 - ⁶² A. A. Bykov, A. V. Goran, and S. A. Vitkalov, *Phys. Rev. B* **81**, 155322 (2010).

- ⁶³ X. L. Lei, Phys. Rev. B **77**, 205309 (2008).
- ⁶⁴ O. E. Raichev, Phys. Rev. B **80**, 075318 (2009).
- ⁶⁵ I. Dmitriev, R. Gellmann, and M. G. Vavilov, Phys. Rev. B **82**, 201311(R) (2010).
- ⁶⁶ O. E. Raichev, Phys. Rev. B **81**, 165319 (2010).
- ⁶⁷ C. L. Yang, M. A. Zudov, J. Zhang, R. R. Du, J. A. Simmons, et al., Physica E (Amsterdam) **12**, 443 (2002).
- ⁶⁸ A. A. Bykov, A. K. Kalagin, and A. K. Bakarov, JETP Lett. **81**, 523 (2005).
- ⁶⁹ I. V. Ponomarev and A. L. Efros, Phys. Rev. B **63**, 165305 (2001).
- ⁷⁰ Longitudinal modes become relevant at higher temperatures than those used in our experiment.
- ⁷¹ For $\epsilon^{1/2} \gg \max\{1, k_F b\}$, the phase of PIRO should approach $\varphi = -\pi/2$, regardless of the quantum well width.
- ⁷² While this condition is not satisfied in our experiment, its violation is not expected to affect the phase.
- ⁷³ Using $h_{14} = 1.2$ V/nm, $\bar{\rho} = 5.3$ g/cm³, $m^* = 0.067m_0$, and $b = 30$ nm, we estimate $g = (eh_{14}/\hbar\omega_s)\sqrt{m^*/2\bar{\rho}b} \simeq 0.03$. Note that this differs from the expression in Ref. 65 considering deformational phonons.