



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Universal features of the excitation spectrum in a generalized Gibbs distribution ensemble

E. Demler and A. M. Tsvelik

Phys. Rev. B **86**, 115448 — Published 27 September 2012

DOI: [10.1103/PhysRevB.86.115448](https://doi.org/10.1103/PhysRevB.86.115448)

Universal Features of the Excitation Spectrum in Generalized Gibbs Distribution Ensemble.

E. Demler and A. M. Tsvelik*

Harvard-MIT Center for Ultracold Atoms, Cambridge, MA

**Department of Condensed Matter Physics and Materials Science,*

Brookhaven National Laboratory, Upton, NY 11973-5000, USA

It is shown that excitation spectra of Generalized Gibbs Ensembles (GGE) of one-dimensional integrable models with isotopic symmetry contain universal features insensitive to details of the distribution. Namely, the low energy limit of the subsystem of isotopic (for instance, spin) excitations is described by the effective action of a ferromagnet at thermodynamic equilibrium with a single temperature and with the stiffness determined by the initial conditions. The condition of universality is that the entropy per excited particle is small.

PACS numbers: 71.10.Pm

I. GENERAL

Dynamics of closed interacting many-body systems has lately become a subject of intense interest. Systems can be prepared in out-of-equilibrium state by any non-adiabatic change of the Hamiltonian (for instance, by quench) or by pumping. In such experiments many-body system is prepared in a state which is not an eigenstate of the Hamiltonian. It is then allowed to evolve coherently with many eigenstates evolving with different energies. In this paper we will focus exclusively on the fate of integrable systems.

After some evolution time, which for integrable systems we expect to be of the order of the energy of the lowest energy excitation, the system can be considered as completely dephased. Then when calculating expectation value of any physical operator, it is reasonable to neglect off-diagonal terms between different eigenstates since such off-diagonal terms oscillate very rapidly in time and cancel each other. To be more rigorous one may consider averaging of expectation values over small time interval. Then it becomes possible to describe the system not as a pure state, but as a density matrix corresponding to some Generalized Gibbs Ensemble (GGE). Physical properties of the resulting GGE are determined by the integrals of motion which expectation values in the initial state fully characterize it^{1, 2, 3, 4, 5}. Conditions for validity of GGE for general integrable systems have been recently discussed in⁶.

A priori there is no reason to expect any universality of resulting GGE. However, in the next sections we will demonstrate that GGEs appearing in quench experiments with one-dimensional integrable models with isotopic symmetry, have a universal structure that are insensitive to details of the distribution. Moreover, the low energy limit of the subsystem of isotopic (for instance, spin) excitations is corresponds to a ferromagnet at thermodynamic equilibrium with a single temperature.

The defining feature of intergrable systems is existence of infinite set of mutually commuting operators I_j (integrals of motion). Alongside with the momentum operator $I_0 = P$ and the Hamiltonian $I_1 = H$ these conserving quantities can be extracted by expanding the logarithm of transfer matrix in spectral parameter θ :

$$\ln T(\theta) = P + i\theta H + (i\theta)^2 I_2/2 + \dots \quad (1)$$

Generalized Gibbs Ensemble (GGE) described above is given by:

$$P = Z^{-1} \exp\left(-\sum_j \beta_j I_j\right), \quad (2)$$

where I_j are integrals of motion and β_j are corresponding Lagrange multipliers which value is determined by the initial values of I_j . It maximizes entropy while taking into account the constraints imposed by the conserved quantities^{1, 7}. This hypothesis has been proven in some particular cases^{5, 8}.

In this paper we consider several typical examples of GGE in relativistic massive field theories possessing some isotopic symmetry. To work with such models is easier because their integrals of motion are known explicitly being fixed by the Lorentz symmetry and requirements of locality. However, as questions may emerge from the point of view of practicality of such models, we will first address prospects of their experimental realization.

II. MODELS

In this Section we give several examples of integrable models which find applications in cold atom physics. Some of them appear in the context of one-dimensional Bose gas. The situation relevant to our discussion is an experiment where a one-dimensional trap containing condensate is abruptly split to become a double-well potential as, for instance, was done in⁹. We will be interested in the situation when the split is not complete such that there is still a substantial inter-well tunneling. In the low energy limit interacting 1D Bose condensate is uniquely characterized by its local phase field $\phi(t, x)$ which dynamics is described by the Tomonaga-Luttinger liquid model (the model of non-interacting bosonic field). When the trap is split there are two condensates and an additional term appears in the Hamiltonian which describes their coupling. As has been demonstrated (see e.g.¹⁰), the Hamiltonian describing a system of two parallel traps consists of two independent parts. One describes the symmetric part of the condensate phase field $\phi_+ = (\phi_1 + \phi_2)/\sqrt{2}$ and the other one describes the asymmetric part $\phi_- = (\phi_1 - \phi_2)/\sqrt{2}$:

$$H = H_+ + H_-, \quad H_+ = \frac{v}{2} \int_0^L dx \left[K(\partial_x \phi_+)^2 + K^{-1}(\partial_x \theta_+)^2 \right] \quad (3)$$

$$H_- = \frac{v}{2} \int_0^L dx \left[K(\partial_x \phi_-)^2 + K^{-1}(\partial_x \theta_-)^2 - \lambda \cos(\sqrt{2\pi} \phi_-) \right]. \quad (4)$$

where v is the phase velocity, K is the Luttinger parameter normalized in such a way that in Tonks-Girardeau limit $K = 1$, and parameter λ is proportional to the inter-trap tunneling. Fields ϕ_a, θ_a obey the standard commutation relations:

$$[\phi_a(x), \theta_b(y)] = \delta_{ab} \Theta_H(x - y), \quad (5)$$

where $\Theta_H(x)$ is the Heaviside function. The low energy limit of the original bosonic fields is given by

$$\psi_{1,2} = \sqrt{\rho_0} e^{i\sqrt{\pi/2}\phi_+} e^{\pm i\sqrt{\pi/2}\phi_-} + \dots, \quad (6)$$

where dots stand for operators with higher dimensions.

Models (3,4) are integrable; the first one describes a free field, the second one is the famous sine-Gordon model. For $K > 1/4$ the cosine term in (4) is relevant and scales to strong coupling. As a result the sine-Gordon model has massive spectrum of the relativistic form:

$$E_n(p) = \sqrt{(vp)^2 + M_n^2}. \quad (7)$$

Its excitations include solitons and antisolitons corresponding to kinks interpolating between neighboring minima of the cosine potential and (for $K > 1/2$) their bound states called breathers. Their masses in term of the soliton mass M_0 are

$$M_n = 2M_0 \sin \left[\frac{\pi n}{2(4K - 1)} \right], \quad n = 1, 2, \dots, [4K - 1]. \quad (8)$$

Sine-Gordon model (4) with an additional chemical potential field also describes the superfluid-insulator transition¹¹. This fact is important for practical applications since in cold atom physics there are well developed techniques for observation of such transition and of its associated features¹².

Another relevant model of ultracold atom physics is the model of SU(N) fermions with a point-like attractive interaction. The corresponding Hamiltonian is

$$H - \mu N = \int_0^L dx \left(\frac{1}{2m} \partial_x \psi_a^\dagger \partial_x \psi_a - \mu \psi_a^\dagger \psi_a - g \sum_{a \neq b} \psi_a^\dagger \psi_a \psi_b^\dagger \psi_b \right) \quad (9)$$

where $a, b = 1, 2, \dots, N$. Experimentally one dimensional systems of SU(2) fermions have been realized by Moritz *et al.*¹³. Fermions with SU(N) symmetry have been recently realized in optical lattices in^{14,15}. Earlier theoretical work on SU(N) fermions focused on their equilibrium properties¹⁶⁻¹⁹ and their applications to quantum information processing²⁰.

In the limit of weak coupling $g \ll (\mu/m)^{1/2}$ model (9) becomes equivalent to another famous integrable model, the so-called Chiral Gross-Neveu one. In this limit one can linearize the spectrum of fermions near the Fermi points and replace the operators:

$$\psi(x) = e^{-ik_F x} R(x) + e^{ik_F x} L(x), \quad (10)$$

where slow fields R, L contains Fourier harmonics with momenta much smaller than the Fermi momentum k_F . Substituting (10) into (9) one obtains

$$H = \int_0^L dx \left(-ivR_\alpha^+ \partial_x R_\alpha + ivL_\alpha^+ \partial_x L_\alpha - gR_\alpha^+ L_\alpha L_\beta^+ R_\beta \right), \quad (11)$$

where $v = k_F/m$.

Both models (9) and its relativistic limit (11) are integrable and together with the sine-Gordon model (4) are among the best studied models of that kind. At $g > 0$ model (11) is asymptotically free with the interaction scaling to strong coupling in the infrared limit. The spectrum of (11) is split into two independent parts, one of which remains gapless and will not be discussed. The spin sector having SU(N) symmetry have massive excitations with spectrum (7) consisting of the fundamental particle with mass $M_0 \sim \mu(g/v)^{1/N} \exp(-2\pi v/Ng)$ and its bound states

$$M_n = M_0 \frac{\sin(\pi n/N)}{\sin(\pi/N)}, \quad n = 1, 2, \dots, N-1. \quad (12)$$

These excitations all carry isotopic indices of the SU(N) group and transform according to its fundamental representations described by vertical column of Yang tableau of length n .

Model (11) is particularly interesting for us since it has a non-Abelian continuous symmetry which plays a principal role in the subsequent discussion.

III. TBA EQUATIONS FOR GGE

Models (4,11) were among the first field theories solved by the Bethe Ansatz (BA). To simplify our consideration we discuss in detail only the simplest case of SU(2) invariant Gross-Neveu (GN) model $N = 2$ and also touch on the sine-Gordon model (4).

A. General facts about Bethe ansatz

Relativistic (Lorentz invariant) models are particularly convenient for our discussion since they admit a simple classification of integrals of motion. This classification becomes particularly transparent in the so-called rapidity representation when energy and momentum of particle of mass M are parametrized as

$$E = M \cosh \theta, \quad P = M \sinh \theta, \quad E^2 - P^2 = M^2, \quad (13)$$

(we set $v = 1$ for simplicity). Then Lorentz transformation (boost) becomes just a shift of rapidities of all particles: $\theta_i \rightarrow \theta_i + \alpha$. Consequently, all integrals of motion can be classified according to their Lorentz spin:

$$E \pm P = M \sum_{i=1}^{\mathcal{N}} e^{\pm \theta_i}, \quad I_{l+1}^{(0)} \pm I_{l+1}^{(1)} = M \sum_{i=1}^{\mathcal{N}} e^{\pm (l+1)\theta_i}, \quad l = 1, 2, \dots \quad (14)$$

Being a Lorentz invariant object the N -body scattering matrix depends only on difference of rapidities of individual particles. For integrable models such S -matrix can be written as a product of two-body scattering matrices. Until the system is somehow restricted (for instance, placed in a box), particles rapidities in (14) are arbitrary. However, as soon as motion of the particles is restricted, this changes.

The Bethe ansatz (BA) equations which determine eigenvalues of all integrals of motion for a model in a box of length L with periodic boundary conditions can be derived from the straightforward solution²¹ or taking the SU(N)-invariant solution of the Yang-Baxter equations for the two-particle scattering matrix and applying the methods of factorized scattering²². The condition for the periodicity of the wave function of relativistic interacting particles of mass M is

$$\exp(iML \sinh \theta_i) \vec{\xi} = \prod_{j \neq i}^{\mathcal{N}} \hat{S}(\theta_i - \theta_j) \vec{\xi}, \quad (15)$$

where $\vec{\xi}$ is a vector depending of spin indices of the particles. The meaning of this equation is straightforward: the i -th particle going around the system scatters on all others (it does it one-by-one which is the condition of integrability)

acquiring a phase factor given by the product of all S -matrices on the right hand side of (15). This phase factor is compensated by $\exp(ip_i L)$ in the left hand side of this equation.

For models with internal (isotopic) symmetry \hat{S} is a tensor and diagonalization of (15) requires some effort. The result of this diagonalization is the so-called nested Bethe ansatz. For the GN and the sine-Gordon models the result is

$$\exp(iML \sinh \theta_i) = \prod_{j \neq i}^{\mathcal{N}} S_0(\theta_i - \theta_j) \prod_{a=1}^{\mathcal{M}} e_1(\theta_i - \lambda_a) \quad (16)$$

$$\prod_{i=1}^{\mathcal{N}} e_1(\lambda_a - \theta_i) = \prod_{b=1}^{\mathcal{M}} e_2(\lambda_a - \lambda_b) \quad (17)$$

where for the GN model

$$e_n(x) = \frac{x - in\pi/2}{x + in\pi/2}$$

and for the sine-Gordon it is

$$e_n(x) = \frac{\sinh \left[\frac{\gamma}{2}(x - in\pi/2) \right]}{\sinh \left[\frac{\gamma}{2}(x + in\pi/2) \right]}, \quad \gamma = (4K - 1)^{-1}.$$

where $S_0(\theta)$ is some known function which exact form is not important for the present discussion. Number \mathcal{N} stands for the number of particles and \mathcal{M} is related to the spin projection: $S^z = \mathcal{N}/2 - \mathcal{M}$. The qualitative difference between the GN and the sine-Gordon model is that in the latter case for $K > 1/2$ function $S_0(\theta)$ have poles on the physical strip and there are bound states (breathers). These breathers carry no spin. We have to add that the integrals (14) with integer l are local (they have integer Lorenz spin l); in principle, one can imagine that quench generates non-local integrals with non-integer l . However, the further discussion does not depend on whether it is true or not.

Generalization of the nested BA equations (16,17) for models with other simple Lie group symmetry follows the standard scheme described, for instance, in²³. For a given simple Lie group one has to modify $S_0(\theta)$ and replace (17) with a hierarchy of coupled algebraic equations for rapidities $\lambda^{(j)}$ ($j=1, \dots, N-1$) where $N-1$ is the dimension of the corresponding Karta subalgebra. The structure of the hierarchy reflects the structure of the Dynkin diagram for the given group.

B. Derivation of the universal dynamics

The emergence of the universal spin dynamics can be ascertained already from (16,17). Since the particles are massive, their number in the ground state is zero: $\mathcal{N} = \mathcal{M} = 0$. However, a nontrivial GGE emerges after a work had been performed on the system resulting in a finite particle density. For the infinite system the values of all integrals of motion are determined by the rapidities θ_i ; the auxiliary variables λ appear only when the system is put in a box. Therefore, at least in the limit of small particle density, one can neglect a feedback on θ 's from λ 's and consider the distribution of θ 's as an independent function.

When the ratios I_j/L are finite in thermodynamic limit the distribution function of rapidities θ must decay sufficiently fast at infinity. Then in the limit of large λ 's one can replace in Eq.(17)

$$\prod_{i=1}^{\mathcal{N}} e_1(\lambda_a - \theta_i) \approx [e_1(\lambda_a)]^{\mathcal{N}} \quad (18)$$

which indicates that the spin sector (described by λ 's) decouples from θ 's. In fact Eq.(17) with substitution (18) resembles the BA equation for a spin $S=1/2$ Heisenberg magnet. In order to figure out whether this is ferro- or antiferromagnet more detailed analysis is needed (see below). As we will show, it is ferromagnet. The difference between the $SU(2)$ GN model and the sine-Gordon model is evident already at this stage: in the former case we have an isotropic magnet and in the latter case it is anisotropic ($U(1)$ or easy plane magnet).

In order to see the emergence of universal spin dynamics and establish its conditions we have to derive Thermodynamic Bethe Ansatz (TBA) equations. This derivation follows the standard scheme. First, we establish that

generically complex solutions of Eqs.(17) in the thermodynamic limit ($L \rightarrow \infty, \mathcal{N}/L, \mathcal{M}/L = \text{finite}$) have only fixed imaginary parts. More specifically, these solutions group into clusters with a common real part (the so-called "strings"):

$$\lambda_{n,j;\alpha} = \theta_\alpha^{(n)} + i\pi(n+1-2j)/2 + O(\exp(-\text{const}L)), \quad n = 1, 2, \dots; \quad j = 1, 2, \dots, n. \quad (19)$$

Then we introduce distribution functions of rapidities of string centers $\rho_n(\theta)$ and the distribution function of particle rapidities $\rho_0(\theta)$. Functions $\tilde{\rho}_n(\theta), \tilde{\rho}_0(\theta)$ describe distribution of unoccupied spaces. Their ratios are parametrized by excitation energy functions ϵ_n :

$$\tilde{\rho}_n(\theta)/\rho_n(\theta) = \exp[-\epsilon_n(\theta)], \quad \tilde{\rho}_0(\theta)/\rho_0(\theta) = \exp[-\epsilon_0(\theta)], \quad (20)$$

$$\mathcal{N}/L = \int d\theta \rho_0(\theta), \quad \mathcal{M}/L = \sum_{n=1}^{\infty} n \int d\theta \rho_n(\theta). \quad (21)$$

The entropy of the state is given by the expression

$$S = L \sum_{n=0}^{\infty} \int d\theta [(\rho_n + \tilde{\rho}_n) \ln(\rho_n + \tilde{\rho}_n) - \rho_n \ln \rho_n - \tilde{\rho}_n \ln \tilde{\rho}_n]. \quad (22)$$

The TBA equations are a result of minimization of the generalized free energy

$$\Omega = \sum_j \beta_j I_j - S = \int d\theta K(\theta) \rho_0(\theta) - S \quad (23)$$

where

$$K(\theta) = \sum_k [\beta_k e^{k\theta} + \bar{\beta}_k e^{-k\theta}], \quad (24)$$

subject to constraints imposed by the equations for the distribution functions:

$$\tilde{\rho}_n + \rho_n = s * (\tilde{\rho}_{n-1} + \tilde{\rho}_{n+1}) + \delta_{n,0} \frac{M}{2\pi} \cosh \theta, \quad (25)$$

The result is

$$\epsilon_n(\theta) = s * \ln[1 + e^{\epsilon_{n-1}(\theta)}][1 + e^{\epsilon_{n+1}(\theta)}] - \delta_{n,0} K(\theta) \quad (26)$$

$$\Omega/L = - \int \frac{d\theta}{2\pi} K(\theta) \ln [1 + e^{\epsilon_0(\theta)}] \quad (27)$$

$$s * f(\theta) = \int_{-\infty}^{\infty} \frac{d\theta' f(\theta')}{2\pi \cosh(\theta - \theta')}$$

From these equations one can restore the integrals of motion:

$$I_j = - \frac{\partial \Omega}{\partial \beta_j}. \quad (28)$$

A peculiar property of TBA equations (25,26) is their quasi-locality: given ϵ_n, ρ_n are related only to their nearest neighbors. Therefore, if ϵ_0, ρ_0 are fixed the TBA for $\epsilon_n, n = 1, 2, \dots$ will have the same form as (26), but with K replaced by $G(\theta) = s * \ln(1 + e^{\epsilon_0(\theta)})$.

It is natural to assume that all integrals of motion are extensive quantities $\sim L$ and hence their densities are finite. It follows then from (27) that $\epsilon_0(\theta) \rightarrow -\infty$ at infinity (changing sign more than ones at finite θ). Then the function $G(\theta)$ is integrable which will be important for what follows. On Fig. 1 we give an example of such function. We have chosen ϵ_0 in such a way, that $G(\theta)$ has some nontrivial structure at small θ . This structure is determined by the initial conditions of the quench. All these details, however, does not make a difference in the asymptotic region of large θ (see Fig. 2) where the behavior of ϵ_n is determined by a single integral characteristic of the θ -distribution (see the derivation below).




FIG. 1: An example of integrable function $G(\theta; \{\beta\})$. Function ϵ_0 changes sign six times going to $-\infty$ at both infinities.




FIG. 2: $\epsilon_n(\theta)$ functions for $n = 1, 2, 3$ corresponding to $G(\theta)$ shown at Fig.1. The asymptotics does not depend on fine details of behavior $G(\theta)$ at finite θ .

Inverting the kernels in (25,26) with $n > 1$ we obtain TBA equations for GGE in the following form:

$$\ln[1 + e^{\epsilon_n(\theta)}] - A_{nm} * \ln[1 + e^{-\epsilon_m(\theta)}] = a_n * G(\theta), \quad n, m = 1, 2, \dots \quad (29)$$

$$\tilde{\rho}_n + A_{nm} * \rho_m = a_n * s * \tilde{\rho}_0 \quad (30)$$

where the Fourier images of the kernels are

$$a_n(\omega) = \exp[-\pi n|\omega|/2], \quad A_{nm}(\omega) = \coth(\pi\omega/2) \left[\exp(-|n-m|\pi|\omega|/2) - \exp(-(n+m)\pi|\omega|/2) \right].$$

Eqs.(29,30) with G and $s * \tilde{\rho}_0$ replaced by the delta functions are precisely TBA's for a spin $S=1/2$ Heisenberg ferromagnet in thermodynamic equilibrium (!)²⁴. When these functions are not delta functions, but just sharp peaks, the analogy with the ferromagnet remains valid just asymptotically. For this analogy to hold it is sufficient that the integral $\int G(y)dy$ converges, but is $\gg 1$. This would correspond to low effective temperature limit of the ferromagnet when the free energy is determined by large rapidities (small momenta) such that details of the dispersion at large momenta are not important. Indeed we have

$$a_n * G(\theta) = n\pi \int dy \frac{G(y)}{(\theta - y)^2 + \pi^2 n^2} \rightarrow \frac{n\pi}{2\theta^2} \int dy G(y)$$

$$J/T = \frac{1}{2} \int dy G(y). \quad (31)$$

A similar condition must be satisfied by the function $\rho_0(\theta)$:

$$\int dy \rho_0(y) = \text{finite}. \quad (32)$$

The condition $\int G(y)dy \gg 1$ means that there is little feedback from ϵ_n , $n \geq 1$ to ϵ_0 and therefore one may use function $G(\theta)$ to characterize the state. This is more convenient than to use the intergrals of motion which themselves can be restored from G via (28). From (27) and (31) it follows that the effective exchange integral of our ferromagnet is of the order of the energy density per particle. Since the entropy per particle in the ferromagnet $(T/J)^{1/2}$, the requirement $T/J \ll 1$ is equivalent to requiring the entropy per particle to be small. Under that condition the universal spin dynamics emerges in GGE which is, perhaps, the most striking result of our derivation. It is also clear that the spin subsystem of GGE is at thermal equilibrium and is described by the ordinary Gibbs ensemble with a single temperature.

IV. CORRELATION FUNCTIONS

To understand how the universal dynamics described in the previous subsection is related to observable quantities one has to consider correlation functions. This is a difficult problem and though we are not in a position to offer a detailed solution, we feel obliged to make some remarks.

To be closer to real experimental systems we consider correlators of the bosonic creation and annihilation operators in the model of two coupled Bose condensates (3,4). In the low energy limit these operators are expressed in terms of the phase fields ϕ_{\pm} (6). Since the two sectors of the model are decoupled, the correlation functions factorize. For instance, for the 2-point one we have:

$$\langle\langle \psi_a(t, x) \psi_b^{\dagger}(0, 0) \rangle\rangle \sim \rho_0 \langle\langle e^{i\sqrt{\pi/2}\phi_+(t, x)} e^{-i\sqrt{\pi/2}\phi_+(0, 0)} \rangle\rangle \langle\langle e^{i\sqrt{\pi/2}\phi_-(t, x)} e^{i(1-2\delta_{ab})\sqrt{\pi/2}\phi_-(0, 0)} \rangle\rangle \quad (33)$$

The correlation functions of bosonic exponents of gapless fields in GGE have been calculated²⁵, but correlators of the sine-Gordon fields ϕ_{\pm} have never been analyzed in this context. Below we will discuss some general features of these correlators.

We will discuss the limit of zero effective temperature $T_{eff} = 0$ when all $\epsilon_n \rightarrow \infty$ in (29), but their ratios remain constant. This is the limit when GGE distribution is reduced to a single "vacuum" state. This state is characterized by some distribution of the particle rapidities and is ferromagnetic, that is has maximal possible spin. Using the Lehmann expansion where Green's functions are expanded in matrix elements over excited states $|n\rangle$:

$$\langle \mathcal{O}(t, x) \mathcal{O}^{\dagger}(0, 0) \rangle = \sum_n | \langle n | \mathcal{O}(0, 0) | vac \rangle |^2 e^{-it(E_n - E_{vac}) - ix(P_n - P_{vac})} \quad (34)$$

In the sine-Gordon model excitations are classified as solitons, anti-solitons and their bound states (see Section II). (Anti)solitons can be described as particles with (negative) positive spin projection ($s = \pm 1/2$). The ferromagnetic state in this context corresponds to the state with only one type of particles (for example, solitons). We denote such state as

$$|(\theta_n, \dots, \theta_1)_S; (0)_{AS}\rangle$$

According to²⁶, the following matrix elements do not vanish:

$$\begin{aligned} & \langle (\theta_1, \dots, \theta_n)_S; (0)_{AS} | e^{i\beta\phi_-} | (\theta'_1, \dots, \theta'_{n+N})_S; (\bar{\theta}_1, \dots, \bar{\theta}_N)_{AS} \rangle = \\ & \langle vac | e^{i\beta\phi_-} | (\theta'_1, \dots, \theta'_{n+N})_S; (\bar{\theta}_1, \dots, \bar{\theta}_N, i\pi + \theta_1, \dots, i\pi + \theta_n)_{AS} \rangle \end{aligned} \quad (35)$$

These matrix elements would correspond to excitation of N magnons and therefore the spectral function of bosons contains multi-magnon processes. Unfortunately, the form of these matrix elements is very cumbersome which makes further calculations difficult. At the moment the moment this is all we can say.

V. OTHER MODELS

The results obtained for the SU(2)-invariant model (11) can be easily generalized for any simple Lie group. Qualitative differences appear only when the S-matrix for physical particles contains the so-called RSOS (Restricted

Solid-on-Solid) component. Such models have an exotic degenerate ground state and excitations with non-Abelian statistics. In view of rarity of such problems we will confine ourselves to a brief discussion.

A typical representative of this class of models is the $SU_k(2)$ Wess-Zumino-Novikov-Witten (WZNW) model perturbed by the current-current interaction term:

$$H = \int_0^L dx \left[\frac{2\pi v}{k+2} \left(: J^a J^a : + : \bar{J}^a \bar{J}^a : \right) + g J^a \bar{J}^a \right] \quad (36)$$

where the current operators satisfy $SU(2)$ Kac-Moody algebra of level k :

$$[J^a(x), J^b(y)] = i\epsilon^{abc} J^c(x)\delta(x-y) + \frac{k}{2\pi} \delta^{ab} \delta'(x-y) \quad (37)$$

which, in fact, coincides with the commutation relations of the fermionic bilinears

$$J^a = \sum_{j=1}^k R_{j\alpha}^+ \sigma_{\alpha\beta}^a R_{j\beta}, \quad \bar{J}^a = \sum_{j=1}^k L_{j\alpha}^+ \sigma_{\alpha\beta}^a L_{j\beta}$$

Model (36) describes the $SU(2)$ -invariant sector of the $SU(2) \times SU(k)$ model of fermions²⁷. The soliton excitations of this model carry zero modes of parafermions; this is the origin of the non-Abelian statistics (see, for instance^{28, 29} for the discussion). This model also emerges as a continuum limit of the lattice model of spin $S = k/2$ integrable magnet with a small Ising-like anisotropy Δ ³⁰:

$$H = \sum_j \mathcal{P}_k \left[(S_j^+ S_{j+1}^- + h.c.), S_j^z S_{j+1}^z; \Delta \right] \quad (38)$$

where

$$\mathcal{P}_k(x, y; \Delta) = \sum_{n+m \leq k} A_{nm}(\Delta) x^n y^m$$

is some known polynomial. The coupling constant in (36) $g \sim \sqrt{\Delta - 1}$.

The TBA equations for this model differ from (26,24,25) only in one respect: the driving term is placed not in the first, but in the k -th equation and the free energy is also determined by ϵ_k ²⁷. Therefore inverting the kernels in TBA yields not one, but two sets of independent equations:

$$\ln[1 + e^{\epsilon_{n+k}(\theta)}] - A_{nm} * \ln[1 + e^{-\epsilon_{m+k}(\theta)}] = a_n * G(\theta), \quad n, m = 1, 2, \dots \quad (39)$$

$$\ln[1 + e^{\epsilon_n(\theta)}] - \mathcal{A}_{nm} * \ln[1 + e^{-\epsilon_m(\theta)}] = \mathcal{B}_n * G(\theta), \quad n, m = 1, 2, \dots, k-1 \quad (40)$$

where

$$\mathcal{B}_n = \frac{\sinh[\pi(k-n)\omega/2]}{\sinh(\pi k\omega/2)}, \quad \mathcal{A}_{nm} = 2 \coth(\pi\omega/2) \frac{\sinh[\pi(k - \max(n, m))\omega/2] \sinh[\pi \min(n, m)\omega/2]}{\sinh(\pi k\omega/2)}$$

and the corresponding equations for the densities are

$$\tilde{\rho}_{n+k} + A_{nm} * \rho_{m+k} = a_n * s * \tilde{\rho}_k \quad (41)$$

$$\tilde{\rho}_n + \mathcal{A}_{nm} * \rho_{m+1} = \mathcal{B}_n * s * \tilde{\rho}_k \quad (42)$$

The set of equations for $n > k$ describes spin $S=1/2$ ferromagnet as before. As far as the equations for $n < k$ are concerned, in the low energy limit they describe a conformal theory. From (40) and (42) it follows that at large $|\theta|$

$$\epsilon_n(\theta) \sim e^{-2|\theta|/N}, \quad \tilde{\rho}_n(\theta) \sim e^{-2|\theta|/N}, \quad (43)$$

and, since $p(\theta) = 2\pi \int^\theta d\theta' \tilde{\rho}(\theta')$, the spectrum is linear. The low energy limit of the second one corresponds to the conformal field theory of Z_k parafermions.

VI. PHYSICAL CONSEQUENCIES

We conclude this paper by reiterating its main result: provided the initial state of our system containing a gas of excited particles is a low entropy state with (small entropy per particle), its spin dynamics is universal. This condition is equivalent to the condition $T \ll J$ (the effective exchange integral (31)). Namely, in the low energy limit GGE spin subsystem looks like a ferromagnet at thermodynamic equilibrium. GGE's of models with non-Abelian statistics also contain a critical sector described by some conformal field theory (for the example given in the main text it was the theory of Z_k parafermions). A qualitative explanation for this result is simple: quench creates particles not fixing the total magnetic moment and the system chooses a state with maximal entropy which is a state with maximal total moment.

There two natural questions to ask. The first is how to prepare such a state, the second is how this universal dynamics reveals itself in observable quantities. One possible answer to the first question is to use pumping with an appropriate field generating pairs of excitations in the spin sector. In the case of ultracold atoms in optical lattices such pumping can be performed with optical lattice modulation. A pulse with duration t_0 and frequency ω may excite pairs of particles of type n momenta provided that $\omega > 2M_n$. Assuming that pumping does not give momentum to the system, excitations will be generated as pairs of quasiparticles with opposite momenta. Energy conservation then gives that such quasiparticles will be excited with rapidities centered around $M_n \cosh \theta \approx \omega$ and the width of energy distribution t_0^{-1} .

Now let us discuss the second question. What are the physical consequences of the effectively "thermal" character of the spin subsystem? Obviously, one cannot argue that the form of generic isospin correlation functions will be that of a corresponding ferromagnet. This is clear from the analysis of Section IV. Therefore even though we find the same distribution of eigenstates as in the appropriate ferromagnet, matrix elements of the operators may be very different. Is it possible then to perform any physical experiments, which would demonstrate the corresponding "thermal" character of the isospin sector?

The first option is to measure fluctuations of the magnetization. For the Chiral Gross-Neveu model the former corresponds to selecting any of the $SU(N)$ operators and measuring the value of this operator in a finite segment of the system^{31,32}. Unlike more generic operators, smooth components of the spin density operators have the same matrix elements for the Chiral Gross-Neveu and ferromagnetic models (analogously $N_- \sim \partial_x \phi_-$ operator for the sine-Gordon has the same matrix elements as the S^z operator for the easy-plane ferromagnet). We emphasize that measurements of the spin operator should be repeated many times so that one could extract not just the average value but all fluctuations of the operator. Putting it differently one can say that results of individual measurements should be combined into distribution function. We predict that the distribution function of magnetization fluctuations of the Chiral Gross-Neveu model after a quench should be the same as in a ferromagnet at finite temperature.

The second way of observing thermal character of the isospin sector is to measure its fluctuations of energy. In a thermal ensemble energy fluctuations are given by the specific heat $\langle \Delta E^2 \rangle = c_V T^{33}$. In equilibrium specific heat is itself a function of temperature. Hence by measuring the average energy and its fluctuations one can effectively measure the equation of state. Our analysis suggests that the average energy and fluctuations of the energy in the Chiral Gross-Neveu model following the quench should be given by the equilibrium equation of state of an $SU(N)$ ferromagnet. Measurements of energy fluctuations in the system (or in a fragment of the full system) can be done experimentally (see e.g. experiments³⁴). In these experiments the average energy of an interacting 1D Bose gas was measured in an array of tubes. Recent experiments allow local resolution of individual 1d systems³⁵⁻³⁷, which should make it possible to measure not only the average energy but also energy fluctuations. Assuming separation of the isospin and density sectors, it should also be possible to separate the isospin part of the energy from the total energy.

AMT is grateful to Robert Konik, Andrew Green, Misha Zvonarev and Fabian Essler for interesting and inspiring discussions. We also thank Galileo Galilei Institute where this work was finished for its hospitality. AMT was supported by US DOE under contract number DE-AC02-98 CH 10886. ED acknowledges support from Harvard-MIT CUA, NSF Grant No. DMR-07-05472, the Army Research Office with funding from the DARPA OLE program, AFOSR Quantum Simulation MURI, the ARO-MURI on Atomtronics.

¹ M. Rigol, V. Dunjko, V. Yurovsky and M. Olshanii, Phys. Rev. Lett. **98**, 050405 (2007); Nature **452**, 854 (2008).

² P. Calabrese and J. L. Cardy, J. Stat. Mech: Th. Exp. **2007**: P06008 (2007).

³ M. A. Cazalilla, A. Iucci, M. C. Chang, Phys. Rev. E **85**, 011133 (2012) M. Cramer *et al.*, Phys. Rev. Lett **100**, 030602 (2008); T. Barthel and Schollwöck, *ibid.* **100**, 100601 (2008); G. Roux, Phys. Rev. A **79**, 021608 (2009).

⁴ J. Mossel and J.-S. Caux, New J. Phys. **12**, 055028 (2010); arXiv:1203.1305.

⁵ P. Calabrese, F. H. L. Essler and M. Fagotti, Phys. Rev. Lett. **106**, 227203 (2011).

⁶ D. Fioretto and G. Mussardo, New J. Phys. **12**, 055015 (2010).

- ⁷ E. T. Jaynes, Phys. Rev. B **106**, 620; *ibid.* **108**, 171 (1957).
- ⁸ A. C. Cassidi, C. W. Clark and M. Rigol, Phys. Rev. Lett. **106**, 140405 (2011).
- ⁹ M. Gring, M. Kuhnert, T. Langen, T. Kitagawa, B. Rauer, M. Schreitl, D. A. Smith, E. Demler, J. Schmiedmayer, arXiv:1112.0013.
- ¹⁰ V. Gritsev, A. Polkovnikov, and E. Demler. Phys. Rev. B **75**, 174511 (2007).
- ¹¹ H. P. Büchler, G. Blatter, and W. Zwerger. Phys. Rev. Lett. **90**, 130401 (2003).
- ¹² I. Bloch, J. Dalibard, and W. Zwerger. Rev. Mod. Phys. **80**, 885 (2008).
- ¹³ H. Moritz, T. Stöferle, K. Günter, M. Köhl, and T. Esslinger. Phys. Rev. Lett. **94**, 210401 (2005).
- ¹⁴ G. K. Campbell, M. M. Boyd, J. W. Thomsen, M. J. Martin, S. Blatt, M. D. Swallows, T. L. Nicholson, T. Fortier, C. W. Oates, S. A. Daddams, N. D. Lemke, P. Naidon, P. Julienne, J. Ye, A. D. Ludlow, Science **32**, 360 (2009).
- ¹⁵ S. Taie, Y. Takasu, S. Sugawa, R. Yamazaki, T. Tsujimoto, R. Murakami, Y. Takahashi, Phys. Rev. Lett **105**, 190401 (2010).
- ¹⁶ C. Wu, J-P. Hu, and S-C. Zhang, Phys. Rev. Lett. **91**, 186402 (2003).
- ¹⁷ M. Hermele, V. Gurarie, and A.-M. Rey. Phys. Rev. Lett. **103**, 135301 (2009).
- ¹⁸ C. Honerkamp and W. Hofstetter, Phys. Rev. Lett. **92**, 170403 (2004).
- ¹⁹ A. V. Gorshkov, M. Hermele, V. Gurarie, C. Xu, P. S. Julienne, J. Ye, P. Zoller, E. Demler, M. D. Lukin, A. M. Rey, Nature Physics **6**, 289 (2010).
- ²⁰ A. V. Gorshkov, A. M. Rey, A. J. Daley, M. M. Boyd, J. Ye, P. Zoller, M. D. Lukin, Phys. Rev. Lett. **102**, 110503 (2009).
- ²¹ N. Andrei, J. H. Lowenstein, Phys. Lett B **90**, 106 (1980); *ibid.*, **91**, 401 (1983).
- ²² A. B. Zamolodchikov and Al. B. Zamolodchikov, Ann. Phys. **120**, 253 (1979).
- ²³ E. Ogievetskii, N. Yu. Reshetikhin and P. B. Wiegmann, Nucl. Phys. **280**, 45 (1987).
- ²⁴ M. Takahashi, Prog. Theor. Phys. **46**, 401 (1971).
- ²⁵ A. Mitra and T. Giamarchi, Phys. Rev. B **85**, 075117 (2012).
- ²⁶ S. Lukyanov, A. Zamolodchikov, Nucl. Phys. **607**, 437 (2001).
- ²⁷ A. M. Tsvelick, ZhETP **93**, 1329 (1987).
- ²⁸ F. A. Smirnov, Int. J. Mod. Phys. A **9**, 5121 (1994).
- ²⁹ P. Fendley and H. Saleur, Phys. Rev. D **65**, 025001 (2002).
- ³⁰ A. B. Zamolodchikov and V. A. Fateev, Sov. J. Nucl. Phys. **32**, 298 (1980).
- ³¹ R. W. Cherng and E. Demler, New Journal of Physics **9**, 7 (2007).
- ³² K. Eckert, L. Zawitkowski, A. Sanpera, M. Lewenstein, and E. S. Polzik, Phys. Rev. Lett. **98**, 100404 (2007).
- ³³ P. K. Pathria, P. D. Beale, "Statistical mechanics", third edition, Elsevier, 2011
- ³⁴ T. Kinoshita, T. Wenger, and D. S. Weiss. *Science*, **305** 1125 (2004).
- ³⁵ R. Folman, P. Krger, J. Schmiedmayer, J. Denschlag, and C. Henkel. volume 48 of *Advances In Atomic, Molecular, and Optical Physics*, pp. 263 – 356. Academic Press, 2002.
- ³⁶ J. F. Sherson, C. Weitenberg, M. Endres, M. Cheneau, I. Bloch, S. Kuhr, Nature **467**, 68 (2010).
- ³⁷ W. S. Bakr, J. I. Gillen, A. Peng, S. Folling, M. Greiner, Nature **462**, 74 (2009).