



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Decay of the Kohn mode in hydrodynamic regime

A. Iqbal, A. Levchenko, and M. Khodas

Phys. Rev. B **92**, 024303 — Published 15 July 2015

DOI: [10.1103/PhysRevB.92.024303](https://doi.org/10.1103/PhysRevB.92.024303)

Decay of the Kohn mode in hydrodynamic regime

A. Iqbal,¹ A. Levchenko,^{2,3} and M. Khodas^{1,4}

¹*Department of Physics and Astronomy, University of Iowa, Iowa City, Iowa 52242, USA*

²*Department of Physics and Astronomy, Michigan State University, East Lansing, Michigan 48824, USA*

³*Department of Physics, University of Wisconsin-Madison, Madison, Wisconsin 53706, USA*

⁴*Racah Institute of Physics, Hebrew University of Jerusalem, Jerusalem 91904, Israel*

(Dated: June 2, 2015)

We develop a hydrodynamic description of the collective modes of interacting liquids in a quasi-one-dimensional confining potential. By solving Navier-Stokes equations we determine analytically excitation spectrum of sloshing oscillations. For parabolic confinement, the lowest frequency eigenmode is not renormalized by interactions and is protected from decay by the Kohn's theorem, which states that center of mass motion decouples from internal dynamics. We find that the combined effect of potential anharmonicity and interactions results in the frequency shift and final lifetime of the Kohn mode. All other excited modes of sloshing oscillations thermalize with the parametrically faster rate. Our results are significant for the interpretation of recent experiments with trapped Fermi gases observed weak violation of the Kohn theorem.

PACS numbers: 67.10.Jn, 72.15.Lh, 72.15.Nj

I. INTRODUCTION

Properties of quantum liquids in one-dimension (1D), as realized experimentally in nanoscale semiconducting wires, carbon nanotubes, laser traps of cold atoms, and edge channels of the quantum Hall effect, continue to attract tremendous attention in the physics research (see Refs.¹⁻³ for recent reviews and references herein). With increasing sophistication in high precision measurements and techniques these systems provide serious tests for the existing theoretical models, such as Luttinger liquid theory⁴⁻⁷, and ultimately challenge their completeness. For example, the powerful approach of the Luttinger liquid formalism allows to account for the interaction effects nonperturbatively. However, this model does not adequately describe relaxation phenomena due to built-in approximation of the linearized quasiparticle dispersion, which by virtue of the kinematic constraints effectively closes the phase space available for inelastic scattering. In certain special cases, the lack of relaxation may be a generic property of the many-body system because of its complete integrability^{8,9}. Alternatively, vanishing relaxation rates may happen because of the reasons prescribed by the Kohn theorem^{10,11}.

Motivated by recent experiments¹²⁻¹⁵ we study relaxation of collective excitations in interacting two-dimensional (2D) systems confined along one of the two spatial dimensions. These systems interpolate between strictly one-dimensional limit of Luttinger liquids and the two-dimensional Fermi liquids. The geometrical confinement in such systems splits the single band spectrum into multiple one-dimensional subbands. The convenient and practically justified model idealization is the interacting particles confined by a harmonic potential. In this case the one-dimensional subbands are equidistantly separated by a frequency of oscillations ω_{\perp} across the channel. Similar to the spectrum linearization in the strictly one-dimensional liquids, harmonic approximation in the

quasi-one-dimensional case on one hand simplifies the dynamics, and at the same time does not allow for proper description of thermalization processes. One necessarily has to account for the confinement anharmonicity, which thermalizes the motion across the channel in much the same way as the spectrum nonlinearity causes the relaxation of charge and spin excitations in one-dimensional quantum wires¹⁶⁻²⁰.

Despite the similarity with 1D, the relaxation of transversal excitations has a few distinct features setting these two problems apart. The kinematical constraints of momentum and energy conservations operational in 1D are less restrictive in quasi-1D. In contrast to the 1D case, which require three-particle scattering processes, the two-body collisions do cause the relaxation via the inter-subband transitions. Thermalization in quasi-1D may nevertheless be prohibited due to the Kohn theorem rather than kinematical restrictions.

This theorem states that the motion of the system as a whole is unaffected by interactions. Classically, it follows as the translationally invariant interaction energy is insensitive to the system displacement as a whole. For the same reason, quantum mechanically, the interaction drops out of the center of mass Heisenberg equation of motion^{11,21}. In a quantum Fermi liquid the Kohn theorem follows from the solution of the kinetic equation on the quasi-particle distribution function²².

In all of the above cases the Kohn theorem states that if the confining potential is harmonic the collective sloshing oscillations proceed without decay. The frequency of the Kohn, or so called sloshing mode, ω_{\perp} , is insensitive to interaction, temperature and particle statistics^{11,21,22}.

This fact makes the observation of the Kohn mode possible in a wide variety of systems. In semiconductor quantum wires the Kohn mode is observed in optical transmission at far infrared^{23,24}. In trapped ultracold Fermi gas of ⁶Li the Kohn mode of a half KHz frequency was excited by sudden displacement of the trap and detected




FIG. 1. (color online) The definition of the displacement field $\phi(z, t)$ in Lagrangian formulation. At the time, shown as vertical axis, $t = 0$ the fluid is contained in the interval $|z| < a$. As time progresses the particles of a fluid move as indicated by narrow arrows pointing upward. The particle located at z at $t = 0$ is shifted to the new position $z + \phi(z, t)$. A fluid volume shown as a narrow (red) horizontal rectangle occupies a segment $[z, z + dz]$ at time $t = 0$. At a later time $t > 0$ this volume is displaced and occupies the segment $[z + \phi(z, t), z + dz + \phi(z + dz, t)]$. The fluid volume expands (shrinks) if $\partial\phi(z, t)/\partial z > (<)0$. The expansion (shrinkage) translates in the decrease (increase) in the density respectively.

by absorption imaging of a released cloud^{14,25}.

The weak violation of the Kohn theorem due to anharmonicity observed in the above three classes of systems plays a key role in the data interpretation. In the case of the semiconductor quantum wires it controls the line broadening and the higher harmonics of light transmission²³. The observed sloshing frequency of an atomic cloud in the optical trap shows systematic deviations from the Kohn theorem prediction²⁶. Such deviations grow with heating as the expanding atomic cloud senses a progressively less parabolic confining potential. Here we concentrate on two fundamental aspects of Kohn theorem violation: (i) frequency shift of the sloshing frequency, and (ii) final lifetime of sloshing oscillations. We approach this problem based on a very general grounds of hydrodynamic theory, which accurately describes most liquids at length scales long compared to the particle-particle mean-free path.

II. HYDRODYNAMIC THEORY

A hydrodynamic description is based on the existence of slow variables associated with locally conserved quantities such as number of particles, momentum and energy. The motion of the liquid is described by the Navier-Stokes equations which in Eulerian continuous field coordinates can be put in the form²⁷

$$\partial_t(\rho v_j) = -\partial_i \Pi_{ij} - \rho \partial_j U, \quad (1)$$

that guaranties the momentum conservation, which holds in ideal and nonideal liquids alike in the absence of confining potential U . The stress tensor of a two-dimensional

fluid giving rise to the Navier-Stokes equation is²⁷

$$\Pi_{ij} = \delta_{ij}P + \rho v_i v_j - \zeta \delta_{ij} \partial_k v_k - \eta (\partial_i v_j + \partial_j v_i - \delta_{ij} \partial_k v_k). \quad (2)$$

Here η, ζ are the first (shear) and second (bulk) viscosities, and P is pressure. In the following we focus on a one-dimensional flow in z -direction of a two-dimensional liquid occupying the strip $|z| < a$, so that velocity vector field can be taken in the form $\mathbf{v} = v(z, t)\mathbf{e}_z$ and Eq. (1) simplifies to

$$\rho(\partial_t v + v \partial_z v) = -\partial_z P - \rho \partial_z U + \partial_z(\eta \partial_z v). \quad (3)$$

When writing this equation we made use of the continuity equation $\partial_t \rho + \partial_z(\rho v) = 0$ and employed standard assumption $\eta \gg \zeta$. For the purpose of our study it will be convenient to use a particle description of Navier-Stokes equation. In this approach the coordinate z labels an equilibrium position of a fluid particle, and its location at later time t is the $z + \phi(z, t)$ such that ϕ is the displacement field. By definition, the density is $\rho(z + \phi(z, t), t) = \rho_0(z)/(1 + \partial_z \phi(z, t))$, the velocity $v(z + \phi(z, t)) = \partial_t \phi(z, t)$. The linearization of Eq. (3) is equivalent to the linearization in ϕ . To the leading order,

$$\rho = \rho_0 + \delta\rho, \quad \delta\rho = -(\rho_0 \dot{\phi})', \quad v = \dot{\phi}, \quad (4)$$

where ρ_0 is a stationary equilibrium density distribution, the notations $f' = \partial f/\partial z$ and $\dot{f} = \partial f/\partial t$ are introduced and the pair of arguments (z, t) common to all the functions is omitted. The parametrization (4) of $\delta\rho$ and v by a single displacement field automatically satisfies the linearized continuity equation, $\delta\dot{\rho} + (\rho_0 \dot{\phi})' = 0$. The concept of the displacement field is further illustrated in Fig. 1. For the solutions of the form $\phi(z, t) = e^{i\omega t} \chi'(z)$ the equation (3) in the parameterization, (4) reads, (see App. A):

$$\omega^2 \chi = -v_s^2 \chi'' + U' \chi' - i\omega \int_{z_0}^z \frac{dz}{\rho_0} (\nu \rho_0 \chi'')', \quad (5)$$

where $\nu(z) = \eta(z)/\rho_0(z)$ is the kinematic viscosity, and

$$v_s = \sqrt{\partial P/\partial \rho_0} \quad (6)$$

has a meaning of a local speed of sound that depends on z only through the equilibrium density ρ_0 . We show that the results are independent on arbitrary z_0 .

The spectrum of collective excitations and their decay rates as given by Eq. (5) depend on the details of the confining potential and dependence of the viscosity on density. Here for definiteness we consider the confining potential per unit mass with a weak quartic anharmonicity,

$$U = \frac{\omega_{\perp}^2 z^2}{2} + \frac{\epsilon z^4}{4m} + \delta U, \quad (7)$$

where m is the mass of an individual particles, and the constant $\delta U \propto \epsilon$ is added in such a way that the spatial extent of the liquid stays the same as for $\epsilon = 0$. In other

words $\delta U + \epsilon a^4/4m = 0$. This choice is not obligatory, yet convenient in the further calculations. The anharmonic part $\propto \epsilon$ of U not only modifies the second term in the right-hand-side of Eq. (5) but also the first one. Indeed, in the parabolic confinement the sound velocity is $v_s^2(z) = 2\pi\rho(z)/m^3 = \omega_\perp^2(a^2 - z^2)/2$, which acquires now a correction $\delta v_s^2(z) = -\epsilon(z^4 - a^4)/4m$. With these observations in mind we multiply Eq. (5) by $2/\omega_\perp^2$, rescale coordinates $z \rightarrow z/a$ and introduce $\lambda_\omega^2 = 2\omega^2/\omega_\perp^2$ to find

$$\lambda_\omega^2 \chi + (1 - z^2)\chi'' - 2z\chi' + V_\epsilon + V_\nu = 0. \quad (8)$$

The perturbation term due to anharmonicity can be cast in the form

$$V_\epsilon = -\frac{\epsilon a^2}{2m\omega_\perp^2} [(z^4 - 1)\chi']'. \quad (9)$$

This formula can be generalized to any confining potential after the change $[(z^4 - 1)\chi']' \rightarrow [(U(z) - U(a))\chi']'$. Consequently, our results for the frequency shift are straightforward to modify for arbitrary confinements. We emphasize that V_ϵ is Hermitian. This property guarantees that the anharmonicity alone causes only the frequency shift but no dissipation. We argue below that only the combination of the anharmonicity and viscosity leads to the dissipation of the Kohn mode. The exact z -dependence of the viscosity $\nu(z)$ is specific to the model. This however only influences the numerical prefactors in the final results and we take the expression for the viscosity from the theory of Fermi liquids, $\eta \propto v_F \rho \ell$, where mean free path is $\ell \propto v_F E_F / T^2$ with v_F and E_F being Fermi velocity and energy, respectively. For this case $\nu(z) = C \rho^2(z)/m^5 T^2$ where C is the numerical factor of the order of unity²⁸. In the above specified dimensionless notations this results in

$$V_\nu = -i\lambda_\omega B \int_{z_0}^z \frac{dz}{1 - z^2} [(1 - z^2)^3 \chi'']', \quad (10)$$

where we have introduced $B = \sqrt{2}\nu_0/a^2\omega_\perp$ with $\nu_0 = \nu(z=0)$. We proceed with the analysis of Eq. (8).

III. RESULTS

As the first step let us discuss the eigenmodes of an ideal fluid confined to a harmonic trap. For that purpose we neglect anharmonicity and interaction effects implicit in the viscosity term of Eq. (8), i.e. we set $V_\epsilon = V_\eta = 0$. What remains is familiar Legendre equation and we therefore immediately read off its solutions $\lambda_{\omega_n}^2 = 2\omega_n^2/\omega_\perp^2 = n(n+1)$ with $n = 0, 1, 2, \dots$ so that the eigenfrequencies are

$$\omega_n = \omega_\perp \sqrt{\frac{n(n+1)}{2}}. \quad (11)$$

The $n = 0$ gives an equilibrium since $\chi_{n=0} = \text{const}$ this velocity is identically zero. The $n = 1$ is a Kohn

mode $\omega_1 = \omega_\perp$. We thus found the whole hierarchy of eigenoscillations, they are Legendre polynomials

$$\chi_n(z) = \sqrt{\frac{2n+1}{2}} P_n(z) \quad (12)$$

and the related velocity fields are $v_n(z) = \partial_z P_n$. Remarkably, the same spectrum of collective modes as (11) was obtained recently for the longitudinal oscillations of the one-dimensional Coulomb chains²⁹. In this systems the collective behavior sets in due to the long range Coulomb forces rather than collisions.

Next we discuss the significance of perturbation terms on the spectrum of collective modes. Let us first consider anharmonicity alone, i.e. we set $V_\eta = 0$ and $V_\epsilon \neq 0$ in Eq. (8). As V_ϵ is Hermitian the spectrum remains real. As a result all the eigenmodes remain undamped as expected in the absence of collisions. To find the frequency shift of the Kohn mode, $\delta^{(1)}\omega_1$ to the leading order in ϵ we write $\lambda_{\omega_1}^2 = 2 + \delta^{(1)}\lambda_{\omega_1}^2$, where the correction term is found from the first order perturbation theory

$$\delta^{(1)}\lambda_{\omega_1}^2 = \frac{3\epsilon a^2}{4m\omega_\perp^2} \langle P_1(z) | (z^4 - 1)\partial_z^2 + 4z^3\partial_z | P_1(z) \rangle. \quad (13)$$

The matrix element gives a factor of $8/5$ which eventually translates into the correction to the Kohn mode frequency

$$\delta^{(1)}\omega_1 = \frac{3\epsilon a^2}{10m\omega_\perp}. \quad (14)$$

We have checked that the result (14) agrees with the result obtained by the method of moments suggested in Ref.²⁵. We note however that the result for the depolarization shift in the collisionless regime differ from Eq. (14) by a nonuniversal numerical prefactor. For instance, for the contact interaction we obtain by direct perturbation theory in Fermions $\delta^{(1)}\omega_1 = 3\epsilon a^2/5m\omega_\perp$ ³⁰. Corrections to all other eigenfrequencies can be computed in the same fashion, which will have the same form as above but with the different numerical coefficients.

Before considering the generic case, it is instructive to verify the Kohn's theorem within our hydrodynamic approach. It amounts to the statement that $\chi_1(z) \propto z$ remains the solution of Eq. (8) with the frequency $\omega_1 = \omega_\perp$ even for $V_\eta \neq 0$ provided only $V_\epsilon = 0$. This in turn will be proven once we show that for any non-negative n , $\langle P_1 | V_\eta | P_n \rangle = \langle P_n | V_\eta | P_1 \rangle = 0$. Clearly $V_\eta | P_1 \rangle = 0$ as $[P_1(z)]'' = 0$. On the other hand from Eq. (5) it follows that

$$\langle P_1 | V_\nu | P_n \rangle \propto \int_{-1}^1 d\bar{z} P_1(\bar{z}) \int_{z_0}^{\bar{z}} \frac{dz}{\rho_0(z)} \{ \nu(z) \rho_0(z) [P_n(z)]' \}'. \quad (15)$$

Realizing that $P_1(\bar{z}) = -\partial_z \rho_0(\bar{z})/2$ and integrating by parts one concludes that $\langle P_1 | V_\nu | P_n \rangle = 0$ for all n thus proving the Kohn theorem in the present context.

It follows that the Kohn mode may acquire a finite lifetime only when the anharmonicity is included. Yet the

perturbation V_ϵ is Hermitian and by itself is insufficient. We therefore consider both perturbations, and look for the corrections that are first order in each of the two. In the second order perturbation theory such a correction is of the form

$$\delta^{(2)}\lambda_{\omega_1}^2 = -i\lambda_\omega B \sum_{n=1}^{\infty} \frac{\sqrt{3/2}\sqrt{(2n+1)/2}}{2-n(n+1)} \\ [\langle P_1|V_\nu|P_n\rangle\langle P_n|V_\epsilon|P_1\rangle + \langle P_1|V_\epsilon|P_n\rangle\langle P_n|V_\nu|P_1\rangle]. \quad (16)$$

As we saw above both terms in this equation are zero as the Kohn mode does not couple to any other mode by a viscosity term V_η , and $\delta^{(2)}\lambda_{\omega_1}^2 = 0$.

Inevitably we have to consider the third order corrections to λ_{ω_1} . The third order correction to the energy E_l of a state $|l\rangle$ is

$$\delta^{(3)}E_l = \sum_{k,m \neq l} \frac{\langle l|V|m\rangle\langle m|V|k\rangle\langle k|V|l\rangle}{(E_m - E_l)(E_k - E_l)} \\ - \langle l|V|l\rangle \sum_{m \neq l} \frac{\langle l|V|m\rangle\langle m|V|l\rangle}{(E_m - E_l)^2}. \quad (17)$$

To apply this expression to our problem we identify $|l\rangle = P_1$ as a Kohn mode and $V = V_\epsilon + V_\nu$. We observe that for the perturbation terms given by Eqs. (9) and (10) the following properties of matrix elements hold $\langle P_1|V_\nu|P_n\rangle = \langle P_n|V_\nu|P_1\rangle = \langle P_1|V_\epsilon|P_1\rangle = 0$ for all n . Furthermore, for the quartic anharmonicity under consideration the only nonzero off-diagonal matrix elements are $\langle P_1|V_\epsilon|P_3\rangle = \langle P_3|V_\epsilon|P_1\rangle$ with the rest of matrix elements $\langle P_n|V_\epsilon|P_1\rangle = 0$ for $n \neq 1, 3$. We thus have, accounting for all the normalization factors of eigenoscillation modes (12)

$$\delta^{(3)}\lambda_1^2 = \frac{147}{8(\lambda_1^2 - \lambda_3^2)^2} \langle P_1|V_\epsilon|P_3\rangle\langle P_3|V_\nu|P_3\rangle\langle P_3|V_\epsilon|P_1\rangle. \quad (18)$$

Using the explicit expressions (9) and (10) we find for the matrix elements

$$\int_{-1}^1 d\bar{z} P_3(\bar{z}) \int_{z_0}^{\bar{z}} \frac{dz}{1-z^2} \partial_z [(1-z^2)^3 \partial_z^2 P_3] = -\frac{40}{21}, \quad (19)$$

$$\int_{-1}^1 dz P_1(z) \partial_z [(z^4 - 1) \partial_z P_3(z)] = \frac{8}{5}, \quad (20)$$

and eventually

$$\delta^{(3)}\lambda_1^2 = i\lambda_\omega \frac{28\sqrt{2}}{125} \frac{\nu_0}{a^2\omega_\perp} \left(\frac{\epsilon a^2}{m\omega_\perp^2} \right)^2. \quad (21)$$

This result enables us to find imaginary part of the Kohn mode $\omega_1 = \omega_\perp + \delta\omega_1 + i\tau_1^{-1}$, which corresponds to its attenuation with the rate

$$\tau_1^{-1} \simeq \frac{\nu_0 \epsilon^2 a^2}{m^2 \omega_\perp^4} \simeq \left(\frac{\delta\omega_1}{\omega_\perp} \right)^2 \frac{\nu_0}{a^2}, \quad (22)$$

where we omitted numerical factors of order unity for brevity. This expression constitutes the main result of

our work and has straightforward interpretation. The higher excitations modes not protected by the Kohn theorem decay with the rate $\sim \nu_0/a^2$. As ν_0 has dimensions of the diffusion coefficient, this is the typical rate of the momentum relaxation. The ratio $(\delta\omega_1/\omega_\perp)^2$ is the probability of finding the system in higher modes.

IV. DISCUSSIONS

Hydrodynamic description requires short equilibration length ℓ . Thus validity of our theory is limited by the condition $\ell \ll a$, which imposes certain restriction on temperature. Specifically, for the Fermi liquids $\ell = v_F \tau_{ee}$ is determined by collisions with the typical rate $\tau_{ee}^{-1} \sim T^2/E_F$. Since $\omega_\perp \sim v_F/a$, hydrodynamic regime is realized at temperatures $T > T_h$ above the crossover scale $T_h \sim \sqrt{\omega_\perp E_F} \sim E_F/\sqrt{N}$, where N is the number of occupied sub-bands of the transversal quantization. It also follows that with necessity hydrodynamics requires $T \gg \omega_\perp$. While this inequality is reasonably satisfied for the cold gases that are confined by a very shallow potential, it obviously breaks in the ultra-cold limit where collisionless regime prevails. In the latter case attenuation coefficient of the Kohn mode is expected to follow quadratic temperature dependence $\tau_1^{-1} \propto \alpha T^2/E_F$ based on the Pauli principle and phase space restrictions argument, whereas in the hydrodynamic regime $\tau_1^{-1} \propto 1/T^2$ in accordance with Eq. (22). The nonmonotonic temperature dependence of the decay rate has been observed experimentally²⁶.

Our hydrodynamic approach has interesting parallels with the Luttinger liquid description of collective modes in confined inhomogeneous one-dimensional gases³¹. The eigenvalue equation for the normal eigenmodes in that case, analogous to our Eq. (5), is given by

$$-\omega_n^2 \chi_n(z) = v(z) K(z) \partial_z \left(\frac{v(z)}{K(z)} \partial_z \chi_n(z) \right) \quad (23)$$

where Luttinger liquid interaction parameter satisfies the relation $v(z)K(z) = \pi\rho(z)/m^2$. This equation is supplemented by the boundary condition $\chi_n(\pm a) = 0$ and normalization condition $\int_{-a}^a dz \chi_j(z) \chi_j(z) / v(z) K(z) = \delta_{ij}$. For the particular choice of $v(z) = v_0 \sqrt{1-z^2/a^2}$ and $K(z) = K_0(1-x^2/a^2)^\gamma$ the solutions $\chi_n(z)$ are obtained in terms of Gegenbauer polynomials with the spectrum of excitations $\omega_n^2 = (v_0/a)^2(n+1)(n+2\gamma+1)^{31-33}$. In the model of $\gamma = 2$, the problem simplifies to the case of Legendre polynomials³⁴ with the spectrum of excitations analogous to our result (11). Another interesting limit is $\gamma = 0$, which corresponds to the case of the Tonks-Girardeau gas, where the Gegenbauer polynomials reduce to Chebyshev polynomials. Inclusion of dissipative terms into Eq. (23) requires consideration of corrections to Luttinger liquid model which account for the inelastic scattering of bosons and ultimately describe equilibration processes. As recently shown such generalization is

possible both in the limit of weak³⁵ and strong³⁶ interactions and application of this formalism to the problem of decay of collective modes is an interesting problem for future research. Along this route one may hope to find a unified description, which interpolates between the quantum³⁶ and classical³⁷ hydrodynamic regimes of Luttinger liquids, and which is broadly applicable for arbitrarily strong interactions.

Note. After the submission of this article for the review we became aware of the related study³⁸ where decay of the Kohn mode has been studied in the context of the famous Quantum Newton's Cradle experiment³⁹.

ACKNOWLEDGMENTS

We thank M. Raikh for discussions that attracted our attention to this interesting problem. We indebted to E. Bettelheim for general discussions of the Navier-Stokes equation in the inhomogeneous geometry, and for pointing out to us importance of terms that make this equation consistent with the boundary conditions. We also thank M. Dykman for the discussion of results. The work of A.L. was supported by NSF Grant No. DMR-1401908. The work of M.K. and A.L. has been supported by the National Science Foundation (Grant No. 2014107). A.I. and M.K. are grateful to the University of Iowa for the support.

Appendix A: Displacement field parametrization of the linearized Navier-Stokes equations

In the parametrization (4), the left hand side of Eq. (3) takes the form,

$$\rho(\partial_t v + v\partial_z v) \approx \rho_0 \ddot{\phi}. \quad (\text{A1})$$

To linearize the right hand side of Eq. (3) we note that the pressure is fixed by the density via the equation of

state such that

$$P(z, t) = P[\rho(z, t)] \approx P[\rho_0(z)] - v_s^2[\rho_0(z)](\rho_0\phi)', \quad (\text{A2})$$

where the velocity v_s is defined in Eq. (6). At equilibrium, Eq. (3) yields

$$v_s^2 \rho_0' = -\rho_0 U'. \quad (\text{A3})$$

We have therefore,

$$-P' - \rho U' \approx [v_s^2(\rho_0\phi)']' + (\rho_0\phi)'U'. \quad (\text{A4})$$

Writing $(\rho_0\phi)'U' = [\rho_0\phi U']' - \rho_0\phi U''$ and using (A3) we obtain,

$$-P' - \rho U' \approx [v_s^2 \rho_0 \phi']' - (\rho_0\phi)U''. \quad (\text{A5})$$

Writing

$$[v_s^2 \rho_0 \phi']' = \rho_0 [v_s^2 \phi']' + \rho_0' [v_s^2 \phi'] \quad (\text{A6})$$

and using (A3) again we obtain

$$-P' - \rho U' \approx \rho_0 [v_s^2 \phi']' - \rho_0 \phi' U' - \rho_0 \phi U''. \quad (\text{A7})$$

The third, viscosity term on the right hand side of Eq. (3) reads

$$\partial_z(\eta\partial_z v) = [\eta\dot{\phi}]'. \quad (\text{A8})$$

Substituting Eqs. (A1), (A7) and (A8) in Eq. (3) we obtain

$$\ddot{\phi} = [v_s^2 \phi']' - \phi' U' - \phi U'' + \rho_0^{-1} [\eta\dot{\phi}]'. \quad (\text{A9})$$

For the solutions of the form $\phi(z, t) = e^{i\omega t} \chi'(z)$ we obtain the equation,

$$-\omega^2 \chi' = [v_s^2 \chi'']' - \chi'' U' - \chi' U'' + (-i\omega) \rho_0^{-1} [\eta \chi''']'. \quad (\text{A10})$$

Integration of Eq. (A10) over z yields Eq. (5).

¹ V. V. Deshpande, M. Bockrath, L. I. Glazman, A. Yacoby, Nature **464**, 209 (2010).
² M. A. Cazalilla, R. Citro, T. Giamarchi, E. Orignac, and M. Rigol, Rev. Mod. Phys. **83**, 1405 (2011).
³ A. Imambekov, T. L. Schmidt, and L. I. Glazman, Rev. Mod. Phys. **84**, 1253 (2012).
⁴ F. D. M. Haldane, J. Phys. C: Solid State Phys., **14**, 2585 (1981).
⁵ M. Stone, *Bosonization*, (World Scientific Publishing Co., 1994).
⁶ A. O. Gogolin, A. A. Nersisyan, and A. M. Tsvelik, *Bosonization and strongly correlated systems*, (Cambridge University Press, 1998).
⁷ T. Giamarchi, *Quantum Physics in One Dimension*, (Clarendon Press, Oxford, 2003).
⁸ D. C. Mattis, *The Many-Body Problem: An Encyclopedia*

of Exactly Solved Models in One Dimension, (World Scientific Publishing, 1992).
⁹ B. Sutherland, *Beautiful models: 70 years of exactly solved quantum many-body problems*, (World Sci. Pub., 2004).
¹⁰ W. Kohn, Phys. Rev. **123**, 1242 (1961).
¹¹ J. F. Dobson, Phys. Rev. Lett. **73**, 2244 (1994).
¹² J. Kinast, S. L. Hemmer, M. E. Gehm, A. Turlapov, and J. E. Thomas, Phys. Rev. Lett. **92**, 150402 (2004).
¹³ M. Bartenstein, A. Altmeyer, S. Riedl, S. Jochim, C. Chin, J. Hecker Denschlag, and R. Grimm, Phys. Rev. Lett. **92**, 203201 (2004).
¹⁴ A. Altmeyer, S. Riedl, C. Kohstall, M. J. Wright, R. Geursen, M. Bartenstein, C. Chin, J. Hecker Denschlag, and R. Grimm, Phys. Rev. Lett. **98**, 040401 (2007).
¹⁵ M. J. Wright, S. Riedl, A. Altmeyer, C. Kohstall, E. R. Sanchez Guajardo, J. Hecker Denschlag, and

- R. Grimm, Phys. Rev. Lett. **99**, 150403 (2007).
- ¹⁶ M. Khodas, M. Pustilnik, A. Kamenev, and L. I. Glazman, Phys. Rev. B **76**, 155402 (2007).
- ¹⁷ G. Barak, H. Steinberg, L. N. Pfeiffer, K. W. West, L. Glazman, F. von Oppen, and A. Yacoby, Nat. Phys. **6**, 489 (2010).
- ¹⁸ T. Karzig, L. I. Glazman, and F. von Oppen, Phys. Rev. Lett. **105**, 226407 (2010).
- ¹⁹ T. Micklitz and A. Levchenko, Phys. Rev. Lett. **106**, 196402 (2011).
- ²⁰ A. Levchenko, Phys. Rev. Lett. **113**, 196401 (2014).
- ²¹ L. Brey, N. F. Johnson, and B. I. Halperin, Phys. Rev. B **40**, 10647 (1989).
- ²² A. Iqbal and M. Khodas, Phys. Rev. B **90**, 155439, (2014).
- ²³ H. Drexler, W. Hansen, J. P. Kotthaus, M. Holland, and S. P. Beaumont, Phys. Rev. B **46**, 12849(R) (1992).
- ²⁴ L. Wendler and R. Haupt, Phys. Rev. B **52**, 9031 (1995).
- ²⁵ P. A. Pantel, D. Davesne, S. Chiacchiera, and M. Urban, Phys. Rev. A **86**, 023635 (2012).
- ²⁶ S. Riedl, E. R. Sanchez Guajardo, C. Kohstall, A. Altmeyer, M. J. Wright, J. Hecker Denschlag, R. Grimm, G. M. Bruun, and H. Smith, Phys. Rev. A **78**, 053609 (2008).
- ²⁷ L. D. Landau and E. M. Lifshitz, *Fluid Mechanics* (Pergamon Press, Oxford, 1987).
- ²⁸ A. A. Abrikosov and I. M. Khalatnikov, Rep. Prog. Phys. **22**, 329 (1959).
- ²⁹ G. Morigi and S. Fishman, Phys. Rev. E **70**, 066141 (2004).
- ³⁰ A. Iqbal, A. Levchenko, and M. Khodas, unpublished.
- ³¹ R. Citro, S. De Palo, E. Orignac, P. Pedri, and M.-L. Chiofalo, New J. Phys. **10**, 045011 (2008).
- ³² C. Menotti and S. Stringari, Phys. Rev. A **66**, 043610 (2002).
- ³³ D. Petrov, D. Gangardt, and G. Shlyapnikov, J. Physique IV **116**, 5 (2004).
- ³⁴ D. S. Petrov, G. V. Shlyapnikov, and J. T. M. Walraven, Phys. Rev. Lett. **85**, 3745 (2000).
- ³⁵ A. Levchenko, T. Micklitz, J. Rech, and K. A. Matveev, Phys. Rev. B **82**, 115413 (2010).
- ³⁶ W. DeGottardi and K. A. Matveev, preprint arXiv:1412.0693.
- ³⁷ A. V. Andreev, S. A. Kivelson, and B. Spivak, Phys. Rev. Lett. **106**, 256804 (2011).
- ³⁸ R. Bamler and A. Rosch, preprint arXiv:1504.03375.
- ³⁹ T. Kinoshita, T. Wenger, and D. S. Weiss, Nature **440**, 900 (2006).