

This is the accepted manuscript made available via CHORUS. The article has been published as:

Reorientation of quantum Hall stripes within a partially filled Landau level

Q. Shi, M. A. Zudov, J. D. Watson, G. C. Gardner, and M. J. Manfra

Phys. Rev. B **93**, 121404 — Published 14 March 2016

DOI: [10.1103/PhysRevB.93.121404](https://doi.org/10.1103/PhysRevB.93.121404)

Reorientation of quantum Hall stripes within a partially filled Landau level

Q. Shi,¹ M. A. Zudov,^{1,*} J. D. Watson,^{2,3} G. C. Gardner,^{3,4} and M. J. Manfra^{2,3,4,5}

¹*School of Physics and Astronomy, University of Minnesota, Minneapolis, Minnesota 55455, USA*

²*Department of Physics and Astronomy, Purdue University, West Lafayette, Indiana 47907, USA*

³*Birck Nanotechnology Center, Purdue University, West Lafayette, Indiana 47907, USA*

⁴*School of Materials Engineering, Purdue University, West Lafayette, Indiana 47907, USA*

⁵*School of Electrical and Computer Engineering,
Purdue University, West Lafayette, Indiana 47907, USA*

We investigate the effect of the filling factor on transport anisotropies, known as stripes, in high Landau levels of a two-dimensional electron gas. We find that at certain in-plane magnetic fields, the stripes orientation is sensitive to the filling factor within a given Landau level. This sensitivity gives rise to the emergence of stripes *away from half-filling* while an orthogonally-oriented, native stripes reside at half-filling. This switching of the anisotropy axes within a single Landau level can be attributed to a strong dependence of the native symmetry breaking potential on the filling factor.

PACS numbers: 73.43.Qt, 73.63.Hs, 73.40.-c

Electronic analogs of liquid crystals, commonly termed electron nematics or stripes, are believed to exist in a wide variety of condensed matter systems¹, such as ruthenates², high temperature superconductors^{3,4}, and heavy fermion systems⁵. However, the first evidence for such nematic phases^{6–11} came from a clean 2D electron gas (2DEG) formed at the interface between GaAs and AlGaAs, which is often the system of choice to study of quantized Hall effects^{12,13}.

One important parameter which is unique to a 2DEG, is the filling factor, $\nu = n_e h / e B$, where n_e is the electron density and B is the magnetic field. Stripes in GaAs manifest themselves in the resistance minima (maxima) in the easy (hard) transport direction when ν is close to $\nu_N^\pm \equiv 2N + 1 \pm 1/2$ ($N \geq 2$), where $+$ ($-$) describes spin-down (spin-up) branches of the N -th Landau level. While stripes exist in a finite filling factor range^{6,7,14–16}, $\nu_N^\pm - 0.1 \lesssim \nu \lesssim \nu_N^\pm + 0.1$, transport studies have focused almost exclusively¹⁷ at half-integer ν where the anisotropy is the strongest.

In a purely perpendicular magnetic field, stripes orient along $\langle 110 \rangle$ crystal direction with very few exceptions^{18,19}. Despite nearly two decades of investigations, the origin of the native symmetry breaking potential responsible for this particular stripes orientation continues to remain elusive^{19–21}. It is well documented that an in-plane magnetic field B_{\parallel} provides an external symmetry breaking potential which competes with and can overcome the native symmetry breaking potential. In particular, when applied along stripes (easy direction), B_{\parallel} can switch the anisotropy axes^{19,22–27} aligning stripes perpendicular to it²⁸. There thus exists a characteristic in-plane magnetic field B_c which renders 2DEG macroscopically isotropic. This field is routinely used to quantify the strength of the native symmetry breaking potential^{18,19,24,29}.

In this Rapid Communication we report on transport studies of stripes in a high-mobility 2DEG focusing on filling factors *away from half-filling*. At certain B_{\parallel} , we observe distinct anisotropic phases which reside both at

and *away from half-filling* within a single Landau level. These stripe phases have orthogonal orientation, indicating that it is largely determined by the filling factor. The observed reorientations of stripes within a single Landau level implies strong sensitivity of the native symmetry breaking potential to the filling factor. Indeed, the characteristic in-plane field B_c drops roughly symmetrically when ν deviates from half-filling in either direction. This dependence on the filling factor is quite significant and should be accounted for by theories attempting to identify the origin of the native symmetry breaking potential.

While similar results have been obtained from samples fabricated from several different wafers, here we present the data from one $\sim 4 \times 4$ mm square sample cleaved from a symmetrically doped, 30 nm-wide GaAs/AlGaAs quantum well. Electron density and mobility were $n_e \approx 2.9 \times 10^{11} \text{ cm}^{-2}$ and $\mu \approx 1.6 \times 10^7 \text{ cm}^2/\text{Vs}$, respectively^{30,31}. Eight indium ohmic contacts were fabricated at the corners and midsides of the sample. The longitudinal resistances, R_{xx} and R_{yy} , were measured using four-terminal, low-frequency lock-in technique; the current (typically 50 nA) was sent through the midside contacts and the voltage drop was measured between the corner contacts. An in-plane magnetic field was introduced by tilting the sample by angle θ about \hat{x} axis, i.e. $B_{\parallel} = B_y = B \sin \theta$. Unless otherwise noted, all the data we acquired at $T \approx 20 \text{ mK}$.

In Fig. 1 we present R_{xx} (solid line) and R_{yy} (dotted line) versus filling factor ν at different θ , from 0° to 12.2° , as marked. At $\theta = 0^\circ$, the data reveal strong anisotropy near $\nu = 9/2$ with $R_{xx} \gg R_{yy}$ [see Fig. 1(a)], i.e., stripes are oriented along \hat{y} direction. Near $\nu \approx 4.28$ and $\nu \approx 4.72$, R_{xx} and R_{yy} reveal isotropic insulating states, reflecting the formation of “bubble” phases^{6,7,32–35}.

At $\theta = 5.7^\circ$ [Fig. 1(b)], new anisotropic states of *orthogonal* orientation (e.g., along \hat{x} -direction) develop near $\nu \approx 4.4$ and $\nu \approx 4.6$, i.e., at the edges of the native stripe range found at $\theta = 0^\circ$. Like the native stripes, which reside near $\nu = 9/2$ ³⁶, these new anisotropic states are represented by high peaks along




FIG. 1. (Color online) R_{xx} (solid line) and R_{yy} (dotted line) versus ν at $T = 25$ mK and θ from (a) 0° to (e) 12.2° , as marked.

one direction and deep minima along another, albeit with $R_{yy} \gg R_{xx}$. As θ increases to 8.1° [Fig. 1(c)], these orthogonal anisotropic states take over larger range of ν , while the native anisotropy at $\nu = 9/2$ becomes noticeably weaker. At $\theta = 8.7^\circ$ [Fig. 1(d)], stripes along \hat{x} -direction occupy almost the whole filling factor range of the native stripes, except near $\nu = 9/2$, where the system becomes isotropic. Finally, at $\theta = 12.2^\circ$ we observe that the B_{\parallel} -induced reorientation of the native stripes is complete at all ν within the native stripe range. The isotropic insulating states representing “bubble” phases remain essentially unchanged.

Before proceeding with the more detailed data analysis, we demonstrate that the evolution shown in Fig. 1 is not unique to $\nu = 9/2$. In Fig. 2 we show R_{xx} (solid line) and R_{yy} (dotted line) at (a) $5 \leq \nu \leq 6$ and (b) $6 \leq \nu \leq 7$ measured at $\theta = 8.1^\circ$ and 8.7° , respectively. In both cases, the data show alternating stripes orientation which is identical to that shown in Fig. 1(c). We thus conclude that the sensitivity of the stripes orientation to the filling factor is a generic feature of stripes in our sample; it appears in both spin-up and spin-down branches and in different ($N = 2, 3$) Landau levels.

To further examine the evolution of the anisotropic states with the tilt angle, we compute the resistance anisotropy $A_R = (R_{xx} - R_{yy}) / (R_{xx} + R_{yy})$ from the data shown in Fig. 1 and present the results in Fig. 3. At $\theta = 0$ [Fig. 3(a)] we see that $A_R \approx 1$ at all filling factors within a band $4.4 \leq \nu \leq 4.6$, reflecting the native stripes orientation along \hat{y} direction. As the sample is tilted, this $A_R \approx 1$ band becomes narrower [Fig. 3(b) and Fig. 3(c)], and eventually vanishes as the system becomes isotropic at $\nu = 9/2$ where $A_R \approx 0$ [Fig. 3(d)]. Concomitant with the narrowing of the $A_R \approx 1$ band (stripes $\parallel \hat{y}$), we observe the emergence of the $A_R \approx -1$ bands (stripes $\parallel \hat{x}$), which germinate at the edges of the native stripe band [Fig. 3(b)]. With increasing θ , these $A_R \approx -1$ bands expand towards each other [Fig. 3(c)] and eventually merge at half-filling [Fig. 3(d)]. At the $\theta = 12.2^\circ$ [Fig. 3(e)], the $A_R \approx -1$ band occupies the whole range of the na-

tive stripes, $4.4 \leq \nu \leq 4.6$, reflecting stripes orientation along \hat{x} direction.

Taken together, the data of Fig. 1, Fig. 2, and Fig. 3 demonstrate that the reorientation of stripes by B_{\parallel} depends sensitively on the filling factor³⁷ and that at a certain fixed θ stripes orientation can vary within a single Landau level. More specifically, we establish that the reorientation first occurs at filling factors near the onset of the native stripes, i.e. at $\nu \approx 4.4$ and $\nu \approx 4.6$, and then the boundary, separating two orthogonal stripe phases, propagates towards half-filling.

We next examine the evolution of the anisotropy with B_{\parallel} at different fixed filling factors in the vicinity of $\nu = 9/2$. In Fig. 4(a), (b), and (c), we present R_{xx} (filled circles) and R_{yy} (open circles) as a function of B_{\parallel} , at $\nu = 4.5$, 4.46 , and 4.43 , respectively. In all three cases, we find that the anisotropy axes are switched when B_{\parallel} reaches a characteristic field B_c , defined such that $R_{xx} \approx R_{yy}$ (cf. \downarrow). We further notice that B_c decreases as ν deviates from half-filling. We summarize these findings in Fig. 4(d) showing B_c as a function of ν . The dashed and dash-dotted lines in Fig. 4(d) represent B_{\parallel} at $\theta = 8.1^\circ$




FIG. 2. (Color online) R_{xx} (solid line) and R_{yy} (dotted line) versus ν around (a) $\nu = 11/2$, $\theta = 8.1^\circ$ and (b) $\nu = 13/2$, $\theta = 8.7^\circ$.




FIG. 3. (Color online) Resistance anisotropy $A_R = (R_{xx} - R_{yy})/(R_{xx} + R_{yy})$ versus ν computed from the data in Fig. 1.

and $\theta = 5.7^\circ$, corresponding to the situation of Fig. 1(c) and Fig. 1(b), respectively. Two crossings of these lines with B_c is reflected in the transport data as switching of the anisotropy axes during the magnetic field sweep.

The characteristic field B_c reaches its maximum value of about 0.43 T at $\nu = 9/2$. As the filling factor deviates from half-filling, B_c gradually decreases to ≈ 0.23 T at both $\nu = 4.41$ or $\nu = 4.59$ ³⁸. The dependence of B_c on ν is roughly symmetric about half-filling and can be reasonably well described by a parabola (dotted line). This parabola crosses zero at filling factors which are very close (within 0.01) to the onset of the native anisotropy [cf. Fig. 3(a)].

Since B_c characterizes the competition between the native symmetry breaking potential and the effect of $B_{||}$ on the orientation of stripes, the decrease of B_c away from half-filling can be due to either a stronger effect of $B_{||}$ or a

weaker native symmetry breaking potential at these filling factors. However, since the effect of $B_{||}$ is predicted to have a very weak dependence on ν ^{25,26}, the relationship between B_c and ν must reflect a weaker native symmetry breaking potential away from half-filling. We note that the observed dependence is quite significant; B_c decreases by nearly 50% as ν is changed from 9/2 to 4.4 (or 4.6).

Finally, we examine the temperature dependence of the anisotropic states away from half-filling. In Fig. 5(a) and Fig. 5(b) we present R_{xx} (solid line) and R_{yy} (dotted line) measured at $\theta = 8.5^\circ$ and $T = 56$ mK and 109 mK, respectively. As seen in Fig. 5(a), R_{yy} (R_{xx}) exhibits strong maxima (deep minima) away from half-filling and the corresponding anisotropy $A_R \approx -1$ (stripes $\parallel \hat{x}$). At $\nu = 9/2$, however, R_{yy} exhibits a minimum, R_{xx} a maximum, and the anisotropy becomes positive. As the temperature is raised, the R_{yy} peaks away from half-filling quickly decay, while the resistances at the corresponding R_{xx} minima become larger [see Fig. 5(b)]. As a result, the anisotropy away from half-filling is significantly reduced. Interestingly, the R_{yy} peaks also move towards each other with increasing T and eventually merge into a single peak at half-filling before disappearing at higher temperatures. This is illustrated in the inset of Fig. 5(b) which shows the




FIG. 4. (Color online) R_{xx} (filled circles) and R_{yy} (open circles) versus B_y (a) $\nu = 4.5$, (b) 4.46 and (c) 4.43. (d) B_c (filled circles) versus ν . Dashed (dash-dotted) line represents B_y at $\theta = 8.1^\circ$ ($\theta = 5.7^\circ$).




FIG. 5. (Color online) R_{xx} (solid line) and R_{yy} (dotted line) versus ν at $\theta = 8.5^\circ$ and (a) $T = 56$ mK and (b) 109 mK. The inset shows the change of ν_m at two R_{yy} peaks versus T .

filling factors of the R_{yy} peaks as a function of temperature. The merging of the R_{yy} peaks indicates that the anisotropy away from half-filling disappears at a lower temperature compared to that at half-filling.

An interesting question is why the filling factor-dependence of B_c has not been seen in previous experiments. We argue that it is the significant native anisotropy away from half-filling, see Fig. 3(a), that made possible a clear observation of this behavior. Indeed, in the absence of an in-plane magnetic field our data show an unusually wide R_{xx} peak and a vanishingly small R_{yy} over a wide filling factor range, $4.4 \leq \nu \leq 4.6$, see Fig. 1(a). It is indeed this wide range which allows significant anisotropy (of the opposite sign) to develop away from half-filling at intermediate tilt angles [see Fig. 1(b-d) and Fig. 3(b-d)]. In contrast, in early studies^{22,23} the native anisotropy rapidly decays away from half-filling, as manifested by narrower resistance maxima and minima, and only weak signatures suggestive of filling factor-driven reorientation were observed [see, e.g. Fig. 2 in Ref. 22].

There is both experimental and theoretical evidence that weakening of the anisotropy away from half-filling is a result of stronger effect of disorder. Indeed, in agreement with the theory³⁹, pinning mode resonances in the ac conductivity¹⁴ revealed higher resonance frequencies away from half-filling. Although the sample in the present study has the mobility comparable to (or even lower than) samples used in earlier experiments, it is well established that the mobility alone is not a good metric of the quality of the transport data^{30,40–43}. We also notice that in contrast to early studies^{22,23}, which utilized conventional single heterointerfaces, our experiments exploit a double heterointerface of a “doping well” design³⁰, which is known to produce better quality transport data revealed in, e.g., larger gaps of the fragile $\nu = 5/2$ fractional quantum Hall state^{42–45}.

In addition to weak anisotropy away from half-filling there exist another argument why the phenomenon reported here can be suppressed in more disordered samples. Experiments on pinning mode resonances under $B_{||}$ ²⁷ suggest that the disorder favors stripes *parallel* to

$B_{||}$. This finding implies that due to stronger effect of disorder away from half-filling^{14,39}, stripes at such filling factors would be more difficult to reorient perpendicular to $B_{||}$. Thus, the effect of disorder would lead to *larger* B_c away from half-filling which may mask the ν -dependence of the native symmetry breaking field observed in the present study.

In summary, we have studied anisotropic phases in high Landau levels in tilted magnetic fields. We have demonstrated the realization of anisotropic phases, distinct by their orthogonal orientation, which reside at and away from half-filling in a single Landau level⁴⁶. The boundaries separating these states propagate towards each other and eventually merge at half-filling with increasing $B_{||}$. The observed switching of the anisotropy axes within a single Landau level can be explained by a strong dependence of the native symmetry breaking potential on the filling factor. This conclusion is corroborated by a monotonic decrease of the characteristic in-plane magnetic field B_c as ν deviates from half-filling. The experimentally observed dependence on the filling factor is quite significant and should be taken into account by theories attempting to identify the nature of the native symmetry breaking potential.

ACKNOWLEDGMENTS

We thank G. Jones, S. Hannas, T. Murphy, J. Park, and A. Suslov for technical support. We thank G. Csathy, L. Engel, H. Fertig, I. Kukushkin, M. Shayegan, and B. Shklovskii for discussions. The work at Minnesota (Purdue) was supported by the U.S. Department of Energy, Office of Science, Basic Energy Sciences, under Award # ER 46640-SC0002567 (DE-SC0006671). Q.S. acknowledges University of Minnesota Doctoral Dissertation Fellowship. Experiments were performed at the National High Magnetic Field Laboratory, which is supported by NSF Cooperative Agreement No. DMR-0654118, by the State of Florida, and by the DOE.

* Corresponding author: zudov@physics.umn.edu

¹ E. Fradkin, S. A. Kivelson, M. J. Lawler, J. P. Eisenstein, and A. P. Mackenzie, *Nematic fermi fluids in condensed matter physics*, Annu. Rev. Condens. Matter Phys. **1**, 153 (2010).

² R. A. Borzi, S. A. Grigera, J. Farrell, R. S. Perry, S. J. S. Lister, S. L. Lee, D. A. Tennant, Y. Maeno, and A. P. Mackenzie, *Formation of a nematic fluid at high fields in $Sr_3Ru_2O_7$* , Science **315**, 214 (2007).

³ R. Daou, J. Chang, D. LeBoeuf, O. Cyr-Choiniere, F. Laliberte, N. Doiron-Leyraud, B. J. Ramshaw, R. Liang, D. A. Bonn, W. N. Hardy, et al., *Broken rotational symmetry in the pseudogap phase of a high- T_c superconductor*, Nature **463**, 519 (2010).

⁴ J.-H. Chu, J. G. Analytis, K. De Greve, P. L. McMahon, Z. Islam, Y. Yamamoto, and I. R. Fisher, *In-plane resistivity anisotropy in an underdoped iron arsenide superconductor*, Science **329**, 824 (2010).

⁵ R. Okazaki, T. Shibauchi, H. J. Shi, Y. Haga, T. D. Matsuda, E. Yamamoto, Y. Onuki, H. Ikeda, and Y. Matsuda, *Rotational symmetry breaking in the hidden-order phase of URu_2Si_2* , Science **331**, 439 (2011).

⁶ M. P. Lilly, K. B. Cooper, J. P. Eisenstein, L. N. Pfeiffer, and K. W. West, *Evidence for an anisotropic state of two-dimensional electrons in high Landau levels*, Phys. Rev. Lett. **82**, 394 (1999).

⁷ R. R. Du, D. C. Tsui, H. L. Stormer, L. N. Pfeiffer, K. W. Baldwin, and K. W. West, *Strongly anisotropic transport*

- in higher two-dimensional Landau levels, *Solid State Commun.* **109**, 389 (1999).
- ⁸ A. A. Koulakov, M. M. Fogler, and B. I. Shklovskii, *Charge density wave in two-dimensional electron liquid in weak magnetic field*, *Phys. Rev. Lett.* **76**, 499 (1996).
 - ⁹ M. M. Fogler, A. A. Koulakov, and B. I. Shklovskii, *Ground state of a two-dimensional electron liquid in a weak magnetic field*, *Phys. Rev. B* **54**, 1853 (1996).
 - ¹⁰ E. Fradkin and S. A. Kivelson, *Liquid-crystal phases of quantum Hall systems*, *Phys. Rev. B* **59**, 8065 (1999).
 - ¹¹ E. Fradkin, S. A. Kivelson, E. Manousakis, and K. Nho, *Nematic phase of the two-dimensional electron gas in a magnetic field*, *Phys. Rev. Lett.* **84**, 1982 (2000).
 - ¹² K. von Klitzing, G. Dorda, and M. Pepper, *New method for high-accuracy determination of the fine-structure constant based on quantized Hall resistance*, *Phys. Rev. Lett.* **45**, 494 (1980).
 - ¹³ D. C. Tsui, H. L. Stormer, and A. C. Gossard, *Two-dimensional magnetotransport in the extreme quantum limit*, *Phys. Rev. Lett.* **48**, 1559 (1982).
 - ¹⁴ G. Sambandamurthy, R. M. Lewis, H. Zhu, Y. P. Chen, L. W. Engel, D. C. Tsui, L. N. Pfeiffer, and K. W. West, *Observation of pinning mode of stripe phases of 2D systems in high Landau levels*, *Phys. Rev. Lett.* **100**, 256801 (2008).
 - ¹⁵ I. V. Kukushkin, V. Umansky, K. von Klitzing, and J. H. Smet, *Collective modes and the periodicity of quantum Hall stripes*, *Phys. Rev. Lett.* **106**, 206804 (2011).
 - ¹⁶ Experiments on pinning mode resonances in the ac conductivity¹⁴ revealed slightly larger resonance frequencies and smaller quality factors away from half-filling, suggesting a stronger effect of disorder and/or a reduced stripes stiffness. Another experiment¹⁵, probing stripes with surface acoustic waves, showed that the frequency of the collective modes decreases as ν is tuned away from half-filling, in general agreement with theoretical proposal⁴⁷.
 - ¹⁷ One exception is Ref. 33 investigating the effects of dc current.
 - ¹⁸ J. Zhu, W. Pan, H. L. Stormer, L. N. Pfeiffer, and K. W. West, *Density-induced interchange of anisotropy axes at half-filled high Landau levels*, *Phys. Rev. Lett.* **88**, 116803 (2002).
 - ¹⁹ J. Pollanen, K. B. Cooper, S. Brandsen, J. P. Eisenstein, L. N. Pfeiffer, and K. W. West, *Heterostructure symmetry and the orientation of the quantum Hall nematic phases*, *Phys. Rev. B* **92**, 115410 (2015).
 - ²⁰ I. Sodemann and A. H. MacDonald, *Theory of native orientational pinning in quantum Hall nematics*, arXiv:1307.5489 (2013).
 - ²¹ S. P. Koduvayur, Y. Lyanda-Geller, S. Khlebnikov, G. Csathy, M. J. Manfra, L. N. Pfeiffer, K. W. West, and L. P. Rokhinson, *Effect of strain on stripe phases in the quantum Hall regime*, *Phys. Rev. Lett.* **106**, 016804 (2011).
 - ²² M. P. Lilly, K. B. Cooper, J. P. Eisenstein, L. N. Pfeiffer, and K. W. West, *Anisotropic states of two-dimensional electron systems in high Landau levels: Effect of an in-plane magnetic field*, *Phys. Rev. Lett.* **83**, 824 (1999).
 - ²³ W. Pan, R. R. Du, H. L. Stormer, D. C. Tsui, L. N. Pfeiffer, K. W. Baldwin, and K. W. West, *Strongly anisotropic electronic transport at Landau level filling factor under a tilted magnetic field*, *Phys. Rev. Lett.* **83**, 820 (1999).
 - ²⁴ K. B. Cooper, M. P. Lilly, J. P. Eisenstein, T. Jungwirth, L. N. Pfeiffer, and K. W. West, *An investigation of orientational symmetry-breaking mechanisms in high Landau levels*, *Solid State Commun.* **119**, 89 (2001).
 - ²⁵ T. Jungwirth, A. H. MacDonald, L. Smrčka, and S. M. Girvin, *Field-tilt anisotropy energy in quantum Hall stripe states*, *Phys. Rev. B* **60**, 15574 (1999).
 - ²⁶ T. D. Stanescu, I. Martin, and P. Phillips, *Finite-temperature density instability at high Landau level occupancy*, *Phys. Rev. Lett.* **84**, 1288 (2000).
 - ²⁷ H. Zhu, G. Sambandamurthy, L. W. Engel, D. C. Tsui, L. N. Pfeiffer, and K. W. West, *Pinning mode resonances of 2D electron stripe phases: Effect of an in-plane magnetic field*, *Phys. Rev. Lett.* **102**, 136804 (2009).
 - ²⁸ There are also examples when B_{\parallel} can induce strong anisotropies in natively isotropic states^{22,23,48–54}, with the hard transport direction perpendicular to it.
 - ²⁹ K. B. Cooper, J. P. Eisenstein, L. N. Pfeiffer, and K. W. West, *Metastable resistance-anisotropy orientation of two-dimensional electrons in high Landau levels*, *Phys. Rev. Lett.* **92**, 026806 (2004).
 - ³⁰ M. J. Manfra, *Molecular beam epitaxy of ultra-high-quality AlGaAs/GaAs heterostructures: Enabling physics in low-dimensional electronic systems*, *Annu. Rev. Condens. Matter Phys.* **5**, 347 (2014).
 - ³¹ The doping wells are 90 nm away from the center of GaAs quantum well which is located 215 nm from the surface. At $T \approx 0.3$ K, the quantum lifetime obtained from the Dingle analysis of microwave-induced resistance oscillations⁵⁵ is about 20 ps.
 - ³² J. P. Eisenstein, K. B. Cooper, L. N. Pfeiffer, and K. W. West, *Insulating and fractional quantum Hall states in the first excited Landau level*, *Phys. Rev. Lett.* **88**, 076801 (2002).
 - ³³ J. Gores, G. Gamez, J. H. Smet, L. Pfeiffer, K. West, A. Yacoby, V. Umansky, and K. von Klitzing, *Current-induced anisotropy and reordering of the electron liquid-crystal phases in a two-dimensional electron system*, *Phys. Rev. Lett.* **99**, 246402 (2007).
 - ³⁴ N. Deng, A. Kumar, M. J. Manfra, L. N. Pfeiffer, K. W. West, and G. A. Csathy, *Collective nature of the reentrant integer quantum Hall states in the second Landau level*, *Phys. Rev. Lett.* **108**, 086803 (2012).
 - ³⁵ N. Deng, J. D. Watson, L. P. Rokhinson, M. J. Manfra, and G. A. Csathy, *Contrasting energy scales of reentrant integer quantum Hall states*, *Phys. Rev. B* **86**, 201301 (2012).
 - ³⁶ The R_{xx} at $\nu = 9/2$ is somewhat higher than at $\theta = 0$. We notice that similar observation was made in Ref. 27.
 - ³⁷ At a given θ , stripes orientation is uniquely determined by ν and is insensitive to the magnetic field sweep direction, in contrast to Ref. 18 and 29.
 - ³⁸ We do not include B_c at smaller or larger ν because of large experimental uncertainty in this regime.
 - ³⁹ H. Yi, H. A. Fertig, and R. Côté, *Stability of the smectic quantum Hall state: A quantitative study*, *Phys. Rev. Lett.* **85**, 4156 (2000).
 - ⁴⁰ V. Umansky, M. Heiblum, Y. Levinson, J. Smet, J. Nübler, and M. Dolev, *MBE growth of ultra-low disorder 2DEG with mobility exceeding 35×10^6 cm²/Vs*, *J. Cryst. Growth* **311**, 1658 (2009).
 - ⁴¹ G. Gamez and K. Muraki, $\nu = 5/2$ fractional quantum Hall state in low-mobility electron systems: Different roles of disorder, *Phys. Rev. B* **88**, 075308 (2013).
 - ⁴² M. Samani, A. V. Rossokhaty, E. Sajadi, S. Lüscher, J. A. Folk, J. D. Watson, G. C. Gardner, and M. J. Manfra, *Low-temperature illumination and annealing of ultrahigh quality quantum wells*, *Phys. Rev. B* **90**, 121405 (2014).

- ⁴³ J. D. Watson, G. A. Csáthy, and M. J. Manfra, *Impact of heterostructure design on transport properties in the second Landau level of In Situ back-gated two-dimensional electron gases*, Phys. Rev. Applied **3**, 064004 (2015).
- ⁴⁴ N. Deng, G. C. Gardner, S. Mondal, E. Kleinbaum, M. J. Manfra, and G. A. Csáthy, $\nu = 5/2$ fractional quantum Hall state in the presence of alloy disorder, Phys. Rev. Lett. **112**, 116804 (2014).
- ⁴⁵ According to Ref. 42, doping well design provides better equilibration of charges among the dopants which improves screening. In addition, the placement of dopants in separate doping wells increases the chance that the dopant energy levels will be shallow, which dramatically increases their ionization fraction and thus reduces the disorder potential due to random ionization states.
- ⁴⁶ While a slight hysteresis is observed upon changing the field sweep direction, it does not affect the orientation of stripes, in contrast to Ref. 18 and 29.
- ⁴⁷ R. Côté and H. A. Fertig, *Collective modes of quantum Hall stripes*, Phys. Rev. B **62**, 1993 (2000).
- ⁴⁸ J. Xia, V. Cvicek, J. P. Eisenstein, L. N. Pfeiffer, and K. W. West, *Tilt-induced anisotropic to isotropic phase transition at $\nu = 5/2$* , Phys. Rev. Lett. **105**, 176807 (2010).
- ⁴⁹ C. Zhang, T. Knuuttila, Y. Dai, R. R. Du, L. N. Pfeiffer, and K. W. West, *$5/2$ fractional quantum Hall effect at 10 T: Implications for the Pfaffian state*, Phys. Rev. Lett. **104**, 166801 (2010).
- ⁵⁰ J. Xia, J. P. Eisenstein, L. N. Pfeiffer, and K. W. West, *Evidence for a fractionally quantized Hall state with anisotropic longitudinal transport*, Nat. Phys. **7**, 845 (2011).
- ⁵¹ M. Mulligan, C. Nayak, and S. Kachru, *Effective field theory of fractional quantized Hall nematics*, Phys. Rev. B **84**, 195124 (2011).
- ⁵² Y. Liu, S. Hasdemir, M. Shayegan, L. N. Pfeiffer, K. W. West, and K. W. Baldwin, *Evidence for a $5/2$ fractional quantum Hall nematic state in parallel magnetic fields*, Phys. Rev. B **88**, 035307 (2013).
- ⁵³ Q. Shi, M. A. Zudov, C. Morrison, and M. Myronov, *Strong transport anisotropy in Ge/SiGe quantum wells in tilted magnetic fields*, Phys. Rev. B **91**, 201301(R) (2015).
- ⁵⁴ Q. Shi, M. A. Zudov, C. Morrison, and M. Myronov, *Transport anisotropy in Ge quantum wells in the absence of quantum oscillations*, Phys. Rev. B **92**, 161405 (2015).
- ⁵⁵ M. A. Zudov, R. R. Du, J. A. Simmons, and J. L. Reno, *Shubnikov-de Haas-like oscillations in millimeter-wave photoconductivity in a high-mobility two-dimensional electron gas*, Phys. Rev. B **64**, 201311(R) (2001).