

This is the accepted manuscript made available via CHORUS. The article has been published as:

Exact conditions on the temperature dependence of density functionals

K. Burke, J. C. Smith, P. E. Grabowski, and A. Pribram-Jones

Phys. Rev. B **93**, 195132 — Published 17 May 2016

DOI: [10.1103/PhysRevB.93.195132](https://doi.org/10.1103/PhysRevB.93.195132)

Exact conditions on the temperature dependence of density functionals

K. Burke,^{1,2} J. C. Smith,¹ P. E. Grabowski,¹ and A. Pribram-Jones^{3,4}

¹*Department of Physics and Astronomy, University of California, Irvine, CA 92697*

²*Department of Chemistry, University of California, Irvine, CA 92697*

³*Lawrence Livermore National Laboratory, 7000 East Avenue, L-413, Livermore, California 94550*

⁴*Department of Chemistry, University of California, Berkeley, CA 94720*

Universal exact conditions guided the construction of most ground-state density functional approximations in use today. We derive the relation between the entropy and Mermin free energy density functionals for thermal density functional theory. Both the entropy and sum of kinetic and electron-electron repulsion functionals are shown to be monotonically increasing with temperature, while the Mermin functional is concave downwards. Analogous relations are found for both exchange and correlation. The importance of these conditions is illustrated in two extremes: the Hubbard dimer and the uniform gas.

PACS numbers: 31.10.+z, 31.15.E-

I. INTRODUCTION

Warm dense matter (WDM) is a rapidly growing multidisciplinary field that spans many branches of physics, including for example astrophysics, geophysics, attosecond physics, and nuclear physics^{1–9}. In the last decade, quantum molecular dynamics, using DFT with electrons at finite temperatures, has been extremely successful at predicting material properties under extreme conditions, and has become a standard simulation tool in this field¹⁰. Almost all such simulations use ground-state exchange-correlation (XC) approximations, even when the electrons are significantly heated. Thermal density functional theory (thDFT) was formalized by Mermin¹¹, when he showed that the reasoning of Hohenberg and Kohn¹² could be extended to the grand canonical potential of electrons coupled to a thermal bath at temperature τ . In recent times, the Mermin-Kohn-Sham (MKS) equations of non-interacting electrons at finite temperature, whose density matches that of the physical system, are being solved to simulate warm dense matter^{13,14}. In most of these calculations, the ground-state approximation (GSA) is made, in which the exchange-correlation (XC) free energy, which typically depends on τ , is approximated by its ground-state value. Accurate results for the uniform gas are still being found^{15–18}, which provide input to a thermal local density approximation, but LDA is insufficiently accurate for many modern applications, and thermal GGA's are being explored¹⁹.

Many useful exact conditions in ground-state DFT (relation between coupling constant and scaling, correlation scaling inequalities, exchange and kinetic scaling equalities, signs of energy components) were first derived²⁰ by studying the variational principle in the form of the Levy constrained search²¹. Most of these conditions are satisfied (by construction) by the local density approximation¹³ and have been used for decades to constrain and/or improve more advanced approximations²². Their finite temperature analogs were derived in Ref.²³ (see also Ref.²⁴), and extended in Ref.²⁵. Because the

kinetic and entropic contributions always appear in the same combination as the so-called kintropic energy (see Eq. (21) and related text), such relations can never be used to extract either component individually.

Many basic thermodynamic relations are proven via quantum statistical mechanics²⁶. However, converting these to conditions on density functionals is neither obvious nor trivial. In the present work, we extend these methods to the dependence of the Mermin functional (i.e., the universal part of the free-energy functional) on the *temperature*, rather than on the coupling constant or the scale of the density. We find several new equalities and inequalities which apply to thDFT of all electronic systems. This allows us to separate entropic and kinetic contributions. We show that the entropy density functional is monotonically increasing with temperature, as is the sum of the kinetic and electron-electron repulsion density functionals, and that the temperature derivative of the Mermin functional is the negative of the entropy functional. Thus the Mermin functional is concave downwards as a function of temperature. Applying these conditions to the MKS system yields conditions on the exchange-correlation free energy functionals. Lastly, we illustrate all our findings in the two extreme cases of the uniform gas and the Hubbard dimer. We find a recent parametrization of the XC free energy of the uniform gas violates our conditions, although only for densities that are so low as to be unlikely to significantly affect any property calculated within thLDA.

II. THEORY

For a given average particle number, define the free energy of a statistical density-matrix Γ as

$$A^\tau[\Gamma] = H[\Gamma] - \tau S[\Gamma], \quad (1)$$

where \hat{H} is the Hamiltonian operator, S extracts the entropy, and we use τ to denote temperature. Define

$$F_1[\Gamma] = T[\Gamma] + V_{ee}[\Gamma], \quad (2)$$

where \hat{T} is the kinetic energy operator and \hat{V}_{ee} the electron-electron repulsion operator. Then

$$F^\tau[\Gamma] = F_I[\Gamma] - \tau S[\Gamma]. \quad (3)$$

The Mermin functional, written in terms of a constrained search, is²³

$$F^\tau[n] = \min_{\Gamma \rightarrow n} F^\tau[\Gamma], \quad (4)$$

where the argument distinguishes functionals of the density from those of the density-matrix. The free energy of a given system can be found from

$$A^\tau = \min_n \left\{ F^\tau[n] + \int d^3r v(\mathbf{r}) n(\mathbf{r}) \right\}. \quad (5)$$

We denote by $\Gamma^\tau[n]$ the statistical density matrix that minimizes \hat{F}^τ and yields density $n(\mathbf{r})$. Then:

$$\frac{dF^\tau[n]}{d\tau} = \frac{\partial F^\tau[\Gamma]}{\partial \tau} + \int d\Gamma \frac{\partial F^\tau[\Gamma]}{\partial \Gamma} \frac{d\Gamma^\tau[n]}{d\tau}, \quad (6)$$

where all are evaluated at $\Gamma^\tau[n]$. Because $\Gamma^\tau[n]$ is the minimizer, its derivative with respect to temperature (or any variable) vanishes. Thus

$$\frac{dF^\tau[n]}{d\tau} = -S^\tau[n]. \quad (7)$$

This is the DFT analog of the standard thermodynamic relation²⁶, and implies

$$F^\tau[n] = F^0[n] - \int_0^\tau d\tau' S^{\tau'}[n], \quad (8)$$

where $F^0[n]$ is the ground-state functional¹². We note that Eq. (7) was derived in⁹, but only within lattice DFT.

Given a Mermin functional (approximate or exact, interacting or not), Eq. (7) defines what the corresponding entropy functional must be. Since coordinate scaling²³ can separate the entropic and potential contributions in F , Eq. (7) allows the entropic and kinetic energy functionals to be separated. Alternatively, given an entropy functional, Eq. (7) defines the temperature-dependence of the corresponding Mermin functional. Since the entropy is always positive,

$$dF^\tau[n]/d\tau \leq 0, \quad (9)$$

i.e., the Mermin functional is monotonically decreasing.

Now consider what happens when, for a given density and temperature τ , we evaluate the Mermin functional on the density matrix for that density but at a different temperature. By the variational principle, Eq. (5),

$$F^\tau[\Gamma^{\tau'}[n]] \geq F^\tau[n], \quad (10)$$

for any value of τ' . Thus

$$F_I[\Gamma^{\tau'}[n]] - \tau S[\Gamma^{\tau'}[n]] \geq F_I^\tau[n] - \tau S^\tau[n], \quad (11)$$

or

$$F_I^{\tau'}[n] - \tau S^{\tau'}[n] \geq F_I^\tau[n] - \tau S^\tau[n]. \quad (12)$$

Since this result is true for any pair of temperatures, we reverse τ and τ' to find:

$$F_I^\tau[n] - \tau' S^\tau[n] \geq F_I^{\tau'}[n] - \tau' S^{\tau'}[n]. \quad (13)$$

Addition of Eqs. (12) and (13) yields

$$(\tau - \tau')(S^\tau[n] - S^{\tau'}[n]) \geq 0, \quad (14)$$

so that the entropy monotonically increases with τ :

$$dS^\tau[n]/d\tau \geq 0. \quad (15)$$

Combining this with Eq. (7) implies

$$d^2 F^\tau[n]/d\tau^2 \leq 0. \quad (16)$$

Thus $F^\tau[n]$ is concave downwards.

We can also isolate the behavior of $F_I^\tau[n]$. If we multiply Eq. (12) by τ' , and Eq. (13) by τ , and add them together, all entropic contributions cancel, yielding

$$(\tau' - \tau)(F_I^{\tau'}[n] - F_I^\tau[n]) \geq 0, \quad dF_I^\tau[n]/d\tau \geq 0. \quad (17)$$

Both $F_I^\tau[n]$ and $S^\tau[n]$ are monotonically increasing, but the net effect is that the Mermin free energy is decreasing.

Applying these conditions to the Mermin-Kohn-Sham electrons¹⁴, we find

$$dF_s^\tau[n]/d\tau = -S_s^\tau[n], \quad (18)$$

and the inequalities

$$\frac{dT_s^\tau[n]}{d\tau}, \frac{dS_s^\tau[n]}{d\tau} \geq 0 \geq \frac{dF_s^\tau[n]}{d\tau}, \frac{d^2 F_s^\tau[n]}{d\tau^2} \quad (19)$$

where subscript s denotes non-interacting, and $F_s^\tau[n] = T_s^\tau[n] - \tau S_s^\tau[n]$. Some of these relations have long been invoked for the uniform and slowly-varying gases and for constructing orbital-free density functionals (see Ref.²⁷ and references therein), but here they have been proven for every inhomogeneous system.

III. ILLUSTRATION

To illustrate these results, we calculate all energy components for an asymmetric Hubbard dimer, i.e. a **two-site Hubbard model with a potential $v_1 = -v_2$** , as described in Ref. ²⁸ for the groundstate and ²⁹ for the thermal system. Here t is the hopping parameter, U the on-site repulsion, and Δn the difference in site occupations **where the difference comes from having an inhomogeneous potential $\Delta v = v_2 - v_1$** . This is the simplest possible model in which one can perform an exact thermal calculation, including the exact thermal correlation components. Fig. 1 shows the energy components, both interacting and




FIG. 1. Energy components for the Hubbard dimer in units of $2t$, where $U = 2t$ and $\Delta n = 0$: F^τ , F_I^τ , S^τ , both interacting (solid) and non-interacting (dashed).




FIG. 2. Temperature dependence of the Mermin functional for spin-unpolarized uniform gas for several values of the Wigner-Seitz radius r_s , using the XC parametrization of Ref.¹⁶, where ϵ_F is the Fermi energy.

non-interacting, as a function of temperature for the [homogeneous system with \$\Delta n = 0\$](#) . All our exact conditions are satisfied for many values of Δn and U .

At the other extreme is the uniform electron gas and a modern parametrization of its free energy¹⁶. In the special case of a uniform density and potential, our formulas become the same as the standard thermodynamic formulas. In Fig. 2, we plot the derivative of the free energy per particle for fixed density (r_s value where $r_s = (3/(4\pi n))^{1/3}$) as a function of temperature, on the scale of the Fermi energy and in atomic units. As $r_s \rightarrow 0$, these curves converge to their well known³⁰ non-interacting value, in which the derivative is negative and decreasing everywhere, in accordance with Eq. (9). Unfortunately, by decreasing the density so that XC effects become relatively more important, we find that the parametrization violates our conditions for $r_s > 10$. Via Eq. (7), this implies that the entropy is unphysically

negative. While such low densities are irrelevant to most practical calculations using thLDA, parametrizations of the uniform gas should build in simple exact conditions such as ours. Note that our restrictions apply only to continuous parametrizations. The QMC data on which Ref.¹⁶ is based¹⁵ is for the XC energy at discrete values of the density, and so does not directly give the entropy.

For extremely high temperatures, sums over KS eigenstates become impractical, and only pure DFT can be applied. Because the uniform gas satisfies our conditions, and because Thomas-Fermi (TF) theory uses local approximations to the kinetic and entropic contributions which satisfy the conditions pointwise, we deduce that TF theory satisfies our conditions. However, recent attempts to go beyond TF theory, such as using generalized gradient approximations for the energy^{19,31,32}, should be tested for satisfaction of these constraints.

IV. EXCHANGE-CORRELATION

In the final section of this paper, we apply this reasoning to the MKS method. The Mermin functional is written in terms of the MKS quantities and a correction:

$$F^\tau[n] = F_s^\tau[n] + U_H[n] + A_{xc}^\tau[n], \quad (20)$$

called the exchange-correlation (XC) free energy. (The Hartree energy, $U_H[n]$, has no explicit temperature dependence). The XC free energy is a sum of three components:

$$A_{xc}^\tau[n] = K_{xc}^\tau[n] + U_{xc}^\tau[n] = T_{xc}^\tau[n] - \tau S_{xc}^\tau[n] + U_{xc}^\tau[n], \quad (21)$$

where U_{xc}^τ is the potential contribution and K_{xc}^τ is the kinetic contribution, which in turn consists of T_{xc}^τ , the kinetic contribution, and $-\tau S_{xc}^\tau$, where S_{xc}^τ is the entropic contribution.

Subtract Eq. (18) from Eq. (7) to find

$$\frac{dA_{xc}^\tau[n]}{d\tau} = -S_{xc}^\tau[n], \quad (22)$$

or

$$A_{xc}^\tau[n] = E_{xc}[n] - \int_0^\tau d\tau' S_{xc}^{\tau'}[n]. \quad (23)$$

All thermal XC effects are contained in the XC contribution to the entropy. This provides an intriguing alternative to the adiabatic connection formula of Ref.²³ or the thermal connection formula of Ref.²⁵. Our inequalities do not yield definite signs for XC quantities, just weak constraints that would be difficult to impose universally on an XC approximation:

$$\frac{dT_{xc}^\tau}{d\tau} \geq -\frac{dT_s^\tau}{d\tau}, \quad \frac{dS_{xc}^\tau}{d\tau} \geq -\frac{dS_s^\tau}{d\tau}. \quad (24)$$

We can also combine these with the coupling-constant derivatives of Ref.²⁵ to find Maxwell-style relations:

$$\left(\frac{\partial U_{xc}}{\partial \tau} \right)_\lambda = -\lambda \left(\frac{\partial S_{xc}}{\partial \lambda} \right)_\tau \quad (25)$$

where λ denotes evaluation at coupling-constant λ , holding the density fixed²³.

Exchange can be isolated by considering the limit of either weak interaction or scaling to the high-density limit²³. The exchange free energy is

$$A_X^\tau[n] = V_{ee}[\Gamma_S^\tau[n]] - U_H[n] \quad (26)$$

in a case of no degeneracies (the only case we consider here). Because Γ_S^τ minimizes the kentropy alone, to first order in λ , kentropy corrections must be zero. Thus

$$K_X^\tau[n] = 0, \quad T_X^\tau[n] = \tau S_X^\tau[n] = -\tau dA_{XC}^\tau[n]/d\tau. \quad (27)$$

It may seem odd to consider a kinetic contribution to exchange (impossible in the ground state), but T_X^τ vanishes as $\tau \rightarrow 0$ in Eq. (27). For a uniform gas, the thermal exchange energy is well-known³⁰. But for our Hubbard dimer²⁹, when $\langle N \rangle = 2$, we find $E_X[n] = -U_H[n]/2$, so that $T_X^\tau = S_X^\tau = 0$.




FIG. 3. Correlation entropy in the Hubbard dimer for several values of Δn as a function of temperature, in units of $2t$, where $U = 2t$.

The results of Eq. (23) apply to correlation alone and can be used in either direction, just as the relation for the full functional. They are well-known for the uniform gas from statistical mechanics^{33–35}. But for an inhomogeneous system, they are non-trivial, and so we illustrate them on the [asymmetric](#) Hubbard dimer. In Fig. 3, we plot the entropic correlation as a function of temperature for several values of Δn , the occupation difference [that arises from the asymmetric potential](#). Eq. (23) is satisfied within numerical precision. The derivative of S_C^τ can change sign, even though both $S^\tau(\Delta n)$ and $S_S^\tau(\Delta n)$ are monotonically increasing (This explains the small dip seen in Fig. 7 of Ref.²⁹).

V. DISCUSSION AND CONCLUSIONS

Finally, we explain the apparent success of the ground-state approximation (GSA) for $A_{XC}^\tau[n]$ in MKS equilibrium calculations. Almost all present-day calculations of WDM use this approximation, and a recent calculation on the Hubbard dimer²⁹ found that GSA worked well

when neither the temperature nor the strength of the correlations were large (the conditions corresponding to most WDM calculations). Now we explain why. Write

$$F^{\tau, \text{GSA}}[n] = F_S^\tau[n] + U_H[n] + E_{XC}[n]. \quad (28)$$

Clearly, all temperature dependence is contained only in the KS part (usually a very dominant piece). Since the KS piece satisfies all the different inequalities and equalities, then so does any GSA calculation. But attempt to add corrections to a GSA calculation by writing

$$A_{XC}^{\tau, \text{GSA}}[n] = E_{XC}^{\text{GSA}}[n] + \Delta A_{XC}^\tau[n]. \quad (29)$$

Only the thermal correction appears in the exact conditions we have derived, since they all contain temperature derivatives. But there is no simple way to know if the corrections will satisfy the exact conditions for all possible systems. The only case would be using local approximations for all temperature-dependent quantities, and then using energy densities from the uniform gas. Thus a TF calculation, with thermal LDA corrections, would satisfy these conditions, since they would be satisfied pointwise, as the uniform gas satisfies these conditions for every density. But in any MKS calculation using approximate thermal XC corrections, this is not guaranteed. Unless special care is taken to guarantee satisfaction of our conditions, *only* GSA automatically does this. This is analogous to the situation in TDDFT (at zero temperature): The adiabatic LDA, which ignores the history dependence that is known to exist in the TDDFT functionals, satisfies most exact conditions, while the time-dependent LDA (the Gross-Kohn approximation³⁶) violates several important constraints³⁷. All this explains why the GSA has been working well in many situations^{1,38}. The GSA appears to be correct in both the low- and high-temperature limits and, at least for model systems, reproduces the exact KS orbitals accurately²⁹. Of course, this depends on the specific property being calculated and the acceptable level of error, and does not preclude moderate deviations, especially between these extremes, i.e., warm dense matter. But any calculation that includes, e.g., semilocal thermal XC corrections, risks violating the exact conditions listed here that GSA automatically satisfies, and should be checked for such violations. On the other hand, the Hartree-Fock approximation (or rather, the DFT equivalent, called EXX³⁹), must satisfy the conditions since any expansion in powers of the coupling constant up to some order must satisfy all our conditions.

To conclude, the formulas presented here are exact conditions applying to every thermal electronic system when treated with DFT, and should guide the future construction of approximate functionals.

ACKNOWLEDGMENTS

The authors acknowledge support from the National Science Foundation (NSF) under grant CHE - 1464795.

J.C.S. acknowledges support through the NSF Graduate Research fellowship program under award # DGE-1321846. P.E.G. acknowledges support from the Department of Energy (DOE) under grant DE14-017426. A.P.J.'s work was performed under the auspices of the

U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344. A.P.J. was supported in part by the University of California President's Postdoctoral Fellowship.

- ¹ M. D. Knudson and M. P. Desjarlais, "Shock compression of quartz to 1.6 TPa: Redefining a pressure standard," *Phys. Rev. Lett.* **103**, 225501 (2009).
- ² B. I. Cho, K. Engelhorn, A. A. Correa, T. Ogitsu, C. P. Weber, H. J. Lee, J. Feng, P. A. Ni, Y. Ping, A. J. Nelson, D. Prendergast, R. W. Lee, R. W. Falcone, and P. A. Heimann, "Electronic structure of warm dense copper studied by ultrafast x-ray absorption spectroscopy," *Phys. Rev. Lett.* **106**, 167601 (2011).
- ³ Michael P. Desjarlais, "First-principles calculation of entropy for liquid metals," *Phys. Rev. E* **88**, 062145 (2013).
- ⁴ B. Militzer and W. B. Hubbard, "Ab initio equation of state for hydrogen-helium mixtures with recalibration of the giant-planet mass-radius relation," *The Astrophysical Journal* **774**, 148 (2013).
- ⁵ Hannes R. Rüter and Ronald Redmer, "*Ab Initio* simulations for the ion-ion structure factor of warm dense aluminum," *Phys. Rev. Lett.* **112**, 145007 (2014).
- ⁶ R F Smith, J H Eggert, R Jeanloz, T S Duffy, D G Braun, J R Patterson, R E Rudd, J Biener, A E Lazicki, A V Hamza, J Wang, T Braun, L X Benedict, P M Celliers, and G W Collins, "Ramp compression of diamond to five terapascals," *Nature* **511**, 330–3 (2014).
- ⁷ P. Sperling, E. J. Gamboa, H. J. Lee, H. K. Chung, E. Galtier, Y. Omarbakiyeva, H. Reinholz, G. Röpke, U. Zastrau, J. Hastings, L. B. Fletcher, and S. H. Glenzer, "Free-electron x-ray laser measurements of collisional-damped plasmons in isochorically heated warm dense matter," *Phys. Rev. Lett.* **115**, 115001 (2015).
- ⁸ M. D. Knudson, M. P. Desjarlais, A. Becker, R. W. Lemke, K. R. Cochrane, M. E. Savage, D. E. Bliss, T. R. Mattsson, and R. Redmer, "Direct observation of an abrupt insulator-to-metal transition in dense liquid deuterium," *Science* **348**, 1455–1460 (2015).
- ⁹ V. L. Campo, "Density-functional-theory approach to the thermodynamics of the harmonically confined one-dimensional hubbard model," *Phys. Rev. A* **92**, 013614 (2015).
- ¹⁰ Frank Graziani, Michael P. Desjarlais, Ronald Redmer, and Samuel B. Trickey, eds., *Frontiers and Challenges in Warm Dense Matter*, Lecture Notes in Computational Science and Engineering, Vol. 96 (Springer International Publishing, 2014).
- ¹¹ N. D. Mermin, "Thermal properties of the inhomogeneous electron gas," *Phys. Rev.* **137**, A: 1441 (1965).
- ¹² P. Hohenberg and W. Kohn, "Inhomogeneous electron gas," *Phys. Rev.* **136**, B864–B871 (1964).
- ¹³ W. Kohn and L. J. Sham, "Self-consistent equations including exchange and correlation effects," *Phys. Rev.* **140**, A1133–A1138 (1965).
- ¹⁴ Aurora Pribram-Jones, Stefano Pittalis, E.K.U. Gross, and Kieron Burke, "Thermal density functional theory in context," in *Frontiers and Challenges in Warm Dense Matter*, Lecture Notes in Computational Science and Engineering, Vol. 96, edited by Frank Graziani, Michael P. Desjarlais, Ronald Redmer, and Samuel B. Trickey (Springer International Publishing, 2014) pp. 25–60.
- ¹⁵ Ethan W. Brown, Bryan K. Clark, Jonathan L. DuBois, and David M. Ceperley, "Path-integral monte carlo simulation of the warm dense homogeneous electron gas," *Phys. Rev. Lett.* **110**, 146405 (2013).
- ¹⁶ Valentin V. Karasiev, Travis Sjostrom, James Dufty, and S. B. Trickey, "Accurate homogeneous electron gas exchange-correlation free energy for local spin-density calculations," *Phys. Rev. Lett.* **112**, 076403 (2014).
- ¹⁷ V. S. Filinov, V. E. Fortov, M. Bonitz, and Zh. Moldabekov, "Fermionic path-integral monte carlo results for the uniform electron gas at finite temperature," *Phys. Rev. E* **91**, 033108 (2015).
- ¹⁸ T. Schoof, S. Groth, J. Vorberger, and M. Bonitz, "*Ab Initio* thermodynamic results for the degenerate electron gas at finite temperature," *Phys. Rev. Lett.* **115**, 130402 (2015).
- ¹⁹ Travis Sjostrom and Jérôme Daligault, "Gradient corrections to the exchange-correlation free energy," *Phys. Rev. B* **90**, 155109 (2014).
- ²⁰ M. Levy and J.P. Perdew, "Hellmann-Feynman, virial, and scaling requisites for the exact universal density functionals. shape of the correlation potential and diamagnetic susceptibility for atoms," *Phys. Rev. A* **32**, 2010 (1985).
- ²¹ Mel Levy, "Universal variational functionals of electron densities, first-order density matrices, and natural spin-orbitals and solution of the *v*-representability problem," *Proceedings of the National Academy of Sciences of the United States of America* **76**, 6062–6065 (1979).
- ²² John P. Perdew, Kieron Burke, and Matthias Ernzerhof, "Generalized gradient approximation made simple," *Phys. Rev. Lett.* **77**, 3865–3868 (1996), *ibid.* **78**, 1396(E) (1997).
- ²³ S. Pittalis, C. R. Proetto, A. Floris, A. Sanna, C. Bersier, K. Burke, and E. K. U. Gross, "Exact conditions in finite-temperature density-functional theory," *Phys. Rev. Lett.* **107**, 163001 (2011).
- ²⁴ James W. Dufty and S. B. Trickey, "Scaling, bounds, and inequalities for the noninteracting density functionals at finite temperature," *Phys. Rev. B* **84**, 125118 (2011).
- ²⁵ A. Pribram-Jones and K. Burke, "Connection formula for thermal density functional theory," *Phys. Rev.* (2016), accepted.
- ²⁶ F. Schwabl, *Quantum Mechanics* (Springer-Verlag, 2007).
- ²⁷ Valentin V. Karasiev, Travis Sjostrom, and S. B. Trickey, "Generalized-gradient-approximation noninteracting free-energy functionals for orbital-free density functional calculations," *Phys. Rev. B* **86**, 115101 (2012).
- ²⁸ D J Carrascal, J Ferrer, J C Smith, and K Burke, "The hubbard dimer: a density functional case study of a many-body problem," *Journal of Physics: Condensed Matter* **27**, 393001 (2015).

- ²⁹ J.C. Smith, A. Pribram-Jones, and K. Burke, “Exact thermal density functional theory for a model system: Correlation components and accuracy of the zero-temperature exchange-correlation approximation,” *Phys. Rev. B* (2016), submitted.
- ³⁰ R. G. Dandrea, N. W. Ashcroft, and A. E. Carlsson, “Electron liquid at any degeneracy,” *Phys. Rev. B* **34**, 2097–2111 (1986).
- ³¹ Valentin V. Karasiev, Debajit Chakraborty, Olga A. Shukruto, and S. B. Trickey, “Nonempirical generalized gradient approximation free-energy functional for orbital-free simulations,” *Phys. Rev. B* **88**, 161108 (2013).
- ³² Travis Sjostrom and Jerome Daligault, “Nonlocal orbital-free noninteracting free-energy functional for warm dense matter,” *Phys. Rev. B* **88**, 195103 (2013).
- ³³ S. Ichimaru, *Rev. Mod. Phys.* **54**, 1017 (1982).
- ³⁴ Francois Perrot and M. W. C. Dharma-wardana, “Exchange and correlation potentials for electron-ion systems at finite temperatures,” *Phys. Rev. A* **30**, 2619–2626 (1984).
- ³⁵ Francois Perrot and M. W. C. Dharma-wardana, “Spin-polarized electron liquid at arbitrary temperatures: Exchange-correlation energies, electron-distribution functions, and the static response functions,” *Phys. Rev. B* **62**, 16536–16548 (2000).
- ³⁶ E.K.U. Gross and W. Kohn, “Local density-functional theory of frequency-dependent linear response,” *Phys. Rev. Lett.* **55**, 2850 (1985).
- ³⁷ John F. Dobson, “Harmonic-potential theorem: Implications for approximate many-body theories,” *Phys. Rev. Lett.* **73**, 2244–2247 (1994).
- ³⁸ M. D. Knudson, M. P. Desjarlais, and D. H. Dolan, “Shock-wave exploration of the high-pressure phases of carbon,” *Science* **322**, 1822–1825 (2008), <http://science.sciencemag.org/content/322/5909/1822.full.pdf>.
- ³⁹ Stephan Kümmel and Leeor Kronik, “Orbital-dependent density functionals: Theory and applications,” *Rev. Mod. Phys.* **80**, 3–60 (2008).