

This is the accepted manuscript made available via CHORUS. The article has been published as:

Classification of (2+1)-dimensional topological order and symmetry-protected topological order for bosonic and fermionic systems with on-site symmetries

Tian Lan, Liang Kong, and Xiao-Gang Wen

Phys. Rev. B **95**, 235140 — Published 22 June 2017

DOI: [10.1103/PhysRevB.95.235140](https://doi.org/10.1103/PhysRevB.95.235140)

Classification of 2+1D topological orders and SPT orders for bosonic and fermionic systems with on-site symmetries

Tian Lan,^{1,2} Liang Kong,^{3,4} and Xiao-Gang Wen^{5,1}

¹*Perimeter Institute for Theoretical Physics, Waterloo, Ontario N2L 2Y5, Canada*

²*Department of Physics and Astronomy, University of Waterloo, Waterloo, Ontario N2L 3G1, Canada*

³*Department of Mathematics and Statistics, University of New Hampshire, Durham, NH 03824, USA*

⁴*Center of Mathematical Sciences and Applications, Harvard University, Cambridge, MA 02138*

⁵*Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

In 2+1-dimensional space-time, gapped quantum states are always gapped quantum liquids (GQL) which include both topologically ordered states (with long range entanglement) and symmetry protected topological (SPT) states (with short range entanglement). In this paper, we propose a classification of 2+1D GQLs for both bosonic and fermionic systems: 2+1D bosonic/fermionic GQLs with finite on-site symmetry are classified by non-degenerate unitary braided fusion categories over a symmetric fusion category (SFC) \mathcal{E} , abbreviated as $\text{UMTC}_{/\mathcal{E}}$, together with their modular extensions and total chiral central charges. In our classification, SFC \mathcal{E} describes the symmetry, which is $\text{Rep}(G)$ for bosonic symmetry G , or $\text{sRep}(G^f)$ for fermionic symmetry G^f . As a special case of the above result, we find that the modular extensions of $\text{Rep}(G)$ classify the 2+1D bosonic SPT states of symmetry G , while the $c = 0$ modular extensions of $\text{sRep}(G^f)$ classify the 2+1D fermionic SPT states of symmetry G^f . Many fermionic SPT states are studied based on the constructions from free-fermion models. But free-fermion constructions cannot produce all fermionic SPT states. Our classification does not have such a drawback. We show that, for interacting 2+1D fermionic systems, there are exactly 16 superconducting phases with no symmetry and no fractional excitations (up to E_8 bosonic quantum Hall states). Also, there are exactly 8 $Z_2 \times Z_2^f$ -SPT phases, 2 Z_8^f -SPT phases, and so on. Besides, we show that two topological orders with identical bulk excitations and central charge always differ by the stacking of the SPT states of the same symmetry.

PACS numbers: 11.15.-q, 11.15.Yc, 02.40.Re, 71.27.+a

CONTENTS

| | | | |
|---|----|---|----|
| | | B. The limitations of the naive calculation | 14 |
| I. Introduction | 2 | VIII. Examples of 2+1D SET orders and SPT orders | 14 |
| II. Gapped quantum liquids, topological order and symmetry | 4 | A. Z_2 bosonic SPT states | 15 |
| A. The finite on-site symmetry and symmetric fusion category | 4 | B. Z_2 -SET orders for bosonic systems | 15 |
| B. Categorical description of topological excitations with symmetry | 4 | C. Two other Z_2 -SET orders for bosonic systems | 16 |
| III. Invertible GQLs and modular extension | 5 | D. The rank $N = 5$ Z_2 -SET orders for bosonic systems | 16 |
| A. Invertible GQLs | 5 | 1. The first entry in Table XXIII | 17 |
| B. Modular extension | 6 | 2. The second entry in Table XXIII | 18 |
| C. Classify 2+1D bosonic SPT states | 6 | 3. The third entry in Table XXIII | 19 |
| D. Classify 2+1D fermionic SPT states | 6 | E. Z_3 , Z_5 , and S_3 SPT orders for bosonic systems | 20 |
| IV. A full classification of 2+1D GQLs with symmetry | 7 | F. Invertible fermionic topological orders | 20 |
| V. Another description of 2+1D GQLs with symmetry | 8 | G. Z_{2n}^f SPT orders for fermionic systems | 20 |
| VI. The stacking operation of GQLs | 9 | IX. Summary | 21 |
| A. Stacking operation | 9 | A. Tables for the solutions of $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$ – imperfect tables for $\text{UMTC}_{/\mathcal{E}}$ | 27 |
| B. The group structure of bosonic SPT states | 9 | 1. Z_2 -SET orders | 27 |
| C. Mathematical construction of the stacking operation | 10 | 2. Z_3 -SET orders | 28 |
| D. Modular extensions of $\text{Rep}(G)$ | 12 | 3. S_3 -SET orders | 28 |
| E. Relation to numerical calculations | 12 | 4. $Z_2 \times Z_2$ -SET orders | 28 |
| VII. How to calculate the modular extension of a $\text{UMTC}_{/\mathcal{E}}$ | 13 | 5. $Z_2 \times Z_2^f$ -SET and Z_4^f -SET orders | 28 |
| A. A naive calculation | 13 | B. Fusion ring for the modular extensions of $\text{Rep}(G)$ or $\text{sRep}(G^f)$ when G or G^f is abelian group | 28 |
| | | C. Conditions to obtain $\text{UMTC}_{/\mathcal{E}}$'s | 31 |
| | | D. Symmetry breaking | 34 |

| | |
|---|----|
| E. Physical and mathematical meaning of UMTC/\mathcal{E} and its modular extensions | 38 |
| References | 40 |

I. INTRODUCTION

Topological order^{1–3} is a new kind of order beyond the symmetry breaking orders⁴ in gapped quantum systems. Topological orders are patterns of *long-range entanglement*⁵ in *gapped quantum liquids* (GQL)⁶. Based on the unitary modular tensor category (UMTC) theory for non-abelian statistics^{7–9}, in Ref. 10 and 11, it is proposed that 2+1D bosonic topological orders are classified by $\{\text{UMTC}\} \times \{\text{iTO}_B\}$, where $\{\text{UMTC}\}$ is the set of UMTCs and $\{\text{iTO}_B\}$ is the set of invertible topological orders (iTO)^{10,12} for 2+1D boson systems. In fact $\{\text{iTO}_B\} = \mathbb{Z}$ which is generated by the E_8 bosonic quantum Hall (QH) state, and a table of UMTCs was obtained in Ref. 11 and 13. Thus, we have a table (and a classification) of 2+1D bosonic topological orders.

In a recent work¹⁴, we show that 2+1D fermionic topological orders are classified by $\{\text{UMTC}/_{\text{sRep}(Z_2^f)}\} \times \{\text{iTO}_F\}$, where $\{\text{UMTC}/_{\text{sRep}(Z_2^f)}\}$ is the set of non-degenerate unitary braided fusion categories (UBFC) over the symmetric fusion category (SFC) $\text{sRep}(Z_2^f)$ (see Definition 3). We also require $\text{UMTC}/_{\text{sRep}(Z_2^f)}$ to have modular extensions. $\{\text{iTO}_F\}$ is the set of invertible topological orders for 2+1D fermion systems. In fact $\{\text{iTO}_F\} = \mathbb{Z}$ which is generated by the $p + ip$ superconductor. In Ref. 14 we computed the table for $\text{UMTC}/_{\text{sRep}(Z_2^f)}$ s, and obtained a table (and a classification) of 2+1D fermionic topological orders.

In Ref. 14, we also point out the importance of modular extensions. If a $\text{UMTC}/_{\text{sRep}(Z_2^f)}$ does not have a modular extension, it means that the fermion-number-parity symmetry is not on-site (*i.e.* anomalous¹⁵). On the other hand, if a $\text{UMTC}/_{\text{sRep}(Z_2^f)}$ does have modular extensions, then the $\text{UMTC}/_{\text{sRep}(Z_2^f)}$ is realizable by a lattice model of fermions. In this case, a given $\text{UMTC}/_{\text{sRep}(Z_2^f)}$ may have several modular extensions. We found that different modular extensions of $\text{UMTC}/_{\text{sRep}(Z_2^f)}$ contain information of iTO_F s.

Our result on fermionic topological orders can be easily generalized to describe bosonic/fermionic topological orders with symmetry. This will be the main topic of this paper. (Some of the results are announced in Ref. 14). In this paper, we will consider symmetric GQL phases for 2+1D bosonic/fermionic systems. The notion of GQL was defined in Ref. 6. The symmetry group of GQL is G (for bosonic systems) or G^f (for fermionic systems). If a symmetric GQL has long-range entanglement (as defined in Ref. 5 and 6), it corresponds to a symmetry enriched topological (SET) order⁵. If a symmetric GQL has short-range entanglement, it corresponds to a sym-




FIG. 1. Bosonic topological orders with symmetry G are classified by three unitary categories: SFC $\mathcal{E} = \text{Rep}(G) \subset \text{UBFC } \mathcal{C} \subset \text{UMTC } \mathcal{M}$, which describe quasiparticle excitations and symmetry-twist defects. The particles connected by lines have non-trivial mutual statistics between them.




FIG. 2. Fermionic topological orders with symmetry G^f are classified by three unitary categories: SFC $\mathcal{E} = \text{sRep}(G^f) \subset \text{UBFC } \mathcal{C} \subset \text{UMTC } \mathcal{M}$.

metry protected trivial (SPT) order [which is also known as symmetry protected topological (SPT) order]^{16–20}.

In this paper, we are going to show that, 2+1D symmetric GQLs are classified by UMTC/\mathcal{E} plus their modular extensions and chiral central charge. In other words, GQLs are labeled by three UBFCs $\mathcal{E} \subset \mathcal{C} \subset \mathcal{M}$ plus the central charge c (see Fig. 1 and 2). Roughly speaking, a UBFC can be viewed as a set of quasiparticle types, plus the data on quasiparticle fusion and braiding:

1. \mathcal{E} is a special kind of UBFC called SFC where all the quasiparticles have trivial mutual statistics between each other. Such a SFC \mathcal{E} describes the local excitations (*i.e.* the excitations that can be created by local operators). The types of those local excitations are described the representations of the symmetry group. Thus \mathcal{E} is given by $\mathcal{E} = \text{Rep}(G)$ for bosonic cases, or $\mathcal{E} = \text{sRep}(G^f)$ for fermionic cases.
2. The UBFC \mathcal{C} contains both local excitations and topological excitations (*i.e.* the excitations that cannot be created by local operators), and thus $\mathcal{E} \subset \mathcal{C}$. Those topological excitations can carry fractional statistics and fractional angular momentum s , which will be called *topological spin*. The topological excitations may also have symmetry fractionalization (such as fractional symmetry quantum numbers). We also require \mathcal{E} to include all the excitations that have trivial mutual statistics with every excitation in \mathcal{C} (which can be viewed as an operational definition of the so called *local excitation*), which leads to a mathematical notion of *UBFC over SFC* \mathcal{E} (denoted as UMTC/\mathcal{E}).

3. The UBFC \mathcal{M} contains both quasiparticle excitations and symmetry-twist defects^{21–23}, and thus $\mathcal{C} \subset \mathcal{M}$. We require that every particle in \mathcal{M} (except the trivial one) has a non-trivial mutual statistics with at least one particle in \mathcal{M} . A UBFC satisfying such a condition is called UMTC, and we call the extension from \mathcal{C} to \mathcal{M} a modular extension. (To be more precise, a modular extension of \mathcal{C} , \mathcal{M} , is a UMTC with a fully faithful embedding $\mathcal{C} \rightarrow \mathcal{M}$. In particular, even if the UMTC \mathcal{M} is fixed, different embeddings correspond to different modular extensions.) The existence of modular extensions for \mathcal{C} is an anomaly-free condition for \mathcal{C} : *the quasiparticles described by \mathcal{C} can be realized by a well defined local lattice model with on-site-symmetry in the same dimension*¹⁵.

The chiral central charge c for the edge states describes the invertible topological orders which have trivial bulk excitations.

We like to remark that symmetry charges by carried topological excitations are in general not well defined. In other words, a topological excitation may not carry a representation of the symmetry group. This phenomenon is called symmetry fractionalization. In general, a topological excitation may not even carry a projective representation of the symmetry group (which corresponds to fractionalized symmetry quantum numbers). In other words, a topological excitations can carry something more exotic than projective representations of the symmetry group. For example, in a gauge theory with gauge group K and symmetry group G , a topological excitation (a gauge charge) may carry a representation of group H which satisfies $H/K = G$. So symmetry fractionalization can be more general than fractionalized quantum numbers and projective representations of the symmetry group.

One example of the classified bosonic SET (see Table VI) is given by the Z_2^{gauge} spin liquid^{24,25} with excitations $1, e, m, f$, where 1 is the trivial excitation, e the Z_2^{gauge} charge, m the Z_2^{gauge} vortex, and f the bound state of e and m . The excitation $1, e, m$ are bosons and f is a fermion. There is also a Z_2^{sym} symmetry which exchanges e and m ^{26–28}. The excitations in such a SET state are labeled by $1_+, 1_-, f_+, f_-, e \oplus m$, which form the UBFC \mathcal{C} . They have topological spins $s_i = 0, 0, \frac{1}{2}, \frac{1}{2}, 0$ and quantum dimensions $d_i = 1, 1, 1, 1, 2$. 1_+ and 1_- are the local excitations with Z_2^{sym} charge 0 and 1. The two excitations 1_+ and 1_- form the SFC $\mathcal{E} = \text{Rep}(Z_2^{\text{sym}})$. f_+ and f_- are topological fermionic excitations with Z_2^{sym} charges 0 and 1. $e \oplus m$ is a doublet excitation that corresponds to degenerate e and m (just like the spin-1/2 doublet that corresponds to degenerate spin-up and spin-down). This is why $e \oplus m$ has a quantum dimension 2. The modular extension is obtained by adding the Z_2^{sym} -symmetry twist defect, as well as its bound states with excitations $f_+, f_-, e \oplus m$. Fig. 1 happens to describe such a SET.

As a second example, Fig. 2 describes the topological order $\mathcal{F}_{(A_1,6)}$ in Table I of Ref. 14, which has a $G^f = Z_2^f$

TABLE I. Some mathematical concepts and their physical correspondences, as well as the meaning of some notations.

| Mathematical term | Physical correspondence |
|--|--|
| UBFC (unitary braided fusion category) \mathcal{C} | Set of excitations that can braid and fuse |
| SFC (symmetric fusion category) \mathcal{E} , which is a special kind of UBFC | Set of local excitations carrying representations of symmetry group |
| UMTC (unitary modular tensor category) \mathcal{M} , which is a special kind of UBFC | Set of excitations such that every non-trivial excitation has a non-trivial mutual statistics with at least one excitation |
| UMTC/ \mathcal{E} (UBFC over \mathcal{E}) a special kind of UBFC | Set of excitations that contain a subset SFC \mathcal{E} , where \mathcal{E} is formed by the excitations that have trivial mutual statistics with all excitations |
| Modular extension | Adding symmetry-twist defects (<i>i.e.</i> gauging the symmetry) |
| Chiral central charge c | The number of right-moving edge modes minus the number of left-moving edge modes (c can be fractional) |
| Topological spin s_i | Fractional part of 2D angular momentum of the quasiparticle i |
| Quantum dimension d_i | The effective dimension of the Hilbert space for the internal degrees of freedom of the quasiparticle i (d_i can be non-integer) |
| N | Number of particle types (also called rank of category) |
| D | $\sqrt{\sum_i d_i^2}$ (total quantum dim.) |
| Θ | $D^{-1} \sum_i e^{2\pi i s_i} d_i^2 = \Theta e^{2\pi i c/8}$ |
| $N_c^{ \Theta }$ | A short label of topological orders |
| N_c^B | When $ \Theta = 1$, rewrite $N_c^{ \Theta }$ as N_c^B |
| ζ_n^m | $\sin \frac{\pi(m+1)}{n+2} / \sin \frac{\pi}{n+2}$ |
| (A_n, k) | Topological order of $SU(n+1)$ level- k Chern-Simons theory |
| (B_n, k) | Topological order of $SO(2n+1)$ level- k Chern-Simons theory |
| (C_n, k) | Topological order of $Sp(2n)$ level- k Chern-Simons theory |
| (D_n, k) | Topological order of $SO(2n)$ level- k Chern-Simons theory |
| \boxtimes | Stacking of two states |
| \otimes | Fusion of two particles |

symmetry. The state has two types of local excitations with Z_2^f -charge 0 (a boson) and 1 (a fermion) that form the SFC $\mathcal{E} = \text{sRep}(Z_2^f)$. They have topological spin $s_i = 0, \frac{1}{2}$. The state also has two types of topological excitations with topological spin $s_i = \frac{1}{4}, -\frac{1}{4}$ and quantum dimension $d_i = 1 + \sqrt{2}, 1 + \sqrt{2}$. The local and topological excitations form the UBFC \mathcal{C} . The modular extension

is obtained by adding the Z_2^f -symmetry twist defect, as well as its bound state with the excitations in \mathcal{C} , which gives rise to three types of symmetry twist defects.

There is another more precise and mathematical way to phrase our result: we find that the structure $\mathcal{E} \hookrightarrow \mathcal{C} \hookrightarrow \mathcal{M}$ (plus the chiral central charge c) classifies the 2+1D GQLs with symmetry \mathcal{E} , where \hookrightarrow represents the embeddings and $\mathcal{E}_{\mathcal{M}}^{\text{cen}} = \mathcal{C}$ (see Definition 2).

As a special case of the above result, we find that bosonic 2+1D SPT phase with symmetry G are classified by the modular extensions of $\text{Rep}(G)$, while fermionic 2+1D SPT phase with symmetry G^f are classified by the modular extensions of $\text{sRep}(G^f)$ that have central charge $c = 0$.

We like to mention that Ref. 29 has classified bosonic GQLs with symmetry G , using G -crossed UMTCs. This paper uses a different approach so that we can classify both bosonic and fermionic GQLs with symmetry. For bosonic systems, the two approaches produces identical classification. We also like to mention that there is a mathematical companion Ref. 30 of this paper, where one can find detailed proof and explanations for related mathematical results.

The paper is organized as the following. In Section II, we review the notion of topological order and introduce category theory as a theory of quasiparticle excitations in a GQL. We will introduce a categorical way to view the symmetry. In Section III, we discuss invertible GQLs and their classification based on modular extensions. In Sections IV and V, we generalize the above results and propose a classification of all GQLs. Section VI investigates the stacking operation from physical and mathematical points of view. Section VII describes how to numerically calculate the modular extensions and Section VIII discusses some simple examples. For people with physics background, one way to read this paper is to start with the Sections II and V, and then go to Section VIII for the examples. The Table I summarizes some important mathematical concepts and their physical correspondences.

II. GAPPED QUANTUM LIQUIDS, TOPOLOGICAL ORDER AND SYMMETRY

A. The finite on-site symmetry and symmetric fusion category

In this paper, we consider physical systems with an on-site symmetry described by a finite group G . For fermionic systems, we further require that G contains a central Z_2 fermion-number-parity subgroup. More precisely, fermionic symmetry group is a pair (G, f) , where G is a finite group, $f \neq 1$ is an element of G satisfying $f^2 = 1, fg = gf, \forall g \in G$. We denote the pair (G, f) as G^f .

There is another way to view the on-site symmetries, which is nicer because bosonic and fermionic symme-

tries can be formulated in the same manner. Consider a bosonic/fermionic product state $|\psi\rangle$ that does not break the symmetry G : $U_g|\psi\rangle = |\psi\rangle, g \in G$. Then the new way to view the symmetry is to use the properties of the excitations above the product state to encode the information of the symmetry G .

The product state contain only local excitations that can be created by acting local operators O on the ground state $O|\psi\rangle$. For any group action $U_g, U_g O|\psi\rangle = U_g O U_g^\dagger U_g |\psi\rangle = U_g O U_g^\dagger |\psi\rangle$ is an excited state with the same energy as $O|\psi\rangle$. Since we assume the symmetry to be on-site, $U_g O U_g^\dagger$ is also a local operator. Therefore, $U_g O U_g^\dagger |\psi\rangle$ and $O|\psi\rangle$ correspond to the degenerate local excitations. We see that local excitations “locally” carry group representations. In other words, different types of local excitations are labeled by irreducible representations of the symmetry group.

By looking at how the local excitations (more precisely, their group representations) fuse and braid with each other, we arrive at the mathematical structure called symmetric fusion categories (SFC). By definition a SFC is a braided fusion category where all the objects (the excitations) have trivial mutual statistics (*i.e.* centralize each other, see next section). A SFC is automatically a unitary braided fusion category.

In fact, there are only two kinds of SFCs: one is representation category of G : $\text{Rep}(G)$, with the usual braiding (all representations are bosonic); the other is $\text{sRep}(G^f)$ where an irreducible representation is bosonic if f is represented trivially (+1), and fermionic if f is represented non-trivially (−1).

It turns out that SFC (or the fusion and braiding properties of the local excitations) fully characterize the symmetry group (which is known as Tannaka duality³¹). Therefore, a finite on-site symmetry is equivalently given by a SFC \mathcal{E} . Also, by checking the braiding in \mathcal{E} we know whether it is bosonic or fermionic. This is the new way, the categorical way, to view the symmetry. Such a categorical view of bosonic/fermionic symmetry allows us to obtain a classification of symmetric topological/SPT orders.

B. Categorical description of topological excitations with symmetry

In symmetric GQLs with topological order (*i.e.* with long range entanglement), there can be particle-like excitations with local energy density, but they cannot be created by local operators. They are known as (non-trivial) topological excitations. Topological excitations do not necessarily carry group representations. Nevertheless, we can still study how they fuse and braid with each other; so we have a unitary braided fusion category (UBFC) to describe the particle-like excitations. To proceed, we need the following key definition on “centralizers.”

Definition 1. The objects X, Y in a UBFC \mathcal{C} are said

to *centralize* (mutually local to) each other if

$$c_{Y,X} \circ c_{X,Y} = \text{id}_{X \otimes Y}, \quad (1)$$

where $c_{X,Y} : X \otimes Y \cong Y \otimes X$ is the braiding in \mathcal{C} .

Physically, we say that X and Y have trivial mutual statistics.

Definition 2. Given a subcategory $\mathcal{D} \subset \mathcal{C}$, its *centralizer* $\mathcal{D}_{\mathcal{C}}^{\text{cen}}$ in \mathcal{C} is the full subcategory of objects in \mathcal{C} that centralize all the objects in \mathcal{D} .

We may roughly view a category as a “set” of particle-like excitations. So the centralizer $\mathcal{D}_{\mathcal{C}}^{\text{cen}}$ is the “subset” of particles in \mathcal{C} that have trivial mutual statistics with all the particles in \mathcal{D} .

Definition 3. A UBFC \mathcal{E} is a *symmetric* fusion category if $\mathcal{E}_{\mathcal{E}}^{\text{cen}} = \mathcal{E}$. A UBFC \mathcal{C} with a fully faithful embedding $\mathcal{E} \hookrightarrow \mathcal{C}_{\mathcal{C}}^{\text{cen}}$ is called a UBFC over \mathcal{E} . Moreover, \mathcal{C} is called a non-degenerate UBFC over \mathcal{E} , or $\text{UMTC}_{/\mathcal{E}}$, if $\mathcal{C}_{\mathcal{C}}^{\text{cen}} = \mathcal{E}$.

Definition 4. Two UBFCs over \mathcal{E} , \mathcal{C} and \mathcal{C}' are equivalent if there is a unitary braided equivalence $F : \mathcal{C} \rightarrow \mathcal{C}'$ such that it preserves the embeddings, i.e., the following diagram commute.

$$\begin{array}{ccc} \mathcal{E} & \hookrightarrow & \mathcal{C} \\ \parallel & & \downarrow F \\ \mathcal{E} & \hookrightarrow & \mathcal{C}' \end{array} \quad (2)$$

We denote the category of unitary braided auto-equivalences of \mathcal{C} by $\text{Aut}(\mathcal{C})$ and its underlining group by $\text{Aut}(\mathcal{C})$.

We recover the usual definition of UMTC when \mathcal{E} is trivial, i.e. the category of Hilbert spaces, denoted by $\text{Vec} = \text{Rep}(\{1\})$. In this case the subscript is omitted.

Physically, a UBFC \mathcal{C} is the collection of all bulk topological excitations plus their fusion and braiding data. Requiring \mathcal{C} to be a $\text{UMTC}_{/\mathcal{E}}$ means: (1) the set of local excitations, \mathcal{E} (which is the set of all the irreducible representations of the symmetry group), is included in \mathcal{C} as a subcategory; (2) \mathcal{C} is anomaly-free, i.e. all the topological excitations (the ones not in \mathcal{E}) can be detected by mutual braiding¹⁰. In other words, every topological excitation must have non-trivial mutual statistics with some excitations. Those excitations that cannot be detected by mutual braiding (i.e., $\mathcal{C}_{\mathcal{C}}^{\text{cen}}$) are exactly the local excitations in \mathcal{E} . Moreover, we want the symmetry to be on-site (gaugeable), which requires the existence of modular extensions (see Definition 6). Such an understanding leads to the following conjecture:

Conjecture 1. Bulk topological excitations of topological orders with symmetry \mathcal{E} are classified by $\text{UMTC}_{/\mathcal{E}}$'s that have modular extensions.

We like to remark that $\text{UMTC}_{/\mathcal{E}}$'s fail to classify topological orders. This is because two different topologically ordered phases may have bulk topological excitations with the same non-abelian statistics (i.e. described by the same $\text{UMTC}_{/\mathcal{E}}$). However, $\text{UMTC}_{/\mathcal{E}}$'s, with modular extensions, do classify topological orders up to invertible ones. See next section for details. The relation between anomaly and modular extension will also be discussed later.

III. INVERTIBLE GQLS AND MODULAR EXTENSION

A. Invertible GQLs

There exist non-trivial topological ordered states that have only trivial topological excitations in the bulk (but non-trivial edge states). They are “invertible” under the stacking operation^{10,12} (see Section VI for details). More generally, we define

Definition 5. A GQL is invertible if its bulk topological excitations are all trivial (i.e. can all be created by local operators).

Consider some invertible GQLs with the same symmetry \mathcal{E} . The bulk excitations of those invertible GQLs are the same which are described by the same SFC \mathcal{E} . Now the question is: How to distinguish those invertible GQLs?

First, we believe that invertible bosonic topological orders with no symmetry are generated by the E_8 QH state (with central charge $c = 8$) via time-reversal and stacking, and form a \mathbb{Z} group. Stacking with an E_8 QH state only changes the central charge by 8, and does not change the bulk excitations or the symmetry. So the only data we need to know to determine the invertible bosonic topological order with no symmetry is the central charge c . The story is parallel for invertible fermionic topological orders with no symmetry, which are believed to be generated by the $p + ip$ superconductor state with central charge $c = 1/2$.

Second, invertible bosonic GQLs with symmetry are generated by bosonic SPT states and invertible bosonic topological orders (i.e. E_8 states) via stacking. We know that the bosonic SPT states with symmetry G are classified by the 3-cocycles in $H^3[G, U(1)]$. Therefore, bosonic invertible GQLs with symmetry G are classified by $H^3[G, U(1)] \times \mathbb{Z}$ (where \mathbb{Z} corresponds to layers of E_8 states).

However, this result and this point of view is not natural to generalize to fermionic cases or non-invertible GQLs. Thus, we introduce an equivalent point of view, which can cover boson, fermion, and non-invertible GQLs in the same fashion.

B. Modular extension

First, we introduce the notion of modular extension of a UMTC $_{/\mathcal{E}}$:

Definition 6. Given a UMTC $_{/\mathcal{E}}$ \mathcal{C} , its *modular extension* is a UMTC \mathcal{M} , together with a fully faithful embedding $\iota_{\mathcal{M}} : \mathcal{C} \hookrightarrow \mathcal{M}$, such that $\mathcal{E}_{\mathcal{M}}^{\text{cen}} = \mathcal{C}$, equivalently $\dim(\mathcal{M}) = \dim(\mathcal{C}) \dim(\mathcal{E})$.

Two modular extensions \mathcal{M} and \mathcal{M}' are equivalent if there is an equivalence between the UMTCs $F : \mathcal{M} \rightarrow \mathcal{M}'$ that preserves the embeddings, i.e., the following diagram commute.

$$\begin{array}{ccc} \mathcal{C} & \hookrightarrow & \mathcal{M} \\ \parallel & & \downarrow F \\ \mathcal{C} & \hookrightarrow & \mathcal{M}' \end{array} \quad (3)$$

We denote the set of equivalent classes of modular extensions of \mathcal{C} by $\mathcal{M}_{\text{ext}}(\mathcal{C})$.

Remark 1. Since the total quantum dimension of modular extensions of a given \mathcal{C} is fixed, there are only finitely many different modular extensions, due to Ref. 32. In principle we can always perform a finite search to exhaust all the modular extensions.

Remember that \mathcal{C} describes the particle-like excitations in our topological state. Some of those excitations are local that have trivial mutual statistics with all other excitations. Those local excitations form $\mathcal{E} \subset \mathcal{C}$. The modular extension \mathcal{M} of \mathcal{C} is obtained as adding particles that have non-trivial mutual statistics with the local excitations in \mathcal{E} , so that every particle in \mathcal{M} will always have non-trivial mutual statistics with some particles in \mathcal{M} . Since the particles in \mathcal{E} carry “charges” (i.e. the irreducible representations of G), the added particles correspond to “flux” (i.e. the symmetry twists of G). So the modular extension correspond to gauging²¹ the on-site symmetry G . Since we can use the gauged symmetry to detect SPT orders²³, we like to propose the following conjecture

Conjecture 2. Invertible bosonic GQLs with symmetry $\mathcal{E} = \text{Rep}(G)$ are classified by (\mathcal{M}, c) where \mathcal{M} is a modular extension of \mathcal{E} and $c = 0 \bmod 8$.

C. Classify 2+1D bosonic SPT states

Invertible bosonic GQLs described by (\mathcal{M}, c) include both bosonic SPT states and bosonic topological orders. Among those, $(\mathcal{M}, c = 0)$ classify bosonic SPT states. In other words:

Corollary 1. 2+1D bosonic SPT states with symmetry G are classified by the modular extensions of $\text{Rep}(G)$ (which always have $c = 0$).

In Ref. 18–20, it was shown that 2+1D bosonic SPT states are classified by $H^3[G, U(1)]$. Such a result agrees with our conjecture, due to the following theorem, which follows immediately from results in Ref. 33.

Theorem 2. The modular extensions of $\text{Rep}(G)$ 1-to-1 correspond to 3-cocycles in $H^3[G, U(1)]$. The central charge of these modular extensions are $c = 0 \bmod 8$.

Remark 2. In Sec. VID, we give more detailed explanation of the 1-to-1 correspondence in Theorem 2. Moreover, we will prove a stronger result in Theorem 11. It turns out that the set $\mathcal{M}_{\text{ext}}(\text{Rep}(G))$ of modular extensions of $\text{Rep}(G)$ is naturally equipped with a physical stacking operation such that $\mathcal{M}_{\text{ext}}(\text{Rep}(G))$ forms an abelian group, which is isomorphic to the group $H^3[G, U(1)]$.

Remark 3. $c/8$ determines the number of layers of the E_8 QH states, which is the topological order part of invertible bosonic symmetric GQLs. In other words

$$\begin{aligned} & \{\text{invertible bosonic symmetric GQLs}\} \\ &= \{\text{bosonic SPT states}\} \times \{\text{layers of } E_8 \text{ states}\}. \end{aligned} \quad (4)$$

D. Classify 2+1D fermionic SPT states

The above approach also apply to fermionic case. Note that, the invertible fermionic GQLs with symmetry G^f have bulk excitations described by SFC $\mathcal{E} = \text{sRep}(G^f)$. So we would like to conjecture that

Conjecture 3. Invertible fermionic GQLs with symmetry G^f are classified by (\mathcal{M}, c) , where \mathcal{M} is a modular extension of $\mathcal{E} = \text{sRep}(G^f)$, and c is the central charge determining the layers of $\nu = 8$ IQH states.

Remark 4. Note that, the central charge $c \bmod 8$ is determined by \mathcal{M} , while $(c - \text{mod}(c, 8))/8$ determines the number of layers of the $\nu = 8$ IQH states.

Remark 5. Invertible fermionic symmetric GQLs include both fermion SPT states and fermionic topological orders. (\mathcal{M}, c) with $c = 0$ classify fermionic SPT states.

In other words,

Corollary 3. 2+1D fermionic SPT states with symmetry G are classified by the $c = 0$ modular extensions of $\text{sRep}(G^f)$.

Remark 6. Unlike the bosonic case, in general

$$\begin{aligned} & \{\text{invertible fermionic symmetric GQLs}\} \\ & \neq \{\text{fermionic SPT states}\} \times \{\text{layers of } p + ip \text{ states}\}. \end{aligned} \quad (5)$$

For example (see Table XV)

$$\begin{aligned} & \{\text{invertible } Z_4^f \text{ fermionic symmetric GQLs}\} \\ &= \{\text{fermionic } Z_4^f \text{-SPT states}\} \times \\ & \quad \{\text{layers of } \nu = 1 \text{ integer quantum Hall states}\}. \end{aligned} \quad (6)$$

But we have

$$\begin{aligned} & \{\text{invertible fermionic symmetric GQLs}\} \\ &= \{\text{invertible fermionic symmetric GQLs with } c \in [0, 8)\} \\ & \quad \times \{\text{layers of } E_8\text{-states}\}. \end{aligned} \quad (7)$$

Or when $G^f = G_b \times Z_2^f$

$$\begin{aligned} & \{\text{invertible fermionic symmetric GQLs}\} \\ &= \{\text{fermionic SPT states}\} \times \{\text{layers of } p + ip \text{ states}\}, \end{aligned} \quad (8)$$

where the fermions in the $p + ip$ states are G_b -invariant.

When there is no symmetry, the invertible fermionic GQLs become the invertible fermionic topological order, which have bulk excitations described by $\mathcal{E} = \text{sRep}(Z_2^f)$. $\text{sRep}(Z_2^f)$ has 16 modular extensions, with central charges $c = n/2, n = 0, 1, 2, \dots, 15$. There is only one modular extension with $c = 0$, which correspond to trivial product state. Thus there is no non-trivial fermionic SPT state when there is no symmetry, as expected.

The modular extensions with $c = n/2$ correspond to invertible fermionic topological order formed by n layers of $p + ip$ states. Since the modular extensions can only determine $c \bmod 8$, in order for the above picture to be consistent, we need to show the following

Theorem 4. The stacking of 16 layers $c = 1/2 p + ip$ states is equivalent to a $\nu = 8$ IQH state, which is in turn equivalent to a E_8 bosonic QH state stacked with a trivial fermionic product state.

Proof. First, two layers of $p + ip$ states is equal to one layer of $\nu = 1$ IQH state. Thus, 16 layers $c = 1/2 p + ip$ states is equivalent to a $\nu = 8$ IQH state. To show $\nu = 8$ IQH state is equivalent to E_8 bosonic QH state stacked with a trivial fermionic product state, we note that the $\nu = 8$ IQH state is described by K -matrix $K_{\nu=8} = I_{8 \times 8}$ which is a 8-by-8 identity matrix. While the E_8 bosonic QH state stacked with a trivial fermionic product state

is described by K -matrix $K_{E_8 \boxtimes \mathcal{F}_0} = K_{E_8} \oplus \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$,

where K_{E_8} is the matrix that describe the E_8 root lattice. We also know that two odd³⁴ K -matrices K_1 and K_2 describe the same fermionic topological order if after direct summing with proper number of $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$'s:

$$\begin{aligned} K'_1 &= K_1 \oplus \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \oplus \dots \\ K'_2 &= K_2 \oplus \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \oplus \dots, \end{aligned} \quad (9)$$

K'_1 and K'_2 become equivalent, *i.e.*

$$K'_1 = UK'_2U^T, \quad U \in SL(N, \mathbb{Z}). \quad (10)$$

Notice that $K_{\nu=8} \oplus \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ and $K_{E_8 \boxtimes \mathcal{F}_0}$ have the same determinant -1 and the same signature. Using the result that odd matrices with ± 1 determinants are equivalent if they have the same signature, we find that $K_{\nu=8} \oplus \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ and $K_{E_8 \boxtimes \mathcal{F}_0}$ are equivalent. Therefore $\nu = 8$ IQH state is equivalent to E_8 bosonic QH state stacked with a trivial fermionic product state. \square

IV. A FULL CLASSIFICATION OF 2+1D GQLS WITH SYMMETRY

We have seen that all invertible GQLs with symmetry G (or G^f) have the same kind of bulk excitations, described by $\text{Rep}(G)$ (or $\text{sRep}(G^f)$). To classify distinct invertible GQLs that shared the same kind of bulk excitations, we need to compute the modular extensions of $\text{Rep}(G)$ (or $\text{sRep}(G^f)$). This result can be generalized to non-invertible topological orders.

In general, the bulk excitations of a 2+1D bosonic/fermionic SET are described by a $\text{UMTC}_{/\mathcal{E}} \mathcal{C}$. However, there can be many distinct SET orders that have the same kind of bulk excitations described by the same \mathcal{C} . To classify distinct invertible SET orders that shared the same kind of bulk excitations \mathcal{C} , we need to compute the modular extensions of \mathcal{C} . This leads to the following

Conjecture 4. 2+1D GQLs with symmetry \mathcal{E} (*i.e.* the 2+1D SET orders) are classified by $(\mathcal{C}, \mathcal{M}, c)$, where \mathcal{C} is a $\text{UMTC}_{/\mathcal{E}}$ describing the bulk topological excitations, \mathcal{M} is a modular extension of \mathcal{C} describing the edge state up to E_8 states, and c is the central charge determining the layers of E_8 states.

Let \mathcal{M} be a modular extension of a $\text{UMTC}_{/\mathcal{E}} \mathcal{C}$. We note that all the simple objects (particles) in \mathcal{C} are contained in \mathcal{M} as simple objects. Assume that the particle labels of \mathcal{M} are $\{i, j, \dots, x, y, \dots\}$, where i, j, \dots correspond to the particles in \mathcal{C} and x, y, \dots the additional particles (not in \mathcal{C}). Physically, the additional particles x, y, \dots correspond to the symmetry twists of the on-site symmetry²². The modular extension \mathcal{M} describes the fusion and the braiding of original particles i, j, \dots with the symmetry twists. In other words, the modular extension \mathcal{M} is the resulting topological order after we gauge the on-site symmetry²¹.

Now, it is clear that the existence of modular extension is closely related to the on-site symmetry (*i.e.* anomaly-free symmetry) which is gaugable (*i.e.* allows symmetry twists). For non-on-site symmetry (*i.e.* anomalous symmetry¹⁵), the modular extension does not exist since the symmetry is not gaugable (*i.e.* does not allow symmetry twists). We also have

Conjecture 5. 2+1D GQLs with anomalous symmetry¹⁵ \mathcal{E} are classified by $\text{UMTC}_{/\mathcal{E}}$'s that have no modular extensions.

It is also important to clarify the equivalence relation between the triples $(\mathcal{C}, \mathcal{M}, c)$. Two triples $(\mathcal{C}, \mathcal{M}, c)$ and $(\mathcal{C}', \mathcal{M}', c')$ are equivalent if: (1) $c = c'$; (2) there exists braided equivalences $F_{\mathcal{C}} : \mathcal{C} \rightarrow \mathcal{C}'$ and $F_{\mathcal{M}} : \mathcal{M} \rightarrow \mathcal{M}'$ such that all the embeddings are preserved, i.e., the following diagram commutes.

$$\begin{array}{ccccc} \mathcal{E} & \hookrightarrow & \mathcal{C} & \hookrightarrow & \mathcal{M} \\ \parallel & & \downarrow F_{\mathcal{C}} & & \downarrow F_{\mathcal{M}} \\ \mathcal{E} & \hookrightarrow & \mathcal{C}' & \hookrightarrow & \mathcal{M}' \end{array} \quad (11)$$

The equivalence classes will be in one-to-one correspondence with GQLs (i.e. SET orders and SPT orders).

Note that the group of the automorphisms of a $\text{UMTC}_{/\mathcal{E}}$ \mathcal{C} , denoted by $\text{Aut}(\mathcal{C})$ (recall Definition 4), naturally acts on the modular extensions $\mathcal{M}_{\text{ext}}(\mathcal{C})$ by changing the embeddings, i.e. $F \in \text{Aut}(\mathcal{C})$ acts as follows:

$$(\mathcal{C} \hookrightarrow \mathcal{M}) \mapsto (\mathcal{C} \xrightarrow{F} \mathcal{C} \hookrightarrow \mathcal{M})$$

For a fixed \mathcal{C} , the above equivalence relation amounts to say that GQLs with bulk excitations described by a fixed \mathcal{C} are in one-to-one correspondence with the quotient $\mathcal{M}_{\text{ext}}(\mathcal{C})/\text{Aut}(\mathcal{C})$ plus a central charge c . When $\mathcal{C} = \mathcal{E}$, the GQLs with bulk excitations described by \mathcal{E} and central charge $c = 0$ are SPT phases. In this case, the group $\text{Aut}(\mathcal{E})$, where \mathcal{E} is viewed as the trivial $\text{UMTC}_{/\mathcal{E}}$, is trivial. Thus, SPT phases are classified by the modular extensions of \mathcal{E} with $c = 0$.

V. ANOTHER DESCRIPTION OF 2+1D GQLS WITH SYMMETRY

Although the above result has a nice mathematical structure, it is hard to implement numerically to produce a table of GQLs. To fix this problem, we propose a different description of 2+1D GQLs. The second description is motivated by a conjecture that the fusion and the spins of the particles, $(\mathcal{N}_K^{IJ}, \mathcal{S}_I)$, completely characterize a UMTC . We conjecture that

Conjecture 6. The data $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i; \mathcal{N}_K^{IJ}, \mathcal{S}_I; c)$, up to some equivalence relations, gives a one-to-one classification of 2+1D GQLs with symmetry G (for boson) or G^f (for fermion), with a restriction that the symmetry group can be fully characterized by the fusion ring of its irreducible representations. The data $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i; \mathcal{N}_K^{IJ}, \mathcal{S}_I; c)$ satisfies the conditions described in Appendix C (see Ref. 11 for UMTC s).

Here $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i; \mathcal{N}_K^{IJ}, \mathcal{S}_I; c)$ is closely related to $(\mathcal{E}; \mathcal{C}; \mathcal{M}; c)$ discussed above. The data $(\tilde{N}_c^{ab}, \tilde{s}_a)$ describes the symmetry (i.e. the SFC \mathcal{E}): $a = 1, \dots, \tilde{N}$

label the irreducible representations and \tilde{N}_c^{ab} are the fusion coefficients of irreducible representations. $\tilde{s}_a = 0$ or $1/2$ depending on if the fermion-number-parity transformation f is represented trivially or non-trivially in the representation a . The data (N_k^{ij}, s_i) describes fusion and the spins of the bulk particles $i = 1, \dots, N$ in the GQL. The data (N_k^{ij}, s_i) contains $(\tilde{N}_c^{ab}, \tilde{s}_a)$ as a subset, where a is identified with the first \tilde{N} particles of the GQL. The data $(\mathcal{N}_K^{IJ}, \mathcal{S}_I)$ describes fusion and the spins of a UMTC , and it includes (N_k^{ij}, s_i) as a subset, where i is identified with the first N particles of the UMTC . Also among all the particles in UMTC , only the first N (i.e. $I = 1, \dots, N$) have trivial mutual statistics with first \tilde{N} particles (i.e. $I = 1, \dots, \tilde{N}$). Last, c is the chiral central charge of the edge state.

If the data $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$ fully characterized the $\text{UMTC}_{/\mathcal{E}}$, then the Conjecture 6 would be equivalent to the Conjecture 4. However, for non-modular tensor category, $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$ fails to fully characterize a $\text{UMTC}_{/\mathcal{E}}$. In other words, there are different $\text{UMTC}_{/\mathcal{E}}$'s that have the same data $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$. We need to include the extra data, such as the F -tensor and the R -tensor, to fully characterize the $\text{UMTC}_{/\mathcal{E}}$.

In Appendix A, we list the data $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$ that satisfy the conditions in Appendix C (without the modular extension condition) in many tables. Those tables include all the $\text{UMTC}_{/\mathcal{E}}$'s (up to certain total quantum dimensions), but the tables are not perfect: (1) some entries in the tables may be fake and do not correspond to any $\text{UMTC}_{/\mathcal{E}}$ (for the conditions are only necessary); (2) some entries in the tables may correspond to more than one $\text{UMTC}_{/\mathcal{E}}$ (since $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$ does not fully characterize a $\text{UMTC}_{/\mathcal{E}}$).

We then continue to compute $(\mathcal{N}_K^{IJ}, \mathcal{S}_I; c)$, the modular extensions of $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$. We find that the modular extensions can fix the imperfectness mentioned above. First, we find that the fake entries do not have modular extensions, and are ruled out. Second, as we will show in Section VI, all $\text{UMTC}_{/\mathcal{E}}$'s have the same numbers of modular extensions (if they exist); therefore, the entry that corresponds to more $\text{UMTC}_{/\mathcal{E}}$'s has more modular extensions. The modular extensions can tell us which entries correspond to multiple $\text{UMTC}_{/\mathcal{E}}$'s. This leads to the conjecture that the full data $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i; \mathcal{N}_K^{IJ}, \mathcal{S}_I; c)$ gives rise to an one-to-one classification of 2+1D GQLs, and allows us to calculate the tables of 2+1D GQLs, which include 2+1D SET states and 2+1D SPT states. Those are given in Section VIII.

As for the equivalence relation, we only need to consider $(\mathcal{N}_K^{IJ}, \mathcal{S}_I; c)$, since the data $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$ is included in $(\mathcal{N}_K^{IJ}, \mathcal{S}_I; c)$. Two such data $(\mathcal{N}_K^{IJ}, \mathcal{S}_I; c)$ and $(\mathcal{N}_K^{IJ}, \mathcal{S}_I; \bar{c})$ are called equivalent if $c = \bar{c}$, and $(\mathcal{N}_K^{IJ}, \mathcal{S}_I)$ and $(\mathcal{N}_K^{IJ}, \mathcal{S}_I)$ are related by two permutations of indices in the range $N_{\mathcal{M}} \geq I > N$ and in the range $N \geq I > \tilde{N}$, where $N_{\mathcal{M}}$ is the range of I . Such an equivalence relation corresponds to the one in eqn. (11) and will be called

the TO-equivalence relation. We use the TO-equivalence relation to count the number of GQL phases (*i.e.* the number of SET orders and SPT orders).

We can also define another equivalence relation, called ME-equivalence relation: we say $(\mathcal{N}_K^{IJ}, \mathcal{S}_I; c)$ and $(\tilde{\mathcal{N}}_K^{IJ}, \tilde{\mathcal{S}}_I; \bar{c})$ to be ME-equivalent if $c = \bar{c}$ and they only differ by a permutation of indices in range $I > N$. The ME-equivalence relation is closely related to the one defined in eqn. (3). We use the ME-equivalence relation to count the number of modular extensions of a *fixed* \mathcal{C} .

Last, let us explain the restriction on the symmetry group. In the Conjecture 6, we try to use the fusion \tilde{N}_c^{ab} of the irreducible representations to characterize the symmetry group. However, it is known that certain different groups may have identical fusion ring for their irreducible representations. So we need to restrict the symmetry group to be the group that can be fully characterized by its fusion ring. Those groups include simple groups and abelian groups³⁵. If we do not impose such a restriction, then the Conjecture 6 give rise to GQLs with a given symmetry fusion ring, instead of a given symmetry group.

VI. THE STACKING OPERATION OF GQLS

A. Stacking operation

Consider two GQLs \mathcal{C}_1 and \mathcal{C}_2 . If we stack them together (without introducing interactions between them), we obtain another GQL, which is denoted by $\mathcal{C}_1 \boxtimes \mathcal{C}_2$. The stacking operation \boxtimes makes the set of GQLs into a monoid. \boxtimes does not makes the set of GQLs into a group, because in general, a GQL \mathcal{C} may not have an inverse under \boxtimes . *i.e.* there is no GQL \mathcal{D} such that $\mathcal{C} \boxtimes \mathcal{D}$ becomes a trivial product state. This is because when a GQL have non-trivial topological excitations, stacking it with another GQL can never cancel out those topological excitations.

When we are considering GQLs with symmetry \mathcal{E} , the simple stacking \boxtimes will “double” the symmetry, leads to a GQL with symmetry $\mathcal{E} \boxtimes \mathcal{E}$ ($\text{Rep}(G \times G)$ or $\text{sRep}(G^f \times G^f)$). In general we allow local interactions between the two layers to break some symmetry such that the resulting system only has the original symmetry \mathcal{E} (In terms of the symmetry group, keep only the subgroup $G \hookrightarrow G \times G$ with the diagonal embedding $g \mapsto (g, g)$). This leads to the stacking between GQLs with symmetry \mathcal{E} , denoted by $\boxtimes_{\mathcal{E}}$. Similarly, $\boxtimes_{\mathcal{E}}$ makes GQLs with symmetry \mathcal{E} a monoid, but in general not all GQLs are invertible.

However, if the bulk excitations of \mathcal{C} are all local (*i.e.* all described by SFC \mathcal{E}), then \mathcal{C} will have an inverse under the stacking operation $\boxtimes_{\mathcal{E}}$, and this is why we call such GQL invertible. Those invertible GQLs include invertible topological orders and SPT states.

B. The group structure of bosonic SPT states

We have proposed that 2+1D SPT states are classified by $c = 0$ modular extensions of the SFC \mathcal{E} that describes the symmetry. Since SPT states are invertible, they form a group under the stacking operation $\boxtimes_{\mathcal{E}}$. This implies that the modular extensions of the SFC should also form a group under the stacking operation. So checking if the modular extensions of the SFC have a group structure is a way to find support for our conjecture.

However, in this section, we will first discuss such stacking operation and group structure from a physical point of view. We will only consider bosonic SPT states.

It has been proposed that the bosonic SPT states are described by group cohomology $\mathcal{H}^{d+1}[G, U(1)]$ ^{18–20}. However, it has not been shown that those bosonic SPT states form a group under stacking operation. Here we will fill this gap. An ideal bosonic SPT state of symmetry G in $d + 1$ D is described the following path integral

$$Z = \sum_{\{g_i\}} \prod_{\{i,j,\dots\}} \nu_{d+1}(g_i, g_j, \dots) \quad (12)$$

where $\nu_{d+1}(g_i, g_j, \dots)$ is a function $G^{d+1} \rightarrow U(1)$, which is a cocycle $\nu_{d+1} \in \mathcal{H}^{d+1}[G, U(1)]$. Here the space-time is a complex whose vertices are labeled by i, j, \dots , and $\prod_{\{i,j,\dots\}}$ is the product over all the simplices of the space-time complex. Also $\sum_{\{g_i\}}$ is a sum over all g_i on each vertex.

Now consider the stacking of two SPT states described by cocycle ν'_{d+1} and ν''_{d+1} :

$$Z = \sum_{\{g'_i, g''_i\}} \prod_{\{i,j,\dots\}} \nu'_{d+1}(g'_i, g'_j, \dots) \nu''_{d+1}(g''_i, g''_j, \dots). \quad (13)$$

Such a stacked state has a symmetry $G \times G$ and is a $G \times G$ SPT state.

Now let us add a term to break the $G \times G$ -symmetry to G -symmetry and consider

$$Z = \sum_{\{g'_i, g''_i\}} \prod_{\{i,j,\dots\}} \nu'_{d+1}(g'_i, g'_j, \dots) \nu''_{d+1}(g''_i, g''_j, \dots) \times \prod_i e^{-U|g'_i - g''_i|^2}, \quad (14)$$

where $|g' - g''|$ is an invariant distance between group elements. As we change $U = 0$ to $U = +\infty$, the stacked system changes into the system for an ideal SPT state described by the cocycle $\nu_{d+1}(g_i, g_j, \dots) = \nu'_{d+1}(g_i, g_j, \dots) \nu''_{d+1}(g_i, g_j, \dots)$. If such a deformation does not cause any phase transition, then we can show that the stacking of a ν'_{d+1} -SPT state with a ν''_{d+1} -SPT state give rise to a $\nu_{d+1} = \nu'_{d+1} \nu''_{d+1}$ -SPT state. Thus, the key to show the stacking operation to give rise to the group structure for the SPT states, is to show the theory eqn. (14) has no phase transition as we change $U = 0$ to $U = +\infty$.

To show there is no phase transition, we put the system on a closed space-time with no boundary, say S^{d+1} . In this case, $\prod_{\{i,j,\dots\}} \nu'_{d+1}(g'_i, g'_j, \dots) \nu''_{d+1}(g''_i, g''_j, \dots) = 1$, since ν'_{d+1} and ν''_{d+1} are cocycles. Thus the path integral (14) is reduced to

$$Z = \sum_{\{g'_i, g''_i\}} \prod_i e^{-U|g'_i - g''_i|^2} = \left(|G| \sum_g e^{-U|1-g|^2}\right)^{N_v}, \quad (15)$$

where N_v is the number of vertices and $|G|$ the order of the symmetry group. We see that the free energy density

$$f = - \lim_{N_v \rightarrow \infty} \ln Z / N_v \quad (16)$$

is a smooth function of U for $U \in [0, \infty)$. There is indeed no phase transition.

The above result is highly non trivial from a categorical point of view. Consider two 2+1D bosonic SPT states described by two modular extensions \mathcal{M}' and \mathcal{M}'' of $\text{Rep}(G)$. The natural tensor product $\mathcal{M}' \boxtimes \mathcal{M}''$ is not a modular extension of $\text{Rep}(G)$, but a modular extension of $\text{Rep}(G) \boxtimes \text{Rep}(G) = \text{Rep}(G \times G)$. So, $\mathcal{M}' \boxtimes \mathcal{M}''$ describes a $G \times G$ -SPT state. According to the above discussion, we need to break the $G \times G$ -symmetry down to the G -symmetry to obtain the G -SPT state. Such a symmetry breaking process correspond to the so call “anyon condensation” in category theory. We will discuss such anyon condensation later. The stacking operation $\boxtimes_{\mathcal{E}}$, with such a symmetry breaking process included, is the correct stacking operation that maintains the symmetry G .

C. Mathematical construction of the stacking operation

We have conjectured that a 2+1D topological order with symmetry \mathcal{E} is classified by $(\mathcal{C}, \mathcal{M}_{\mathcal{C}}, c)$, where \mathcal{C} is a UMTC/ \mathcal{E} , $\mathcal{M}_{\mathcal{C}}$ is a modular extension of \mathcal{C} , and c is the central charge. If we have another topological order of the same symmetry \mathcal{E} described by $(\mathcal{C}', \mathcal{M}_{\mathcal{C}'}, c')$, stacking $(\mathcal{C}, \mathcal{M}_{\mathcal{C}}, c)$ and $(\mathcal{C}', \mathcal{M}_{\mathcal{C}'}, c')$ should give a third topological order described by similar data $(\mathcal{C}'', \mathcal{M}_{\mathcal{C}''}, c'')$:

$$(\mathcal{C}, \mathcal{M}_{\mathcal{C}}, c) \boxtimes_{\mathcal{E}} (\mathcal{C}', \mathcal{M}_{\mathcal{C}'}, c') = (\mathcal{C}'', \mathcal{M}_{\mathcal{C}''}, c'') \quad (17)$$

In this section, we will show that such a stacking operation can be defined mathematically. This is an evidence supporting our Conjecture 4. We like to point out that a special case of the above result for $\mathcal{C} = \mathcal{C}' = \mathcal{C}'' = \mathcal{E} = \text{Rep}(G)$ was discussed in section VIB.

To define $\boxtimes_{\mathcal{E}}$ mathematically, first, we like to introduce

Definition 7. A *condensable algebra* in a UBFC \mathcal{C} is a triple (A, m, η) , $A \in \mathcal{C}$, $m : A \otimes A \rightarrow A$, $\eta : \mathbf{1} \rightarrow A$ satisfying

- Associative: $m(\text{id}_A \otimes m) = m(m \otimes \text{id}_A)$

- Unit: $m(\eta \otimes \text{id}_A) = m(\text{id}_A \otimes \eta) = \text{id}_A$
- Isometric: $mm^\dagger = \text{id}_A$
- Connected: $\text{Hom}(\mathbf{1}, A) = \mathbb{C}$
- Commutative: $mc_{A,A} = m$

Physically, such an condensable algebra A is a composite self-bosonic anyon satisfies additional conditions such that one can condense A to obtain another topological phase.

Definition 8. A (left) *module* over a condensable algebra (A, m, η) in \mathcal{C} is a pair (X, ρ) , $X \in \mathcal{C}$, $\rho : A \otimes X \rightarrow X$ satisfying

$$\begin{aligned} \rho(\text{id}_A \otimes \rho) &= \rho(m \otimes \text{id}_M), \\ \rho(\eta \otimes \text{id}_M) &= \text{id}_M. \end{aligned} \quad (18)$$

It is further a *local* module if

$$\rho c_{M, A \otimes A, M} = \rho.$$

We denote the category of left A modules by \mathcal{C}_A . A left module (X, ρ) is turned into a right module via the braiding, $(X, \rho c_{X,A})$ or $(X, \rho c_{A,X}^{-1})$, and thus an A - A bimodule. The relative tensor functor \otimes_A of bimodules then turns \mathcal{C}_A into a fusion category. (This is known as α -induction in subfactor context.) In general there can be two monoidal structures on \mathcal{C}_A , since there are two ways to turn a left module into a bimodule (usually we pick one for definiteness when considering \mathcal{C}_A as a fusion category). The two monoidal structures coincide for the fusion subcategory \mathcal{C}_A^0 of local A modules. Moreover, \mathcal{C}_A^0 inherited the braiding from \mathcal{C} and is also a UBFC. The local modules are nothing but the anyons in the topological phases after condensing A .

Lemma 1 (DMNO³⁶).

$$\dim(\mathcal{C}_A) = \frac{\dim(\mathcal{C})}{\dim(A)}.$$

If \mathcal{C} is a UMTC, then so is \mathcal{C}_A^0 , and

$$\dim(\mathcal{C}_A^0) = \frac{\dim(\mathcal{C})}{\dim(A)^2}.$$

A non-commutative algebra A is also of interest. We have the left center A_l of A , the maximal subalgebra such that $mc_{A_l, A} = m$, and the right center A_r , the maximal subalgebra such that $mc_{A, A_r} = m$. A_l and A_r are commutative subalgebras, thus condensable.

Theorem 5 (FFRS³⁷). There is a canonical equivalence between the categories of local modules over the left and right centers, $\mathcal{C}_{A_l}^0 = \mathcal{C}_{A_r}^0$.

Definition 9. The Drinfeld center $Z(\mathcal{A})$ of a monoidal category \mathcal{A} is a monoidal category with objects as pairs $(X \in \mathcal{A}, b_{X,-})$, where $b_{X,-} : X \otimes - \rightarrow - \otimes X$ are half-braidings that satisfy similar conditions as braidings. Morphisms and the tensor product are naturally defined.

$Z(\mathcal{A})$ is a braided monoidal category. There is a forgetful tensor functor $for_{\mathcal{A}} : Z(\mathcal{A}) \rightarrow \mathcal{A}$, $(X, b_{X,-}) \mapsto X$ that forgets the half-braidings.

Theorem 6 (Müger³⁸). $Z(\mathcal{A})$ is a UMTC if \mathcal{A} is a fusion category and $\dim(Z(\mathcal{A})) = \dim(\mathcal{A})^2$.

Definition 10. Let \mathcal{C} be a braided fusion category and \mathcal{A} a fusion category, a tensor functor $F : \mathcal{C} \rightarrow \mathcal{A}$ is called a central functor if it factorizes through $Z(\mathcal{A})$, i.e., there exists a braided tensor functor $F' : \mathcal{C} \rightarrow Z(\mathcal{A})$ such that $F = F' for_{\mathcal{A}}$.

Lemma 2 (DMNO³⁶). Let $F : \mathcal{C} \rightarrow \mathcal{A}$ be a central functor, and $R : \mathcal{A} \rightarrow \mathcal{C}$ the right adjoint functor of F . Then the object $A = R(\mathbf{1}) \in \mathcal{C}$ has a canonical structure of condensable algebra. \mathcal{C}_A is monoidally equivalent to the image of F , i.e. the smallest fusion subcategory of \mathcal{A} containing $F(\mathcal{C})$.

Example 1. If \mathcal{C} is a UBFC, it is naturally embedded into $Z(\mathcal{C})$, so is $\bar{\mathcal{C}}$. Therefore, $\mathcal{C} \boxtimes \bar{\mathcal{C}} \hookrightarrow Z(\mathcal{C})$. Compose this embedding with the forgetful functor $for_{\mathcal{C}} : Z(\mathcal{C}) \rightarrow \mathcal{C}$ we get a central functor

$$\begin{aligned} \mathcal{C} \boxtimes \bar{\mathcal{C}} &\rightarrow \mathcal{C} \\ X \boxtimes Y &\mapsto X \otimes Y. \end{aligned}$$

Let R be its right adjoint functor, we obtain a condensable algebra $L_{\mathcal{C}} := R(\mathbf{1}) \cong \oplus_i (i \boxtimes \bar{i}) \in \mathcal{C} \boxtimes \bar{\mathcal{C}}$ (\bar{i} denotes the dual object, or anti-particle of i) and $\mathcal{C} = (\mathcal{C} \boxtimes \bar{\mathcal{C}})_{L_{\mathcal{C}}}$, $\dim(L_{\mathcal{C}}) = \dim(\mathcal{C})$. In particular, for a symmetric category \mathcal{E} , $L_{\mathcal{E}}$ is a condensable algebra in $\mathcal{E} \boxtimes \mathcal{E}$, and $\mathcal{E} = (\mathcal{E} \boxtimes \mathcal{E})_{L_{\mathcal{E}}} = (\mathcal{E} \boxtimes \mathcal{E})_{L_{\mathcal{E}}}^0$ for \mathcal{E} is symmetric, all $L_{\mathcal{E}}$ -modules are local. Condensing $L_{\mathcal{E}}$ is nothing but breaking the symmetry from $\mathcal{E} \boxtimes \mathcal{E}$ to \mathcal{E} .

Now, we are ready to define the stacking operation for UMTC/ \mathcal{E} 's as well as their modular extensions.

Definition 11. Let \mathcal{C}, \mathcal{D} be UMTC/ \mathcal{E} 's, and $\mathcal{M}_{\mathcal{C}}, \mathcal{M}_{\mathcal{D}}$ their modular extensions. The stacking is defined by:

$$\mathcal{C} \boxtimes_{\mathcal{E}} \mathcal{D} := (\mathcal{C} \boxtimes \mathcal{D})_{L_{\mathcal{E}}}^0, \quad \mathcal{M}_{\mathcal{C}} \boxtimes_{\mathcal{E}} \mathcal{M}_{\mathcal{D}} := (\mathcal{M}_{\mathcal{C}} \boxtimes \mathcal{M}_{\mathcal{D}})_{L_{\mathcal{E}}}^0$$

Note that in Ref. 39, the tensor product $\boxtimes_{\mathcal{E}}$ for UMTC/ \mathcal{E} 's is defined as $(\mathcal{C} \boxtimes \mathcal{D})_{L_{\mathcal{E}}}$. For UMTC/ \mathcal{E} 's the two definitions coincide $(\mathcal{C} \boxtimes \mathcal{D})_{L_{\mathcal{E}}}^0 = (\mathcal{C} \boxtimes \mathcal{D})_{L_{\mathcal{E}}}$, for $L_{\mathcal{E}}$ lies in the centralizer of $\mathcal{C} \boxtimes \mathcal{D}$ which is $\mathcal{E} \boxtimes \mathcal{E}$. But for the modular extensions we have to take the unusual definition above.

Theorem 7. $\mathcal{C} \boxtimes_{\mathcal{E}} \mathcal{D}$ is a UMTC/ \mathcal{E} , and $\mathcal{M}_{\mathcal{C}} \boxtimes_{\mathcal{E}} \mathcal{M}_{\mathcal{D}}$ is a modular extension of $\mathcal{C} \boxtimes_{\mathcal{E}} \mathcal{D}$.

Proof. The embeddings $\mathcal{E} = (\mathcal{E} \boxtimes \mathcal{E})_{L_{\mathcal{E}}}^0 \hookrightarrow (\mathcal{C} \boxtimes \mathcal{D})_{L_{\mathcal{E}}}^0 = \mathcal{C} \boxtimes_{\mathcal{E}} \mathcal{D} \hookrightarrow (\mathcal{M}_{\mathcal{C}} \boxtimes \mathcal{M}_{\mathcal{D}})_{L_{\mathcal{E}}}^0 = \mathcal{M}_{\mathcal{C}} \boxtimes_{\mathcal{E}} \mathcal{M}_{\mathcal{D}}$ are obvious. So $\mathcal{C} \boxtimes_{\mathcal{E}} \mathcal{D}$ is a UBFC over \mathcal{E} . Also

$$\dim(\mathcal{C} \boxtimes_{\mathcal{E}} \mathcal{D}) = \frac{\dim(\mathcal{C} \boxtimes \mathcal{D})}{\dim(L_{\mathcal{E}})} = \frac{\dim(\mathcal{C}) \dim(\mathcal{D})}{\dim(\mathcal{E})},$$

and $\mathcal{M}_{\mathcal{C}} \boxtimes_{\mathcal{E}} \mathcal{M}_{\mathcal{D}}$ is a UMTC,

$$\dim(\mathcal{M}_{\mathcal{C}} \boxtimes_{\mathcal{E}} \mathcal{M}_{\mathcal{D}}) = \frac{\dim(\mathcal{M}_{\mathcal{C}} \boxtimes \mathcal{M}_{\mathcal{D}})}{\dim(L_{\mathcal{E}})^2} = \dim(\mathcal{C}) \dim(\mathcal{D}).$$

Thus, $\mathcal{M}_{\mathcal{C}} \boxtimes_{\mathcal{E}} \mathcal{M}_{\mathcal{D}}$ is a modular extension of $\mathcal{C} \boxtimes_{\mathcal{E}} \mathcal{D}$. \square

Take $\mathcal{D} = \mathcal{E}$. Note that $\mathcal{C} \boxtimes_{\mathcal{E}} \mathcal{E} = \mathcal{C}$. Therefore, for any modular extension $\mathcal{M}_{\mathcal{E}}$ of \mathcal{E} , $\mathcal{M}_{\mathcal{C}} \boxtimes_{\mathcal{E}} \mathcal{M}_{\mathcal{E}}$ is still a modular extension of \mathcal{C} . In the following we want to show the inverse, that one can extract the “difference”, a modular extension of \mathcal{E} , between two modular extensions of \mathcal{C} .

Lemma 3. We have $(\mathcal{C} \boxtimes \bar{\mathcal{C}})_{L_{\mathcal{C}}}^0 = \mathcal{C}_{\mathcal{C}}^{\text{cen}}$.

Proof. $(\mathcal{C} \boxtimes \bar{\mathcal{C}})_{L_{\mathcal{C}}}$ is equivalent to \mathcal{C} (as a fusion category). Moreover, for $X \in \mathcal{C}$ the equivalence gives the free module $L_{\mathcal{C}} \otimes (X \boxtimes \mathbf{1}) \cong L_{\mathcal{C}} \otimes (\mathbf{1} \boxtimes X)$. $L_{\mathcal{C}} \otimes (X \boxtimes \mathbf{1})$ is a local $L_{\mathcal{C}}$ module if and only if $X \boxtimes \mathbf{1}$ centralize $L_{\mathcal{C}}$. This is the same as $X \in \mathcal{C}_{\mathcal{C}}^{\text{cen}}$. Therefore we have $(\mathcal{C} \boxtimes \bar{\mathcal{C}})_{L_{\mathcal{C}}}^0 = \mathcal{C}_{\mathcal{C}}^{\text{cen}}$. \square

Theorem 8. let \mathcal{M} and \mathcal{M}' be two modular extensions of the UMTC/ \mathcal{E} \mathcal{C} . There exists a unique $\mathcal{K} \in \mathcal{M}_{\text{ext}}(\mathcal{E})$ such that $\mathcal{K} \boxtimes_{\mathcal{E}} \mathcal{M} = \mathcal{M}'$. Such \mathcal{K} is given by

$$\mathcal{K} = (\mathcal{M}' \boxtimes \bar{\mathcal{M}})_{L_{\mathcal{C}}}^0.$$

Proof. \mathcal{K} is a modular extension of \mathcal{E} . This follows Lemma 3, that $\mathcal{E} = \mathcal{C}_{\mathcal{C}}^{\text{cen}} = (\mathcal{C} \boxtimes \bar{\mathcal{C}})_{L_{\mathcal{C}}}^0$ is a full subcategory of \mathcal{K} . \mathcal{K} is a UMTC by construction, and $\dim(\mathcal{K}) = \frac{\dim(\mathcal{M}) \dim(\mathcal{M}')}{\dim(L_{\mathcal{C}})^2} = \dim(\mathcal{E})^2$.

To show that $\mathcal{K} = (\mathcal{M}' \boxtimes \bar{\mathcal{M}})_{L_{\mathcal{C}}}$ satisfies $\mathcal{M}' = \mathcal{K} \boxtimes_{\mathcal{E}} \mathcal{M}$, note that $\mathcal{M}' = \mathcal{M}' \boxtimes \text{Vec} = \mathcal{M}' \boxtimes (\bar{\mathcal{M}} \boxtimes \mathcal{M})_{L_{\mathcal{M}}}^0$. It suffices that

$$\begin{aligned} (\mathcal{M}' \boxtimes \bar{\mathcal{M}} \boxtimes \mathcal{M})_{L_{\mathcal{M}}}^0 &= [(\mathcal{M}' \boxtimes \bar{\mathcal{M}})_{L_{\mathcal{C}}}^0 \boxtimes \mathcal{M}]_{L_{\mathcal{E}}}^0 \\ &= (\mathcal{M}' \boxtimes \bar{\mathcal{M}} \boxtimes \mathcal{M})_{(L_{\mathcal{C}} \boxtimes \mathbf{1}) \otimes (\mathbf{1} \boxtimes L_{\mathcal{E}})}. \end{aligned}$$

This follows that $\mathbf{1} \boxtimes L_{\mathcal{M}}$ and $(L_{\mathcal{C}} \boxtimes \mathbf{1}) \otimes (\mathbf{1} \boxtimes L_{\mathcal{E}})$ are left and right centers of the algebra $(L_{\mathcal{C}} \boxtimes \mathbf{1}) \otimes (\mathbf{1} \boxtimes L_{\mathcal{M}})$.

If $\mathcal{M}' = \mathcal{K} \boxtimes_{\mathcal{E}} \mathcal{M} = (\mathcal{K} \boxtimes \mathcal{M})_{L_{\mathcal{E}}}^0$, then

$$\begin{aligned} \mathcal{K} &= (\mathcal{K} \boxtimes \mathcal{M} \boxtimes \bar{\mathcal{M}})_{L_{\mathcal{M}}}^0 = (\mathcal{K} \boxtimes \mathcal{M} \boxtimes \bar{\mathcal{M}})_{(L_{\mathcal{E}} \boxtimes \mathbf{1}) \otimes (\mathbf{1} \boxtimes L_{\mathcal{C}})}^0 \\ &= [(\mathcal{K} \boxtimes_{\mathcal{E}} \mathcal{M}) \boxtimes \bar{\mathcal{M}}]_{L_{\mathcal{C}}}^0 = (\mathcal{M}' \boxtimes \bar{\mathcal{M}})_{L_{\mathcal{C}}}^0. \end{aligned}$$

It is similar here that $\mathbf{1} \boxtimes L_{\mathcal{M}}$ and $(L_{\mathcal{E}} \boxtimes \mathbf{1}) \otimes (\mathbf{1} \boxtimes L_{\mathcal{C}})$ are the left and right centers of the algebra $(L_{\mathcal{E}} \boxtimes \mathbf{1}) \otimes (\mathbf{1} \boxtimes L_{\mathcal{M}})$. This proves the uniqueness of \mathcal{K} . \square

Let us list several consequences of Theorem 8.

Theorem 9. $\mathcal{M}_{\text{ext}}(\mathcal{E})$ forms a finite abelian group.

Proof. Firstly, there exists at least one modular extension of a symmetric fusion category \mathcal{E} , the Drinfeld center $Z(\mathcal{E})$. So the set $\mathcal{M}_{\text{ext}}(\mathcal{E})$ is not empty. The multiplication is given by the stacking $\boxtimes_{\mathcal{E}}$. It is easy to verify

that the stacking $\boxtimes_{\mathcal{E}}$ for modular extensions is associative and commutative. To show that they form a group we only need to find out the identity and inverse. In this case $\mathcal{K} = (\mathcal{M}' \boxtimes_{\mathcal{E}} \overline{\mathcal{M}})_{L_{\mathcal{E}}}^0 = \mathcal{M}' \boxtimes_{\mathcal{E}} \overline{\mathcal{M}}$, Theorem 8 becomes $\mathcal{M}' \boxtimes_{\mathcal{E}} \overline{\mathcal{M}} \boxtimes_{\mathcal{E}} \mathcal{M} = \mathcal{M}'$, for any modular extensions $\mathcal{M}, \mathcal{M}'$ of \mathcal{E} . Thus, $\overline{\mathcal{M}} \boxtimes_{\mathcal{E}} \mathcal{M}' = \overline{\mathcal{M}} \boxtimes_{\mathcal{E}} \mathcal{M}' \boxtimes_{\mathcal{E}} \overline{\mathcal{M}} \boxtimes_{\mathcal{E}} \mathcal{M} = \overline{\mathcal{M}} \boxtimes_{\mathcal{E}} \mathcal{M}$, i.e. $\overline{\mathcal{M}} \boxtimes_{\mathcal{E}} \mathcal{M}$, is the same category for any extension \mathcal{M} , which turns out to be $Z(\mathcal{E})$. It is exactly the identity element. It is then obvious that the inverse of \mathcal{M} is $\overline{\mathcal{M}}$. The finiteness follows from Ref. 32. \square

Example 2. For bosonic case we find that $\mathcal{M}_{ext}(\text{Rep}(G)) = H^3(G, U(1))$, which is discussed in more detail in the next subsection. For fermionic case a general group cohomological classification is still lacking. We know some simple ones such as $\mathcal{M}_{ext}(\text{sRep}(\mathbb{Z}_2^f)) = \mathbb{Z}_{16}$, which agrees with Kitaev's 16-fold way⁹.

Theorem 10. For a UMTC/ \mathcal{E} \mathcal{C} , if the modular extensions exist, $\mathcal{M}_{ext}(\mathcal{C})$ form a $\mathcal{M}_{ext}(\mathcal{E})$ -torsor. In particular, $|\mathcal{M}_{ext}(\mathcal{C})| = |\mathcal{M}_{ext}(\mathcal{E})|$.

Proof. The action is given by the stacking $\boxtimes_{\mathcal{E}}$. For any two extensions $\mathcal{M}, \mathcal{M}'$, there is a unique extension \mathcal{K} of \mathcal{E} , such that $\mathcal{M} \boxtimes_{\mathcal{E}} \mathcal{K} = \mathcal{M}'$. To see $Z(\mathcal{E})$ acts trivially, note that $\mathcal{M}' \boxtimes_{\mathcal{E}} Z(\mathcal{E}) = \mathcal{M} \boxtimes_{\mathcal{E}} \mathcal{K} \boxtimes_{\mathcal{E}} Z(\mathcal{E}) = \mathcal{M} \boxtimes_{\mathcal{E}} \mathcal{K} = \mathcal{M}'$ holds for any \mathcal{M}' . Due to uniqueness we also know that only $Z(\mathcal{E})$ acts trivially. Thus, the action is free and transitive. \square

This means that for any modular extension of \mathcal{C} , stacking with a nontrivial modular extensions of \mathcal{E} , one always obtains a different modular extension of \mathcal{C} ; on the other hand, starting with a particular modular extension of \mathcal{C} , all the other modular extensions can be generated by staking modular extensions of \mathcal{E} (in other words, there is only one orbit). However, in general, there is no preferred choice of the starting modular extension, unless \mathcal{C} is the form $\mathcal{C}_0 \boxtimes_{\mathcal{E}}$ where \mathcal{C}_0 is a UMTC.

D. Modular extensions of $\text{Rep}(G)$

We set $\mathcal{E} = \text{Rep}(G)$ throughout this subsection. Let $(\mathcal{M}, \iota_{\mathcal{M}})$ be a modular extension of $\text{Rep}(G)$. $\iota_{\mathcal{M}}$ is the embedding $\iota_{\mathcal{M}} : \mathcal{E} \hookrightarrow \mathcal{M}$ that we need to consider explicitly in this subsection. The algebra $A = \text{Fun}(G)$ is a condensable algebra in $\text{Rep}(G)$ and also a condensable algebra in \mathcal{M} . Moreover, A is a Lagrangian algebra in \mathcal{M} because $(\dim A)^2 = |G|^2 = (\dim \text{Rep}(G))^2 = \dim \mathcal{M}$. Therefore, $\mathcal{M} \simeq Z(\mathcal{M}_A)$, where \mathcal{M}_A is the category of right A -modules in \mathcal{M} . In other words, \mathcal{M} describes the bulk excitations in a 2+1D topological phase with a gapped boundary (see Fig. 3). Moreover, the fusion category \mathcal{M}_A is pointed and equipped with a canonical fully faithful G -grading³³, which means that

$$\mathcal{M}_A = \bigoplus_{g \in G} (\mathcal{M}_A)_g, \quad (\mathcal{M}_A)_g \simeq \text{Vec}, \quad \forall g \in G,$$




FIG. 3. Consider a physical situation in which the excitations in the 2 + 1D bulk are given by a modular extension \mathcal{M} of $\text{Rep}(G)$, and those on the gapped boundary by the UFC \mathcal{M}_A . Consider a simple particle $e \in \text{Rep}(G)$ in the bulk moving toward the boundary. The bulk-to-boundary map is given by the central functor $- \otimes A : \mathcal{M} \rightarrow \mathcal{M}_A$, which restricted to $\text{Rep}(G)$ is nothing but the forgetful functor $F : \text{Rep}(G) \rightarrow \text{Vec}$. Let x be a simple excitation in \mathcal{M}_A sitting next to $F(e)$. We move $F(e)$ along the semicircle γ_1 (defined by the half-braiding), then move along the semicircle γ_2 (defined by the symmetric braiding in the trivial phase Vec).

$$\text{and } \otimes : (\mathcal{M}_A)_g \boxtimes (\mathcal{M}_A)_h \xrightarrow{\sim} (\mathcal{M}_A)_{gh}.$$

Let us recall the construction of this G -grading. The physical meaning of acquiring a G -grading on \mathcal{M}_A after condensing the algebra $A = \text{Fun}(G)$ in \mathcal{M} is depicted in Figure 3. The process in Figure 3 defines the isomorphism $F(e) \otimes_A x \xrightarrow{z_{e,x}} x \otimes_A F(e) = F(e) \otimes_A x$, which further gives a monoidal automorphism $\phi(x) \in \text{Aut}(F) = G$ of the fiber functor $F : \text{Rep}(G) \rightarrow \text{Vec}$.

Since ϕ is an isomorphism, the associator of the monoidal category \mathcal{M}_A determines a unique $\omega_{(\mathcal{M}, \iota_{\mathcal{M}})} \in H^3(G, U(1))$ such that $\mathcal{M}_A \simeq \text{Vec}_G^\omega$ as G -graded fusion categories.

Theorem 11. The map $(\mathcal{M}, \iota_{\mathcal{M}}) \mapsto \omega_{(\mathcal{M}, \iota_{\mathcal{M}})}$ defines a group isomorphism $\mathcal{M}_{ext}(\text{Rep}(G)) \simeq H^3(G, U(1))$. In particular, we have

$$(Z(\text{Vec}_G^{\omega_1}), \iota_{\omega_1}) \boxtimes_{\mathcal{E}} (Z(\text{Vec}_G^{\omega_2}), \iota_{\omega_2}) \simeq (Z(\text{Vec}_G^{\omega_1 + \omega_2}), \iota_{\omega_1 + \omega_2}).$$

For the proof and more related details, see also Ref. 30.

E. Relation to numerical calculations

In Section V we proposed another way to characterise GQIs, using the data $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i; \mathcal{N}_K^{IJ}, \mathcal{S}_I; c)$ which is more friendly in numerical calculations. We would like to investigate how to calculate the stacking operation in terms of these data.

Assuming that \mathcal{C} and \mathcal{C}' can be characterized by data (N_k^{ij}, s_i) and $(N'_k{}^{ij}, s'_i)$. Let $(N_k^{\mathcal{D}, ij}, s_i^{\mathcal{D}})$ be the data that characterizes the stacked UMTC/ \mathcal{E} $\mathcal{D} = \mathcal{C} \boxtimes_{\mathcal{E}} \mathcal{C}'$.

To calculate $(N_k^{\mathcal{D},ij}, s_i^{\mathcal{D}})$, let us first construct

$$N_{kk'}^{ii',jj'} = N_k^{ij} N_{k'}^{i'j'}, \quad s_{ii'} = s_i + s_{i'}. \quad (19)$$

Note that, the above data describes a $\text{UMTC}_{/\mathcal{E} \boxtimes \mathcal{E}} \mathcal{D}' = \mathcal{C} \boxtimes \mathcal{C}'$ (*i.e.* with centralizer $\mathcal{E} \boxtimes \mathcal{E}$), which is not what we want. We need reduce centralizer from $\mathcal{E} \boxtimes \mathcal{E}$ to \mathcal{E} . This is the $G \times G$ to G process and \mathcal{C} - \mathcal{C}' coupling, or condensing the $L_{\mathcal{E}}$ algebra, as discussed above

To do the $\mathcal{E} \boxtimes \mathcal{E}$ to \mathcal{E} reduction (*i.e.* to obtain the real stacking operation $\boxtimes_{\mathcal{E}}$), we can introduce an equivalence relation. Noting that the excitations in $\mathcal{D}' = \mathcal{C} \boxtimes \mathcal{C}'$ are labeled by $ii' = i \boxtimes i'$, the equivalence relation is

$$ii' \sim jj', \quad \text{if } ii' \otimes L_{\mathcal{E}} = jj' \otimes L_{\mathcal{E}}. \quad (20)$$

where $L_{\mathcal{E}} = \oplus_a a \bar{a}$, $a \in \mathcal{E}$. In the simple case of abelian groups, where all the a 's are abelian particles, the equivalence relation reduces to

$$(a \otimes i)i' \sim i(a \otimes i'), \quad \forall i \in \mathcal{C}, i' \in \mathcal{C}', a \in \mathcal{E}. \quad (21)$$

Mathematically, this amounts to consider only the free local $L_{\mathcal{E}}$ modules. The equivalent classes $[ii']$ are then some composite anyons in $\mathcal{D} = \mathcal{C} \boxtimes_{\mathcal{E}} \mathcal{C}'$

$$[ii'] = k \oplus l \oplus \dots, \quad \text{for some } k, l, \dots \in \mathcal{D}. \quad (22)$$

In other words, they form a fusion sub ring of \mathcal{D} . Moreover, the spin of ii' is the same as the direct summands

$$s_{ii'} = s_k^{\mathcal{D}} = s_l^{\mathcal{D}} = \dots \quad (23)$$

Since it is limited to a subset of data of $\text{UMTC}_{/\mathcal{E}}$'s, we can only give these necessary conditions. However, as we already give a large list of GQLs in terms of these data, they are usually enough to pick the resulting $\mathcal{C} \boxtimes_{\mathcal{E}} \mathcal{C}'$ from the list.

VII. HOW TO CALCULATE THE MODULAR EXTENSION OF A $\text{UMTC}_{/\mathcal{E}}$

A. A naive calculation

How do we calculate the modular extension \mathcal{M} of $\text{UMTC}_{/\mathcal{E}} \mathcal{C}$ from the data of \mathcal{C} ? Actually, we do not know how to do that. So here, we will follow a closely related Conjecture 6, and calculate instead $(\mathcal{N}_K^{IJ}, \mathcal{S}_I, c)$ (that fully characterize \mathcal{M}) from the data $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$ (that partially characterize \mathcal{C}). In this section, we will describe such a calculation.

We note that all the simple objects (particles) in \mathcal{C} are contained in \mathcal{M} as simple objects, and \mathcal{M} may contain some extra simple objects. Assume that the particle labels of \mathcal{M} are $\{I, J, \dots\} = \{i, j, \dots, x, y, \dots\}$, where we use i, j, \dots to label the particles in \mathcal{C} and x, y, \dots to label the additional particles (not in \mathcal{C}). Also let us use a, b, \dots to label the simple objects in the centralizer of \mathcal{C} : $\mathcal{E} = \mathcal{C}_{\mathcal{E}}^{\text{cen}}$. Let \mathcal{N}_K^{IJ} , \mathcal{S}_I be the fusion coefficients

and the spins for \mathcal{M} , and N_k^{ij} , s_i be the fusion coefficients and the spins for \mathcal{C} . The idea is to find as many conditions on $(\mathcal{N}_K^{IJ}, \mathcal{S}_I)$ as possible, and use those conditions to solve for $(\mathcal{N}_K^{IJ}, \mathcal{S}_I)$. Since the data $(\mathcal{N}_K^{IJ}, \mathcal{S}_I)$ describe the $\text{UMTC} \mathcal{M}$, they should satisfy all the conditions discussed in Ref. 11. On the other hand, as a modular extension of \mathcal{C} , $(\mathcal{N}_K^{IJ}, \mathcal{S}_I)$ also satisfy some additional conditions. Here, we will discuss those additional conditions.

First, the modular extension \mathcal{M} has a fixed total quantum dimension.

$$\dim(\mathcal{M}) = \dim(\mathcal{E}) \dim(\mathcal{C}). \quad (24)$$

In other words

$$\sum_{I \in \mathcal{M}} d_I^2 = \sum_{a \in \mathcal{E}} d_a^2 \sum_{i \in \mathcal{C}} d_i^2. \quad (25)$$

Physically, the modular extension \mathcal{M} is obtained by “gauging” the symmetry \mathcal{E} in \mathcal{C} (*i.e.* adding the symmetry twists of \mathcal{E}). So the additional particles x, y, \dots correspond to the symmetry twists. Fusing an original particle $i \in \mathcal{C}$ to a symmetry twist $x \notin \mathcal{C}$ still give us a symmetry twist. Thus

$$\mathcal{N}_j^{ix} = \mathcal{N}_j^{xi} = \mathcal{N}_x^{ij} = 0. \quad (26)$$

Therefore, \mathcal{N}_i for $i \in \mathcal{C}$ is block diagonal:

$$\mathcal{N}_i = N_i \oplus \hat{N}_i, \quad (27)$$

where $(N_i)_{jk} = \mathcal{N}_k^{ij} = N_k^{ij}$ and $(\hat{N}_i)_{xy} = \mathcal{N}_x^{iy}$.

If we pick a charge conjugation for the additional particles $x \mapsto \bar{x}$, the conditions for fusion rules reduce to

$$\begin{aligned} \mathcal{N}_y^{ix} &= \mathcal{N}_y^{xi} = \mathcal{N}_i^{\bar{x}y} = \mathcal{N}_{\bar{x}}^{iy}, \\ \sum_{k \in \mathcal{C}} N_k^{ij} \mathcal{N}_y^{kx} &= \sum_{z \notin \mathcal{C}} \mathcal{N}_x^{iz} \mathcal{N}_z^{jy}. \end{aligned} \quad (28)$$

With a choice of charge conjugation, it is enough to construct (or search for) the matrices \hat{N}_i and \mathcal{N}_z^{xy} to determine all the extended fusion rules \mathcal{N}_K^{IJ} .

Besides the general condition (28), there are also some simple constraints on \hat{N}_i that may speed up the numerical search. Firstly, observe that (28) is the same as

$$\hat{N}_i \hat{N}_j = \sum_{k \in \mathcal{C}} N_k^{ij} \hat{N}_k, \quad (29)$$

where $i, j, k \in \mathcal{F}$. This means that \hat{N}_i satisfy the same fusion algebra as N_i , and $N_k^{ij} = \mathcal{N}_k^{ij}$ is the structure constant; therefore, the eigenvalues of \hat{N}_i must be a subset of the eigenvalues of N_i .

Secondly, since $\sum_{y \notin \mathcal{C}} \mathcal{N}_y^{ix} d_y = d_i d_x$, by Perron-Frobenius theorem, we know that d_i is the largest eigenvalue of \hat{N}_i , with eigenvector $v, v_x = d_x$. (d_i is also the largest absolute values of the eigenvalues of \hat{N}_i .) Note that $\hat{N}_i \hat{N}_i = \hat{N}_i \hat{N}_i$, $\hat{N}_i = \hat{N}_i^\dagger$. Thus, d_i^2 is the largest

eigenvalue of the positive semi-definite Hermitian matrix $\tilde{N}_i^\dagger \tilde{N}_i$. For any unit vector v we have $v^\dagger \tilde{N}_i^\dagger \tilde{N}_i v \leq d_i^2$, in particular,

$$(\tilde{N}_i^\dagger \tilde{N}_i)_{xx} = \sum_y (\mathcal{N}_y^{ix})^2 \leq d_i^2. \quad (30)$$

The above result is very helpful to reduce the scope of numerical search.

Once we find the fusion rules, \mathcal{N}_K^{IJ} , we can then use the rational conditions and other conditions to determine the spins \mathcal{S}_I (for details, see Ref. 11). The set of data $(\mathcal{N}_K^{IJ}, \mathcal{S}_I)$ that satisfy all the conditions give us the set of modular extensions.

The above proposed calculation for modular extensions is quite expensive. If the quantum dimensions of the particles in \mathcal{C} are all equal to 1: $d_i = 1$, then there is another much cheaper way to calculate the fusion coefficient \mathcal{N}_K^{IJ} of the modular extension \mathcal{M} . Such an approach is explained in Appendix B. We will also use such an approach in our calculation.

Last, we would like to mention that two sets of data $(\mathcal{N}_K^{IJ}, \mathcal{S}_I)$ and $(\tilde{\mathcal{N}}_K^{IJ}, \tilde{\mathcal{S}}_I)$ describe the same modular extension of \mathcal{C} , if they only differ by a permutation of indices $x \in \mathcal{M}$ but $x \notin \mathcal{C}$. So some times, two sets of data $(\mathcal{N}_K^{IJ}, \mathcal{S}_I)$ and $(\tilde{\mathcal{N}}_K^{IJ}, \tilde{\mathcal{S}}_I)$ can describe different modular extensions, even through they describe the same UMTC. (Two sets of data $(\mathcal{N}_K^{IJ}, \mathcal{S}_I)$ and $(\tilde{\mathcal{N}}_K^{IJ}, \tilde{\mathcal{S}}_I)$ describe the same UMTC, if they are only different by a permutation of indices $I \in \mathcal{M}$.)

Why we use such a permutation in the calculation of modular extensions. (which is the ME-equivalence relation discussed before)? This is because when we considering modular extensions, the particle $x \in \mathcal{M}$ but $x \notin \mathcal{C}$ correspond to symmetry twists. They are extrinsic excitations that do not appear in the finite energy spectrum of the Hamiltonian. While the particle $i \in \mathcal{C}$ are intrinsic excitations that do appear in the finite energy spectrum of the Hamiltonian. So $x \notin \mathcal{C}$ and $i \in \mathcal{C}$ are physically distinct and we do not allow permutations that mix them. Also we should not permute the particles $a \in \mathcal{E}$, because they correspond to symmetries. We should not mix, for example, the Z_2 symmetry of exchange layers and the Z_2 symmetry of 180° spin rotation.

B. The limitations of the naive calculation

Since a UMTC/ \mathcal{E} \mathcal{C} is not modular, the data $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$ may not fully characterize \mathcal{C} . To fully characterize \mathcal{C} , we need to use additional data, such as the F -tensor and the R -tensor^{9,11}.

In this paper, we will not use those additional data. As a result, the data $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$ may correspond to several different UMTC/ \mathcal{E} \mathcal{C} 's. In other words, $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$ is a one-to-many labeling of UMTC/ \mathcal{E} 's.

So in our naive calculation, when we calculate the modular extensions of $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$, we may actually calculate the modular extension of several different \mathcal{C} 's that

TABLE II. The bottom two rows correspond to the two modular extensions of $\text{Rep}(Z_2)$ (denoted by $N_c^{|\Theta|} = 2_0^{\zeta_2^1}$). Thus we have two different trivial topological orders with Z_2 symmetry in 2+1D (i.e. two Z_2 SPT states). We use $N_c^{|\Theta|}$ to label UMTC/ \mathcal{E} 's, where $\Theta = D^{-1} \sum_i e^{2\pi i s_i} d_i^2 = |\Theta| e^{2\pi i c/8}$ and $D^2 = \sum_i d_i^2$.

| $N_c^{ \Theta }$ | D^2 | d_1, d_2, \dots | s_1, s_2, \dots | comment |
|-------------------|-------|-------------------|----------------------------------|-------------------|
| $2_0^{\zeta_2^1}$ | 2 | 1, 1 | 0, 0 | $\text{Rep}(Z_2)$ |
| 4_0^B | 4 | 1, 1, 1, 1 | 0, 0, 0, $\frac{1}{2}$ | Z_2 gauge |
| 4_0^B | 4 | 1, 1, 1, 1 | 0, 0, $\frac{1}{4}, \frac{3}{4}$ | double semion |

TABLE III. The two modular extensions of $N_c^{|\Theta|} = 3_2^{\zeta_2^1} \cdot 3_2^{\zeta_2^2}$ has a centralizer $\text{Rep}(Z_2)$. Thus we have two topological orders with Z_2 symmetry in 2+1D which has only one type of spin-1/3 topological excitations.

| $N_c^{ \Theta }$ | D^2 | d_1, d_2, \dots | s_1, s_2, \dots | comment |
|-------------------|-------|--|---|---|
| $3_2^{\zeta_2^1}$ | 6 | 1, 1, 2 | 0, 0, $\frac{1}{3}$ | $\kappa = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$ |
| 5_2^B | 12 | 1, 1, 2, $\zeta_4^1, \zeta_4^1 = \sqrt{3}$ | 0, 0, $\frac{1}{3}, \frac{1}{8}, \frac{5}{8}$ | $(A_1, 4)$ |
| 5_2^B | 12 | 1, 1, 2, ζ_4^1, ζ_4^1 | 0, 0, $\frac{1}{3}, \frac{3}{8}, \frac{7}{8}$ | |

are described by the same data $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$. But for UMTC/ \mathcal{E} 's that can be fully characterized by the data $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$, our calculation produce the modular extensions of a single \mathcal{C} . For example, the naive calculation can obtain the correct modular extensions of $\mathcal{C} = \text{Rep}(G)$ and $\mathcal{C} = \text{sRep}(G^f)$, when G and G^f are abelian groups, or simple finite groups³⁵.

If the $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$ happen to describe two different UMTC/ \mathcal{E} 's, we find that our naive calculation will produce the modular extensions for both of UMTC/ \mathcal{E} 's (see Section VIII D). So by computing the modular extensions of $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$, we can tell if $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$ corresponds to none, one, two, etc UMTC/ \mathcal{E} 's. This leads to the Conjecture 6 that $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i, \mathcal{N}_K^{IJ}, \mathcal{S}_I; c)$ can fully and one-to-one classify GQLs in 2+1D.

VIII. EXAMPLES OF 2+1D SET ORDERS AND SPT ORDERS

In this section, we will discuss simple examples of UMTC/ \mathcal{E} \mathcal{C} 's, and their modular extensions \mathcal{M} . The triple $(\mathcal{C}, \mathcal{M}, c)$ describe a topologically ordered or SPT phase. A single UMTC/ \mathcal{E} \mathcal{C} only describes the set of bulk topological excitations, which correspond to topologically ordered states up to invertible ones.

However, in this section we will not discuss examples of UMTC/ \mathcal{E} \mathcal{C} . What we really do is to discuss examples of the solutions $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$ (which are not really UMTC/ \mathcal{E} 's, but closely related). We will also discuss the modular extensions $(\mathcal{N}_K^{IJ}, \mathcal{S}_I; c)$ of $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$. $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$ will correspond to

UMTC/ \mathcal{C} if it has modular extensions $(\mathcal{N}_K^{IJ}, \mathcal{S}_I; c)$. This allows us to classify GQLs in terms of the data $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i, \mathcal{N}_K^{IJ}, \mathcal{S}_I; c)$.

A. Z_2 bosonic SPT states

Tables [XXII](#), [XXIII](#), and [XXIV](#) list the solutions $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$ when $(\tilde{N}_c^{ab}, \tilde{s}_a)$ describes a SFC $\text{Rep}(Z_2)$. The table contains all UMTC/ $\text{Rep}(Z_2)$'s but may contain extra fake entries. Physically, they describe possible sets of bulk excitations for Z_2 -SET orders of bosonic systems. The sets of bulk excitations are listed by their quantum dimensions d_i and spins s_i .

For example, let us consider the entry $N_c^{|\Theta|} = 2_0^{\zeta_2^1}$ in Table [XXII](#). Such an entry has a central charge $c = 0$. Also $N = 2$, hence the Z_2 -SET state has two types of bulk excitations both with $d_i = 1$ and $s_i = 0$. Both types of excitations are local excitations; one is the trivial type and the other carries an Z_2 charge.

The first question that we like to ask is that “is such an entry a fake entry, or it corresponds to some Z_2 -symmetric GQL's?” If it corresponds to some Z_2 -symmetric GQL's, how many distinct Z_2 -symmetric GQL phases that it corresponds to? In other word, how many distinct Z_2 -symmetric GQL phases are there, that share the same set of bulk topological excitations described by the entry $2_0^{\zeta_2^1}$?

Both questions can be answered by computing the modular extensions of $2_0^{\zeta_2^1}$ (which is also denoted as $\text{Rep}(Z_2)$). We find that the modular extensions exist, and thus $\text{Rep}(Z_2)$ does correspond to some Z_2 -symmetric GQL's. In fact, one of the Z_2 -symmetric GQL's is the trivial product state with Z_2 symmetry. Other Z_2 -symmetric GQL's are Z_2 SPT states.

After a numerical calculation, we find that there are only two different modular extensions of $\text{Rep}(Z_2)$ (see Table [II](#)). Thus there are two distinct Z_2 -symmetric GQL phases whose bulk excitations are described by the $\text{Rep}(Z_2)$. The first one corresponds to the trivial product states whose modular extension is the Z_2 gauge theory which has four types of particles with $(d_i, s_i) = (1, 0), (1, 0), (1, 0), (1, \frac{1}{2})$. (Gauging the Z_2 symmetry of the trivial product state gives rise to a Z_2 gauge theory.) The second one corresponds to the only non-trivial Z_2 bosonic SPT state in 2+1D, whose modular extension is the double-semion theory which has four types of particles with $(d_i, s_i) = (1, 0), (1, 0), (1, \frac{1}{4}), (1, -\frac{1}{4})$. (Gauging the Z_2 symmetry of the Z_2 -SPT state gives rise to a double-semion theory²¹.) So the Z_2 -SPT phases are classified by \mathbb{Z}_2 , reproducing the group cohomology result^{18–20}. In general, the modular extensions of $\text{Rep}(G)$ correspond to the bosonic SPT states in 2+1D with symmetry G .

TABLE IV. The fusion rule of the $N_c^{|\Theta|} = 3_2^{\zeta_2^1}$ Z_2 -SET order. The particle **1** carries the Z_2 -charge 0, and the particle s carries the Z_2 -charge 1. From the table, we see that $\sigma \otimes \sigma = \mathbf{1} \oplus s \oplus \sigma$.

| | | | |
|-------------------|----------|----------|-------------------------------------|
| s_i | 0 | 0 | $\frac{1}{3}$ |
| d_i | 1 | 1 | 2 |
| $3_2^{\zeta_2^1}$ | 1 | s | σ |
| 1 | 1 | s | σ |
| s | s | 1 | σ |
| σ | σ | σ | $\mathbf{1} \oplus s \oplus \sigma$ |

TABLE V. The fusion rules of the two $N_c^{|\Theta|} = 4_1^{\zeta_2^1}$ Z_2 symmetry enriched topological orders with identical d_i and s_i . We see that one has a $Z_2 \times Z_2$ fusion rule and the other has a Z_4 fusion rule.

| | | | | |
|-------------------|-----------|-----------|---------------|---------------|
| s_i | 0 | 0 | $\frac{1}{4}$ | $\frac{1}{4}$ |
| d_i | 1 | 1 | 1 | 1 |
| $4_1^{\zeta_2^1}$ | 00 | 01 | 10 | 11 |
| 00 | 00 | 01 | 10 | 11 |
| 01 | 01 | 00 | 11 | 10 |
| 10 | 10 | 11 | 00 | 01 |
| 11 | 11 | 10 | 01 | 00 |

| | | | | |
|-------------------|----------|----------|---------------|---------------|
| s_i | 0 | 0 | $\frac{1}{4}$ | $\frac{1}{4}$ |
| d_i | 1 | 1 | 1 | 1 |
| $4_1^{\zeta_2^1}$ | 0 | 2 | 1 | 3 |
| 0 | 0 | 2 | 1 | 3 |
| 2 | 2 | 0 | 3 | 1 |
| 1 | 1 | 3 | 2 | 0 |
| 3 | 3 | 1 | 0 | 2 |

B. Z_2 -SET orders for bosonic systems

The entry $N_c^{|\Theta|} = 3_2^{\zeta_2^1}$ in Table [XXII](#) corresponds to more non-trivial UMTC/ $\text{Rep}(Z_2)$. It describes the bulk excitations of Z_2 -SET orders which has only one type of non-trivial topological excitation (with quantum dimension $d = 2$ and spin $s = 1/3$, see Table [IV](#)). The other two types of excitations are local excitations with Z_2 -charge 0 and 1. We find that $3_2^{\zeta_2^1}$ has modular extensions and hence is not a fake entry.

To see how many SET orders that have such set of bulk excitations, we need to compute how many modular extensions are there for $3_2^{\zeta_2^1}$. We find that $3_2^{\zeta_2^1}$ has two modular extensions (see Table [III](#)). Thus there are two Z_2 -SET orders with the above mentioned bulk excitations. It is not an accident that the number of Z_2 -SET orders with the same set of bulk excitations is the same as the number of Z_2 SPT states. This is because the different Z_2 -SET orders with a fixed set of bulk excitations are generated by stacking with Z_2 SPT states.

We would like to point out that for any G -SET state, if we break the symmetry, the G -SET state will reduce to a topologically ordered state described by a UMTC. In fact, the different G -SET states described by the same UMTC/ \mathcal{C} (*i.e.* with the same set of bulk excitations) will reduce to the same topologically ordered state (*i.e.* the same UMTC). In Appendix [D](#), we discussed such a symmetry breaking process and how to compute UMTC from UMTC/ \mathcal{C} . We found that the two Z_2 -SET

orders from $3_2^{\zeta_2^1}$ reduce to an abelian topological order described by a K -matrix $\begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$. This is indicated by $\text{SB}:K = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$ in the comment column of Table

XXII. In other place, we use $\text{SB}:N_c^B$ or $\text{SB}:N_c^F(a_b)$ to indicate the reduced topological order after the symmetry breaking (for bosonic or fermionic cases). (The topological orders described by N_c^B or $N_c^F(a_b)$ are given by the tables in Ref. 11 or Ref. 14.)

As we have mentioned, there are two Z_2 -SET orders with the same bulk excitations. But how to realize those Z_2 -SET orders? We find that one of the Z_2 -SET orders is the double layer FQH state with K -matrix $\begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$ (same as the reduced topological order after symmetry breaking), where the Z_2 symmetry is the layer-exchange symmetry. The quasiparticles are labeled by the l -vectors $l = \begin{pmatrix} l_1 \\ l_2 \end{pmatrix}$. The two non-trivial quasiparticles are given by

$$l = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad (31)$$

whose spins are all equal to $\frac{1}{3}$.

Since the layer-exchange Z_2 symmetry exchanges l_1 and l_2 , we see that the two excitations $\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ always have the same energy. Despite the Z_2 symmetry has no 2-dim irreducible representations, the above spin-1/3 topological excitations has an exact two-fold degeneracy due to the Z_2 layer-exchange symmetry. This effect is an interplay between the long-range entanglement and the symmetry: *degeneracy in excitations may not always arise from high dimensional irreducible representations of the symmetry*.

Such two degenerate excitations are viewed as one type of topological excitations with quantum dimension $d = 2$ (for the two-fold degeneracy) and spin $s = \frac{1}{3}$ (see Table **XXII**). The Z_2 symmetry twist in such a double-layer state carry a non-abelian statistics with quantum dimension $d = \sqrt{3}$. In fact, there are two such Z_2 symmetry twists whose spin differ by $1/2$.

The other Z_2 -SET order can be viewed as the above double layer FQH state $K = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$ stacked with a Z_2 SPT state.

C. Two other Z_2 -SET orders for bosonic systems

The fourth and fifth entries in Table **XXII** describe the bulk excitations of two other Z_2 -SET orders. Those

TABLE VI. The four modular extensions of $N_c^{|\Theta|} = 5_0^{\zeta_2^1}$ with $Z_2 \times Z_2$ fusion. $5_0^{\zeta_2^1}$ has a centralizer $\text{Rep}(Z_2)$. The first pair and the second pair turns out to be equivalent.

| $N_c^{ \Theta }$ | D^2 | d_1, d_2, \dots | s_1, s_2, \dots | comment |
|-------------------|-------|-------------------------------------|---|----------------------------------|
| $5_0^{\zeta_2^1}$ | 8 | $1 \times 4, 2$ | $0, 0, \frac{1}{2}, \frac{1}{2}, 0$ | |
| 9_0^B | 16 | $1 \times 4, 2, \zeta_2^1 \times 4$ | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, \frac{15}{16}, \frac{1}{16}, \frac{7}{16}, \frac{9}{16}$ | $3_{-1/2}^B \boxtimes 3_{1/2}^B$ |
| 9_0^B | 16 | $1 \times 4, 2, \zeta_2^1 \times 4$ | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, \frac{3}{16}, \frac{13}{16}, \frac{11}{16}, \frac{5}{16}$ | $3_{3/2}^B \boxtimes 3_{-3/2}^B$ |
| 9_0^B | 16 | $1 \times 4, 2, \zeta_2^1 \times 4$ | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, \frac{1}{16}, \frac{15}{16}, \frac{9}{16}, \frac{7}{16}$ | $3_{1/2}^B \boxtimes 3_{-1/2}^B$ |
| 9_0^B | 16 | $1 \times 4, 2, \zeta_2^1 \times 4$ | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, \frac{13}{16}, \frac{3}{16}, \frac{5}{16}, \frac{11}{16}$ | $3_{-3/2}^B \boxtimes 3_{3/2}^B$ |

TABLE VII. The four modular extensions of $N_c^{|\Theta|} = 5_1^{\zeta_2^1}$ with $Z_2 \times Z_2$ fusion. $5_1^{\zeta_2^1}$ has a centralizer $\text{Rep}(Z_2)$.

| $N_c^{ \Theta }$ | D^2 | d_1, d_2, \dots | s_1, s_2, \dots | comment |
|-------------------|-------|-------------------------------------|---|----------------------------------|
| $5_1^{\zeta_2^1}$ | 8 | $1 \times 4, 2$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}$ | |
| 9_1^B | 16 | $1 \times 4, 2, \zeta_2^1 \times 4$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}, \frac{1}{16}, \frac{1}{16}, \frac{9}{16}, \frac{9}{16}$ | $3_{1/2}^B \boxtimes 3_{1/2}^B$ |
| 9_1^B | 16 | $1 \times 4, 2, \zeta_2^1 \times 4$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}, \frac{13}{16}, \frac{13}{16}, \frac{5}{16}, \frac{5}{16}$ | $3_{-3/2}^B \boxtimes 3_{5/2}^B$ |
| 9_1^B | 16 | $1 \times 4, 2, \zeta_2^1 \times 4$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}, \frac{15}{16}, \frac{3}{16}, \frac{7}{16}, \frac{11}{16}$ | $3_{-1/2}^B \boxtimes 3_{3/2}^B$ |
| 9_1^B | 16 | $1 \times 4, 2, \zeta_2^1 \times 4$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}, \frac{3}{16}, \frac{15}{16}, \frac{11}{16}, \frac{7}{16}$ | $3_{3/2}^B \boxtimes 3_{-1/2}^B$ |

bulk excitations have identical s_i and d_i , but they have different fusion rules N_k^{ij} (see Table **V**).

Both entries have two modular extensions, and correspond to two SET orders. Among the two SET orders for the $Z_2 \times Z_2$ fusion rule, one of them is obtained by stacking a Z_2 neutral $\nu = 1/2$ Laughlin state with a trivial Z_2 product state. The other is obtained by stacking a Z_2 neutral $\nu = 1/2$ Laughlin state with a non-trivial Z_2 SPT state.

The entry with Z_4 fusion rule also correspond to two SET orders. They are obtained by stacking a Z_2 charged $\nu = 1/2$ Laughlin state with a trivial or a non-trivial Z_2 SPT state. Here, *charged* means that the particles forming the $\nu = 1/2$ Laughlin state carry Z_2 -charge 1. In this case, the anyon in the $\nu = 1/2$ Laughlin state carries a fractional Z_2 -charge $1/2$. So the fusion of two such anyons give us a Z_2 -charge 1 excitation instead of a trivial neutral excitation. This leads to the Z_4 fusion rule.

D. The rank $N = 5$ Z_2 -SET orders for bosonic systems

The first and the second entries in Table **XXIII** describe two $N = 5$ UMTC/ $\text{Rep}(Z_2)$'s. They describe two different sets of bulk excitations for Z_2 -SET orders. Those bulk excitations have identical s_i and d_i , but they have different fusion rules N_k^{ij} : the 4 $d = 1$ particles have a $Z_2 \times Z_2$ fusion rule for the first entry, and they have a Z_4 fusion rule for the second entry (as indicated by $\text{F}:Z_2 \times Z_2$ or $\text{F}:Z_4$ in the comment column of Table **XXIII**).

TABLE VIII. The first and the third entries in Table VI have different fusion rules, despite they have the same (d_i, s_i) .

| | | | | | | | | | |
|---------|---|---|---------------|---------------|--------------------------------|----------------|----------------|----------------|-----------------|
| s_i | 0 | 0 | $\frac{1}{2}$ | $\frac{1}{2}$ | 0 | $\frac{1}{16}$ | $\frac{7}{16}$ | $\frac{9}{16}$ | $\frac{15}{16}$ |
| d_i | 1 | 1 | 1 | 1 | 2 | ζ_2^1 | ζ_2^1 | ζ_2^1 | ζ_2^1 |
| 9_0^1 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| 1 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| 2 | 2 | 1 | 4 | 3 | 5 | 8 | 9 | 6 | 7 |
| 3 | 3 | 4 | 1 | 2 | 5 | 8 | 7 | 6 | 9 |
| 4 | 4 | 3 | 2 | 1 | 5 | 6 | 9 | 8 | 7 |
| 5 | 5 | 5 | 5 | 5 | $1 \oplus 2 \oplus 3 \oplus 4$ | $7 \oplus 9$ | $6 \oplus 8$ | $7 \oplus 9$ | $6 \oplus 8$ |
| 6 | 6 | 8 | 8 | 6 | $7 \oplus 9$ | $1 \oplus 4$ | 5 | $2 \oplus 3$ | 5 |
| 7 | 7 | 9 | 7 | 9 | $6 \oplus 8$ | 5 | $1 \oplus 3$ | 5 | $2 \oplus 4$ |
| 8 | 8 | 6 | 6 | 8 | $7 \oplus 9$ | $2 \oplus 3$ | 5 | $1 \oplus 4$ | 5 |
| 9 | 9 | 7 | 9 | 7 | $6 \oplus 8$ | 5 | $2 \oplus 4$ | 5 | $1 \oplus 3$ |

| | | | | | | | | | |
|---------|---|---|---------------|---------------|--------------------------------|----------------|----------------|----------------|-----------------|
| s_i | 0 | 0 | $\frac{1}{2}$ | $\frac{1}{2}$ | 0 | $\frac{1}{16}$ | $\frac{7}{16}$ | $\frac{9}{16}$ | $\frac{15}{16}$ |
| d_i | 1 | 1 | 1 | 1 | 2 | ζ_2^1 | ζ_2^1 | ζ_2^1 | ζ_2^1 |
| 9_0^1 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| 1 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| 2 | 2 | 1 | 4 | 3 | 5 | 8 | 9 | 6 | 7 |
| 3 | 3 | 4 | 1 | 2 | 5 | 6 | 9 | 8 | 7 |
| 4 | 4 | 3 | 2 | 1 | 5 | 8 | 7 | 6 | 9 |
| 5 | 5 | 5 | 5 | 5 | $1 \oplus 2 \oplus 3 \oplus 4$ | $7 \oplus 9$ | $6 \oplus 8$ | $7 \oplus 9$ | $6 \oplus 8$ |
| 6 | 6 | 8 | 6 | 8 | $7 \oplus 9$ | $1 \oplus 3$ | 5 | $2 \oplus 4$ | 5 |
| 7 | 7 | 9 | 9 | 7 | $6 \oplus 8$ | 5 | $1 \oplus 4$ | 5 | $2 \oplus 3$ |
| 8 | 8 | 6 | 8 | 6 | $7 \oplus 9$ | $2 \oplus 4$ | 5 | $1 \oplus 3$ | 5 |
| 9 | 9 | 7 | 7 | 9 | $6 \oplus 8$ | 5 | $2 \oplus 3$ | 5 | $1 \oplus 4$ |

1. The first entry in Table XXIII

Let us compute the modular extensions of the first entry (*i.e.* $5_0^{\zeta_2^1}$ with $Z_2 \times Z_2$ fusion). Since the total quantum dimension of the modular extensions is $D^2 = 16$, the modular extensions must have rank $N = 13$ or less (since quantum dimension $d \geq 1$).

Now we would like to show $N = 13$ is not possible. If a modular extension has $N = 13$, then it must have 12 particles (labeled by $a = 1, \dots, 12$) with quantum dimension $d_a = 1$, and one particle (labeled by x) with quantum dimension $d_x = 2$, so that $12 \times 1^2 + 2^2 = D^2 = 16$. In this case, we must have the fusion rule

$$a \otimes x = x, \quad x \otimes x = 1 \oplus 2 \oplus 3 \oplus 4. \quad (32)$$

where $x \otimes x$ is determined by the fusion rule of the $\text{UMTC}/\text{Rep}(Z_2)$. The above determines the fusion matrix N_x defined as $(N_x)_{ij} \equiv N_j^{xi}$. The largest eigenvalue of N_x should be 2, the quantum dimension of x . Indeed, we find that the largest eigenvalue of N_x is 2. But we also require that N_x can be diagonalized by a unitary matrix (which happens to be the S -matrix). N_x fails such a test. So N cannot be 13.

N also cannot be 12. If $N = 12$, then the modular extension will have 10 particles (labeled by $a = 1, \dots, 10$)

TABLE IX. The third and the fourth entries in Table VII have different fusion rules, despite they have the same (d_i, s_i) .

| | | | | | | | | | |
|---------|---|---|---------------|---------------|--------------------------------|----------------|----------------|-----------------|-----------------|
| s_i | 0 | 0 | $\frac{1}{2}$ | $\frac{1}{2}$ | $\frac{1}{8}$ | $\frac{3}{16}$ | $\frac{7}{16}$ | $\frac{11}{16}$ | $\frac{15}{16}$ |
| d_i | 1 | 1 | 1 | 1 | 2 | ζ_2^1 | ζ_2^1 | ζ_2^1 | ζ_2^1 |
| 9_1^1 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| 1 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| 2 | 2 | 1 | 4 | 3 | 5 | 8 | 9 | 6 | 7 |
| 3 | 3 | 4 | 1 | 2 | 5 | 8 | 7 | 6 | 9 |
| 4 | 4 | 3 | 2 | 1 | 5 | 6 | 9 | 8 | 7 |
| 5 | 5 | 5 | 5 | 5 | $1 \oplus 2 \oplus 3 \oplus 4$ | $7 \oplus 9$ | $6 \oplus 8$ | $7 \oplus 9$ | $6 \oplus 8$ |
| 6 | 6 | 8 | 8 | 6 | $7 \oplus 9$ | $1 \oplus 4$ | 5 | $2 \oplus 3$ | 5 |
| 7 | 7 | 9 | 7 | 9 | $6 \oplus 8$ | 5 | $1 \oplus 3$ | 5 | $2 \oplus 4$ |
| 8 | 8 | 6 | 6 | 8 | $7 \oplus 9$ | $2 \oplus 3$ | 5 | $1 \oplus 4$ | 5 |
| 9 | 9 | 7 | 9 | 7 | $6 \oplus 8$ | 5 | $2 \oplus 4$ | 5 | $1 \oplus 3$ |

| | | | | | | | | | |
|---------|---|---|---------------|---------------|--------------------------------|----------------|----------------|-----------------|-----------------|
| s_i | 0 | 0 | $\frac{1}{2}$ | $\frac{1}{2}$ | $\frac{1}{8}$ | $\frac{3}{16}$ | $\frac{7}{16}$ | $\frac{11}{16}$ | $\frac{15}{16}$ |
| d_i | 1 | 1 | 1 | 1 | 2 | ζ_2^1 | ζ_2^1 | ζ_2^1 | ζ_2^1 |
| 9_1^1 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| 1 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| 2 | 2 | 1 | 4 | 3 | 5 | 8 | 9 | 6 | 7 |
| 3 | 3 | 4 | 1 | 2 | 5 | 6 | 9 | 8 | 7 |
| 4 | 4 | 3 | 2 | 1 | 5 | 8 | 7 | 6 | 9 |
| 5 | 5 | 5 | 5 | 5 | $1 \oplus 2 \oplus 3 \oplus 4$ | $7 \oplus 9$ | $6 \oplus 8$ | $7 \oplus 9$ | $6 \oplus 8$ |
| 6 | 6 | 8 | 6 | 8 | $7 \oplus 9$ | $1 \oplus 3$ | 5 | $2 \oplus 4$ | 5 |
| 7 | 7 | 9 | 9 | 7 | $6 \oplus 8$ | 5 | $1 \oplus 4$ | 5 | $2 \oplus 3$ |
| 8 | 8 | 6 | 8 | 6 | $7 \oplus 9$ | $2 \oplus 4$ | 5 | $1 \oplus 3$ | 5 |
| 9 | 9 | 7 | 7 | 9 | $6 \oplus 8$ | 5 | $2 \oplus 3$ | 5 | $1 \oplus 4$ |

with quantum dimension $d_a = 1$, one particle (labeled by x) with quantum dimension $d_x = 2$, and one particle (labeled by y) with quantum dimension $d_y = \sqrt{2}$. The fusion of 10 $d_a = 1$ particles is described by an abelian group Z_{10} or $Z_2 \times Z_5$. None of them contain $Z_2 \times Z_2$ as subgroup. Thus $N = 12$ is incompatible with the $Z_2 \times Z_2$ fusion of the first four $d_a = 1$ particles.

We searched the modular extensions with N up to 11. We find four $N = 9$ modular extensions (see Table VI), and thus the first entry corresponds to valid Z_2 -SET states.

In fact one of the Z_2 -SET states is the Z_2 gauge theory with a Z_2 global symmetry, where the Z_2 symmetry action exchange the Z_2 -charge e and the Z_2 -vortex m . The degenerate e and m give rise to the $(d, s) = (2, 0)$ particle (the fifth particle in the table). The bound state of e and m is a fermion f . It may carry the Z_2 -charge 0 or 1, which correspond to the third and the fourth particle with $(d, s) = (1, 1/2)$ in the table.

However, from the discussion in the last few sections, we know that a $\text{UMTC}/\text{Rep}(Z_2)$ always has 2 modular extensions, corresponding to the 2 bosonic Z_2 -SPT states in 2+1D. This seems contradictory with the above result that the Z_2 -SET state, $5_0^{\zeta_2^1}$ with $Z_2 \times Z_2$ fusion, has four different modular extensions.

TABLE X. The fusion rules of the first and the second entries in Table VII.

| s_i | 0 | 0 | $\frac{1}{2}$ | $\frac{1}{2}$ | $\frac{1}{8}$ | $\frac{1}{16}$ | $\frac{1}{16}$ | $\frac{9}{16}$ | $\frac{9}{16}$ |
|---------|---|---|---------------|---------------|--------------------------------|----------------|----------------|----------------|----------------|
| d_i | 1 | 1 | 1 | 1 | 2 | ζ_2^1 | ζ_2^1 | ζ_2^1 | ζ_2^1 |
| 9_1^1 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| 1 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| 2 | 2 | 1 | 4 | 3 | 5 | 8 | 9 | 6 | 7 |
| 3 | 3 | 4 | 1 | 2 | 5 | 8 | 7 | 6 | 9 |
| 4 | 4 | 3 | 2 | 1 | 5 | 6 | 9 | 8 | 7 |
| 5 | 5 | 5 | 5 | 5 | $1 \oplus 2 \oplus 3 \oplus 4$ | $7 \oplus 9$ | $6 \oplus 8$ | $7 \oplus 9$ | $6 \oplus 8$ |
| 6 | 6 | 8 | 8 | 6 | $7 \oplus 9$ | $1 \oplus 4$ | 5 | $2 \oplus 3$ | 5 |
| 7 | 7 | 9 | 7 | 9 | $6 \oplus 8$ | 5 | $1 \oplus 3$ | 5 | $2 \oplus 4$ |
| 8 | 8 | 6 | 6 | 8 | $7 \oplus 9$ | $2 \oplus 3$ | 5 | $1 \oplus 4$ | 5 |
| 9 | 9 | 7 | 9 | 7 | $6 \oplus 8$ | 5 | $2 \oplus 4$ | 5 | $1 \oplus 3$ |

| s_i | 0 | 0 | $\frac{1}{2}$ | $\frac{1}{2}$ | $\frac{1}{8}$ | $\frac{5}{16}$ | $\frac{5}{16}$ | $\frac{13}{16}$ | $\frac{13}{16}$ |
|---------|---|---|---------------|---------------|--------------------------------|----------------|----------------|-----------------|-----------------|
| d_i | 1 | 1 | 1 | 1 | 2 | ζ_2^1 | ζ_2^1 | ζ_2^1 | ζ_2^1 |
| 9_1^1 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| 1 | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 |
| 2 | 2 | 1 | 4 | 3 | 5 | 8 | 9 | 6 | 7 |
| 3 | 3 | 4 | 1 | 2 | 5 | 8 | 7 | 6 | 9 |
| 4 | 4 | 3 | 2 | 1 | 5 | 6 | 9 | 8 | 7 |
| 5 | 5 | 5 | 5 | 5 | $1 \oplus 2 \oplus 3 \oplus 4$ | $7 \oplus 9$ | $6 \oplus 8$ | $7 \oplus 9$ | $6 \oplus 8$ |
| 6 | 6 | 8 | 8 | 6 | $7 \oplus 9$ | $1 \oplus 4$ | 5 | $2 \oplus 3$ | 5 |
| 7 | 7 | 9 | 7 | 9 | $6 \oplus 8$ | 5 | $1 \oplus 3$ | 5 | $2 \oplus 4$ |
| 8 | 8 | 6 | 6 | 8 | $7 \oplus 9$ | $2 \oplus 3$ | 5 | $1 \oplus 4$ | 5 |
| 9 | 9 | 7 | 9 | 7 | $6 \oplus 8$ | 5 | $2 \oplus 4$ | 5 | $1 \oplus 3$ |

In fact, there is no contradiction. Here, we only use (N_k^{ij}, s_i) to label different entries. However, a UMTC/\mathcal{E} is fully characterized by (N_k^{ij}, s_i) plus the F -tensors and the R -tensors. To see this point, we note that the Ising-like UMTC $N_c^B = 3_{m/2}^B$, $m = 1, 3, \dots, 15$ (with central charge $c = m/2$) has three particles: 1, f with $(d_f, s_f) = (1, 1/2)$, and σ with $(d_\sigma, s_\sigma) = (\sqrt{2}, m/16)$. Its R -tensor is given by⁹

$$\begin{aligned} R_1^{ff} &= -1, & R_\sigma^{f\sigma} &= R_\sigma^{f\sigma} = -i^m, \\ R_1^{\sigma\sigma} &= (-1)^{\frac{m^2-1}{8}} e^{-i\frac{\pi}{8}m}, & R_f^{\sigma\sigma} &= (-1)^{\frac{m^2-1}{8}} e^{i\frac{3\pi}{8}m}, \end{aligned} \quad (33)$$

and some components of the F -tensor are given by

$$F_{f;1}^{f\sigma\sigma;\sigma} = F_{f;1}^{\sigma\sigma f;\sigma} = 1. \quad (34)$$

The values of R_σ^{ff} and $R_\sigma^{f\sigma}$ are not gauge invariant. But if we fix the values of the F -tensor to be the ones given above, this will fix the gauge, and we can treat R_σ^{ff} and $R_\sigma^{f\sigma}$ as if they are gauge invariant quantities.

If we stack $N_c^B = 3_{m/2}^B$ and $N_{c'}^B = 3_{m'/2}^B$ together, the induced UMTC $3_{m/2}^B \boxtimes 3_{m'/2}^B$ contains particles $\mathbf{1} = (1, 1)$, $\mathbf{2} = (f, f')$, $\mathbf{3} = (f, 1)$, $\mathbf{4} = (1, f')$, $\mathbf{5} = (\sigma, \sigma')$. Those 5 particles are closed under the fusion, and correspond to

the 5 particles in $\text{UMTC}/\text{Rep}(Z_2)$ $5_{m+m'}^{\zeta_2^1}$. We note that some components of the R -tensor of $3_{m/2}^B \boxtimes 3_{m'/2}^B$ are given by

$$\begin{aligned} R_{(\sigma,\sigma')}^{(f,1),(\sigma,\sigma')} &= R_{(\sigma,\sigma')}^{(\sigma,\sigma'),(f,1)} = -i^m, \\ R_{(\sigma,\sigma')}^{(1,f'),(\sigma,\sigma')} &= R_{(\sigma,\sigma')}^{(\sigma,\sigma'),(1,f')} = -i^{m'}. \end{aligned} \quad (35)$$

Taking $(m, m') = (-1, 1)$ and $(1, -1)$, it is clear the $3_{-1/2}^B \boxtimes 3_{1/2}^B$ and $3_{1/2}^B \boxtimes 3_{-1/2}^B$ give rise to two different R -tensors that have identical (N_k^{ij}, s_i) . So the first entry in Table XXIII (*i.e.* $5_0^{\zeta_2^1}$ with $Z_2 \times Z_2$ fusion) split into two different entries if we include the R -tensors. Each give rise to two modular extensions, and this is why we got four modular extensions. In Table VI, the first two modular extensions have the same (N_k^{ij}, s_i) , F -tensor and R -tensors when restricted to the first 5 particles. The second pair of modular extensions also have the same (N_k^{ij}, s_i) , F -tensor and R -tensor when restricted to the first 5 particles, but their R -tensor is different from that of the first pair. However, note that under the exchange of the two fermions, the R -tensor of the first pair becomes that of the second pair.

We like to stress that Table VI is obtained using the ME-equivalence relation, *i.e.* the different entries are different under the ME-equivalence relation (see Section V). We see that for each fixed $\text{UMTC}/\text{Rep}(Z_2)$ (*i.e.* for each fixed set of (N_k^{ij}, s_i) , F -tensor and R -tensor), there are two modular extensions, which agrees with our general result for modular extensions. However, if we ignore F -tensor and R -tensor, then for each fixed set of (N_k^{ij}, s_i) , we get four modular extensions. This is because (N_k^{ij}, s_i) is only a partial description of a $\text{UMTC}/\text{Rep}(Z_2)$, and as discussed above, in this case there are two ways to assign F -tensor and R -tensor to them. This is why each fixed (N_k^{ij}, s_i) has four modular extensions, while each fixed (N_k^{ij}, s_i, F, R) has only two modular extensions.

On the other hand, under the TO-equivalence relation (see Section V), the two ways to assign F -tensor and R -tensor are actually equivalent (related by exchanging the two fermions), and the first entry in Table XXIII corresponds to only one $\text{UMTC}/\text{Rep}(Z_2)$. Thus, the first entry is equivalent to the third entry, and the second entry is equivalent to the fourth entry in Table VI. So the four entries of Table VI in fact represent only two distinct Z_2 -SET orders.

One of the two Z_2 -SET orders have been studied extensively. It corresponds to Z_2 gauge theory with a Z_2 global symmetry that exchanges the Z_2 -gauge-charge e and the Z_2 -gauge-vortex m ^{26,27}.

2. The second entry in Table XXIII

Next, we compute the modular extensions of the second entry in Table XXIII (*i.e.* $5_0^{\zeta_2^1}$ with Z_4 fusion). Again,

we can use the same argument to show that modular extensions of rank 12 and above do not exist. We searched the modular extensions with N up to 11, and find that there is no modular extensions. So the second entry is not realizable and does not correspond to any valid bosonic Z_2 -SET in 2+1D. This is indicated by NR in the comment column of Table XXIII.

Naively, the (none existing) state from the second entry is very similar to that from the first entry. It is also a Z_2 gauge theory with a Z_2 global symmetry that exchange e and m . However, for the second entry, the f particles (the third and the fourth particles) are assigned fraction Z_2 -charge of $\pm 1/2$. This leads to the Z_4 fusion rule. Our result implies that such an assignment is not realizable (or is illegal). It turns out that all the $5_c^{\zeta_2^1}$'s with Z_4 fusion do not have modular extensions. They are not realizable, and do not correspond to any 2+1D bosonic Z_2 -SET orders.

3. The third entry in Table XXIII

Third, let us compute the modular extensions of the third entry in Table XXIII (i.e. $5_1^{\zeta_2^1}$ with $Z_2 \times Z_2$ fusion). We find that the entry has four modular extensions. In fact, the entry corresponds to two different $\text{UMTC}_{/\text{Rep}(Z_2)}$ s, each with two modular extensions, as implied by the two Z_2 -SPT states. The two $\text{UMTC}_{/\text{Rep}(Z_2)}$ s have identical (N_k^{ij}, s_i, c) , but different F -tensors and R -tensors. Sometimes two different $\text{UMTC}_{/\mathcal{E}}$'s (with different F -tensors and the R -tensors) can have the same (N_k^{ij}, s_i) 's. The third, seventh, ..., entries of Table XXIII provide such examples. We like to stress that this is different from the first entry in Table XXIII which corresponds to one $\text{UMTC}_{/\text{Rep}(Z_2)}$.

To see those different F -tensors and R -tensors, we note that one of the two $5_1^{\zeta_2^1}$ with $Z_2 \times Z_2$ fusion has modular extensions given by $3_{1/2}^B \boxtimes 3_{1/2}^B$ and $3_{-3/2}^B \boxtimes 3_{5/2}^B$. We find the R -tensor for this first $5_1^{\zeta_2^1}$ with $Z_2 \times Z_2$ fusion is given by

$$\begin{aligned} R_{(\sigma, \sigma')}^{(f, 1), (\sigma, \sigma')} &= R_{(\sigma, \sigma')}^{(\sigma, \sigma'), (f, 1)} = -i, \\ R_{(\sigma, \sigma')}^{(1, f'), (\sigma, \sigma')} &= R_{(\sigma, \sigma')}^{(\sigma, \sigma'), (1, f')} = -i. \end{aligned} \quad (36)$$

The second $5_1^{\zeta_2^1}$ with $Z_2 \times Z_2$ fusion has modular extensions given by $3_{-1/2}^B \boxtimes 3_{3/2}^B$ and $3_{3/2}^B \boxtimes 3_{-1/2}^B$. We find the R -tensor for the second $5_1^{\zeta_2^1}$ with $Z_2 \times Z_2$ fusion is given by

$$\begin{aligned} R_{(\sigma, \sigma')}^{(f, 1), (\sigma, \sigma')} &= R_{(\sigma, \sigma')}^{(\sigma, \sigma'), (f, 1)} = i, \\ R_{(\sigma, \sigma')}^{(1, f'), (\sigma, \sigma')} &= R_{(\sigma, \sigma')}^{(\sigma, \sigma'), (1, f')} = i. \end{aligned} \quad (37)$$

We see that the two $5_1^{\zeta_2^1}$'s with $Z_2 \times Z_2$ fusion are really different $\text{UMTC}_{/\text{Rep}(Z_2)}$. Each $5_1^{\zeta_2^1}$ has two modular ex-

TABLE XI. The three modular extensions of $\text{Rep}(Z_3)$.

| $N_c^{ \Theta }$ | D^2 | d_1, d_2, \dots | s_1, s_2, \dots | comment |
|-------------------|-------|-------------------|---|-------------------|
| $3_0^{\zeta_4^1}$ | 3 | 1, 1, 1 | 0, 0, 0 | $\text{Rep}(Z_3)$ |
| 9_0^B | 9 | 1×9 | $0, 0, 0, 0, 0, \frac{1}{3}, \frac{1}{3}, \frac{2}{3}, \frac{2}{3}$ | Z_3 gauge |
| 9_0^B | 9 | 1×9 | $0, 0, 0, \frac{1}{9}, \frac{1}{9}, \frac{4}{9}, \frac{4}{9}, \frac{7}{9}, \frac{7}{9}$ | |
| 9_0^B | 9 | 1×9 | $0, 0, 0, \frac{2}{9}, \frac{2}{9}, \frac{5}{9}, \frac{5}{9}, \frac{8}{9}, \frac{8}{9}$ | |

TABLE XII. The six modular extensions of $\text{Rep}(S_3)$.

| $N_c^{ \Theta }$ | D^2 | d_1, d_2, \dots | s_1, s_2, \dots | comment |
|------------------|-------|------------------------|--|-------------------|
| $3_0^{\sqrt{6}}$ | 6 | 1, 1, 2 | 0, 0, 0 | $\text{Rep}(S_3)$ |
| 8_0^B | 36 | 1, 1, 2, 2, 2, 2, 3, 3 | $0, 0, 0, 0, \frac{1}{3}, \frac{2}{3}, 0, \frac{1}{2}$ | S_3 gauge |
| 8_0^B | 36 | 1, 1, 2, 2, 2, 2, 3, 3 | $0, 0, 0, 0, \frac{1}{3}, \frac{2}{3}, \frac{1}{4}, \frac{3}{4}$ | |
| 8_0^B | 36 | 1, 1, 2, 2, 2, 2, 3, 3 | $0, 0, 0, \frac{1}{9}, \frac{4}{9}, \frac{7}{9}, 0, \frac{1}{2}$ | |
| 8_0^B | 36 | 1, 1, 2, 2, 2, 2, 3, 3 | $0, 0, 0, \frac{1}{9}, \frac{4}{9}, \frac{7}{9}, \frac{1}{4}, \frac{3}{4}$ | $(B_4, 2)$ |
| 8_0^B | 36 | 1, 1, 2, 2, 2, 2, 3, 3 | $0, 0, 0, \frac{2}{9}, \frac{5}{9}, \frac{8}{9}, 0, \frac{1}{2}$ | |
| 8_0^B | 36 | 1, 1, 2, 2, 2, 2, 3, 3 | $0, 0, 0, \frac{2}{9}, \frac{5}{9}, \frac{8}{9}, \frac{1}{4}, \frac{3}{4}$ | |

tensions, and that is why we have four entries in Table VII.

Again, Table VII is obtained using the ME-equivalence relation, and is not a table of GQLs. Under the TO-equivalence relation, the third entry is equivalent to the fourth entry of Table VII. So the four entries in Table VII actually describe *three* different Z_2 -SET orders. This has a very interesting consequence: *The Z_2 -SET state described by the third (or fourth) entry in VII, after stacked with an Z_2 -SPT state, still remains in the same phase.* This is an example of the following general statement made previously: *The GQLs with bulk excitations described by \mathcal{C} are in one-to-one correspondence with the quotient $\mathcal{M}_{\text{ext}}(\mathcal{C})/\text{Aut}(\mathcal{C})$ plus a central charge c .* In such an example $\text{Aut}(\mathcal{C})$ is non-trivial.

It is worth noting here that for the second $5_1^{\zeta_2^1}$, two modular extensions $3_{-1/2}^B \boxtimes 3_{3/2}^B$ and $3_{3/2}^B \boxtimes 3_{-1/2}^B$ are actually equivalent UMTCs. This is an example that different embeddings leads to different modular extensions. For $3_{-1/2}^B \boxtimes 3_{3/2}^B$ the first fermion in $5_1^{\zeta_2^1}$ is embedded into $3_{-1/2}^B$ and the second fermion is embedded into $3_{3/2}^B$, while for $3_{3/2}^B \boxtimes 3_{-1/2}^B$ the first fermion is embedded into $3_{3/2}^B$ and the second fermion is embedded into $3_{-1/2}^B$. The equivalence between $3_{-1/2}^B \boxtimes 3_{3/2}^B$ and $3_{3/2}^B \boxtimes 3_{-1/2}^B$ that exchanges both fermions and symmetry twists fails to relate the two embeddings, as they differ by a non-trivial automorphism of $5_1^{\zeta_2^1}$ that exchanges only the two fermions. This is an example that the $\text{Aut}(\mathcal{C})$ action permutes the modular extensions, as discussed in Section IV.

TABLE XIII. The 16 modular extensions of $\text{sRep}(Z_2^f)$.

| $N_c^{ \Theta }$ | D^2 | d_1, d_2, \dots | s_1, s_2, \dots | comment |
|------------------|-------|-------------------|--|----------------------|
| 2_0^0 | 2 | 1, 1 | $0, \frac{1}{2}$ | $\text{sRep}(Z_2^f)$ |
| 4_0^B | 4 | 1, 1, 1, 1 | $0, \frac{1}{2}, 0, 0$ | Z_2 gauge |
| 4_1^B | 4 | 1, 1, 1, 1 | $0, \frac{1}{2}, \frac{1}{8}, \frac{1}{8}$ | F: Z_4 |
| 4_2^B | 4 | 1, 1, 1, 1 | $0, \frac{1}{2}, \frac{1}{4}, \frac{1}{4}$ | F: $Z_2 \times Z_2$ |
| 4_3^B | 4 | 1, 1, 1, 1 | $0, \frac{1}{2}, \frac{3}{8}, \frac{3}{8}$ | F: Z_4 |
| 4_4^B | 4 | 1, 1, 1, 1 | $0, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}$ | F: $Z_2 \times Z_2$ |
| 4_{-3}^B | 4 | 1, 1, 1, 1 | $0, \frac{1}{2}, \frac{5}{8}, \frac{5}{8}$ | F: Z_4 |
| 4_{-2}^B | 4 | 1, 1, 1, 1 | $0, \frac{1}{2}, \frac{3}{4}, \frac{3}{4}$ | F: $Z_2 \times Z_2$ |
| 4_{-1}^B | 4 | 1, 1, 1, 1 | $0, \frac{1}{2}, \frac{7}{8}, \frac{7}{8}$ | F: Z_4 |
| $3_{1/2}^B$ | 4 | 1, 1, ζ_2^1 | $0, \frac{1}{2}, \frac{1}{16}$ | $p + ip$ SC |
| $3_{3/2}^B$ | 4 | 1, 1, ζ_2^1 | $0, \frac{1}{2}, \frac{3}{16}$ | |
| $3_{5/2}^B$ | 4 | 1, 1, ζ_2^1 | $0, \frac{1}{2}, \frac{5}{16}$ | |
| $3_{7/2}^B$ | 4 | 1, 1, ζ_2^1 | $0, \frac{1}{2}, \frac{7}{16}$ | |
| $3_{-7/2}^B$ | 4 | 1, 1, ζ_2^1 | $0, \frac{1}{2}, \frac{9}{16}$ | |
| $3_{-5/2}^B$ | 4 | 1, 1, ζ_2^1 | $0, \frac{1}{2}, \frac{11}{16}$ | |
| $3_{-3/2}^B$ | 4 | 1, 1, ζ_2^1 | $0, \frac{1}{2}, \frac{13}{16}$ | |
| $3_{-1/2}^B$ | 4 | 1, 1, ζ_2^1 | $0, \frac{1}{2}, \frac{15}{16}$ | |

E. Z_3 , Z_5 , and S_3 SPT orders for bosonic systems

We also find that $\text{Rep}(Z_3)$ has 3 modular extensions (see Table XI), $\text{Rep}(Z_5)$ has 5 modular extensions (see Table XIV), and $\text{Rep}(S_3)$ has 6 modular extensions (see Table XII). They correspond to the 3 Z_3 -SPT states, the 5 Z_5 -SPT states and the 6 S_3 -SPT states respectively. These results agree with those from group cohomology theory¹⁹.

We note that for $\text{Rep}(Z_2)$, $\text{Rep}(Z_3)$, and $\text{Rep}(S_3)$, their modular extensions all correspond to distinct UMTCs. However, for $\text{Rep}(Z_5)$, its 5 modular extensions only correspond to 3 distinct UMTCs. $\text{Rep}(Z_5)$ has 5 modular extensions because $\text{Rep}(Z_5)$ can be embedded into the same UMTC in different ways. The different embeddings correspond to different modular extensions.

F. Invertible fermionic topological orders

We find that $\text{sRep}(Z_2^f)$ has 16 modular extensions (see Table XIII) which correspond to invertible fermionic topological orders in 2+1D. One might thought that the invertible fermionic topological orders are classified by \mathbb{Z}_{16} . But in fact, the invertible fermionic topological orders are classified by \mathbb{Z} , obtained by stacking the $c = 1/2$ $p + ip$ states. The discrepancy is due to the fact that the modular extensions cannot see the $c = 8$ E_8 states. The 16 modular extensions exactly correspond to the invertible fermionic topological orders modulo the E_8 states.

We also find that the modular extensions with $c = \text{even}$ have a $Z_2 \times Z_2$ fusion rule, while the modular extensions with $c = \text{odd}$ have a Z_4 fusion rule (indicated by F: $Z_2 \times Z_2$

or F: Z_4 in the comment column of Table).

The Z_2^f -SPT states for fermions is given by the modular extensions with zero central charge. We see that there is only one modular extension with central charge $c = 0$. Thus there is no non-trivial 2+1D fermionic SPT states with Z_2^f symmetry. In general, the modular extensions of $\text{sRep}(G^f)$ with zero central charge correspond to the fermionic SPT states in 2+1D with symmetry G^f .

To calculate the $Z_2 \times Z_2^f$ SPT orders for fermionic systems, we first compute the modular extensions for $\text{sRep}(Z_2 \times Z_2^f)$. We note that $\text{sRep}(Z_2 \times Z_2^f) = \text{sRep}(Z_2^f \times \tilde{Z}_2^f)$. Thus, the modular extensions for $\text{sRep}(Z_2 \times Z_2^f)$ is the modular extensions of $\text{sRep}(Z_2^f \times \tilde{Z}_2^f)$. Some of the modular extensions of $\text{sRep}(Z_2^f \times \tilde{Z}_2^f)$ are given by the modular extensions of $\text{sRep}(Z_2^f)$ stacked (under \boxtimes) with the modular extensions of $\text{sRep}(\tilde{Z}_2^f)$. Some of the modular extensions of $\text{sRep}(Z_2 \times Z_2^f)$ are given by the modular extensions for $\text{Rep}(Z_2)$ stacked (under \boxtimes) with the modular extensions of $\text{sRep}(Z_2^f)$.

The above mathematical statements correspond to the following physical picture: Some fermionic GQLs with $Z_2 \times Z_2^f$ symmetry can be viewed as bosonic GQLs with Z_2 symmetry stacked with fermionic GQLs with Z_2^f symmetry. Also some fermionic GQLs with $Z_2^f \times \tilde{Z}_2^f$ symmetry can be viewed as fermionic GQLs with Z_2^f symmetry stacked with fermionic GQLs with \tilde{Z}_2^f symmetry.

Using eqn. (B12), we find that the modular extensions for $Z_2 \times Z_2^f$ symmetry must have ranks 7, 9, 10, 12, 16. By direct search for those ranks, we find that the modular extensions of $\text{sRep}(Z_2 \times Z_2^f)$ are given by Tables XVII, XVIII, XIX and XX. The $N = 9$ modular extensions of $\text{sRep}(Z_2 \times Z_2^f)$ in Table XVII are given by the stacking of the $N = 3$ modular extensions of $\text{sRep}(Z_2^f)$ and the $N = 3$ modular extensions of $\text{sRep}(\tilde{Z}_2^f)$. The $N = 16$ modular extensions of $\text{sRep}(Z_2 \times Z_2^f)$ in Table XX are given by the stacking of the $N = 4$ modular extensions of $\text{sRep}(Z_2^f)$ and the $N = 4$ modular extensions of $\text{sRep}(\tilde{Z}_2^f)$. There are also 64 $N = 12$ modular extensions of $\text{sRep}(Z_2 \times Z_2^f)$ given by the stacking of the $N = 4$ ($N = 3$) modular extensions of $\text{sRep}(Z_2^f)$ and the $N = 3$ ($N = 4$) modular extensions of $\text{sRep}(\tilde{Z}_2^f)$.

Many of the modular extensions have non-trivial topological orders since the central charge c is non-zero. There are eight modular extensions for each central charge $c = 0, 1/2, 1, 3/2, \dots, 15/2$, and in total $8 \times 16 = 128$ modular extensions. Those eight with $c = 0$ correspond to the $Z_2 \times Z_2^f$ fermionic SPT states. Those are all the $Z_2 \times Z_2^f$ fermionic SPT states⁴⁰.

G. Z_{2n}^f SPT orders for fermionic systems

We also find the modular extensions for $\text{sRep}(Z_4^f)$, $\text{sRep}(Z_6^f)$, and $\text{sRep}(Z_8^f)$ (see Tables XV, XXI, and XVI).

TABLE XIV. The five modular extensions of $\text{Rep}(Z_5)$.

| $N_c^{ \Theta }$ | D^2 | d_1, d_2, \dots | s_1, s_2, \dots | comment |
|------------------|-------|-------------------|---|-------------------------|
| $5_0^{\sqrt{5}}$ | 5 | 1×5 | $0, 0, 0, 0, 0$ | |
| 25_0^B | 25 | 1×25 | $0, 0, 0, 0, 0, 0, 0, 0, 0, \frac{1}{5}, \frac{1}{5}, \frac{1}{5}, \frac{1}{5}, \frac{2}{5}, \frac{2}{5}, \frac{2}{5}, \frac{2}{5}, \frac{3}{5}, \frac{3}{5}, \frac{3}{5}, \frac{3}{5}, \frac{4}{5}, \frac{4}{5}, \frac{4}{5}, \frac{4}{5}$ | $5_0^B \boxtimes 5_0^B$ |
| 25_0^B | 25 | 1×25 | $0, 0, 0, 0, 0, \frac{1}{25}, \frac{1}{25}, \frac{4}{25}, \frac{4}{25}, \frac{6}{25}, \frac{6}{25}, \frac{9}{25}, \frac{9}{25}, \frac{11}{25}, \frac{11}{25}, \frac{14}{25}, \frac{14}{25}, \frac{16}{25}, \frac{16}{25}, \frac{19}{25}, \frac{19}{25}, \frac{21}{25}, \frac{21}{25}, \frac{24}{25}, \frac{24}{25}$ | |
| 25_0^B | 25 | 1×25 | $0, 0, 0, 0, 0, \frac{1}{25}, \frac{1}{25}, \frac{4}{25}, \frac{4}{25}, \frac{6}{25}, \frac{6}{25}, \frac{9}{25}, \frac{9}{25}, \frac{11}{25}, \frac{11}{25}, \frac{14}{25}, \frac{14}{25}, \frac{16}{25}, \frac{16}{25}, \frac{19}{25}, \frac{19}{25}, \frac{21}{25}, \frac{21}{25}, \frac{24}{25}, \frac{24}{25}$ | |
| 25_0^B | 25 | 1×25 | $0, 0, 0, 0, 0, \frac{2}{25}, \frac{2}{25}, \frac{3}{25}, \frac{3}{25}, \frac{7}{25}, \frac{7}{25}, \frac{8}{25}, \frac{8}{25}, \frac{12}{25}, \frac{12}{25}, \frac{13}{25}, \frac{13}{25}, \frac{17}{25}, \frac{17}{25}, \frac{18}{25}, \frac{18}{25}, \frac{22}{25}, \frac{22}{25}, \frac{23}{25}, \frac{23}{25}$ | |
| 25_0^B | 25 | 1×25 | $0, 0, 0, 0, 0, \frac{2}{25}, \frac{2}{25}, \frac{3}{25}, \frac{3}{25}, \frac{7}{25}, \frac{7}{25}, \frac{8}{25}, \frac{8}{25}, \frac{12}{25}, \frac{12}{25}, \frac{13}{25}, \frac{13}{25}, \frac{17}{25}, \frac{17}{25}, \frac{18}{25}, \frac{18}{25}, \frac{22}{25}, \frac{22}{25}, \frac{23}{25}, \frac{23}{25}$ | |

TABLE XV. All the 8 modular extensions of $\text{sRep}(Z_4^f)$.

| $N_c^{ \Theta }$ | D^2 | d_1, d_2, \dots | s_1, s_2, \dots | comment |
|------------------|-------|--|--|-------------------------------|
| 4_0^0 | 4 | $1, 1, 1, 1$ | $0, 0, \frac{1}{2}, \frac{1}{2}$ | $\text{sRep}(Z_4^f)$ |
| 16_0^B | 16 | $1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, 0, 0, 0, 0, \frac{1}{4}, \frac{1}{4}, \frac{1}{2}, \frac{1}{2}, \frac{3}{4}, \frac{3}{4}$ | |
| 16_1^B | 16 | $1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{32}, \frac{1}{32}, \frac{1}{8}, \frac{1}{8}, \frac{1}{8}, \frac{1}{8}, \frac{9}{32}, \frac{9}{32}, \frac{17}{32}, \frac{17}{32}, \frac{25}{32}, \frac{25}{32}$ | $8_1^B \boxtimes 2_1^B$ |
| 16_2^B | 16 | $1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{16}, \frac{1}{16}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{5}{16}, \frac{5}{16}, \frac{9}{16}, \frac{9}{16}, \frac{13}{16}, \frac{13}{16}$ | |
| 16_3^B | 16 | $1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{32}, \frac{3}{32}, \frac{3}{32}, \frac{3}{32}, \frac{8}{8}, \frac{8}{8}, \frac{8}{8}, \frac{8}{8}, \frac{32}{32}, \frac{32}{32}, \frac{27}{32}, \frac{27}{32}$ | $4_3^B \boxtimes 4_1^B$ |
| 16_4^B | 16 | $1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}, \frac{1}{8}, \frac{1}{8}, \frac{1}{8}, \frac{1}{2}, \frac{1}{2}, \frac{5}{8}, \frac{5}{8}, \frac{7}{8}, \frac{7}{8}, \frac{7}{8}, \frac{7}{8}$ | |
| 16_{-3}^B | 16 | $1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{5}{32}, \frac{5}{32}, \frac{13}{32}, \frac{13}{32}, \frac{5}{8}, \frac{5}{8}, \frac{5}{8}, \frac{5}{8}, \frac{21}{32}, \frac{21}{32}, \frac{29}{32}, \frac{29}{32}$ | $8_{-1}^B \boxtimes 2_{-1}^B$ |
| 16_{-2}^B | 16 | $1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{16}, \frac{3}{16}, \frac{7}{16}, \frac{7}{16}, \frac{11}{16}, \frac{11}{16}, \frac{3}{4}, \frac{3}{4}, \frac{3}{4}, \frac{3}{4}, \frac{15}{16}, \frac{15}{16}$ | |
| 16_{-1}^B | 16 | $1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{7}{32}, \frac{7}{32}, \frac{15}{32}, \frac{15}{32}, \frac{23}{32}, \frac{23}{32}, \frac{7}{8}, \frac{7}{8}, \frac{7}{8}, \frac{7}{8}, \frac{31}{32}, \frac{31}{32}$ | |

Again, many of them has non-trivial topological orders since the central charge c is non-zero.

For Z_4^f group, only one of them have $c = 0$. So there is no non-trivial Z_4^f fermionic SPT states. For Z_6^f group, only three of them have $c = 0$. So, the Z_6^f fermionic SPT states are described by Z_3 . For Z_8^f group, only two of them have $c = 0$. So, the Z_8^f fermionic SPT states are described by Z_2 . Those results are consistent with the results in Ref. 41 and 42. However, the calculation present here is more complete.

IX. SUMMARY

GQLs contain both topologically ordered states and SPT states. In this paper, we present a theory that classify GQLs in 2+1D for bosonic/fermionic systems with symmetry.

We propose that the possible non-abelian statistics (or sets of bulk quasiparticles excitations) in 2+1D GQLs are classified by UMTC/\mathcal{E} , where $\mathcal{E} = \text{Rep}(G)$ or $\text{sRep}(G^f)$ describing the symmetry in bosonic or fermionic systems. However, UMTC/\mathcal{E} 's fail to classify GQLs, since different GQL phases can have identical non-abelian statistics, which correspond to identical UMTC/\mathcal{E} .

To fix this problem, we introduce the notion of modular extensions for a UMTC/\mathcal{E} . We propose to use the triple $(\mathcal{C}, \mathcal{M}, c)$ to classify 2+1D GQLs with symmetry G (for boson) or G^f (for fermion). Here \mathcal{C} is a UMTC/\mathcal{E} with $\mathcal{E} = \text{Rep}(G)$ or $\text{sRep}(G^f)$, \mathcal{M} is a modular extension of \mathcal{C} and c is the chiral central charge of the edge state. We

show that the modular extensions of a UMTC/\mathcal{E} has a one-to-one correspondence with the modular extensions of \mathcal{E} . So the number of the modular extensions is solely determined by the symmetry \mathcal{E} . Also, the $c = 0$ modular extensions of a \mathcal{E} ($\mathcal{E} = \text{Rep}(G)$ or $\text{sRep}(G^f)$) classify the 2+1D SPT states for bosons or fermions with symmetry G or G^f .

Although the above result has a nice mathematical structure, it is hard to implement numerically to produce a table of GQLs. To fix this problem, we propose a different description of 2+1D GQLs. We propose to use the data $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i; \mathcal{N}_K^{IJ}, \mathcal{S}_I; c)$, up to some permutations of the indices, to describe 2+1D GQLs with symmetry G (for boson) or G^f (for fermion), with a restriction that the symmetry group G can be fully characterized by the fusion ring of its irreducible representations (for example, for simple groups or abelian groups). Here the data $(\tilde{N}_c^{ab}, \tilde{s}_a)$ describe the symmetry and the data (N_k^{ij}, s_i) describes fusion and the spins of the bulk particles in the GQL. The modular extensions are obtained by “gauging” the symmetry G or G^f . The data $(\mathcal{N}_K^{IJ}, \mathcal{S}_I)$ describes fusion and the spins of the bulk particles in the “gauged” theory. Last, c is the chiral central charge of the edge state.

In this paper (see Appendix C) and in Ref. 11, we list the necessary and the sufficient conditions on the data $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i; \mathcal{N}_K^{IJ}, \mathcal{S}_I; c)$, which allow us to obtain a list of GQLs. However, in this paper, we did not give the list of GQLs directly. We first give a list of $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$, which is an imperfect list of UMTC/\mathcal{E} 's. We then compute the modular extensions

TABLE XVIII. The first 32 modular extensions of $\text{sRep}(Z_2 \times Z_2^f)$ with $N = 12$.

| $N_c^{ \Theta }$ | D^2 | d_1, d_2, \dots | s_1, s_2, \dots | comment |
|------------------|-------|--|---|---------------------------------|
| 4_0^0 | 4 | 1, 1, 1, 1 | $0, 0, \frac{1}{2}, \frac{1}{2}$ | $\text{sRep}(Z_2 \times Z_2^f)$ |
| $12_{1/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{16}, \frac{1}{16}, \frac{1}{16}, \frac{9}{16}$ | $4_0^B \boxtimes 3_{1/2}^B$ |
| $12_{1/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{16}, \frac{1}{16}, \frac{1}{16}, \frac{9}{16}$ | $4_0^B \boxtimes 3_{1/2}^B$ |
| $12_{1/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}, \frac{1}{8}, \frac{5}{8}, \frac{5}{8}, \frac{1}{16}, \frac{1}{16}, \frac{7}{16}, \frac{15}{16}$ | $4_{-3}^B \boxtimes 3_{7/2}^B$ |
| $12_{1/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}, \frac{1}{8}, \frac{5}{8}, \frac{5}{8}, \frac{1}{16}, \frac{1}{16}, \frac{7}{16}, \frac{15}{16}$ | $4_{-3}^B \boxtimes 3_{7/2}^B$ |
| $12_{1/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{1}{4}, \frac{3}{4}, \frac{3}{4}, \frac{1}{16}, \frac{1}{16}, \frac{5}{16}, \frac{13}{16}$ | $6_{-1/2}^B \boxtimes 2_1^B$ |
| $12_{1/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{1}{4}, \frac{3}{4}, \frac{3}{4}, \frac{1}{16}, \frac{1}{16}, \frac{5}{16}, \frac{13}{16}$ | $6_{-1/2}^B \boxtimes 2_1^B$ |
| $12_{1/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{8}, \frac{3}{8}, \frac{7}{8}, \frac{7}{8}, \frac{1}{16}, \frac{1}{16}, \frac{3}{16}, \frac{11}{16}$ | $4_{-1}^B \boxtimes 3_{3/2}^B$ |
| $12_{1/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{8}, \frac{3}{8}, \frac{7}{8}, \frac{7}{8}, \frac{1}{16}, \frac{1}{16}, \frac{3}{16}, \frac{11}{16}$ | $4_{-1}^B \boxtimes 3_{3/2}^B$ |
| $12_{3/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{16}, \frac{3}{16}, \frac{3}{16}, \frac{11}{16}$ | $4_0^B \boxtimes 3_{3/2}^B$ |
| $12_{3/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{16}, \frac{3}{16}, \frac{3}{16}, \frac{11}{16}$ | $4_0^B \boxtimes 3_{3/2}^B$ |
| $12_{3/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}, \frac{1}{8}, \frac{5}{8}, \frac{5}{8}, \frac{1}{16}, \frac{1}{16}, \frac{3}{16}, \frac{9}{16}$ | $4_1^B \boxtimes 3_{1/2}^B$ |
| $12_{3/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}, \frac{1}{8}, \frac{5}{8}, \frac{5}{8}, \frac{1}{16}, \frac{1}{16}, \frac{3}{16}, \frac{9}{16}$ | $4_1^B \boxtimes 3_{1/2}^B$ |
| $12_{3/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{1}{4}, \frac{3}{4}, \frac{3}{4}, \frac{1}{16}, \frac{1}{16}, \frac{7}{16}, \frac{15}{16}$ | $6_{1/2}^B \boxtimes 2_1^B$ |
| $12_{3/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{1}{4}, \frac{3}{4}, \frac{3}{4}, \frac{1}{16}, \frac{1}{16}, \frac{7}{16}, \frac{15}{16}$ | $6_{1/2}^B \boxtimes 2_1^B$ |
| $12_{3/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{8}, \frac{3}{8}, \frac{7}{8}, \frac{7}{8}, \frac{3}{16}, \frac{3}{16}, \frac{5}{16}, \frac{13}{16}$ | $4_{-1}^B \boxtimes 3_{5/2}^B$ |
| $12_{3/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{8}, \frac{3}{8}, \frac{7}{8}, \frac{7}{8}, \frac{3}{16}, \frac{3}{16}, \frac{5}{16}, \frac{13}{16}$ | $4_{-1}^B \boxtimes 3_{5/2}^B$ |
| $12_{5/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{5}{16}, \frac{5}{16}, \frac{5}{16}, \frac{13}{16}$ | $4_0^B \boxtimes 3_{5/2}^B$ |
| $12_{5/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{5}{16}, \frac{5}{16}, \frac{5}{16}, \frac{13}{16}$ | $4_0^B \boxtimes 3_{5/2}^B$ |
| $12_{5/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}, \frac{1}{8}, \frac{5}{8}, \frac{5}{8}, \frac{3}{16}, \frac{3}{16}, \frac{5}{16}, \frac{11}{16}$ | $4_1^B \boxtimes 3_{3/2}^B$ |
| $12_{5/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}, \frac{1}{8}, \frac{5}{8}, \frac{5}{8}, \frac{3}{16}, \frac{3}{16}, \frac{5}{16}, \frac{11}{16}$ | $4_1^B \boxtimes 3_{3/2}^B$ |
| $12_{5/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{1}{4}, \frac{3}{4}, \frac{3}{4}, \frac{1}{16}, \frac{1}{16}, \frac{5}{16}, \frac{9}{16}$ | $6_{3/2}^B \boxtimes 2_1^B$ |
| $12_{5/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{1}{4}, \frac{3}{4}, \frac{3}{4}, \frac{1}{16}, \frac{1}{16}, \frac{5}{16}, \frac{9}{16}$ | $6_{3/2}^B \boxtimes 2_1^B$ |
| $12_{5/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{8}, \frac{3}{8}, \frac{7}{8}, \frac{7}{8}, \frac{5}{16}, \frac{5}{16}, \frac{7}{16}, \frac{15}{16}$ | $4_{-1}^B \boxtimes 3_{7/2}^B$ |
| $12_{5/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{8}, \frac{3}{8}, \frac{7}{8}, \frac{7}{8}, \frac{5}{16}, \frac{5}{16}, \frac{7}{16}, \frac{15}{16}$ | $4_{-1}^B \boxtimes 3_{7/2}^B$ |
| $12_{7/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{7}{16}, \frac{7}{16}, \frac{7}{16}, \frac{15}{16}$ | $4_0^B \boxtimes 3_{7/2}^B$ |
| $12_{7/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{7}{16}, \frac{7}{16}, \frac{7}{16}, \frac{15}{16}$ | $4_0^B \boxtimes 3_{7/2}^B$ |
| $12_{7/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}, \frac{1}{8}, \frac{5}{8}, \frac{5}{8}, \frac{5}{16}, \frac{5}{16}, \frac{7}{16}, \frac{13}{16}$ | $4_1^B \boxtimes 3_{5/2}^B$ |
| $12_{7/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}, \frac{1}{8}, \frac{5}{8}, \frac{5}{8}, \frac{5}{16}, \frac{5}{16}, \frac{7}{16}, \frac{13}{16}$ | $4_1^B \boxtimes 3_{5/2}^B$ |
| $12_{7/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{1}{4}, \frac{3}{4}, \frac{3}{4}, \frac{3}{16}, \frac{3}{16}, \frac{7}{16}, \frac{11}{16}$ | $6_{5/2}^B \boxtimes 2_1^B$ |
| $12_{7/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{1}{4}, \frac{3}{4}, \frac{3}{4}, \frac{3}{16}, \frac{3}{16}, \frac{7}{16}, \frac{11}{16}$ | $6_{5/2}^B \boxtimes 2_1^B$ |
| $12_{7/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{8}, \frac{3}{8}, \frac{7}{8}, \frac{7}{8}, \frac{1}{16}, \frac{1}{16}, \frac{7}{16}, \frac{9}{16}$ | $4_3^B \boxtimes 3_{1/2}^B$ |
| $12_{7/2}^B$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, $\zeta_2^1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{8}, \frac{3}{8}, \frac{7}{8}, \frac{7}{8}, \frac{1}{16}, \frac{1}{16}, \frac{7}{16}, \frac{9}{16}$ | $4_3^B \boxtimes 3_{1/2}^B$ |

Acknowledgement: We like to thank Pavel Etingof, Dmitri Nikshych, Chenjie Wang, and Zhenghan Wang for many helpful discussions. This research is supported by NSF Grant No. DMR-1506475, and NSFC 11274192. It is also supported by the John Templeton Foundation No. 39901. Research at Perimeter Institute is supported by the Government of Canada through Industry Canada and by the Province of Ontario through the Ministry of Research. LK is supported by the Center of Mathematical Sciences and Applications at Harvard University.

[illegible]

[illegible]

TABLE XXI. All the modular extensions of $\text{sRep}(Z_6^f) = \text{sRep}(Z_3 \times Z_2^f)$.

[illegible]

TABLE XXII. Z_2 -SET orders (or $\text{UMTC}_{/\text{Rep}(Z_2)}$) for bosonic systems labeled in terms of sets of topological excitations. The list contains all topological orders with $N = 3, 4$ and $D^2 \leq 100$. All the topological orders in this list are anomaly free (*i.e.* have modular extensions), and are realizable by 2+1D bosonic systems. We use $N_c^{|\Theta|}$ to label $\text{UMTC}_{/\mathcal{E}}$'s, where $\Theta = D^{-1} \sum_i e^{2\pi i s_i} d_i^2 = |\Theta| e^{2\pi i c/8}$ and $D^2 = \sum_i d_i^2$.

| $N_c^{ \Theta }$ | D^2 | d_1, d_2, \dots | s_1, s_2, \dots | comment |
|-------------------------|--------|------------------------------|----------------------------------|---|
| $2_0^{\zeta_2^1}$ | 2 | 1, 1 | 0, 0 | $\mathcal{E} = \text{Rep}(Z_2)$ |
| $3_2^{\zeta_2^1}$ | 6 | 1, 1, 2 | $0, 0, \frac{1}{3}$ | $\text{SB}: K = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$ |
| $3_{-2}^{\zeta_2^1}$ | 6 | 1, 1, 2 | $0, 0, \frac{2}{3}$ | $\text{SB}: K = \begin{pmatrix} -2 & 1 \\ 1 & -2 \end{pmatrix}$ |
| $4_1^{\zeta_2^1}$ | 4 | 1, 1, 1, 1 | $0, 0, \frac{1}{4}, \frac{1}{4}$ | $2_1^B \boxtimes \text{Rep}(Z_2)$ |
| $4_1^{\zeta_2^1}$ | 4 | 1, 1, 1, 1 | $0, 0, \frac{1}{4}, \frac{1}{4}$ | $2_1^B \boxtimes^t \text{Rep}(Z_2)$ |
| $4_{-1}^{\zeta_2^1}$ | 4 | 1, 1, 1, 1 | $0, 0, \frac{3}{4}, \frac{3}{4}$ | $2_{-1}^B \boxtimes \text{Rep}(Z_2)$ |
| $4_{-1}^{\zeta_2^1}$ | 4 | 1, 1, 1, 1 | $0, 0, \frac{3}{4}, \frac{3}{4}$ | $2_{-1}^B \boxtimes^t \text{Rep}(Z_2)$ |
| $4_{14/5}^{\zeta_2^1}$ | 7.2360 | $1, 1, \zeta_3^1, \zeta_3^1$ | $0, 0, \frac{2}{5}, \frac{2}{5}$ | $2_{14/5}^B \boxtimes \text{Rep}(Z_2)$ |
| $4_{-14/5}^{\zeta_2^1}$ | 7.2360 | $1, 1, \zeta_3^1, \zeta_3^1$ | $0, 0, \frac{3}{5}, \frac{3}{5}$ | $2_{-14/5}^B \boxtimes \text{Rep}(Z_2)$ |
| $4_0^{\zeta_2^1}$ | 10 | 1, 1, 2, 2 | $0, 0, \frac{1}{5}, \frac{4}{5}$ | $\text{SB}: K = \begin{pmatrix} 2 & -3 \\ -3 & 2 \end{pmatrix}$ |
| $4_4^{\zeta_2^1}$ | 10 | 1, 1, 2, 2 | $0, 0, \frac{2}{5}, \frac{3}{5}$ | $\text{SB}: K = \begin{pmatrix} 2 & 1 & 0 & 0 \\ 1 & 2 & 0 & 1 \\ 0 & 0 & 2 & 1 \\ 0 & 1 & 1 & 2 \end{pmatrix}$ |

Appendix A: Tables for the solutions of $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$ – imperfect tables for $\text{UMTC}_{/\mathcal{E}}$

In this appendix, we list $\text{UMTC}_{/\mathcal{E}}$'s for various symmetry \mathcal{E} , which can also be viewed as the list of 2+1D SET orders (up to invertible ones) with symmetry \mathcal{E} . Those lists are created using a naive calculation, by checking the necessary conditions on the data $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$ (for details, see Appendix C). So those lists should contain all $\text{UMTC}_{/\mathcal{E}}$'s (*i.e.* all SET orders). But since the conditions are only known to be necessary, the lists may contain fake entries that do not correspond to any $\text{UMTC}_{/\mathcal{E}}$ (or any SET order). In other words, some entries in the lists have no modular extensions and those entries do not correspond any real 2+1D SET order.

The entries with known decomposition $N_c^B \boxtimes \text{Rep}(G)$ or $N_c^B \boxtimes \text{sRep}(G^f)$, or with given K -matrix in the comment column all correspond to existing 2+1D SET orders. (The topological orders described by N_c^B are given by the tables in Ref. 11.) Other entries may or may not correspond to existing 2+1D SET orders, which need to be determined by checking the existence of modular extensions.

Even for the entries that have modular extensions, some times they may correspond to more than one $\text{UMTC}_{/\mathcal{E}}$'s. This is because $(\tilde{N}_c^{ab}, \tilde{s}_a; N_k^{ij}, s_i)$ cannot distinguish all different $\text{UMTC}_{/\mathcal{E}}$'s.

TABLE XXIII. Z_2 -SET orders for bosonic systems labeled in terms of sets of topological excitations. The list contains all topological orders with $N = 5$ and $D^2 \leq 100$.

| $N_c^{ \Theta }$ | D^2 | d_1, d_2, \dots | s_1, s_2, \dots | comment |
|-------------------------|--------|---|---|---|
| $2_0^{\zeta_2^1}$ | 2 | 1, 1 | 0, 0 | $\mathcal{E} = \text{Rep}(Z_2)$ |
| $5_0^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, 0$ | $\text{SB}: 4_0^B \text{ F}: Z_2 \times Z_2$ |
| $5_0^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, 0$ | $\text{SB}: 4_0^B \text{ F}: Z_4 \text{ NR}$ |
| $5_1^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}$ | $\text{SB}: 4_1^B \text{ F}: Z_2 \times Z_2$ |
| $5_1^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}$ | $\text{SB}: 4_1^B \text{ F}: Z_4 \text{ NR}$ |
| $5_2^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}$ | $\text{SB}: 4_2^B \text{ F}: Z_2 \times Z_2$ |
| $5_2^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}$ | $\text{SB}: 4_2^B \text{ F}: Z_4 \text{ NR}$ |
| $5_3^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{8}$ | $\text{SB}: 4_3^B \text{ F}: Z_2 \times Z_2$ |
| $5_3^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{8}$ | $\text{SB}: 4_3^B \text{ F}: Z_4 \text{ NR}$ |
| $5_4^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}$ | $\text{SB}: 4_4^B \begin{pmatrix} 2 & 1 & 1 & 1 \\ 1 & 2 & 0 & 0 \\ 1 & 0 & 2 & 0 \\ 1 & 0 & 0 & 2 \end{pmatrix}$ |
| $5_4^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}$ | $\text{SB}: 4_4^B \text{ F}: Z_4 \text{ NR}$ |
| $5_{-3}^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{5}{8}$ | $\text{SB}: 4_{-3}^B \text{ F}: Z_2 \times Z_2$ |
| $5_{-3}^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{5}{8}$ | $\text{SB}: 4_{-3}^B \text{ F}: Z_4 \text{ NR}$ |
| $5_{-2}^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{4}$ | $\text{SB}: 4_{-2}^B \text{ F}: Z_2 \times Z_2$ |
| $5_{-2}^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{4}$ | $\text{SB}: 4_{-2}^B \text{ F}: Z_4 \text{ NR}$ |
| $5_{-1}^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{7}{8}$ | $\text{SB}: 4_{-1}^B \text{ F}: Z_2 \times Z_2$ |
| $5_{-1}^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{7}{8}$ | $\text{SB}: 4_{-1}^B \text{ F}: Z_4 \text{ NR}$ |
| $5_{-2}^{\zeta_2^1}$ | 14 | 1, 1, 2, 2, 2 | $0, 0, \frac{1}{7}, \frac{2}{7}, \frac{4}{7}$ | $\text{SB}: 7_2^B$ |
| $5_{-2}^{\zeta_2^1}$ | 14 | 1, 1, 2, 2, 2 | $0, 0, \frac{3}{7}, \frac{5}{7}, \frac{6}{7}$ | $\text{SB}: 7_{-2}^B$ |
| $5_{12/5}^{\zeta_2^1}$ | 26.180 | $1, 1, \zeta_8^2, \zeta_8^2, \zeta_8^4$ | $0, 0, \frac{1}{5}, \frac{1}{5}, \frac{3}{5}$ | $\text{SB}: 4_{12/5}^B$ |
| $5_{-12/5}^{\zeta_2^1}$ | 26.180 | $1, 1, \zeta_8^2, \zeta_8^2, \zeta_8^4$ | $0, 0, \frac{4}{5}, \frac{4}{5}, \frac{2}{5}$ | $\text{SB}: 4_{-12/5}^B$ |

1. Z_2 -SET orders

Tables XXII, XXIII, and XXIV list the Z_2 -SET orders (up to invertible ones) for 2+1D bosonic systems. For bosonic systems the central charge is determined up to 8 by the bulk excitations. The $3_2^{\zeta_2^1}$ states and the two $4_1^{\zeta_2^1}$ states in Table XXII are discussed in the main text.

All the Z_2 -SET orders in Table XXII are realizable. Some of the them are realized as $N_c^B \boxtimes \text{Rep}(Z_2)$, as indicated in the comment column. Here N_c^B describes a neutral bosonic topological order (which was denoted as N_c^B in Ref. 11) with rank N and central charge c , which does not transform under the Z_2 symmetry. For example 2_1^B is the $\nu = 1/2$ bosonic Laughlin state, and $2_{14/5}^B$ is the bosonic Fibonacci state¹¹. $\text{Rep}(Z_2)$ describes a product state with Z_2 symmetry of Z_2 charged bosons. $N_c^B \boxtimes \text{Rep}(Z_2)$ is simply the stacking of the neutral bosonic topological order N_c^B with the Z_2 symmetric product state.

We also introduced $N_c^B \boxtimes^t \text{Rep}(Z_2)$ which describe a state similar to $N_c^B \boxtimes \text{Rep}(Z_2)$, except here the bosons that form the topological order N_c^B also carries a Z_2

charge. The $3_2^{\zeta_1}$ state can be realized by double-layer FQH state with K -matrix $\begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$, which is discussed in the main text.

Since we did not use the condition of the existence of modular extensions when we calculate the tables, some the entries in the tables may not be realizable by any 2+1D bosonic systems. We use NR in the comment column to indicate such entries (see Table XXIII).

2. Z_3 -SET orders

Table XXV lists the Z_3 -SET orders (up to invertible ones) for 2+1D bosonic systems.

The Z_3 -SET state $4_4^{\zeta_1}$ in the table becomes the $K = \begin{pmatrix} 2 & 1 & 1 & 1 \\ 1 & 2 & 0 & 0 \\ 1 & 0 & 2 & 0 \\ 1 & 0 & 0 & 2 \end{pmatrix}$ 4-layer FQH state after we break the Z_3 -symmetry. We can add the Z_3 -symmetry back to obtain the Z_3 -SET state. The Z_3 -symmetry is the cyclic permutation of the second, the third, and the fourth layers.

Without the symmetry, the $K = \begin{pmatrix} 2 & 1 & 1 & 1 \\ 1 & 2 & 0 & 0 \\ 1 & 0 & 2 & 0 \\ 1 & 0 & 0 & 2 \end{pmatrix}$ state has four types of particles, a trivial boson and three non-trivial fermions. With the symmetry, the three fermions become degenerate and is combined into the $d = 3$ particle (the fourth particle) for the $4_4^{\zeta_1}$ state. The first three particles for the $4_4^{\zeta_1}$ state all come from the trivial boson. They carry different Z_3 charges: 0, 1, 2, in the presence of the symmetry.

3. S_3 -SET orders

Tables XXVII and XXVIII list the S_3 -SET orders (up to invertible ones) for 2+1D bosonic systems.

Table XXVII has three $5_4^{\sqrt{6}}$ entries that have identical (d_i, s_i) . But the three entries have different fusion rules (see Table XXVI). If we break the symmetry, the three entries all reduce to the $K = \begin{pmatrix} 2 & 1 & 1 & 1 \\ 1 & 2 & 0 & 0 \\ 1 & 0 & 2 & 0 \\ 1 & 0 & 0 & 2 \end{pmatrix}$ 4-layer state. So we expect the S_3 symmetry is the permutation symmetry of the second, the third, and the fourth layers.

The second $5_4^{\sqrt{6}}$ entry can be realized by the $K = \begin{pmatrix} 2 & 1 & 1 & 1 \\ 1 & 2 & 0 & 0 \\ 1 & 0 & 2 & 0 \\ 1 & 0 & 0 & 2 \end{pmatrix}$ 4-layer state. The two $d = 3$ fermions are the direct-sum of the three degenerate fermions in the $K = \begin{pmatrix} 2 & 1 & 1 & 1 \\ 1 & 2 & 0 & 0 \\ 1 & 0 & 2 & 0 \\ 1 & 0 & 0 & 2 \end{pmatrix}$ state. They carry the following S_3 representations

$$\sigma \rightarrow \mathbf{1} \oplus b, \quad \tau \rightarrow a \oplus b. \quad (\text{A1})$$

It is strange that two different irreducible representations are degenerate in energy. But this can happen for topological excitations in the presence of symmetry.

Such an assignment of the S_3 -representations (or S_3 “charges”) is consistent with the fusion rule (see the second table in Table XXVI). For example

$$\begin{aligned} \sigma \otimes \sigma &\rightarrow \mathbf{1} \oplus 2b \oplus b \otimes b = \mathbf{1} \oplus 2b \oplus (\mathbf{1} \oplus a \oplus b) \\ &\rightarrow \mathbf{1} \oplus b \oplus \sigma \oplus \tau \end{aligned} \quad (\text{A2})$$

This is why we say that the second $5_4^{\sqrt{6}}$ entry can be realized by the $K = \begin{pmatrix} 2 & 1 & 1 & 1 \\ 1 & 2 & 0 & 0 \\ 1 & 0 & 2 & 0 \\ 1 & 0 & 0 & 2 \end{pmatrix}$ state.

However, the S_3 -charge assignment eqn. (A1) does not work for the first and the third $5_4^{\sqrt{6}}$ entries (*i.e.* inconsistent with fusion rules in the first and the third tables in Table XXVI). In fact, none of the S_3 -charge assignment works. This means that the $d = 3$ fermions in the first and the third $5_4^{\sqrt{6}}$ entries must carry fractionalized S_3 -charges or fractionalized S_3 -representations. It is not clear if such fractionalized S_3 -charges are realizable or not, since we cannot calculate the modular extensions for those entries (due to the limitation of computer power).

4. $Z_2 \times Z_2$ -SET orders

Tables XXIX, XXX, and XXXI list the $Z_2 \times Z_2$ -SET orders (up to invertible ones) for 2+1D bosonic systems.

Table XXXII list the fusion rules for some $Z_2 \times Z_2$ -SET orders. We see that the 5_1^2 state is a $\nu = 1/2$ bosonic Laughlin state with $Z_2 \times Z_2$ symmetry, where the only topological excitation carries the projective representation of $Z_2 \times Z_2$. We also see that the $5_{14/2}^2$ state is a bosonic Fibonacci state with $Z_2 \times Z_2$ symmetry, where the only non-abelian topological excitation carries the projective representation of $Z_2 \times Z_2$.

5. $Z_2 \times Z_2^f$ -SET and Z_4^f -SET orders

Table XXXIII lists the $Z_2 \times Z_2^f$ -SET orders (up to invertible ones) for 2+1D fermionic systems. Table XXXIV lists the Z_4^f -SET orders (up to invertible ones) for 2+1D fermionic systems. For fermionic systems the central charge is determined up to c_{\min} by the bulk excitations, where c_{\min} is the smallest positive central charge of the modular extensions of $\text{sRep}(G^f)$, for example, $c_{\min} = 1/2$ for $Z_2^f, Z_2 \times Z_2^f, Z_6^f$, $c_{\min} = 1$ for Z_4^f, Z_8^f .

Appendix B: Fusion ring for the modular extensions of $\text{Rep}(G)$ or $\text{sRep}(G^f)$ when G or G^f is abelian group

When the symmetry group G is abelian, the different irreducible representations, under the fusion, form the

TABLE XXIV. Z_2 -SET orders for bosonic systems labeled in terms of sets of topological excitations. The list contains all topological orders with $N = 6$ $D^2 \leq 50$.

| $N_c^{ \Theta }$ | D^2 | d_1, d_2, \dots | s_1, s_2, \dots | comment |
|-------------------------|--------|--|--|--|
| $2_0^{\zeta_2^1}$ | 2 | 1, 1 | 0, 0 | $\mathcal{E} = \text{Rep}(Z_2)$ |
| $6_2^{\zeta_2^1}$ | 6 | 1, 1, 1, 1, 1, 1 | $0, 0, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}$ | $3_2^B \boxtimes \text{Rep}(Z_2)$ |
| $6_{-2}^{\zeta_2^1}$ | 6 | 1, 1, 1, 1, 1, 1 | $0, 0, \frac{2}{3}, \frac{2}{3}, \frac{2}{3}, \frac{2}{3}$ | $3_{-2}^B \boxtimes \text{Rep}(Z_2)$ |
| $6_{1/2}^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, ζ_2^1, ζ_2^1 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{16}, \frac{1}{16}$ | $3_{1/2}^B \boxtimes \text{Rep}(Z_2)$ |
| $6_{1/2}^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, ζ_2^1, ζ_2^1 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{16}, \frac{1}{16}$ | SB: $3_{1/2}^B$ |
| $6_{3/2}^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, ζ_2^1, ζ_2^1 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{16}, \frac{3}{16}$ | $3_{3/2}^B \boxtimes \text{Rep}(Z_2)$ |
| $6_{3/2}^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, ζ_2^1, ζ_2^1 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{16}, \frac{3}{16}$ | SB: $3_{3/2}^B$ |
| $6_{5/2}^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, ζ_2^1, ζ_2^1 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{5}{16}, \frac{5}{16}$ | $3_{5/2}^B \boxtimes \text{Rep}(Z_2)$ |
| $6_{5/2}^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, ζ_2^1, ζ_2^1 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{5}{16}, \frac{5}{16}$ | SB: $3_{5/2}^B$ |
| $6_{7/2}^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, ζ_2^1, ζ_2^1 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{7}{16}, \frac{7}{16}$ | $3_{7/2}^B \boxtimes \text{Rep}(Z_2)$ |
| $6_{7/2}^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, ζ_2^1, ζ_2^1 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{7}{16}, \frac{7}{16}$ | SB: $3_{7/2}^B$ |
| $6_{-7/2}^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, ζ_2^1, ζ_2^1 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{9}{16}, \frac{9}{16}$ | $3_{-7/2}^B \boxtimes \text{Rep}(Z_2)$ |
| $6_{-7/2}^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, ζ_2^1, ζ_2^1 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{9}{16}, \frac{9}{16}$ | SB: $3_{-7/2}^B$ |
| $6_{-5/2}^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, ζ_2^1, ζ_2^1 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{11}{16}, \frac{11}{16}$ | $3_{-5/2}^B \boxtimes \text{Rep}(Z_2)$ |
| $6_{-5/2}^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, ζ_2^1, ζ_2^1 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{11}{16}, \frac{11}{16}$ | SB: $3_{-5/2}^B$ |
| $6_{-3/2}^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, ζ_2^1, ζ_2^1 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{13}{16}, \frac{13}{16}$ | $3_{-3/2}^B \boxtimes \text{Rep}(Z_2)$ |
| $6_{-3/2}^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, ζ_2^1, ζ_2^1 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{13}{16}, \frac{13}{16}$ | SB: $3_{-3/2}^B$ |
| $6_{-1/2}^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, ζ_2^1, ζ_2^1 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{15}{16}, \frac{15}{16}$ | $3_{-1/2}^B \boxtimes \text{Rep}(Z_2)$ |
| $6_{-1/2}^{\zeta_2^1}$ | 8 | 1, 1, 1, 1, ζ_2^1, ζ_2^1 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{15}{16}, \frac{15}{16}$ | SB: $3_{-1/2}^B$ |
| $6_1^{\zeta_2^1}$ | 12 | 1, 1, 1, 1, 2, 2 | $0, 0, \frac{3}{4}, \frac{3}{4}, \frac{1}{12}, \frac{1}{3}$ | SB: 6_1^B |
| $6_3^{\zeta_2^1}$ | 12 | 1, 1, 1, 1, 2, 2 | $0, 0, \frac{1}{4}, \frac{1}{4}, \frac{1}{3}, \frac{7}{12}$ | SB: 6_3^B |
| $6_{-3}^{\zeta_2^1}$ | 12 | 1, 1, 1, 1, 2, 2 | $0, 0, \frac{3}{4}, \frac{3}{4}, \frac{5}{12}, \frac{2}{3}$ | SB: 6_{-3}^B |
| $6_{-1}^{\zeta_2^1}$ | 12 | 1, 1, 1, 1, 2, 2 | $0, 0, \frac{1}{4}, \frac{1}{4}, \frac{2}{3}, \frac{11}{12}$ | SB: 6_{-1}^B |
| $6_0^{\zeta_2^1}$ | 18 | 1, 1, 2, 2, 2, 2 | $0, 0, 0, 0, \frac{1}{3}, \frac{2}{3}$ | SB: 9_0^B |
| $6_0^{\zeta_2^1}$ | 18 | 1, 1, 2, 2, 2, 2 | $0, 0, 0, \frac{1}{9}, \frac{4}{9}, \frac{7}{9}$ | SB: 9_0^B |
| $6_0^{\zeta_2^1}$ | 18 | 1, 1, 2, 2, 2, 2 | $0, 0, 0, \frac{2}{9}, \frac{5}{9}, \frac{8}{9}$ | SB: 9_0^B |
| $6_4^{\zeta_2^1}$ | 18 | 1, 1, 2, 2, 2, 2 | $0, 0, \frac{1}{3}, \frac{1}{3}, \frac{2}{3}, \frac{2}{3}$ | SB: 9_4^B |
| $6_{8/7}^{\zeta_2^1}$ | 18.591 | 1, 1, $\zeta_5^1, \zeta_5^1, \zeta_5^2, \zeta_5^2$ | $0, 0, \frac{6}{7}, \frac{6}{7}, \frac{2}{7}, \frac{2}{7}$ | $3_{8/7}^B \boxtimes \text{Rep}(Z_2)$ |
| $6_{-8/7}^{\zeta_2^1}$ | 18.591 | 1, 1, $\zeta_5^1, \zeta_5^1, \zeta_5^2, \zeta_5^2$ | $0, 0, \frac{1}{7}, \frac{1}{7}, \frac{5}{7}, \frac{5}{7}$ | $3_{-8/7}^B \boxtimes \text{Rep}(Z_2)$ |
| $6_{4/5}^{\zeta_2^1}$ | 21.708 | 1, 1, $\zeta_3^1, \zeta_3^1, 2, \zeta_8^4$ | $0, 0, \frac{2}{5}, \frac{2}{5}, \frac{2}{3}, \frac{1}{15}$ | $2_{14/5}^B \boxtimes 3_{-2}^{\zeta_2^1}$ |
| $6_{16/5}^{\zeta_2^1}$ | 21.708 | 1, 1, $\zeta_3^1, \zeta_3^1, 2, \zeta_8^4$ | $0, 0, \frac{3}{5}, \frac{3}{5}, \frac{2}{3}, \frac{4}{15}$ | $2_{-14/5}^B \boxtimes 3_{-2}^{\zeta_2^1}$ |
| $6_{-16/5}^{\zeta_2^1}$ | 21.708 | 1, 1, $\zeta_3^1, \zeta_3^1, 2, \zeta_8^4$ | $0, 0, \frac{2}{5}, \frac{2}{5}, \frac{1}{3}, \frac{11}{15}$ | $2_{14/5}^B \boxtimes 3_2^{\zeta_2^1}$ |
| $6_{-4/5}^{\zeta_2^1}$ | 21.708 | 1, 1, $\zeta_3^1, \zeta_3^1, 2, \zeta_8^4$ | $0, 0, \frac{3}{5}, \frac{3}{5}, \frac{1}{3}, \frac{14}{15}$ | $2_{-14/5}^B \boxtimes 3_2^{\zeta_2^1}$ |

same group G . Thus different irreducible representations can be labeled by the group elements: (q) , $q \in G$. The different symmetry twists are also labeled by the group elements: $[g]$, $g \in G$. More general symmetry twists may carry some charge. We denote such charge carrying symmetry twists by $[g, q]$ where $q \in G$. In fact we can identify (q) as $[1, q]$. Those irreducible representations and charged symmetry twists are particles in the modular extensions of $\text{Rep}(G)$ or $\text{sRep}(G^f)$.

Since the group is abelian, the symmetry twists do not break the symmetry. Thus, we have the following fusion rule

$$[1, q] \otimes [g, q'] = [g, qq'] \quad (\text{B1})$$

This means that $[g, q']$ and $[g, qq']$ differ by charge q . We also have

$$[g, q] \otimes [g', q'] = [gg', qq'] \quad (\text{B2})$$

TABLE XXV. Z_3 -SET orders for bosonic systems labeled in terms of sets of topological excitations. The list contains all topological orders with $N = 4, 5, 6$ $D^2 \leq 100$, $N = 7$ $D^2 \leq 60$, $N = 8$ $D^2 \leq 40$, and $N = 9$ $D^2 \leq 28$.

| $N_c^{ \Theta }$ | D^2 | d_1, d_2, \dots | s_1, s_2, \dots | comment |
|-------------------------|--------|---|---|---|
| $3_0^{\zeta_4^1}$ | 3 | 1, 1, 1 | 0, 0, 0 | $\mathcal{E} = \text{Rep}(Z_3)$ |
| $4_4^{\zeta_4^1}$ | 12 | 1, 1, 1, 3 | $0, 0, 0, \frac{1}{2}$ | $\text{SB}: K = \begin{pmatrix} 2 & 1 & 1 & 1 \\ 1 & 2 & 0 & 0 \\ 1 & 0 & 2 & 0 \\ 1 & 0 & 0 & 2 \end{pmatrix}$ |
| $6_1^{\zeta_4^1}$ | 6 | 1, 1, 1, 1, 1, 1 | $0, 0, 0, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}$ | $2_1^B \boxtimes \text{Rep}(Z_3)$ |
| $6_{-1}^{\zeta_4^1}$ | 6 | 1, 1, 1, 1, 1, 1 | $0, 0, 0, \frac{3}{4}, \frac{3}{4}, \frac{3}{4}$ | $2_{-1}^B \boxtimes \text{Rep}(Z_3)$ |
| $6_{14/5}^{\zeta_4^1}$ | 10.854 | $1, 1, 1, \zeta_3^1, \zeta_3^1, \zeta_3^1$ | $0, 0, 0, \frac{2}{5}, \frac{2}{5}, \frac{2}{5}$ | $2_{14/5}^B \boxtimes \text{Rep}(Z_3)$ |
| $6_{-14/5}^{\zeta_4^1}$ | 10.854 | $1, 1, 1, \zeta_3^1, \zeta_3^1, \zeta_3^1$ | $0, 0, 0, \frac{3}{5}, \frac{3}{5}, \frac{3}{5}$ | $2_{-14/5}^B \boxtimes \text{Rep}(Z_3)$ |
| $8_3^{\zeta_4^1}$ | 24 | 1, 1, 1, 1, 1, 1, 3, 3 | $0, 0, 0, \frac{3}{4}, \frac{3}{4}, \frac{3}{4}, \frac{1}{4}, \frac{1}{2}$ | $2_{-1}^B \boxtimes 4_4^{\zeta_4^1}$ |
| $8_{-3}^{\zeta_4^1}$ | 24 | 1, 1, 1, 1, 1, 1, 3, 3 | $0, 0, 0, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{2}, \frac{3}{4}$ | $2_1^B \boxtimes 4_4^{\zeta_4^1}$ |
| $8_{6/5}^{\zeta_4^1}$ | 43.416 | $1, 1, 1, \zeta_3^1, \zeta_3^1, \zeta_3^1, 3, \frac{3+\sqrt{45}}{2}$ | $0, 0, 0, \frac{3}{5}, \frac{3}{5}, \frac{3}{5}, \frac{1}{2}, \frac{1}{10}$ | $2_{-14/5}^B \boxtimes 4_4^{\zeta_4^1}$ |
| $8_{-6/5}^{\zeta_4^1}$ | 43.416 | $1, 1, 1, \zeta_3^1, \zeta_3^1, \zeta_3^1, 3, \frac{3+\sqrt{45}}{2}$ | $0, 0, 0, \frac{2}{5}, \frac{2}{5}, \frac{2}{5}, \frac{1}{2}, \frac{9}{10}$ | $2_{14/5}^B \boxtimes 4_4^{\zeta_4^1}$ |
| $9_2^{\zeta_4^1}$ | 9 | 1, 1, 1, 1, 1, 1, 1, 1, 1 | $0, 0, 0, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}$ | $\text{SB}: 3_2^B \text{ F}: Z_9$ |
| $9_2^{\zeta_4^1}$ | 9 | 1, 1, 1, 1, 1, 1, 1, 1, 1 | $0, 0, 0, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}$ | $3_2^B \boxtimes \text{Rep}(Z_3) \text{ F}: Z_3 \times Z_3$ |
| $9_{-2}^{\zeta_4^1}$ | 9 | 1, 1, 1, 1, 1, 1, 1, 1, 1 | $0, 0, 0, \frac{2}{3}, \frac{2}{3}, \frac{2}{3}, \frac{2}{3}, \frac{2}{3}, \frac{2}{3}$ | $\text{SB}: 3_{-2}^B \text{ F}: Z_9$ |
| $9_{-2}^{\zeta_4^1}$ | 9 | 1, 1, 1, 1, 1, 1, 1, 1, 1 | $0, 0, 0, \frac{2}{3}, \frac{2}{3}, \frac{2}{3}, \frac{2}{3}, \frac{2}{3}, \frac{2}{3}$ | $3_{-2}^B \boxtimes \text{Rep}(Z_3) \text{ F}: Z_3 \times Z_3$ |
| $9_{1/2}^{\zeta_4^1}$ | 12 | $1, 1, 1, 1, 1, 1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{16}, \frac{1}{16}, \frac{1}{16}$ | $3_{1/2}^B \boxtimes \text{Rep}(Z_3)$ |
| $9_{3/2}^{\zeta_4^1}$ | 12 | $1, 1, 1, 1, 1, 1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{3}{16}, \frac{3}{16}, \frac{3}{16}$ | $3_{3/2}^B \boxtimes \text{Rep}(Z_3)$ |
| $9_{5/2}^{\zeta_4^1}$ | 12 | $1, 1, 1, 1, 1, 1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{5}{16}, \frac{5}{16}, \frac{5}{16}$ | $3_{5/2}^B \boxtimes \text{Rep}(Z_3)$ |
| $9_{7/2}^{\zeta_4^1}$ | 12 | $1, 1, 1, 1, 1, 1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{7}{16}, \frac{7}{16}, \frac{7}{16}$ | $3_{7/2}^B \boxtimes \text{Rep}(Z_3)$ |
| $9_{-7/2}^{\zeta_4^1}$ | 12 | $1, 1, 1, 1, 1, 1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{9}{16}, \frac{9}{16}, \frac{9}{16}$ | $3_{-7/2}^B \boxtimes \text{Rep}(Z_3)$ |
| $9_{-5/2}^{\zeta_4^1}$ | 12 | $1, 1, 1, 1, 1, 1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{11}{16}, \frac{11}{16}, \frac{11}{16}$ | $3_{-5/2}^B \boxtimes \text{Rep}(Z_3)$ |
| $9_{-3/2}^{\zeta_4^1}$ | 12 | $1, 1, 1, 1, 1, 1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{13}{16}, \frac{13}{16}, \frac{13}{16}$ | $3_{-3/2}^B \boxtimes \text{Rep}(Z_3)$ |
| $9_{-1/2}^{\zeta_4^1}$ | 12 | $1, 1, 1, 1, 1, 1, \zeta_2^1, \zeta_2^1, \zeta_2^1$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{15}{16}, \frac{15}{16}, \frac{15}{16}$ | $3_{-1/2}^B \boxtimes \text{Rep}(Z_3)$ |
| $9_{8/7}^{\zeta_4^1}$ | 27.887 | $1, 1, 1, \zeta_5^1, \zeta_5^1, \zeta_5^1, \zeta_5^2, \zeta_5^2, \zeta_5^2$ | $0, 0, 0, \frac{6}{7}, \frac{6}{7}, \frac{6}{7}, \frac{2}{7}, \frac{2}{7}, \frac{2}{7}$ | $3_{8/7}^B \boxtimes \text{Rep}(Z_3)$ |
| $9_{-8/7}^{\zeta_4^1}$ | 27.887 | $1, 1, 1, \zeta_5^1, \zeta_5^1, \zeta_5^1, \zeta_5^2, \zeta_5^2, \zeta_5^2$ | $0, 0, 0, \frac{1}{7}, \frac{1}{7}, \frac{1}{7}, \frac{5}{7}, \frac{5}{7}, \frac{5}{7}$ | $3_{-8/7}^B \boxtimes \text{Rep}(Z_3)$ |

However, the above fusion rule is too restrictive. Although $[g, q']$ and $[g, qq']$ differ by charge q , we do not know the net charge of $[g, q']$ when $g \neq 1$. Thus the more general fusion rule that still preserves charge conservation is

$$[g, q] \otimes [g', q'] = [gg', \omega_2(g, g')qq'], \quad \omega_2(g, g') \in G. \quad (\text{B3})$$

From

$$\begin{aligned} & ([g_1, q_1] \otimes [g_2, q_2]) \otimes [g_3, q_3] \\ &= [g_1 g_2 g_3, \omega(g_1, g_2)\omega(g_1 g_2, g_3)q_1 q_2 q_3] \\ &= [g_1, q_1] \otimes ([g_2, q_2] \otimes [g_3, q_3]) \\ &= [g_1 g_2 g_3, \omega(g_1, g_2 g_3)\omega(g_2, g_3)q_1 q_2 q_3] \end{aligned} \quad (\text{B4})$$

we see that

$$\omega(g_1, g_2)\omega(g_1 g_2, g_3) = \omega(g_1, g_2 g_3)\omega(g_2, g_3). \quad (\text{B5})$$

i.e. $\omega(g_1, g_2)$ is a group 2-cocycle in $\mathcal{H}^2(G, G)$.

In the above, we have assumed that the modular extension is abelian (*i.e.* all the particles in the modular extension have a quantum dimension 1). We see that the fusion rules of abelian modular extensions are labeled by 2-cocycles in $\mathcal{H}^2(G, G)$.

However, sometimes the modular extension can be non-abelian, such as the modular extension of $\text{sRep}(Z_2^f)$ and $\text{Rep}(Z_2 \times Z_2 \times Z_2)$. To allow such a possibility, we allow $[g, q]$ to be a many-to-one label of the particle, and define a subgroup $H_g \subset G$:

$$H_g = \{h | [g, q] = [g, hq], h \in G\}. \quad (\text{B6})$$

The mapping $g \rightarrow H_g$ is an important data to describe the fusion. H_g represents the charge ambiguity of the symmetry twist $[g, q]$. To get an one-to-one label, we can use

$$[g, qH_g]. \quad (\text{B7})$$

Note that, when g is an identity: $g = 1$, H_g is trivial: $H_1 = 1$.

TABLE XXVI. The fusion rules for the three $5_4^{\sqrt{6}}$ entries in Table XXVII. The three entries have identical (d_i, s_i) but different fusions rules. $\mathbf{1}$, a , b are the three irreducible representations of S_3 with dimension 1, 1, 2.

| | | | | | |
|------------------|--------------|--------------|--------------------------------|--------------------------------------|--------------------------------------|
| s_i | 0 | 0 | 0 | $\frac{1}{2}$ | $\frac{1}{2}$ |
| d_i | 1 | 1 | 2 | 3 | 3 |
| $5_4^{\sqrt{6}}$ | $\mathbf{1}$ | a | b | σ | τ |
| $\mathbf{1}$ | $\mathbf{1}$ | a | b | σ | τ |
| a | a | $\mathbf{1}$ | b | τ | σ |
| b | b | b | $\mathbf{1} \oplus a \oplus b$ | $\sigma \oplus \tau$ | $\sigma \oplus \tau$ |
| σ | σ | τ | $\sigma \oplus \tau$ | $\mathbf{1} \oplus b \oplus 2\sigma$ | $a \oplus b \oplus 2\tau$ |
| τ | τ | σ | $\sigma \oplus \tau$ | $a \oplus b \oplus 2\tau$ | $\mathbf{1} \oplus b \oplus 2\sigma$ |

| | | | | | |
|------------------|--------------|--------------|--------------------------------|---|---|
| s_i | 0 | 0 | 0 | $\frac{1}{2}$ | $\frac{1}{2}$ |
| d_i | 1 | 1 | 2 | 3 | 3 |
| $5_4^{\sqrt{6}}$ | $\mathbf{1}$ | a | b | σ | τ |
| $\mathbf{1}$ | $\mathbf{1}$ | a | b | σ | τ |
| a | a | $\mathbf{1}$ | b | τ | σ |
| b | b | b | $\mathbf{1} \oplus a \oplus b$ | $\sigma \oplus \tau$ | $\sigma \oplus \tau$ |
| σ | σ | τ | $\sigma \oplus \tau$ | $\mathbf{1} \oplus b \oplus \sigma \oplus \tau$ | $a \oplus b \oplus \sigma \oplus \tau$ |
| τ | τ | σ | $\sigma \oplus \tau$ | $a \oplus b \oplus \sigma \oplus \tau$ | $\mathbf{1} \oplus b \oplus \sigma \oplus \tau$ |

| | | | | | |
|------------------|--------------|--------------|--------------------------------|---|---|
| s_i | 0 | 0 | 0 | $\frac{1}{2}$ | $\frac{1}{2}$ |
| d_i | 1 | 1 | 2 | 3 | 3 |
| $5_4^{\sqrt{6}}$ | $\mathbf{1}$ | a | b | σ | τ |
| $\mathbf{1}$ | $\mathbf{1}$ | a | b | σ | τ |
| a | a | $\mathbf{1}$ | b | τ | σ |
| b | b | b | $\mathbf{1} \oplus a \oplus b$ | $\sigma \oplus \tau$ | $\sigma \oplus \tau$ |
| σ | σ | τ | $\sigma \oplus \tau$ | $a \oplus b \oplus \sigma \oplus \tau$ | $\mathbf{1} \oplus b \oplus \sigma \oplus \tau$ |
| τ | τ | σ | $\sigma \oplus \tau$ | $\mathbf{1} \oplus b \oplus \sigma \oplus \tau$ | $a \oplus b \oplus \sigma \oplus \tau$ |

The fusion of $[1, q']$ and $[g, qH_g]$ is still given by

$$[1, q'] \otimes [g, qH_g] = [g, q'qH_g]. \quad (\text{B8})$$

We also have $H_g = H_{g^{-1}}$ and

$$[g, qH_g] \otimes [g^{-1}, q'H_g] = \oplus_{h \in qq'H_g} [1, h] \quad (\text{B9})$$

We see that the quantum dimension of $[g, qH_g]$ is $d = \sqrt{|H_g|}$.

The fusion rule should satisfy

$$\begin{aligned} & [1, q] \otimes ([g_1, q_1 H_{g_1}] \otimes [g_2, q_2 H_{g_2}]) \\ &= ([1, q] \otimes [g_1, q_1 H_{g_1}]) \otimes [g_2, q_2 H_{g_2}] \\ &= [g_1, q_1 H_{g_1}] \otimes ([1, q] \otimes [g_2, q_2 H_{g_2}]) \end{aligned} \quad (\text{B10})$$

We find that the following ansatz satisfy the above condition

$$\begin{aligned} [g_1, q_1 H_{g_1}] \otimes [g_2, q_2 H_{g_2}] &= \frac{m^{g_1 g_2}}{|(H_{g_1} \vee H_{g_2}) \cap H_{g_1 g_2}|} \\ &\oplus_{q \in \omega(g_1, g_2) q_1 q_2 H_{g_1} \vee H_{g_2}} [g_1 g_2, q H_{g_1 g_2}] \end{aligned} \quad (\text{B11})$$

where $m^{g_1 g_2} \in \mathbb{Z}$ and $H_{g_1} \vee H_{g_2}$ is the subgroup generated by H_{g_1} and H_{g_2} . The above implies that

$$\sqrt{|H_{g_1}|} \sqrt{|H_{g_2}|} = m^{g_1 g_2} \frac{|H_{g_1} \vee H_{g_2}|}{|(H_{g_1} \vee H_{g_2}) \cap H_{g_1 g_2}|} \sqrt{|H_{g_1 g_2}|} \quad (\text{B12})$$

We see that different fusion rules are labeled by $\omega(g_1, g_2)$ and H_g .

It is much easier to find all the H_g 's that satisfy eqn. (B12) and all the $\omega(g_1, g_2)$ that satisfy eqn. (B5). From those solutions, we can directly construct the fusion rule from eqn. (B11).

Appendix C: Conditions to obtain UMTC/ \mathcal{E} 's

In our simplified theory, a UMTC/ \mathcal{E} is described by an integer tensor N_k^{ij} and a mod-1 rational vector s_i , where i, j, k run from 1 to N and N is called the rank of the UMTC/ \mathcal{E} . We may simply denote a UMTC/ \mathcal{E} (the collection of data (N_k^{ij}, s_i)) by \mathcal{C} , a particle i in \mathcal{C} by $i \in \mathcal{C}$. Sometimes it is more convenient to use abstract labels rather than 1 to N ; we may also abuse \mathcal{C} as the set of labels (particles).

Not all (N_k^{ij}, s_i) describe a valid UMTC/ \mathcal{E} \mathcal{C} with modular extensions. In order to describe a valid \mathcal{C} , (N_k^{ij}, s_i) must satisfy the following conditions: [13,43–46](#)

1. Fusion ring:

N_k^{ij} for the UMTC/ \mathcal{E} \mathcal{C} are non-negative integers that satisfy

$$\begin{aligned} N_k^{ij} &= N_k^{ji}, \quad N_j^{1i} = \delta_{ij}, \quad \sum_{k=1}^N N_1^{ik} N_1^{kj} = \delta_{ij}, \quad (\text{C1}) \\ \sum_m N_m^{ij} N_l^{mk} &= \sum_n N_l^{in} N_n^{jk} \text{ or } \sum_m N_m^{ij} N_m = N_i N_j \end{aligned}$$

where the matrix N_i is given by $(N_i)_{kj} = N_k^{ij}$, and the indices i, j, k run from 1 to N . In fact N_1^{ij} defines a charge conjugation $i \rightarrow \bar{i}$:

$$N_1^{ij} = \delta_{\bar{i}j}. \quad (\text{C2})$$

N_k^{ij} satisfying the above conditions define a fusion ring which is viewed as the set (of simple objects)

$$\{1, 2, \dots, N\}. \quad (\text{C3})$$

2. Charge conjugation condition:

$$\begin{aligned} N_k^{ij} &= N_{\bar{i}}^{j\bar{k}} = N_{\bar{j}}^{\bar{k}i} \\ &= N_j^{\bar{i}k} = N_i^{k\bar{j}} = N_{\bar{k}}^{\bar{j}\bar{i}}. \end{aligned} \quad (\text{C4})$$

TABLE XXVII. S_3 -SET orders for bosonic systems labeled in terms of sets of topological excitations. The list contains all topological orders with $N = 4, 5, 6$ $D^2 \leq 100$, $N = 7$ $D^2 \leq 60$, and $N = 8$ $D^2 \leq 40$. (In fact, we fail to find any bosonic S_3 -SET orders with $N = 4, 7, 8$.)

| $N_c^{ \Theta }$ | D^2 | d_1, d_2, \dots | s_1, s_2, \dots | comment |
|------------------------|--------|--|--|--|
| $3_0^{\sqrt{6}}$ | 6 | 1, 1, 2 | 0, 0, 0 | $\mathcal{E} = \text{Rep}(S_3)$ |
| $5_4^{\sqrt{6}}$ | 24 | 1, 1, 2, 3, 3 | $0, 0, 0, \frac{1}{2}, \frac{1}{2}$ | SB: 4_4^B |
| $5_4^{\sqrt{6}}$ | 24 | 1, 1, 2, 3, 3 | $0, 0, 0, \frac{1}{2}, \frac{1}{2}$ | SB: $4_4^B \begin{pmatrix} 2 & 1 & 1 & 1 \\ 1 & 2 & 0 & 0 \\ 1 & 0 & 2 & 0 \\ 1 & 0 & 0 & 2 \end{pmatrix}$ |
| $5_4^{\sqrt{6}}$ | 24 | 1, 1, 2, 3, 3 | $0, 0, 0, \frac{1}{2}, \frac{1}{2}$ | SB: 4_4^B |
| $6_1^{\sqrt{6}}$ | 12 | 1, 1, 2, 1, 1, 2 | $0, 0, 0, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}$ | $2_1^B \boxtimes \text{Rep}(S_3)$ |
| $6_1^{\sqrt{6}}$ | 12 | 1, 1, 2, 1, 1, 2 | $0, 0, 0, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}$ | SB: 2_1^B |
| $6_{-1}^{\sqrt{6}}$ | 12 | 1, 1, 2, 1, 1, 2 | $0, 0, 0, \frac{3}{4}, \frac{3}{4}, \frac{3}{4}$ | $2_{-1}^B \boxtimes \text{Rep}(S_3)$ |
| $6_{-1}^{\sqrt{6}}$ | 12 | 1, 1, 2, 1, 1, 2 | $0, 0, 0, \frac{3}{4}, \frac{3}{4}, \frac{3}{4}$ | SB: 2_{-1}^B |
| $6_2^{\sqrt{6}}$ | 18 | 1, 1, 2, 2, 2, 2 | $0, 0, 0, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}$ | SB: 3_2^B |
| $6_2^{\sqrt{6}}$ | 18 | 1, 1, 2, 2, 2, 2 | $0, 0, 0, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}$ | SB: 3_2^B |
| $6_{-2}^{\sqrt{6}}$ | 18 | 1, 1, 2, 2, 2, 2 | $0, 0, 0, \frac{2}{3}, \frac{2}{3}, \frac{2}{3}$ | SB: 3_{-2}^B |
| $6_{-2}^{\sqrt{6}}$ | 18 | 1, 1, 2, 2, 2, 2 | $0, 0, 0, \frac{2}{3}, \frac{2}{3}, \frac{2}{3}$ | SB: 3_{-2}^B |
| $6_{14/5}^{\sqrt{6}}$ | 21.708 | $1, 1, 2, \zeta_3^1, \zeta_3^1, \zeta_8^4$ | $0, 0, 0, \frac{2}{5}, \frac{2}{5}, \frac{2}{5}$ | $2_{14/5}^B \boxtimes \text{Rep}(S_3)$ |
| $6_{-14/5}^{\sqrt{6}}$ | 21.708 | $1, 1, 2, \zeta_3^1, \zeta_3^1, \zeta_8^4$ | $0, 0, 0, \frac{3}{5}, \frac{3}{5}, \frac{3}{5}$ | $2_{-14/5}^B \boxtimes \text{Rep}(S_3)$ |

3. Rational condition:

N_k^{ij} and s_i for \mathcal{C} satisfy^{45,47–49}

$$\sum_r V_{ijkl}^r s_r = 0 \bmod 1 \quad (\text{C5})$$

where

$$V_{ijkl}^r = N_r^{ij} N_{\bar{r}}^{kl} + N_r^{il} N_{\bar{r}}^{jk} + N_r^{ik} N_{\bar{r}}^{jl} - (\delta_{ir} + \delta_{jr} + \delta_{kr} + \delta_{lr}) \sum_m N_m^{ij} N_{\bar{m}}^{kl} \quad (\text{C6})$$

4. Verlinde fusion characters:

Let the topological S -matrix be [see eqn. (223) in Ref. 9]

$$S_{ij} = \frac{1}{D} \sum_k N_k^{ij} e^{2\pi i (s_i + s_j - s_k)} d_k, \quad (\text{C7})$$

where d_i (called quantum dimension) is the largest eigenvalue of the matrix N_i and $D = \sqrt{\sum_i d_i^2}$ (called the total quantum dimension). Then⁵⁰:

$$\frac{S_{il} S_{jl}}{S_{1l}} = \sum_k N_k^{ij} S_{kl}. \quad (\text{C8})$$

5. Weak modularity:

Let the topological T -matrix be

$$T_{ij} = \delta_{ij} e^{2\pi i s_i}. \quad (\text{C9})$$

Then [see eqn. (232) in Ref. 9]

$$S^\dagger T S = \Theta T^\dagger S^\dagger T^\dagger, \quad \Theta = D^{-1} \sum_i e^{2\pi i s_i} d_i^2 = |\Theta| e^{2\pi i c/8}. \quad (\text{C10})$$

The parameter $c \bmod 8$ is defined via Θ , if $|\Theta| \neq 0$.

6. Charge conjugation symmetry:

$$S_{ij} = S_{i\bar{j}}^*, \quad s_i = s_{\bar{i}}, \quad \text{or } S = S^\dagger C, \quad T = TC, \quad (\text{C11})$$

where the charge conjugation matrix C is given by $C_{ij} = N_1^{ij} = \delta_{i\bar{j}}$.

7. The centralizer describes the symmetry:

Let the centralizer of \mathcal{C} , $\mathcal{C}_{\mathcal{C}}^{\text{cen}}$, be the subset of the particle labels:

$$\mathcal{C}_{\mathcal{C}}^{\text{cen}} = \{i \mid S_{ij} = \frac{d_i d_j}{D}, \forall j \in \mathcal{C}\}. \quad (\text{C12})$$

Then, $\mathcal{C}_{\mathcal{C}}^{\text{cen}} = \mathcal{E}$.

8. The second Frobenius-Schur indicator:

Let

$$\nu_k = D^{-2} \sum_{ij} N_k^{ij} d_i d_j \cos(4\pi(s_i - s_j)), \quad (\text{C13})$$

then $\nu_k \in \mathbb{Z}$ if $k = \bar{k}$ ⁵¹.

9. Symmetry breaking:

There is a symmetry breaking induced map $\mathcal{C} \rightarrow \mathcal{C}_0$, where \mathcal{C}_0 is a UMTC if $\mathcal{E} = \text{Rep}(G)$ or a UMTC/ $\text{sRep}(Z_2)$ if $\mathcal{E} = \text{sRep}(G^f)$. See Appendix D for details.

10. Modular extension:

The UMTC/ \mathcal{E} \mathcal{C} has modular extensions.

The above conditions are necessary and sufficient (due to the condition 10) for (N_k^{ij}, s_i) to describe a UMTC/ \mathcal{E} \mathcal{C} with modular extensions.

TABLE XXVIII. S_3 -SET orders for bosonic systems labeled in terms of sets of topological excitations. The list contains all topological orders with $N = 9$ $D^2 \leq 30$.

| $N_c^{ \Theta }$ | D^2 | d_1, d_2, \dots | s_1, s_2, \dots | comment |
|-----------------------|--------|---|---|--|
| $3_0^{\sqrt{6}}$ | 6 | 1, 1, 2 | 0, 0, 0 | $\mathcal{E} = \text{Rep}(S_3)$ |
| $9_2^{\sqrt{6}}$ | 18 | 1, 1, 2, 1, 1, 1, 1, 2, 2 | $0, 0, 0, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}$ | $3_2^B \boxtimes \text{Rep}(S_3)$ |
| $9_{-2}^{\sqrt{6}}$ | 18 | 1, 1, 2, 1, 1, 1, 1, 2, 2 | $0, 0, 0, \frac{2}{3}, \frac{2}{3}, \frac{2}{3}, \frac{2}{3}, \frac{2}{3}, \frac{2}{3}$ | $3_{-2}^B \boxtimes \text{Rep}(S_3)$ |
| $9_0^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, 2, 2, 2, 2 | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, 0, \frac{1}{2}$ | SB:4 ₀ ^B |
| $9_0^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, 2, 2, 2, 2 | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, 0, \frac{1}{2}$ | SB:4 ₀ ^B |
| $9_1^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, 2, 2, 2, 2 | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}, \frac{1}{8}, \frac{1}{8}, \frac{1}{2}$ | SB:4 ₁ ^B |
| $9_1^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, 2, 2, 2, 2 | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}, \frac{1}{8}, \frac{1}{8}, \frac{1}{2}$ | SB:4 ₁ ^B |
| $9_2^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, 2, 2, 2, 2 | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{2}$ | SB:4 ₂ ^B |
| $9_2^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, 2, 2, 2, 2 | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{2}$ | SB:4 ₂ ^B |
| $9_3^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, 2, 2, 2, 2 | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{8}, \frac{3}{8}, \frac{3}{8}, \frac{1}{2}$ | SB:4 ₃ ^B |
| $9_3^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, 2, 2, 2, 2 | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{8}, \frac{3}{8}, \frac{3}{8}, \frac{1}{2}$ | SB:4 ₃ ^B |
| $9_4^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, 2, 2, 2, 2 | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}$ | SB:4 ₄ ^B |
| $9_4^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, 2, 2, 2, 2 | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \frac{1}{2}$ | SB:4 ₄ ^B |
| $9_{-3}^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, 2, 2, 2, 2 | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{5}{8}, \frac{5}{8}, \frac{5}{8}, \frac{1}{2}$ | SB:4 ₋₃ ^B |
| $9_{-3}^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, 2, 2, 2, 2 | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{5}{8}, \frac{5}{8}, \frac{5}{8}, \frac{1}{2}$ | SB:4 ₋₃ ^B |
| $9_{-2}^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, 2, 2, 2, 2 | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{4}, \frac{3}{4}, \frac{3}{4}, \frac{1}{2}$ | SB:4 ₋₂ ^B |
| $9_{-2}^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, 2, 2, 2, 2 | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{4}, \frac{3}{4}, \frac{3}{4}, \frac{1}{2}$ | SB:4 ₋₂ ^B |
| $9_{-1}^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, 2, 2, 2, 2 | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{7}{8}, \frac{7}{8}, \frac{7}{8}, \frac{1}{2}$ | SB:4 ₋₁ ^B |
| $9_{-1}^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, 2, 2, 2, 2 | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{7}{8}, \frac{7}{8}, \frac{7}{8}, \frac{1}{2}$ | SB:4 ₋₁ ^B |
| $9_{5/2}^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, $\zeta_2^1, \zeta_2^1, 2, \sqrt{8}$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{5}{16}, \frac{5}{16}, \frac{1}{2}, \frac{5}{16}$ | $3_{5/2}^B \boxtimes \text{Rep}(S_3)$ |
| $9_{5/2}^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, $\zeta_2^1, \zeta_2^1, 2, \sqrt{8}$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{5}{16}, \frac{5}{16}, \frac{1}{2}, \frac{5}{16}$ | SB:3 _{5/2} ^B |
| $9_{1/2}^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, $\zeta_2^1, \zeta_2^1, 2, \sqrt{8}$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{16}, \frac{1}{16}, \frac{1}{2}, \frac{1}{16}$ | $3_{1/2}^B \boxtimes \text{Rep}(S_3)$ |
| $9_{1/2}^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, $\zeta_2^1, \zeta_2^1, 2, \sqrt{8}$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{16}, \frac{1}{16}, \frac{1}{2}, \frac{1}{16}$ | SB:3 _{1/2} ^B |
| $9_{3/2}^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, $\zeta_2^1, \zeta_2^1, 2, \sqrt{8}$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{16}, \frac{3}{16}, \frac{1}{2}, \frac{3}{16}$ | $3_{3/2}^B \boxtimes \text{Rep}(S_3)$ |
| $9_{3/2}^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, $\zeta_2^1, \zeta_2^1, 2, \sqrt{8}$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{16}, \frac{3}{16}, \frac{1}{2}, \frac{3}{16}$ | SB:3 _{3/2} ^B |
| $9_{7/2}^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, $\zeta_2^1, \zeta_2^1, 2, \sqrt{8}$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{7}{16}, \frac{7}{16}, \frac{1}{2}, \frac{7}{16}$ | $3_{7/2}^B \boxtimes \text{Rep}(S_3)$ |
| $9_{7/2}^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, $\zeta_2^1, \zeta_2^1, 2, \sqrt{8}$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{7}{16}, \frac{7}{16}, \frac{1}{2}, \frac{7}{16}$ | SB:3 _{7/2} ^B |
| $9_{-7/2}^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, $\zeta_2^1, \zeta_2^1, 2, \sqrt{8}$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{9}{16}, \frac{9}{16}, \frac{1}{2}, \frac{9}{16}$ | $3_{-7/2}^B \boxtimes \text{Rep}(S_3)$ |
| $9_{-7/2}^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, $\zeta_2^1, \zeta_2^1, 2, \sqrt{8}$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{9}{16}, \frac{9}{16}, \frac{1}{2}, \frac{9}{16}$ | SB:3 _{-7/2} ^B |
| $9_{-5/2}^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, $\zeta_2^1, \zeta_2^1, 2, \sqrt{8}$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{11}{16}, \frac{11}{16}, \frac{1}{2}, \frac{11}{16}$ | $3_{-5/2}^B \boxtimes \text{Rep}(S_3)$ |
| $9_{-5/2}^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, $\zeta_2^1, \zeta_2^1, 2, \sqrt{8}$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{11}{16}, \frac{11}{16}, \frac{1}{2}, \frac{11}{16}$ | SB:3 _{-5/2} ^B |
| $9_{-3/2}^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, $\zeta_2^1, \zeta_2^1, 2, \sqrt{8}$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{13}{16}, \frac{13}{16}, \frac{1}{2}, \frac{13}{16}$ | $3_{-3/2}^B \boxtimes \text{Rep}(S_3)$ |
| $9_{-3/2}^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, $\zeta_2^1, \zeta_2^1, 2, \sqrt{8}$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{13}{16}, \frac{13}{16}, \frac{1}{2}, \frac{13}{16}$ | SB:3 _{-3/2} ^B |
| $9_{-1/2}^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, $\zeta_2^1, \zeta_2^1, 2, \sqrt{8}$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{15}{16}, \frac{15}{16}, \frac{1}{2}, \frac{15}{16}$ | $3_{-1/2}^B \boxtimes \text{Rep}(S_3)$ |
| $9_{-1/2}^{\sqrt{6}}$ | 24 | 1, 1, 2, 1, 1, $\zeta_2^1, \zeta_2^1, 2, \sqrt{8}$ | $0, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{15}{16}, \frac{15}{16}, \frac{1}{2}, \frac{15}{16}$ | SB:3 _{-1/2} ^B |
| $9_0^{\sqrt{6}}$ | 30 | 1, 1, 2, 2, 2, 2, 2, 2, 2 | $0, 0, 0, \frac{1}{5}, \frac{1}{5}, \frac{1}{5}, \frac{4}{5}, \frac{4}{5}, \frac{4}{5}$ | SB:5 ₀ ^B |
| $9_4^{\sqrt{6}}$ | 30 | 1, 1, 2, 2, 2, 2, 2, 2, 2 | $0, 0, 0, \frac{2}{5}, \frac{2}{5}, \frac{2}{5}, \frac{3}{5}, \frac{3}{5}, \frac{3}{5}$ | SB:5 ₄ ^B |
| $9_{8/7}^{\sqrt{6}}$ | 55.775 | 1, 1, 2, $\zeta_5^1, \zeta_5^1, \zeta_5^2, \zeta_5^2, 2\zeta_5^1, \zeta_{12}^6$ | $0, 0, 0, \frac{6}{7}, \frac{6}{7}, \frac{2}{7}, \frac{2}{7}, \frac{6}{7}, \frac{2}{7}$ | $3_{8/7}^B \boxtimes \text{Rep}(S_3)$ |
| $9_{-8/7}^{\sqrt{6}}$ | 55.775 | 1, 1, 2, $\zeta_5^1, \zeta_5^1, \zeta_5^2, \zeta_5^2, 2\zeta_5^1, \zeta_{12}^6$ | $0, 0, 0, \frac{1}{7}, \frac{1}{7}, \frac{5}{7}, \frac{5}{7}, \frac{1}{7}, \frac{5}{7}$ | $3_{-8/7}^B \boxtimes \text{Rep}(S_3)$ |

However, when we calculate the tables in Appendix A, we do not use the condition 10. So the used conditions are only necessary. As a result, the tables may contain fake entries that have no modular extensions.

To numerically solve the above conditions to obtain the classification tables, we first search for N_k^{ij} 's that satisfy

the condition 1 and 2. Then for each N_k^{ij} , we calculate s_i 's that satisfy the condition 3 via the Smith normal form of integer matrix V_{ijkl}^r , where $ijkl$ is viewed as a single index. Last, from the obtained N_k^{ij} , s_i 's, we select those that satisfy all the conditions.

TABLE XXIX. $Z_2 \times Z_2$ -SET orders for bosonic systems labeled in terms of sets of topological excitations. The list contains all topological orders with $N = 5$ $D^2 \leq 100$ and $N = 6$ $D^2 \leq 200$.

| $N_c^{ \Theta }$ | D^2 | d_1, d_2, \dots | s_1, s_2, \dots | comment |
|------------------|--------|--|--|--|
| 4_0^2 | 4 | 1, 1, 1, 1 | 0, 0, 0, 0 | $\mathcal{E} = \text{Rep}(Z_2 \times Z_2)$ |
| 5_1^2 | 8 | 1, 1, 1, 1, 2 | 0, 0, 0, 0, $\frac{1}{4}$ | SB: 2_1^B |
| 5_{-1}^2 | 8 | 1, 1, 1, 1, 2 | 0, 0, 0, 0, $\frac{3}{4}$ | SB: 2_{-1}^B |
| $5_{14/5}^2$ | 14.472 | 1, 1, 1, 1, ζ_8^4 | 0, 0, 0, 0, $\frac{2}{5}$ | SB: $2_{14/5}^B$ |
| $5_{-14/5}^2$ | 14.472 | 1, 1, 1, 1, ζ_8^4 | 0, 0, 0, 0, $\frac{3}{5}$ | SB: $2_{-14/5}^B$ |
| 6_2^2 | 12 | 1, 1, 1, 1, 2, 2 | 0, 0, 0, 0, $\frac{1}{3}, \frac{1}{3}$ | SB: 3_2^B |
| 6_2^2 | 12 | 1, 1, 1, 1, 2, 2 | 0, 0, 0, 0, $\frac{1}{3}, \frac{1}{3}$ | SB: 3_2^B |
| 6_2^2 | 12 | 1, 1, 1, 1, 2, 2 | 0, 0, 0, 0, $\frac{1}{3}, \frac{1}{3}$ | SB: 3_2^B |
| 6_2^2 | 12 | 1, 1, 1, 1, 2, 2 | 0, 0, 0, 0, $\frac{1}{3}, \frac{1}{3}$ | SB: 3_2^B |
| 6_{-2}^2 | 12 | 1, 1, 1, 1, 2, 2 | 0, 0, 0, 0, $\frac{2}{3}, \frac{2}{3}$ | SB: 3_{-2}^B |
| 6_{-2}^2 | 12 | 1, 1, 1, 1, 2, 2 | 0, 0, 0, 0, $\frac{2}{3}, \frac{2}{3}$ | SB: 3_{-2}^B |
| 6_{-2}^2 | 12 | 1, 1, 1, 1, 2, 2 | 0, 0, 0, 0, $\frac{2}{3}, \frac{2}{3}$ | SB: 3_{-2}^B |
| 6_{-2}^2 | 12 | 1, 1, 1, 1, 2, 2 | 0, 0, 0, 0, $\frac{2}{3}, \frac{2}{3}$ | SB: 3_{-2}^B |
| $6_{1/2}^2$ | 16 | 1, 1, 1, 1, 2, $\sqrt{8}$ | 0, 0, 0, 0, $\frac{1}{2}, \frac{1}{16}$ | SB: $3_{1/2}^B$ |
| $6_{3/2}^2$ | 16 | 1, 1, 1, 1, 2, $\sqrt{8}$ | 0, 0, 0, 0, $\frac{1}{2}, \frac{3}{16}$ | SB: $3_{3/2}^B$ |
| $6_{5/2}^2$ | 16 | 1, 1, 1, 1, 2, $\sqrt{8}$ | 0, 0, 0, 0, $\frac{1}{2}, \frac{5}{16}$ | SB: $3_{5/2}^B$ |
| $6_{7/2}^2$ | 16 | 1, 1, 1, 1, 2, $\sqrt{8}$ | 0, 0, 0, 0, $\frac{1}{2}, \frac{7}{16}$ | SB: $3_{7/2}^B$ |
| $6_{-7/2}^2$ | 16 | 1, 1, 1, 1, 2, $\sqrt{8}$ | 0, 0, 0, 0, $\frac{1}{2}, \frac{9}{16}$ | SB: $3_{-7/2}^B$ |
| $6_{-5/2}^2$ | 16 | 1, 1, 1, 1, 2, $\sqrt{8}$ | 0, 0, 0, 0, $\frac{1}{2}, \frac{11}{16}$ | SB: $3_{-5/2}^B$ |
| $6_{-3/2}^2$ | 16 | 1, 1, 1, 1, 2, $\sqrt{8}$ | 0, 0, 0, 0, $\frac{1}{2}, \frac{13}{16}$ | SB: $3_{-3/2}^B$ |
| $6_{-1/2}^2$ | 16 | 1, 1, 1, 1, 2, $\sqrt{8}$ | 0, 0, 0, 0, $\frac{1}{2}, \frac{15}{16}$ | SB: $3_{-1/2}^B$ |
| 6_4^2 | 36 | 1, 1, 1, 1, 4, 4 | 0, 0, 0, 0, $\frac{1}{3}, \frac{2}{3}$ | SB: 9_4^B |
| $6_{8/7}^2$ | 37.183 | 1, 1, 1, 1, $2\zeta_5^1, \zeta_{12}^6$ | 0, 0, 0, 0, $\frac{6}{7}, \frac{2}{7}$ | SB: $3_{8/7}^B$ |
| $6_{-8/7}^2$ | 37.183 | 1, 1, 1, 1, $2\zeta_5^1, \zeta_{12}^6$ | 0, 0, 0, 0, $\frac{1}{7}, \frac{5}{7}$ | SB: $3_{-8/7}^B$ |

Appendix D: Symmetry breaking

A UMTC/ \mathcal{E} \mathcal{C} describes a SET with symmetry \mathcal{E} (up to invertible GQLs). If we break the symmetry \mathcal{E} , then the UMTC/ \mathcal{E} will become a UMTC \mathcal{C}_0 if $\mathcal{E} = \text{Rep } G$ or become a UMTC/ Z_2^f \mathcal{C}_0 if $\mathcal{E} = \text{sRep } G^f$. So there is a natural mapping from UMTC/ \mathcal{E} 's to UMTCs or UMTC/ Z_2^f : $\mathcal{C} \rightarrow \mathcal{C}_0$. Requiring the existence of such map can give us some additional conditions on (N_k^{ij}, s_i) of \mathcal{C} .

To understand such a map, we note that \mathcal{C} can be viewed as a subcategory of \mathcal{C}_0 , in the sense that the simple objects in \mathcal{C} can be viewed as the simple or composite objects in \mathcal{C}_0 :

$$i \rightarrow \oplus_I M^{iI} I, \quad i \in \mathcal{C}, \quad I \in \mathcal{C}_0. \quad (\text{D1})$$

Physical, if we just pretend the symmetry is not there, then every particle in \mathcal{C} can also be viewed as a particle in \mathcal{C}_0 . However, a particle in \mathcal{C} may be the direct sum of several degenerate particles in \mathcal{C}_0 , where the degeneracy is due to the symmetry, as described by eqn. (D1).

In the following, we will obtain some conditions on M^{iI} , which will help us to calculate it. Let us label the particles in \mathcal{C} as $\{i\} = \{1, a, b, \dots, x, y, \dots\}$. Here a, b, \dots label the *bosonic* part of \mathcal{E} , and x, y, \dots label

the fermionic part of \mathcal{E} (if any) and the rest of non-trivial topological excitations. We have also used I to label the particles in \mathcal{C}_0 . Clearly, the bosonic part of \mathcal{E} are local excitations and are direct sums of $\mathbf{1} \in \mathcal{C}_0$:

$$a \rightarrow d_a \mathbf{1}, \quad \text{or} \quad M^{aI} = d_a \delta_{1I}, \quad (\text{D2})$$

(Here $\mathbf{1}$ is the trivial particle in \mathcal{C}_0 .) By computing $i \otimes j$ in two different ways, we find that M^{iI} must also satisfy

$$\sum_{IJ} M^{iI} M^{jJ} N_K^{IJ} = \sum_k N_k^{ij} M^{kK} \quad (\text{D3})$$

Taking $K = \mathbf{1}$, we obtain

$$\sum_I M^{iI} M^{j\bar{I}} = \sum_a N_a^{ij} d_a \quad (\text{D4})$$

Assuming the charge conjugation symmetry: $M^{iI} = M^{i\bar{I}}$, we can rewrite the above as

$$\sum_I M^{iI} M^{jI} = \sum_a N_a^{i\bar{j}} d_a, \quad (\text{D5})$$

which implies that

$$\sum_I (M^{iI})^2 = \sum_a N_a^{i\bar{i}} d_a. \quad (\text{D6})$$

TABLE XXX. $Z_2 \times Z_2$ -SET orders for bosonic systems labeled in terms of sets of topological excitations. The list contains all topological orders with $N = 7$ $D^2 \leq 120$.

| $N_c^{ \Theta }$ | D^2 | d_1, d_2, \dots | s_1, s_2, \dots | comment |
|------------------|--------|--|---|--|
| 4_0^2 | 4 | 1, 1, 1, 1 | 0, 0, 0, 0 | $\mathcal{E} = \text{Rep}(Z_2 \times Z_2)$ |
| 7_0^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, 0, 0, $\frac{1}{2}$ | SB:4 ₀ ^B |
| 7_0^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, 0, $\frac{1}{4}, \frac{3}{4}$ | SB:4 ₀ ^B |
| 7_1^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{1}{8}, \frac{1}{8}, \frac{1}{2}$ | SB:4 ₁ ^B |
| 7_1^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{1}{8}, \frac{1}{8}, \frac{1}{2}$ | SB:4 ₁ ^B |
| 7_1^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{1}{8}, \frac{1}{8}, \frac{1}{2}$ | SB:4 ₁ ^B |
| 7_1^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{1}{8}, \frac{1}{8}, \frac{1}{2}$ | SB:4 ₁ ^B |
| 7_1^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{1}{8}, \frac{1}{8}, \frac{1}{2}$ | SB:4 ₁ ^B |
| 7_1^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{1}{8}, \frac{1}{8}, \frac{1}{2}$ | SB:4 ₁ ^B |
| 7_1^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{1}{8}, \frac{1}{8}, \frac{1}{2}$ | SB:4 ₁ ^B |
| 7_1^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{1}{8}, \frac{1}{8}, \frac{1}{2}$ | SB:4 ₁ ^B |
| 7_2^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{1}{4}, \frac{1}{4}, \frac{1}{2}$ | SB:4 ₂ ^B |
| 7_3^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{3}{8}, \frac{3}{8}, \frac{1}{2}$ | SB:4 ₃ ^B |
| 7_3^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{3}{8}, \frac{3}{8}, \frac{1}{2}$ | SB:4 ₃ ^B |
| 7_3^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{3}{8}, \frac{3}{8}, \frac{1}{2}$ | SB:4 ₃ ^B |
| 7_3^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{3}{8}, \frac{3}{8}, \frac{1}{2}$ | SB:4 ₃ ^B |
| 7_3^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{3}{8}, \frac{3}{8}, \frac{1}{2}$ | SB:4 ₃ ^B |
| 7_3^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{3}{8}, \frac{3}{8}, \frac{1}{2}$ | SB:4 ₃ ^B |
| 7_3^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{3}{8}, \frac{3}{8}, \frac{1}{2}$ | SB:4 ₃ ^B |
| 7_3^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{3}{8}, \frac{3}{8}, \frac{1}{2}$ | SB:4 ₃ ^B |
| 7_4^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{1}{2}, \frac{1}{2}, \frac{1}{2}$ | SB:4 ₄ ^B |
| 7_{-3}^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{5}{8}, \frac{5}{8}, \frac{1}{2}$ | SB:4 ₋₃ ^B |
| 7_{-3}^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{5}{8}, \frac{5}{8}, \frac{1}{2}$ | SB:4 ₋₃ ^B |
| 7_{-3}^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{5}{8}, \frac{5}{8}, \frac{1}{2}$ | SB:4 ₋₃ ^B |
| 7_{-3}^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{5}{8}, \frac{5}{8}, \frac{1}{2}$ | SB:4 ₋₃ ^B |
| 7_{-3}^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{5}{8}, \frac{5}{8}, \frac{1}{2}$ | SB:4 ₋₃ ^B |
| 7_{-3}^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{5}{8}, \frac{5}{8}, \frac{1}{2}$ | SB:4 ₋₃ ^B |
| 7_{-3}^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{5}{8}, \frac{5}{8}, \frac{1}{2}$ | SB:4 ₋₃ ^B |
| 7_{-2}^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{1}{2}, \frac{3}{4}, \frac{3}{4}$ | SB:4 ₋₂ ^B |
| 7_{-1}^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{1}{2}, \frac{7}{8}, \frac{7}{8}$ | SB:4 ₋₁ ^B |
| 7_{-1}^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{1}{2}, \frac{7}{8}, \frac{7}{8}$ | SB:4 ₋₁ ^B |
| 7_{-1}^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{1}{2}, \frac{7}{8}, \frac{7}{8}$ | SB:4 ₋₁ ^B |
| 7_{-1}^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{1}{2}, \frac{7}{8}, \frac{7}{8}$ | SB:4 ₋₁ ^B |
| 7_{-1}^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{1}{2}, \frac{7}{8}, \frac{7}{8}$ | SB:4 ₋₁ ^B |
| 7_{-1}^2 | 16 | 1, 1, 1, 1, 2, 2, 2 | 0, 0, 0, 0, $\frac{1}{2}, \frac{7}{8}, \frac{7}{8}$ | SB:4 ₋₁ ^B |
| $7_{9/5}^2$ | 28.944 | 1, 1, 1, 1, 2, ζ_8^4, ζ_8^4 | 0, 0, 0, 0, $\frac{3}{4}, \frac{3}{20}, \frac{2}{5}$ | SB:4 _{9/5} ^B |
| $7_{19/5}^2$ | 28.944 | 1, 1, 1, 1, 2, ζ_8^4, ζ_8^4 | 0, 0, 0, 0, $\frac{1}{4}, \frac{2}{5}, \frac{13}{20}$ | SB:4 _{19/5} ^B |
| $7_{-19/5}^2$ | 28.944 | 1, 1, 1, 1, 2, ζ_8^4, ζ_8^4 | 0, 0, 0, 0, $\frac{3}{4}, \frac{7}{20}, \frac{3}{5}$ | SB:4 _{-19/5} ^B |
| $7_{-9/5}^2$ | 28.944 | 1, 1, 1, 1, 2, ζ_8^4, ζ_8^4 | 0, 0, 0, 0, $\frac{1}{4}, \frac{3}{5}, \frac{17}{20}$ | SB:4 _{-9/5} ^B |
| 7_0^2 | 52.360 | 1, 1, 1, 1, $\zeta_8^4, \zeta_8^4, 3 + \sqrt{5}$ | 0, 0, 0, 0, $\frac{2}{5}, \frac{3}{5}, 0$ | SB:4 ₀ ^B |
| $7_{12/5}^2$ | 52.360 | 1, 1, 1, 1, $\zeta_8^4, \zeta_8^4, 3 + \sqrt{5}$ | 0, 0, 0, 0, $\frac{3}{5}, \frac{3}{5}, \frac{1}{5}$ | SB:4 _{12/5} ^B |
| $7_{-12/5}^2$ | 52.360 | 1, 1, 1, 1, $\zeta_8^4, \zeta_8^4, 3 + \sqrt{5}$ | 0, 0, 0, 0, $\frac{2}{5}, \frac{2}{5}, \frac{4}{5}$ | SB:4 _{-12/5} ^B |
| $7_{10/3}^2$ | 76.937 | 1, 1, 1, 1, $2\zeta_7^1, 2\zeta_7^2, \zeta_{16}^8$ | 0, 0, 0, 0, $\frac{1}{3}, \frac{2}{9}, \frac{2}{3}$ | SB:4 _{10/3} ^B |
| $7_{-10/3}^2$ | 76.937 | 1, 1, 1, 1, $2\zeta_7^1, 2\zeta_7^2, \zeta_{16}^8$ | 0, 0, 0, 0, $\frac{2}{3}, \frac{7}{9}, \frac{1}{3}$ | SB:4 _{-10/3} ^B |

TABLE XXXI. $Z_2 \times Z_2$ -SET orders for bosonic systems labeled in terms of sets of topological excitations. The list contains all topological orders with $N = 8$ $D^2 \leq 60$.

| $N_c^{ \Theta }$ | D^2 | d_1, d_2, \dots | s_1, s_2, \dots | comment |
|------------------|--------|--|---|--|
| 4_0^2 | 4 | 1, 1, 1, 1 | 0, 0, 0, 0 | $\mathcal{E} = \text{Rep}(Z_2 \times Z_2)$ |
| 8_1^2 | 8 | 1, 1, 1, 1, 1, 1, 1, 1 | 0, 0, 0, 0, $\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}$ | $2_1^B \boxtimes \text{Rep}(Z_2 \times Z_2)$ |
| 8_1^2 | 8 | 1, 1, 1, 1, 1, 1, 1, 1 | 0, 0, 0, 0, $\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}$ | SB: 2_1^B |
| 8_1^2 | 8 | 1, 1, 1, 1, 1, 1, 1, 1 | 0, 0, 0, 0, $\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}$ | SB: 2_1^B |
| 8_1^2 | 8 | 1, 1, 1, 1, 1, 1, 1, 1 | 0, 0, 0, 0, $\frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}$ | SB: 2_1^B |
| 8_{-1}^2 | 8 | 1, 1, 1, 1, 1, 1, 1, 1 | 0, 0, 0, 0, $\frac{3}{4}, \frac{3}{4}, \frac{3}{4}, \frac{3}{4}$ | $2_{-1}^B \boxtimes \text{Rep}(Z_2 \times Z_2)$ |
| 8_{-1}^2 | 8 | 1, 1, 1, 1, 1, 1, 1, 1 | 0, 0, 0, 0, $\frac{3}{4}, \frac{3}{4}, \frac{3}{4}, \frac{3}{4}$ | SB: 2_{-1}^B |
| 8_{-1}^2 | 8 | 1, 1, 1, 1, 1, 1, 1, 1 | 0, 0, 0, 0, $\frac{3}{4}, \frac{3}{4}, \frac{3}{4}, \frac{3}{4}$ | SB: 2_{-1}^B |
| 8_{-1}^2 | 8 | 1, 1, 1, 1, 1, 1, 1, 1 | 0, 0, 0, 0, $\frac{3}{4}, \frac{3}{4}, \frac{3}{4}, \frac{3}{4}$ | SB: 2_{-1}^B |
| $8_{14/5}^2$ | 14.472 | 1, 1, 1, 1, $\zeta_3^1, \zeta_3^1, \zeta_3^1, \zeta_3^1$ | 0, 0, 0, 0, $\frac{2}{5}, \frac{2}{5}, \frac{2}{5}, \frac{2}{5}$ | $2_{14/5}^B \boxtimes \text{Rep}(Z_2 \times Z_2)$ |
| $8_{-14/5}^2$ | 14.472 | 1, 1, 1, 1, $\zeta_3^1, \zeta_3^1, \zeta_3^1, \zeta_3^1$ | 0, 0, 0, 0, $\frac{3}{5}, \frac{3}{5}, \frac{3}{5}, \frac{3}{5}$ | $2_{-14/5}^B \boxtimes \text{Rep}(Z_2 \times Z_2)$ |
| 8_0^2 | 20 | 1, 1, 1, 1, 2, 2, 2, 2 | 0, 0, 0, 0, $\frac{1}{5}, \frac{1}{5}, \frac{4}{5}, \frac{4}{5}$ | SB: 5_0^B |
| 8_0^2 | 20 | 1, 1, 1, 1, 2, 2, 2, 2 | 0, 0, 0, 0, $\frac{1}{5}, \frac{1}{5}, \frac{4}{5}, \frac{4}{5}$ | SB: 5_0^B |
| 8_0^2 | 20 | 1, 1, 1, 1, 2, 2, 2, 2 | 0, 0, 0, 0, $\frac{1}{5}, \frac{1}{5}, \frac{4}{5}, \frac{4}{5}$ | SB: 5_0^B |
| 8_0^2 | 20 | 1, 1, 1, 1, 2, 2, 2, 2 | 0, 0, 0, 0, $\frac{1}{5}, \frac{1}{5}, \frac{4}{5}, \frac{4}{5}$ | SB: 5_0^B |
| 8_4^2 | 20 | 1, 1, 1, 1, 2, 2, 2, 2 | 0, 0, 0, 0, $\frac{2}{5}, \frac{2}{5}, \frac{3}{5}, \frac{3}{5}$ | SB: 5_4^B |
| 8_4^2 | 20 | 1, 1, 1, 1, 2, 2, 2, 2 | 0, 0, 0, 0, $\frac{2}{5}, \frac{2}{5}, \frac{3}{5}, \frac{3}{5}$ | SB: 5_4^B |
| 8_4^2 | 20 | 1, 1, 1, 1, 2, 2, 2, 2 | 0, 0, 0, 0, $\frac{2}{5}, \frac{2}{5}, \frac{3}{5}, \frac{3}{5}$ | SB: 5_4^B |
| 8_4^2 | 20 | 1, 1, 1, 1, 2, 2, 2, 2 | 0, 0, 0, 0, $\frac{2}{5}, \frac{2}{5}, \frac{3}{5}, \frac{3}{5}$ | SB: 5_4^B |
| 8_2^2 | 48 | 1, 1, 1, 1, 2, $\sqrt{12}, \sqrt{12}, 4$ | 0, 0, 0, 0, 0, $\frac{1}{8}, \frac{5}{8}, \frac{1}{3}$ | SB: 5_2^B |
| 8_2^2 | 48 | 1, 1, 1, 1, 2, $\sqrt{12}, \sqrt{12}, 4$ | 0, 0, 0, 0, 0, $\frac{3}{8}, \frac{7}{8}, \frac{1}{3}$ | SB: 5_2^B |
| 8_{-2}^2 | 48 | 1, 1, 1, 1, 2, $\sqrt{12}, \sqrt{12}, 4$ | 0, 0, 0, 0, 0, $\frac{1}{8}, \frac{5}{8}, \frac{2}{3}$ | SB: 5_{-2}^B |
| 8_{-2}^2 | 48 | 1, 1, 1, 1, 2, $\sqrt{12}, \sqrt{12}, 4$ | 0, 0, 0, 0, 0, $\frac{3}{8}, \frac{7}{8}, \frac{2}{3}$ | SB: 5_{-2}^B |
| $8_{16/11}^2$ | 138.58 | 1, 1, 1, 1, $2\zeta_9^1, 2\zeta_9^2, 2\zeta_9^3, \zeta_{20}^{10}$ | 0, 0, 0, 0, $\frac{9}{11}, \frac{2}{11}, \frac{1}{11}, \frac{6}{11}$ | SB: $5_{16/11}^B$ |
| $8_{-16/11}^2$ | 138.58 | 1, 1, 1, 1, $2\zeta_9^1, 2\zeta_9^2, 2\zeta_9^3, \zeta_{20}^{10}$ | 0, 0, 0, 0, $\frac{2}{11}, \frac{9}{11}, \frac{10}{11}, \frac{5}{11}$ | SB: $5_{-16/11}^B$ |
| $8_{18/7}^2$ | 141.36 | 1, 1, 1, 1, $\zeta_{12}^6, \zeta_{12}^6, 2\zeta_{12}^2, 2\zeta_{12}^4$ | 0, 0, 0, 0, $\frac{6}{7}, \frac{6}{7}, \frac{1}{7}, \frac{3}{7}$ | SB: $5_{18/7}^B$ |
| $8_{-18/7}^2$ | 141.36 | 1, 1, 1, 1, $\zeta_{12}^6, \zeta_{12}^6, 2\zeta_{12}^2, 2\zeta_{12}^4$ | 0, 0, 0, 0, $\frac{1}{7}, \frac{1}{7}, \frac{6}{7}, \frac{4}{7}$ | SB: $5_{-18/7}^B$ |

TABLE XXXII. The fusion rules for some $Z_2 \times Z_2$ -SET orders.

| | | | | | |
|----------|----------|----------|----------|----------|---|
| s_i | 0 | 0 | 0 | 0 | $\frac{1}{4}$ |
| d_i | 1 | 1 | 1 | 1 | 2 |
| 5_1^2 | 1 | <i>a</i> | <i>b</i> | <i>c</i> | ϕ |
| 1 | 1 | <i>a</i> | <i>b</i> | <i>c</i> | ϕ |
| <i>a</i> | <i>a</i> | 1 | <i>c</i> | <i>b</i> | ϕ |
| <i>b</i> | <i>b</i> | <i>c</i> | 1 | <i>a</i> | ϕ |
| <i>c</i> | <i>c</i> | <i>b</i> | <i>a</i> | 1 | ϕ |
| ϕ | ϕ | ϕ | ϕ | ϕ | $\mathbf{1} \oplus a \oplus b \oplus c$ |

| | | | | | |
|--------------|----------|----------|----------|----------|--|
| s_i | 0 | 0 | 0 | 0 | $\frac{2}{5}$ |
| d_i | 1 | 1 | 1 | 1 | $2\zeta_3^1$ |
| $5_{14/5}^2$ | 1 | <i>a</i> | <i>b</i> | <i>c</i> | η |
| 1 | 1 | <i>a</i> | <i>b</i> | <i>c</i> | η |
| <i>a</i> | <i>a</i> | 1 | <i>c</i> | <i>b</i> | η |
| <i>b</i> | <i>b</i> | <i>c</i> | 1 | <i>a</i> | η |
| <i>c</i> | <i>c</i> | <i>b</i> | <i>a</i> | 1 | η |
| η | η | η | η | η | $\mathbf{1} \oplus a \oplus b \oplus c \oplus 2\eta$ |

To obtain more properties of M^{iI} and to solve the above conditions on M^{iI} , let us consider the fusion with a particles:

$$a \otimes x = \oplus_y N_y^{ax} y. \quad (\text{D7})$$

We define x to be equivalent to y if there exists a such that $N_y^{ax} \neq 0$. Let $[x]$ be the equivalent class of x . Clearly $[1] = [a]$.

First, we like to pointed out that if i and j are equivalent, then i and j are formed by the same combination

of I 's, up to an overall factor, such as

$$i \rightarrow I_1 \oplus 2I_2, \quad j \rightarrow 3I_1 \oplus 6I_2. \quad (\text{D8})$$

This is because a particles in \mathcal{C} is mapped to the direct-sum of identity in \mathcal{C}_0 . Since i and j is related by fusing a or identity in \mathcal{C}_0 , then i and j must be formed by the same combination of I 's.

Second, if i and j are not equivalent, then the I 's that enter i do not overlap with the I 's that enter j . This

TABLE XXXIII. $Z_2 \times Z_2^f$ -SET orders (up to invertible ones) for fermionic systems. The list contains all topological orders with $N = 6$ $D^2 \leq 300$, $N = 8$ $D^2 \leq 60$, and $N = 10$ $D^2 \leq 20$.

| $N_c^{ \Theta }$ | D^2 | d_1, d_2, \dots | s_1, s_2, \dots | comment |
|--|--------|--|--|--|
| $4_0^0(\frac{2}{0})$ | 4 | 1, 1, 1, 1 | $0, 0, \frac{1}{2}, \frac{1}{2}$ | $\mathcal{E} = \text{sRep}(Z_2 \times Z_2^f)$ |
| 6_0^0 | 12 | 1, 1, 1, 1, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{6}, \frac{2}{3}$ | SB: $K = \begin{pmatrix} -1 & -2 \\ -2 & -1 \end{pmatrix}$ |
| 6_0^0 | 12 | 1, 1, 1, 1, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{3}, \frac{5}{6}$ | SB: $K = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$ |
| $8_0^0(\frac{0}{0})$ | 8 | 1, 1, 1, 1, 1, 1, 1, 1 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{1}{4}, \frac{3}{4}, \frac{3}{4}$ | $2_1^B \boxtimes \text{sRep}(Z_2 \times Z_2^f)$ |
| $8_0^0(\frac{0}{0})$ | 8 | 1, 1, 1, 1, 1, 1, 1, 1 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{1}{4}, \frac{3}{4}, \frac{3}{4}$ | SB: $4_0^F(\frac{0}{0})$ |
| $8_{-14/5}^0(\frac{\zeta_8^4}{3/20})$ | 14.472 | 1, 1, 1, 1, $\zeta_3^1, \zeta_3^1, \zeta_3^1, \zeta_3^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{10}, \frac{1}{10}, \frac{3}{5}, \frac{3}{5}$ | $2_{-14/5}^B \boxtimes \text{sRep}(Z_2 \times Z_2^f)$ |
| $8_{14/5}^0(\frac{\zeta_8^4}{-3/20})$ | 14.472 | 1, 1, 1, 1, $\zeta_3^1, \zeta_3^1, \zeta_3^1, \zeta_3^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{2}{5}, \frac{2}{5}, \frac{9}{10}, \frac{9}{10}$ | $2_{14/5}^B \boxtimes \text{sRep}(Z_2 \times Z_2^f)$ |
| $8_0^0(\frac{2}{0})$ | 20 | 1, 1, 1, 1, 2, 2, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{10}, \frac{2}{5}, \frac{3}{5}, \frac{9}{10}$ | SB: $10_0^F(\frac{\zeta_2^1}{0})$ |
| $8_0^0(\frac{2}{1/2})$ | 20 | 1, 1, 1, 1, 2, 2, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{5}, \frac{3}{10}, \frac{7}{10}, \frac{4}{5}$ | SB: $10_0^F(\frac{\zeta_2^1}{1/2})$ |
| $8_{1/4}^0(\frac{\zeta_2^3 \zeta_6^3}{1/2})$ | 27.313 | 1, 1, 1, 1, $\zeta_6^2, \zeta_6^2, \zeta_6^2, \zeta_6^2$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{1}{4}, \frac{3}{4}, \frac{3}{4}$ | SB: $4_{1/4}^F(\frac{\zeta_3^3}{1/2})$ |
| $8_{1/4}^0(\frac{\zeta_2^3 \zeta_6^3}{1/2})$ | 27.313 | 1, 1, 1, 1, $\zeta_6^2, \zeta_6^2, \zeta_6^2, \zeta_6^2$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{1}{4}, \frac{3}{4}, \frac{3}{4}$ | SB: $4_{1/4}^F(\frac{\zeta_3^3}{1/2})$ |
| $10_0^0(\frac{4}{0})$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, 0, \frac{1}{2}$ | SB: $8_0^F(\frac{\sqrt{8}}{0})$ |
| $10_0^0(\frac{4}{0})$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, 0, \frac{1}{2}$ | SB: $8_0^F(\frac{\sqrt{8}}{0})$ |
| $10_0^0(\frac{\sqrt{8}}{1/8})$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}, \frac{5}{8}$ | SB: $8_0^F(\frac{2}{1/8})$ |
| $10_0^0(\frac{\sqrt{8}}{1/8})$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}, \frac{5}{8}$ | SB: $8_0^F(\frac{2}{1/8})$ |
| $10_0^0(\frac{0}{0})$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{3}{4}$ | SB: $8_0^F(\frac{0}{0})$ |
| $10_0^0(\frac{0}{0})$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{3}{4}$ | SB: $8_0^F(\frac{0}{0})$ |
| $10_0^0(\frac{\sqrt{8}}{-1/8})$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{8}, \frac{7}{8}$ | SB: $8_0^F(\frac{2}{-1/8})$ |
| $10_0^0(\frac{\sqrt{8}}{-1/8})$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{8}, \frac{7}{8}$ | SB: $8_0^F(\frac{2}{-1/8})$ |

is a consequence of eqn. (D5). The right hand side of eqn. (D5) will vanish if i and j are not equivalent.

Third, the I 's that appear in i must have the same quantum dimensions and spins. This is because those I 's must be degenerate. This can only happen if they have the same quantum dimensions and spins.

Fourth, the I 's that appears in i must each enter with an equal weight, such as

$$i \rightarrow 2I_1 \oplus 2I_2. \quad (\text{D9})$$

Again, this is because those I 's must be degenerate. This can only happen if they can be mapped into each other by symmetry transformations. Since the symmetry transformations only permute I 's, each I enters with an equal weight.

Combine the above results, we see that M^{iI} has the following block structure. We can divide the index I into groups $[I]$, such that there is one-to-one correspondence between $[i]$ and $[I]$: $[i] \leftrightarrow [I]_{[i]}$, and

$$\begin{aligned} M^{iI} &= 0 & \text{if } i \in [i], I \notin [I]_{[i]}, \\ M^{iI} &= m_i > 0 & \text{if } i \in [i], I \in [I]_{[i]}. \end{aligned} \quad (\text{D10})$$

Therefore, we have

$$m_i^2 n_{[i]} = \sum_a N_a^{i\bar{i}} d_a, \quad (\text{D11})$$

where $n_{[i]}$ is the size of the set $[I]_{[i]}$. Since

$$i = \oplus_{I \in [I]_{[i]}} m_i I, \quad (\text{D12})$$

we have

$$m_i m_j n_{[i]} = \sum_a N_a^{i\bar{j}} d_a, \quad i, j \in [i] \quad (\text{D13})$$

In other words, the matrix \tilde{N} with elements $\tilde{N}_{ij} = \sum_a N_a^{i\bar{j}} d_a$ is block diagonal. Each block is formed by particles in an equivalent class $[i]$, and is given by the above expression. We see that, for $i, j \in [i]$, $\sum_a N_a^{i\bar{j}} d_a$ must be a symmetric matrix with a single non-zero eigenvalue $n_{[i]} \sum_{j \in [i]} m_j^2$ and eigenvector (m_j) .

We also find that

$$d_i = m_i n_{[i]} d_I, \quad (\text{D14})$$

or

$$d_I = \frac{m_i d_i}{\sum_a N_a^{i\bar{i}} d_a} \quad \forall \quad I \in [I]_{[i]}. \quad (\text{D15})$$

Using the fact $s_i = s_j = s_I, \forall i, j \in [i], I \in [I]_{[i]}$, we can obtain (d_I, s_I) of \mathcal{C}_0 from (N_k^{ij}, s_i) of \mathcal{C} . The resulting (d_I, s_I) must be the quantum dimensions and the spins of a UMTC. This gives us some extra conditions on (N_k^{ij}, s_i) .

TABLE XXXIV. Z_4^f -SET orders for fermionic systems. The list contains all topological orders with $N = 6$ $D^2 \leq 100$, $N = 8$ $D^2 \leq 60$, and $N = 10$ $D^2 \leq 20$.

| $N_c^{ \Theta }$ | D^2 | d_1, d_2, \dots | s_1, s_2, \dots | comment |
|---------------------------------|--------|--|--|--|
| 4_0^0 | 4 | 1, 1, 1, 1 | $0, 0, \frac{1}{2}, \frac{1}{2}$ | $\mathcal{E} = \text{sRep}(Z_4^f)$ |
| 6_0^0 | 12 | 1, 1, 1, 1, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{6}, \frac{2}{3}$ | $\kappa = -\begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$ |
| 6_0^0 | 12 | 1, 1, 1, 1, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{3}, \frac{5}{6}$ | $\kappa = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$ |
| 8_0^0 | 8 | 1, 1, 1, 1, 1, 1, 1 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{1}{4}, \frac{3}{4}, \frac{3}{4}$ | $2_{-1}^B \boxtimes \text{sRep}(Z_4^f)$ |
| 8_0^0 | 8 | 1, 1, 1, 1, 1, 1, 1 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{1}{4}, \frac{3}{4}, \frac{3}{4}$ | $2_1^B \boxtimes \text{sRep}(Z_4^f)$ |
| $8_{-14/5}^0$ | 14.472 | $1, 1, 1, 1, \zeta_3^1, \zeta_3^1, \zeta_3^1, \zeta_3^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{10}, \frac{1}{10}, \frac{3}{5}, \frac{3}{5}$ | $2_{-14/5}^B \boxtimes \text{sRep}(Z_4^f)$ |
| $8_{14/5}^0$ | 14.472 | $1, 1, 1, 1, \zeta_3^1, \zeta_3^1, \zeta_3^1, \zeta_3^1$ | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{2}{5}, \frac{2}{5}, \frac{9}{10}, \frac{9}{10}$ | $2_{14/5}^B \boxtimes \text{sRep}(Z_4^f)$ |
| 8_0^0 | 20 | 1, 1, 1, 1, 2, 2, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{10}, \frac{2}{5}, \frac{3}{5}, \frac{9}{10}$ | SB: $10_0^F(\zeta_5^1)$ |
| 8_0^0 | 20 | 1, 1, 1, 1, 2, 2, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{5}, \frac{3}{10}, \frac{7}{10}, \frac{4}{5}$ | SB: $10_0^F(\zeta_5^2)$ |
| $10_0^0(\frac{4}{0})$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, 0, \frac{1}{2}$ | SB: $8_0^F(\sqrt[8]{0})$ |
| $10_0^0(\frac{4}{0})$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, 0, \frac{1}{2}$ | SB: $8_0^F(\sqrt[8]{0})$ |
| $10_0^0(\frac{\sqrt{8}}{1/8})$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}, \frac{5}{8}$ | SB: $8_0^F(\frac{2}{1/8})$ |
| $10_0^0(\frac{\sqrt{8}}{1/8})$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{8}, \frac{5}{8}$ | SB: $8_0^F(\frac{2}{1/8})$ |
| $10_0^0(\frac{0}{0})$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{3}{4}$ | SB: $8_0^F(\frac{0}{0})$ |
| $10_0^0(\frac{0}{0})$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{1}{4}, \frac{3}{4}$ | SB: $8_0^F(\frac{0}{0})$ |
| $10_0^0(\frac{\sqrt{8}}{-1/8})$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{8}, \frac{7}{8}$ | SB: $8_0^F(\frac{2}{-1/8})$ |
| $10_0^0(\frac{\sqrt{8}}{-1/8})$ | 16 | 1, 1, 1, 1, 1, 1, 1, 1, 2, 2 | $0, 0, \frac{1}{2}, \frac{1}{2}, 0, 0, \frac{1}{2}, \frac{1}{2}, \frac{3}{8}, \frac{7}{8}$ | SB: $8_0^F(\frac{2}{-1/8})$ |

Appendix E: Physical and mathematical meaning of $\text{UMTC}_{/\mathcal{E}}$ and its modular extensions

In the main text of the paper, we have explained why $\text{UMTC}_{/\mathcal{E}}$ describes the bulk particle-like excitations. We also explained the motivation of modular extension via “gauging” the symmetry. In this section, we will discuss a deeper meaning of $\text{UMTC}_{/\mathcal{E}}$ and its modular extensions.

We know that $\text{UMTC}_{/\mathcal{E}}$ is a very abstract way to describe the non-abelian statistics of the excitations. It is not clear at all that why the excitations described by $\text{UMTC}_{/\mathcal{E}}$ can be realized by a local lattice model with on-site symmetry. In physics, we mainly concern about local lattice models and their properties. It appears that there is a big gap between the $\text{UMTC}_{/\mathcal{E}}$ studied in this paper and local lattice models that physicists want to study. In fact, the two are closely related. Here, we will try to explain such a connection between lattice models and $\text{UMTC}_{/\mathcal{E}}$ (with their modular extensions).

We know that the fusion-braiding properties of particles within a 2-dimensional open disk can be described by a unitary braided fusion category. From this point of view, a unitary braided fusion category is a *local* theory that only encode the local properties of the fusion and braiding (*i.e.* on an open disk). We want to promote fusion-braiding properties to be integrable to any 2-dimensional manifolds because we want those fusion-braiding properties to be realizable by some local lattice models, which can always be defined on any 2-dimensional manifolds. Therefore, the integrability of fusion-braiding properties to any 2-dimensional mani-

folds is necessary for the fusion-braiding properties to be realized by some local lattice models.

Now we assume that “all 2-dimensional manifolds” are the most powerful probes. This means that the integrability of the local fusion-braiding properties to global invariants (on all 2-dimensional manifolds), satisfying natural physically required properties, is also sufficient for those properties to be realizable by some local lattice models.

The process of integrating the local fusion-braiding properties of particles (described by a UBFC \mathcal{C}) to give global invariants is defined by the so-called factorization homology.^{52,53} In order to be free of framing anomaly, we need a spherical structure, which is guaranteed by the unitarity of a UBFC.⁹ For general UBFCs, although the global invariants are well-defined by factorization homology,⁵³ they do not have nice properties that allow us to give them a natural physical meaning. A stronger *integrability condition* needs to be imposed in order for the global invariants to have natural physical meanings.

For example, if \mathcal{C} is assumed to be non-degenerate (*i.e.* UMTC), it was shown in Ref. 54 that factorization homology of a UMTC \mathcal{C} over a closed 2-dimensional manifold is given by the category of finite dimensional Hilbert spaces. If one inserts a finite number of particle-like excitations x_1, \dots, x_r on the closed surface, one simply obtain the Hilbert space $\text{hom}_{\mathcal{C}}(\mathbf{1}, x_1 \otimes \dots \otimes x_r)$, which is also the space of degenerate ground states. This result remains to be true for all closed 2-dimensional manifolds with topological gapped defects and with 2-cells decorated by different phases.⁵⁴ This includes the cases that the topological order is defined on any surfaces with

boundaries. Therefore, the non-degeneracy is certainly a sufficient integrability condition, which is too strong for the purpose of this work.

In this paper, we consider something more complicated – the fusion-braiding properties of particles with symmetry. By “with symmetry”, we mean to include local excitations that carry representations of the symmetry group. Mathematically, this means that the unitary braided fusion category \mathcal{C} contain a SFC \mathcal{E} as its Müger center, i.e. a UMTC $_{/\mathcal{E}}$. We know that either $\mathcal{E} = \text{Rep}(G)$ or $\mathcal{E} = \text{sRep}(G^f)$, where G or G^f is the symmetry group. In this case, we must find a proper integrability condition that is weaker than the non-degeneracy of UBFC.

In order for the factorization homology of \mathcal{C} on a surface, a unitary category denoted by \mathcal{C}_Σ , to have a physical meaning, we suspect that we should be able to interpret its object as finite dimensional Hilbert spaces in a natural way. This suggests that the category \mathcal{C}_Σ should be equipped with a natural functor to the category of finite dimensional Hilbert spaces, which is a factorization homology \mathcal{M}_Σ of a UMTC \mathcal{M} .⁵⁴ So we expect that we should be able to embed \mathcal{C} into a UMTC \mathcal{M} such that the embedding naturally descends to a functor $\mathcal{C}_\Sigma \rightarrow \mathcal{M}_\Sigma$ on factorization homologies. An arbitrary UMTC such as the Drinfeld center $Z(\mathcal{C})$ of \mathcal{C} can not do the job because there is no canonical way to identify \mathcal{C} in \mathcal{M} (with a fixed symmetry \mathcal{E}) so that it is unlikely that it can be compatible with the integration process. So we expect that the condition $\mathcal{E}_\mathcal{M}^{\text{cen}} = \mathcal{C}$ is a natural integrability condition that replace the non-degeneracy condition in this case. This flow of thinking leads us to the concept of the modular extension of \mathcal{C} . It also suggests that the non-existence of the modular extension of a given \mathcal{C} means that \mathcal{C} is somewhat inconsistent globally or not integrable to all 2-dimensional manifolds with natural physical meanings.

This can also be viewed from a different point of view. If we require each particle to be non-trivial in some sense, then we must only consider the non-degenerate unitary braided fusion category over SFC \mathcal{E} . In this case, for particles not in \mathcal{E} , we know they are non-trivial because their non-trivial double braiding (or non-trivial mutual statistics) with some particles. But we still have trouble to know why the particles in \mathcal{E} are non-trivial? From their fusion and braiding properties, they just behave like the identity or a composite of identities.

To fix this problem, we put our particles on any 2-dimensional manifolds. In this case, we can find a way to understand the non-trivialness of the particle in \mathcal{E} . This requires us to twist the symmetry G or G^f on the 2-dimensional manifold. In other words, we equip the 2-dimensional manifold with a flat G -connection. Since the particles in \mathcal{E} all carry irreducible representations of G , as we move the particles along a non-contractile loop, the flat G -connection will induce a G transformation on the particle (or more precisely, on the hom space of the particles). This allows us to probe the particles in \mathcal{E} and detect their non-trivialness.

Therefore, as we put particles on a 2-dimensional man-




FIG. 4. (a) A torus with a flat G -connection (described by a symmetry twist along the dashed loop). The thin solid loop is a braiding path. (b) A handle is deformed into a very thin one. (c) A very thin handle can be viewed as two defects, and each defect corresponds to the added particle in the modular extension.

ifold, it is important to allow any flat G -connection on the manifold. Now we ask, in this case, can a non-degenerate unitary braided fusion category \mathcal{C} over a SFC \mathcal{E} describes the fusion-braiding properties of particles that are consistent on any 2-dimensional manifolds with any flat G -connections?

In this paper, we propose that the answer is no. We also propose that the answer is yes iff the \mathcal{C} over \mathcal{E} has modular extensions, which are the categorical ways of gauging the symmetry \mathcal{E} . So, non-degenerate unitary braided fusion categories over SFC can describe the consistent local fusion and braiding on an open disk. Only the ones with modular extensions can describe the consistent fusion and braiding on any manifolds (with any flat G -connections).

The intuition for the above conjecture is explained in the Fig. 4. Fig. 4(a) describes a braiding of particles on a torus with flat G -connection. As we deform a handle into a very thin one, we may view the above braiding on torus as a braiding around the added particles in the modular extension. So the consistent fusion and braiding on any manifolds with any flat G -connection must be closely related to the consistent fusion and braiding on a sphere with the added particles in the modular extension. So, the mathematical meaning of the modular extension is to make the fusion and braiding to be consistent on any manifolds with any flat G -connection.

For a given \mathcal{C} over \mathcal{E} , there can be several modular extensions \mathcal{M} . We believe that those different modular extensions describe the different structures at the boundary. This picture leads to the physical conjecture that the triple $(\mathcal{C}, \mathcal{M}, c)$ classify the 2+1D topological/SPT orders with symmetry \mathcal{E} .

- ¹ X.-G. Wen, Phys. Rev. B **40**, 7387 (1989).
- ² X.-G. Wen, Int. J. Mod. Phys. B **4**, 239 (1990).
- ³ X.-G. Wen and Q. Niu, Phys. Rev. B **41**, 9377 (1990).
- ⁴ L. D. Landau, Phys. Z. Sowjetunion **11**, 26 (1937).
- ⁵ X. Chen, Z.-C. Gu, and X.-G. Wen, Phys. Rev. B **82**, 155138 (2010), [arXiv:1004.3835](#).
- ⁶ B. Zeng and X.-G. Wen, Phys. Rev. B **91**, 125121 (2015), [arXiv:1406.5090](#).
- ⁷ G. Moore and N. Seiberg, Communications in Mathematical Physics **123**, 177 (1989).
- ⁸ B. Bakalov and J. A. Kirillov, *Lectures on Tensor Categories and Modular Functors* (Amer. Math. Soc., 2001).
- ⁹ A. Kitaev, Annals of Physics **321**, 2 (2006), [cond-mat/0506438](#).
- ¹⁰ L. Kong and X.-G. Wen, (2014), [arXiv:1405.5858](#).
- ¹¹ X.-G. Wen, Natl. Sci. Rev. **3**, 68 (2016), [arXiv:1506.05768](#).
- ¹² D. S. Freed, (2014), [arXiv:1406.7278](#).
- ¹³ E. Rowell, R. Stong, and Z. Wang, Comm. Math. Phys. **292**, 343 (2009), [arXiv:0712.1377](#).
- ¹⁴ T. Lan, L. Kong, and X.-G. Wen, Phys. Rev. B **94**, 155113 (2016), [arXiv:1507.04673](#).
- ¹⁵ X.-G. Wen, Phys. Rev. D **88**, 045013 (2013), [arXiv:1303.1803](#).
- ¹⁶ Z.-C. Gu and X.-G. Wen, Phys. Rev. B **80**, 155131 (2009), [arXiv:0903.1069](#).
- ¹⁷ F. Pollmann, E. Berg, A. M. Turner, and M. Oshikawa, Phys. Rev. B **85**, 075125 (2012), [arXiv:0909.4059](#).
- ¹⁸ X. Chen, Z.-X. Liu, and X.-G. Wen, Phys. Rev. B **84**, 235141 (2011), [arXiv:1106.4752](#).
- ¹⁹ X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen, Phys. Rev. B **87**, 155114 (2013), [arXiv:1106.4772](#).
- ²⁰ X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen, Science **338**, 1604 (2012), [arXiv:1301.0861](#).
- ²¹ M. Levin and Z.-C. Gu, Phys. Rev. B **86**, 115109 (2012), [arXiv:1202.3120](#).
- ²² X.-G. Wen, Phys. Rev. B **89**, 035147 (2014), [arXiv:1301.7675](#).
- ²³ L.-Y. Hung and X.-G. Wen, Phys. Rev. B **89**, 075121 (2014), [arXiv:1311.5539](#).
- ²⁴ N. Read and S. Sachdev, Phys. Rev. Lett. **66**, 1773 (1991).
- ²⁵ X.-G. Wen, Phys. Rev. B **44**, 2664 (1991).
- ²⁶ X.-G. Wen, Phys. Rev. Lett. **90**, 016803 (2003), [quant-ph/0205004](#).
- ²⁷ S.-P. Kou, M. Levin, and X.-G. Wen, Phys. Rev. B **78**, 155134 (2008), [arXiv:0803.2300](#).
- ²⁸ C. Heinrich, F. Burnell, L. Fidkowski, and M. Levin, Phys. Rev. B **94**, 235136 (2016), [arXiv:1606.07816](#).
- ²⁹ M. Barkeshli, P. Bonderson, M. Cheng, and Z. Wang, (2014), [arXiv:1410.4540](#).
- ³⁰ T. Lan, L. Kong, and X.-G. Wen, Commun. Math. Phys. **351**, 709 (2017), [arXiv:1602.05936](#).
- ³¹ T. Tannaka, Tohoku Math. J. **45**, 1 (1938).
- ³² P. Bruillard, S.-H. Ng, E. C. Rowell, and Z. Wang, (2013), [arXiv:1310.7050 \[math.QA\]](#).
- ³³ V. Drinfeld, S. Gelaki, D. Nikshych, and V. Ostrik, (2007), [0704.0195](#).
- ³⁴ An odd matrix is a symmetric integer matrix with at least one of its diagonal elements being odd.
- ³⁵ Q. Yuan, <http://mathoverflow.net/questions/229568> (2016).
- ³⁶ A. Davydov, M. Müger, D. Nikshych, and V. Ostrik, J. Reine Angew. Math. **677**, 135 (2013), [1009.2117](#).
- ³⁷ J. Fröhlich, J. Fuchs, I. Runkel, and C. Schweigert, arXiv Mathematics e-prints (2003), [math/0309465](#).
- ³⁸ M. Müger, arXiv Mathematics e-prints (2001), [math/0111205](#).
- ³⁹ A. Davydov, D. Nikshych, and V. Ostrik, Selecta Math **19**, 237 (2011), [arXiv:1109.5558](#).
- ⁴⁰ Z.-C. Gu and M. Levin, (2013), [arXiv:1304.4569](#).
- ⁴¹ A. Kapustin, R. Thorngren, A. Turzillo, and Z. Wang, Journal of High Energy Physics **2015**, 52 (2015), [arXiv:1406.7329](#).
- ⁴² C. Wang, (2016), [arXiv:1601.02028](#).
- ⁴³ E. Witten, Comm. Math. Phys. **121**, 351 (1989).
- ⁴⁴ D. Gepner and A. Kapustin, Physics Letters B **349**, 71 (1995), [hep-th/9410089](#).
- ⁴⁵ B. Bakalov and A. Kirillov, *Lectures on Tensor Categories and Modular Functors*, University Lecture Series, Vol. 21 (American Mathematical Society, 2001) p. 221.
- ⁴⁶ Z. Wang, *Topological Quantum Computation* (CBMS Regional Conference Series in Mathematics, 2010).
- ⁴⁷ C. Vafa, Phys. Lett. B **206**, 421 (1988).
- ⁴⁸ G. Andersen and G. Moore, Commun. Math. Phys. **117**, 441 (1988).
- ⁴⁹ P. Etingof, Mathematical Research Letters **9**, 651 (2002), [math/0207007](#).
- ⁵⁰ E. Verlinde, Nuclear Physics B **300**, 360 (1988).
- ⁵¹ P. Bruillard, (2012), [arXiv:1204.4836](#).
- ⁵² J. Lurie, Higher algebra, a book available at <http://www.math.harvard.edu/~lurie/> (2014).
- ⁵³ D. Ben-Zvi, A. Brochier, and D. Jordan, (2015), [arXiv:1501.04652](#).
- ⁵⁴ Y. Ai, L. Kong, and H. Zheng, (2016), [in preparation](#).