This is the accepted manuscript made available via CHORUS. The article has been published as:

Magnetic correlations in the two-dimensional repulsive Fermi-Hubbard model

Fedor Šimkovic, IV, Youjin Deng, N. V. Prokof'ev, B. V. Svistunov, I. S. Tupitsyn, and Evgeny Kozik

Phys. Rev. B **96**, 081117 — Published 29 August 2017

DOI: [10.1103/PhysRevB.96.081117](https://doi.org/10.1103/PhysRevB.96.081117)

Magnetic Correlations in the Two-dimensional Repulsive Fermi Hubbard Model

Fedor Šimkovic IV.,¹ Youjin Deng,² N. V. Prokof'ev,^{3,4} B. V. Svistunov,^{3,4} I. Tupitsyn,^{3,4} and Evgeny Kozik¹

¹*Department of Physics, King's College London, Strand, London WC2R 2LS, UK*

²*Hefei National Laboratory for Physical Sciences at Microscale and Department of Modern Physics, University of Science and Technology of China, Hefei, Anhui 230026, China*

³*Department of Physics, University of Massachusetts, Amherst, MA 01003, USA*

⁴*National Research Center "Kurchatov Institute," 123182 Moscow, Russia*

(Dated: August 14, 2017)

The repulsive Fermi Hubbard model on a square lattice has a rich phase diagram near half-filling ($n = 1$): for $n = 1$ the ground state is an antiferromagnetic insulator, at $0.6 < n \lesssim 0.8$, the ground state is a $d_{x^2-y^2}$ -wave superfluid (at least for moderately strong interactions $U \lesssim 4$), and the region $1 - n \ll 1$ is likely subject to phase separation. Much less is known about the nature of strong magnetic fluctuations at finite temperature and how they change with the doping level. Recent experiments on ultra-cold atoms have now reached this interesting fluctuation regime. In this work we employ the skeleton diagrammatic method to quantify the characteristic temperature scale $T_M(n)$ for the onset of magnetic fluctuations with large correlation length.

PACS numbers: 71.10.Fd, 74.72.A, 74.25.Dw, 74.72.Ek

The fermionic Hubbard model [1–4], defined by the square lattice Hamiltonian

$$H = -t \sum_{\langle i,j \rangle \sigma} \hat{c}_{i,\sigma}^\dagger \hat{c}_{j,\sigma} + U \sum_i \hat{n}_{i,\uparrow} \hat{n}_{i,\downarrow} - \mu \sum_{i,\sigma} \hat{n}_{i,\sigma} \quad (1)$$

has for years played a crucial role in studies of correlated electrons in solids; it is regarded as one of the “standard models” of condensed matter physics to introduce and discuss Mott insulating phases, antiferromagnetic (AFM) correlations, novel mechanisms of superconducting pairing, non-Fermi-liquid behavior, etc. In Eq. (1) and in what follows, the nearest-neighbor hopping amplitude t is set to be the energy and temperature unit (distances are measured in units of the lattice constant); U is the on-site repulsive coupling constant; μ is the chemical potential; $\hat{c}_{i,\sigma}^\dagger$ and $\hat{c}_{i,\sigma}$ create and annihilate (respectively) a fermion of the spin component $\sigma \in \{\uparrow, \downarrow\}$ on the site i , and $\hat{n}_{i,\sigma}$ counts the number of fermions of a particular spin on a given lattice site.

On the one hand, Eq. (1) involves a number of crucial simplifications that make it qualitatively different from real materials, such as high- T_c superconductors. The most important ones include (i) two dimensional (2D) as opposed to the strongly anisotropic 3D geometry, (ii) neglect of long-range Coulomb interactions, (iii) suppression of hopping matrix elements beyond the nearest-neighbor ones ($t' = 0$), (iv) single-band approximation, and (v) absence of electron-phonon coupling. Correspondingly, the model (1) cannot feature an ordered AFM phase at finite temperature, but is allowed to have a first-order transition between phases with different electron densities, not to mention that $t' = 0$ leads to the Fermi surface nesting and particle-hole symmetry at $n = \langle n_{i\uparrow} + n_{i\downarrow} \rangle = 1$. As a result, the schematic phase diagram of (1) in the doping-temperature plane shown in Fig. 1 (see discussion below) is distinct from the

Figure 1. (color online). Schematic phase diagram of the repulsive Fermi-Hubbard model on the square lattice near half-filling, showing the Fermi liquid (FL), superfluid (SF), and putative phase separation (PS) regimes, as well as the antiferromagnetic ground state (AFM) at half-filling. Strong magnetic correlations with large correlation length $\xi \gg a$ are observed below the dashed red line.

“canonical” picture of high- T_c -type materials [4]. On the other hand, advances in ultra-cold atomic experiments have made it possible to accurately emulate the model (1) on optical lattices [5–14]. This remarkable progress brings ultra-cold atom experiments into the region of the phase diagram (Fig. 1), where they can cooperate with the state-of-the-art numerical methods to reveal the underlying physics. Numerical results can now be directly compared to experiments (and *vice versa*), which dramatically increases the importance of producing reliable data sets.

Recent years have seen a remarkable progress in unveiling the $T = 0$ phase diagram of (1). Well-understood

regions include the limit of vanishing densities $n \rightarrow 0$ [15–21] and vanishing interaction strength [22–25]. For densities $n \leq 0.7$ and coupling strength $U \leq 4$, the ground state is a BCS superfluid (with the $d_{x^2-y^2}$ -wave symmetry at density $n > 0.6$) [26]. At half-filling $n = 1$, the ground state is an AFM insulator for any U [27–31]. Being a qualitative property, AFM order can only disappear (with doping) by a quantum phase transition. The simplest scenario is the first-order one, implying phase separation (PS) into the insulating AFM and a superfluid is a natural—but still not the only possible—way of the doping-driven transitioning between the two ordered ground states. The first PS state was proposed to be a mixture of AFM and ferromagnetic (FM) orders in the region of small doping $\delta = 1 - n \ll 1$ and large U [32–34]; this conjecture has been later supported numerically for $U > 25$ [35]. The instability of the model towards incommensurate AFM and domain wall formation was also reported in Refs. [36, 37]. Recently, PS for small values of U was observed in Auxiliary-field Quantum Monte Carlo [38, 39] and variational [40] studies.

Much less is known conclusively about the finite-temperature behavior. Given that the correlation length ξ for AFM correlations at $n = 1$ diverges exponentially fast when $T \rightarrow 0$, there exists a relatively high temperature T_M below which magnetic correlations extend over many lattice sites and electronic degrees of freedom are getting locked in collective magnetic modes. The characteristic temperature scale $T_M(n)$ is supposed to decrease with increasing the doping level. Since magnetic correlations and fluctuations are considered to be the prime reason for PS and BCS phenomena near half-filling, quantifying their behavior is paramount to understanding the finite temperature phase diagram of the Fermi Hubbard model in 2D.

In this communication, we employ the skeleton diagrammatic approach to quantify the $T_M(n)$ scale, see Fig. 2, and shed light on the structure of dominant fluctuations. The most recent setups with ultra-cold ^6Li fermions have now reached temperatures low enough to directly observe magnetic correlations in the model (1) [10, 12, 13], with the lowest temperatures attained on the scale of $T \sim 0.25$ [14].

Method. The imaginary-time spin correlation function $\chi(i, \tau) = \langle S_z(\tau, i) S_z(0, 0) \rangle$ and its Fourier transform $\chi(\omega_m, \vec{q})$ at bosonic Matsubara frequencies $\omega_m = 2\pi T m$, where m is integer, was computed within the skeleton diagrammatic framework based on four fully dressed channels. We start with noticing that in any diagram based on bare on-site coupling U every interaction vertex can be “promoted” by adding an infinite sequence based on a particular geometric series, see Fig. 3. This leads to the diagrammatic elements from which an arbitrary Feynman diagram may be constructed by taking any number of diagrammatic elements and connecting their incoming and outgoing ends with propagator lines [see upper and

Figure 2. (color online). Onset of strong (incommensurate) magnetic fluctuations as calculated by the lowest-order GGGW method (see text) for values of $1 \leq U \leq 6$. The region of strong fluctuations increases with U and reaches its maximum at around $U = 4$ at $n = 1$.

Figure 3. (color online). Simple geometric series that can be constructed by connecting bare interaction vertexes using various pairs of bare single-particle propagators (Green's functions) $G_\sigma^{(0)}$. Red and black colors stand for spin-up and spin-down propagators.

lower panels in Fig. 4]. Note that the particle-particle ladders, $G_{pp}^{(0)}$, particle-hole ladders, $G_{ph}^{(0)}$, and screened interaction, $W_{\sigma\sigma'}^{(0)}$, all have the same structure as single-particle propagators (Green's functions) $G_\sigma^{(0)}$; i.e., thanks to the local nature of the on-site Hubbard interaction U they depend only the lattice coordinate and time. In what follows we refer to all G and W functions (bare or fully renormalized) as “lines” and call the U element a “point.” The scheme can be abbreviated as GGGW to underline four renormalization channels.

The necessity of treating the bare interaction point separately originates from precise rules for avoiding double counting when all diagram lines are assumed to be

Figure 4. (color online). Diagrammatic elements based on ladders and screened interactions (upper panel) and an example of a particular diagram for the single-particle self-energy Σ based on them.

dressed with the corresponding self-energy insertions. In the skeleton technique, one computes self-energies for all lines in the graph: Σ_σ for G_σ , Σ_{pp} for G_{pp} , Σ_{ph} for G_{ph} , and $\Pi_{\sigma\sigma'}$ for $W_{\sigma\sigma'}$ and uses them to “dress”/renormalize the corresponding lines using Dyson equations:

$$G_\sigma = \frac{G_\sigma^{(0)}}{1 - G_\sigma^{(0)}\Sigma_\sigma}, \quad W = \frac{U}{1 - U\Pi} - U, \quad (2)$$

$$G_{pp} = \frac{U}{1 - U\Sigma_{pp}} - U, \quad G_{ph} = \frac{U}{1 - U\Sigma_{ph}} - U. \quad (3)$$

These are simple algebraic relations in the momentum/frequency representation. The spin-spin correlation function is directly related to the polarization Π by

$$\chi(\omega_n, \vec{q}) = \text{Tr} S_z \frac{\Pi(\omega_n, \vec{q})}{1 - \Pi(\omega_n, \vec{q})U} S_z, \quad (4)$$

with the trace taken over the spin index.

The self-energy diagrams are computed self-consistently by using the solutions of (2) and (3) for all diagram lines. To avoid double counting, these diagrams have to be from the so-called skeleton set: they remain connected after cutting any two lines of the same kind. This is identical to enforcing the rule that no two lines in the free-energy graph have the same momentum; skeleton self-energy diagrams are then obtained by removing one of the lines from the closed free-energy graph. In addition, one has to enforce two rules concerning dots:

- (1) no two dots can be connected by two Green’s functions, and
- (2) dots cannot be connected to the same end of either G_{pp} , or G_{ph} , or W by two Green’s functions in a manner that reproduces their geometrical-series structure.

This is necessary because, by construction, such diagrams are already accounted for in the line definitions when considering their first-order self-energies. By admitting that the bare interaction vertex can be interpreted in three different ways (with the corresponding momentum going along the line) as depicted in Fig. 5, we observe that these additional rules are accounted

for automatically by the “no two lines carry the same momentum” rule. Finally, there is one exception from the rule: to avoid triple counting of the same diagram contributing to the lowest-order Σ , one has to perform subtraction of the diagram based on two points, see Fig. 6.

Figure 5. (color online). Three different ways for “interpreting” the bare interaction vertex in the context of different channels. In each interpretation the interaction line carries a different momentum.

$$\Sigma_\sigma^{(1)} = \text{diagram 1} + \text{diagram 2} + \text{diagram 3} + \text{diagram 4} - 2 \times \text{diagram 5}$$

$$\Sigma_{pp}^{(1)} = \text{diagram 6}, \quad \Sigma_{ph}^{(1)} = \text{diagram 7}, \quad \Pi_{\sigma\sigma'}^{(1)} = \delta_{\sigma\sigma'} \text{diagram 8}$$

Figure 6. (color online). First-order skeleton graphs for all self-energies.

All results in this work are based on the lowest-order diagrams shown in Fig. 6 and solving Dyson equations (2) and (3) iteratively. The Diagrammatic Monte Carlo scheme for simulating higher-order skeleton graphs (based on standard principles described in previous work [26, 41, 42] with obvious modifications required to reflect a larger set of diagrammatic elements and self energies) was only used to establish that the lowest-order scheme is producing accurate thermodynamic results at the level of few percent. We have also verified this level of accuracy for spin correlations by benchmarking results at $n = 1$ against the sign-free Determinant Diagrammatic Monte Carlo method [43].

Results. The very notion of T_M as the crossover temperature between the high-temperature regime with $\xi \leq 1$ and the regime of strong magnetic correlations with $\xi \gg 1$, implies that its definition is not unique. To define the onset of strong magnetic fluctuations we examine the momentum dependence of the static magnetic susceptibility $\chi(0, \vec{q})$ and the development of a narrow peak structure. More precisely, T_M is the highest temperature that results in the peak amplitude to be an order of magnitude larger than the minimum value of χ over the Brillouin zone, $\chi_{\max}/\chi_{\min} = 10$. Our results are summarized in Fig. 2.

As expected, the largest values of T_M (at fixed U) are observed at half-filling where the crossover temperature can be as high as $T \sim 0.25$ (or about 1000 K in units

Figure 7. (color online). *left*: Static magnetic susceptibility calculated from the lowest-order GGGW diagrams and plotted over the trace $(0,0) \rightarrow (\pi,0) \rightarrow (\pi,\pi) \rightarrow (0,0)$ in the Brillouin zone for $T = 0.05$, $U = 4$ and $n = 0.8085$. Two peaks are observed at the incommensurate vectors (π, Q) and (Q, Q) , where Q is some value other than π . *right*: The corresponding diagonally striped “carpet” pattern is plotted in real space with colors reflecting the sign of the average spin correlation. Whether this pattern survives or changes as $T \rightarrow 0$ remains an open question.

representative of the CuO_2 superconductors with hopping amplitude $t \sim 0.3 \text{ eV}$ [44]). As a function of interaction, T_M goes to zero at both large and small values of U , and within the considered range of $1 \leq U \leq 6$ features a smooth maximum around $U \sim 4$. This appears to be the optimal spot for experimental studies of magnetism in (1) where reaching low temperatures remains challenging. The magnetic crossover scale goes to zero with doping but remains relatively high for intermediate values of U even at doping levels $\delta \in (0.15, 0.25)$. We did not see evidence for PS at T_M , meaning that the PS dome takes place within the magnetic region, see Fig. 1.

The character of spin correlations undergoes a dramatic transformation with doping. A mismatch between the largest momentum transfer at the Fermi surface and the reciprocal lattice vector results in the incommensurate spin-wave fluctuations that take the form of AFM domains and *diagonal* domain walls. In Fig. 7, we show a typical example of the emerging structure (for $T = 0.05$, $U = 4$ and $n = 0.8085$). In the left panel, we see that the otherwise dominant peak around the commensurate vector (π, π) is split and features a minimum at (π, π) surrounded by two maxima at the incommensurate vectors. The real-space spin texture behind this split-peak signal is shown in the right panel with different colors corresponding to the sign of $\chi(0, \vec{r})$. For brevity, we will call it “diagonal stripes.” It is plausible that in the putative PS region, see Fig. 1, the AFM order is intermixed with diagonal stripes, and the mechanism for the $d_{x^2-y^2}$ -wave pairing is based on coupling to these spin fluctuations.

Conclusions. We discussed the finite-temperature phase diagram of the repulsive Fermi Hubbard model on a square lattice and identified the overarching dome defining the onset of strong magnetic correlations that change their structure from commensurate antiferromag-

netism to incommensurate diagonal stripes as the doping level is increased. Given relatively high values of T_M that extend well into the doping region where optimal values for transition temperatures to the $d_{x^2-y^2}$ superfluidity are expected to take place, magnetic correlations appear to be the key ingredient behind both the PS and BCS phenomena near half-filling.

Further development of the Diagrammatic Monte Carlo approach is required to obtain controllable results at temperatures below T_M where convergence of the diagrammatic expansion becomes problematic. Large magnetic correlation length and phase separation are two factors that have to be treated with extreme care by any numerical method based on finite-cluster simulations because they impose restrictions on the minimal acceptable cluster sizes and question the homogeneity of the solution. In particular, superfluid states proposed in [44–46] can not exist in the parameter space of phase separation. To find high- T_c regions one has to avoid PS near half-filling or modify the model to include non-zero values of the next-nearest-neighbor hopping $t' > 0$ [45].

Ultra-cold atom experiments are expected to have a major impact on revealing the finite-temperature phase diagram. Current experiments have already reached temperatures $T \sim T_M$ [14], and are well positioned to explore the structure of strong magnetic correlations. The detection and characterization of the PS state requires lower temperature scales but we do not see any reason for the PS dome to take place at $T \ll T_M$ given that known correlations are getting saturated below T_M .

Quantifying magnetic correlations is also of significant interest in relation with Copper oxides [47] as neutron scattering experiments have revealed the coexistence of commensurate and incommensurate magnetic structures at finite doping. For $\text{La}_{2-p}\text{Sr}_p\text{Cu}_4$ an incommensurate state with a magnetic structure wave vector was found at small dopings; for $\text{YBa}_2\text{Cu}_3\text{O}_{6+y}$ a wide doping window exists where commensurate AFM fluctuations are observed at low temperatures [48].

The authors are grateful to Shiwei Zhang, Sandro Sorella, Luca F. Tocchio, and Markus Greiner for useful discussions of their results. The authors would also like to thank Evgeni Burowski for permitting the use and further development of his original DDMC code. Fruitful exchanges with Ulrich Schollwöck about the possibility of phase separation at larger U ’s are acknowledged. Fedor Šimkovic would like to thank USTC Hefei for generous hospitality during a period when parts of this paper were written. This work was supported by the Simons Collaboration on the Many Electron Problem, National Science Foundation under the grant PHY-1314735, the MURI Program “New Quantum Phases of Matter” from AFOSR, and the Swiss National Science Foundation, NSFC Grant No. 11625522, CAS, and NKBRSC Grant No. 2016YFA0301600.

-
- [1] J. Hubbard, in *Proceedings of the royal society of london a: mathematical, physical and engineering sciences*, Vol. 276 (The Royal Society, 1963) pp. 238–257.
- [2] P. W. Anderson, *Solid state physics* **14**, 99 (1963).
- [3] P. W. Anderson *et al.*, *The theory of superconductivity in the high-T_c cuprate superconductors*, Vol. 446 (Princeton University Press Princeton, NJ, 1997).
- [4] P. A. Lee, N. Nagaosa, and X.-G. Wen, *Rev. Mod. Phys.* **78**, 17 (2006).
- [5] D. Jaksch, C. Bruder, J. I. Cirac, C. W. Gardiner, and P. Zoller, *Physical Review Letters* **81**, 3108 (1998).
- [6] M. Lewenstein, A. Sanpera, V. Ahufinger, B. Damski, A. Sen, and U. Sen, *Advances in Physics* **56**, 243 (2007).
- [7] M. Köhl, H. Moritz, T. Stöferle, K. Günter, and T. Esslinger, *Physical Review Letters* **94**, 080403 (2005).
- [8] R. Jördens, N. Strohmaier, K. Günter, H. Moritz, and T. Esslinger, *Nature* **455**, 204 (2008).
- [9] U. Schneider, L. Hackermüller, S. Will, T. Best, I. Bloch, T. Costi, R. W. Helmes, D. Rasch, and A. Rosch, *Science* **322**, 1520 (2008).
- [10] R. G. Hulet, P. M. Duarte, R. A. Hart, and T.-L. Yang, *arXiv preprint arXiv:1512.05311* (2015).
- [11] R. A. Hart, P. M. Duarte, T.-L. Yang, X. Liu, T. Paiva, E. Khatami, R. T. Scalettar, N. Trivedi, D. A. Huse, and R. G. Hulet, *Nature* **519**, 211 (2015).
- [12] D. Greif, G. Jotzu, M. Messer, R. Desbuquois, and T. Esslinger, *Physical Review Letters* **115**, 260401 (2015).
- [13] M. F. Parsons, A. Mazurenko, C. S. Chiu, G. Ji, D. Greif, and M. Greiner, *Science* **353**, 1253 (2016).
- [14] A. Mazurenko, C. S. Chiu, G. Ji, M. F. Parsons, M. Karsz-Nagy, R. Schmidt, F. Grusdt, E. Demler, D. Greif, and M. Greiner, *Nature* **545**, 462 (2017).
- [15] M. Y. Kagan and A. V. Chubukov, *Soviet Journal of Experimental and Theoretical Physics Letters* **50**, 517 (1989).
- [16] M. A. Baranov and M. Y. Kagan, *Zeitschrift für Physik B Condensed Matter* **86**, 237 (1992).
- [17] A. V. Chubukov and J. P. Lu, *Physical Review B* **46**, 11163 (1992).
- [18] A. V. Chubukov, *Physical Review B* **48**, 1097 (1993).
- [19] D. Zanchi and H. J. Schulz, *Physical Review B* **54**, 9509 (1996).
- [20] C. J. Halboth and W. Metzner, *Physical Review Letters* **85**, 5162 (2000).
- [21] H. Fukazawa and K. Yamada, *Journal of the Physical Society of Japan* **71**, 1541 (2002).
- [22] R. Hlubina, *Physical Review B* **59**, 9600 (1999).
- [23] S. Raghu, S. A. Kivelson, and D. J. Scalapino, *Physical Review B* **81**, 224505 (2010).
- [24] A. T. Rømer, A. Kreisel, I. Eremin, M. A. Malakhov, T. A. Maier, P. J. Hirschfeld, and B. M. Andersen, *Physical Review B* **92**, 104505 (2015).
- [25] F. Šimkovic IV, X.-W. Liu, Y. Deng, and E. Kozik, *Physical Review B* **94**, 085106 (2016).
- [26] Y. Deng, E. Kozik, N. V. Prokof'ev, and B. V. Svistunov, *EPL (Europhysics Letters)* **110**, 57001 (2015).
- [27] J. E. Hirsch, *Physical Review B* **31**, 4403 (1985).
- [28] J. E. Hirsch and S. Tang, *Physical Review Letters* **62**, 591 (1989).
- [29] S. R. White, D. J. Scalapino, R. L. Sugar, E. Y. Loh, J. E. Gubernatis, and R. T. Scalettar, *Physical Review B* **40**, 506 (1989).
- [30] N. Furukawa and M. Imada, *Journal of the Physical Society of Japan* **61**, 3331 (1992).
- [31] C. N. Varney, C.-R. Lee, Z. J. Bai, S. Chiesa, M. Jarrell, and R. T. Scalettar, *Physical Review B* **80**, 075116 (2009).
- [32] D. R. Penn, *Physical Review* **142**, 350 (1966).
- [33] Y. Nagaoka, *Physical Review* **147**, 392 (1966).
- [34] H. Tasaki, *Progress of Theoretical Physics* **99**, 489 (1998).
- [35] R. Zitzler, T. Pruschke, and R. Bulla, *The European Physical Journal B-Condensed Matter and Complex Systems* **27**, 473 (2002).
- [36] H. J. Schulz, *Physical Review Letters* **64**, 1445 (1990).
- [37] H. Q. Lin, *Physical Review B* **44**, 7151 (1991).
- [38] C.-C. Chang and S. Zhang, *Physical Review B* **78**, 165101 (2008).
- [39] S. Sorella, *Phys. Rev. B* **91**, 241116 (2015).
- [40] L. F. Tocchio, F. Becca, and S. Sorella, *Phys. Rev. B* **94**, 195126 (2016).
- [41] K. Van Houcke, E. Kozik, N. Prokofev, and B. Svistunov, *Physics Procedia* **6**, 95 (2010).
- [42] S. A. Kulagin, N. Prokof'ev, O. A. Starykh, B. Svistunov, and C. N. Varney, *Phys. Rev. B* **87**, 024407 (2013).
- [43] E. Burovski, N. Prokofev, B. Svistunov, and M. Troyer, *Physical Review Letters* **96**, 160402 (2006).
- [44] E. Gull, O. Parcollet, and A. J. Millis, *Physical Review Letters* **110**, 216405 (2013).
- [45] X. Chen, J. P. F. LeBlanc, and E. Gull, *Physical Review Letters* **115**, 116402 (2015).
- [46] B.-X. Zheng and G. K.-L. Chan, *Physical Review B* **93**, 035126 (2016).
- [47] M. A. Kastner, R. J. Birgeneau, G. Shirane, and Y. Endoh, *Reviews of Modern Physics* **70**, 897 (1998).
- [48] M. Matsuda, M. Fujita, K. Yamada, R. J. Birgeneau, Y. Endoh, and G. Shirane, *Physical Review B* **65**, 134515 (2002).