



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Out-of-time-order correlators in finite open systems

S. V. Syzranov, A. V. Gorshkov, and V. Galitski

Phys. Rev. B **97**, 161114 — Published 26 April 2018

DOI: [10.1103/PhysRevB.97.161114](https://doi.org/10.1103/PhysRevB.97.161114)

Out-of-time-order correlators in finite open systems

S.V. Syzranov,^{1,2,3} A.V. Gorshkov,^{2,4} and V. Galitski²

¹*Physics Department, University of California, Santa Cruz, CA 95064, USA*

²*Joint Quantum Institute, NIST/University of Maryland, College Park, MD 20742, USA*

³*School of Physics and Astronomy, Monash University, Victoria 3800, Australia*

⁴*Joint Center for Quantum Information and Computer Science, NIST/University of Maryland, College Park, MD 20742, USA*

(Dated: April 4, 2018)

We study out-of-time-order correlators (OTOCs) of the form $\langle \hat{A}(t)\hat{B}(0)\hat{C}(t)\hat{D}(0) \rangle$ for a quantum system weakly coupled to a dissipative environment. Such an open system may serve as a model of, e.g., a small region in a disordered interacting medium coupled to the rest of this medium considered as an environment. We demonstrate that for a system with discrete energy levels the OTOC saturates exponentially $\propto \sum a_i e^{-t/\tau_i} + \text{const}$ to a constant value at $t \rightarrow \infty$, in contrast with quantum-chaotic systems which exhibit exponential growth of OTOCs. Focussing on the case of a two-level system, we calculate microscopically the decay times τ_i and the value of the saturation constant. Because some OTOCs are immune to dephasing processes and some are not, such correlators may decay on two sets of parametrically different time scales related to inelastic transitions between the system levels and to pure dephasing processes, respectively. In the case of a classical environment, the evolution of the OTOC can be mapped onto the evolution of the density matrix of two systems coupled to the same dissipative environment.

Quantum information spreading in a quantum system is often described by out-of-time-order correlators (OTOCs) of the form

$$K(t) = \left\langle \hat{A}(t)\hat{B}(0)\hat{C}(t)\hat{D}(0) \right\rangle, \quad (1)$$

where \hat{A} , \hat{B} , \hat{C} and \hat{D} are Hermitian operators, and $\langle \dots \rangle$ is the average with respect to the initial state of the system. Correlators of such form have been first introduced by A. Larkin and Y.N. Ovchinnikov¹ in the context of disordered conductors, where the correlator $\langle [p_z(t), p_z(0)]^2 \rangle$ of particle momenta p_z has been demonstrated to grow exponentially $\propto e^{2\lambda t}$ on times $\tau_0 \ll t \ll t_E$, where the exponent λ characterises the rate of divergence of two classical electron trajectories with slightly different initial conditions, τ_0 is the elastic scattering time and t_E is the Ehrenfest time, characterising the crossover between classical and quantum dynamics².

The concept of OTOC has revived³ recently in the context of quantum information scrambling and black holes, motivating further studies of such quantities (see, e.g., Refs. 4–8). Despite not being measurable observables⁹, OTOCs (1) characterise the spreading of quantum information and the sensitivity of the system to the change of the initial conditions. At present, the exponential growth of commutators of the form $\langle [\hat{W}(t), \hat{V}(0)]^2 \rangle$ in a sufficiently large interval of time is often used as a definition of quantum chaos (for a discussion of alternative definitions see, e.g., Ref.¹⁰), and the Lyapunov exponent λ serves as a measure of quantum chaotic behaviour in a system. It is also expected that OTOCs may be used^{5,11–14} to identify the many-body-localisation transition¹⁵.

So far the studies of quantum chaos and information scrambling have been focussing on closed quantum systems. In reality, however, each system is coupled to

a noisy environment, which leads to decoherence and affects information spreading. Moreover, a sufficiently strongly disordered interacting system may be separated into a small subsystem, of the size of the single-particle localisation length or a region of quasi-localised states, coupled to the rest of the system considered as environment. In this paper we analyse out-of-time-order correlators in a quantum system weakly coupled to a dissipative environment.

Summary of the results. We demonstrate that, for a system with discrete non-degenerate levels E_n , correlator (1) at long times t exponentially saturates to a constant value, $K(t) \propto \sum a_n e^{i\omega_n t} e^{-t/\tau_n} + \text{const}$, in contrast with chaotic systems, where OTOCs contain exponentially growing contributions. The times τ_n describe the rates of information scrambling in the system and are distinct from the dephasing and decoherence times which describe the decay rates of the density matrix. We calculate microscopically the value of the saturation constant and the relaxation times τ_n as a function of the environment spectral function and the matrix elements of the




FIG. 1. Time dependence of the out-of-time-order correlator (1) in a system with significantly larger dephasing rate than relaxation. τ_{deph} and τ_{rel} are the (longest) characteristic times of dephasing and inelastic relaxation.

system-environment coupling. Depending on the choice of the operators \hat{A} , \hat{B} , \hat{C} and \hat{D} , the saturation value may be finite or zero. OTOCs relax due to both inelastic transitions between the system's levels and pure dephasing processes caused by slow fluctuations of the energies E_n . While some OTOCs are immune to dephasing processes, a generic correlator has components both sensitive and insensitive to dephasing and thus decays on two sets of parametrically different scales related to dephasing and relaxation respectively, as shown in Fig. 1.

Our results suggest, in particular, that a disordered system of interacting particles cannot exhibit quantum chaotic behaviour if the typical single-particle level splitting δ_ξ in a volume of linear size ξ (single-particle localisation length) exceeds the dephasing rate and the rate of inelastic transitions due to interactions and/or phonons. Thus, chaotic behaviour in a disordered interacting system requires either a sufficiently large single-particle localisation length or strong interactions or, e.g., a phonon bath, which would lead to the quasiparticle decay rate exceeding the level spacing δ_ξ .

For a classical environment, the evolution of an OTOC (1) in an open system may be mapped onto the evolution of the density matrix of two systems coupled to the same environment, which allows one to measure OTOCs by observing the correlations between two systems in a noisy environment, such as spins in a random time-dependent magnetic field.

Model. We consider a system with discrete *non-degenerate* energy levels E_n coupled to a dissipative environment and described by the Hamiltonian

$$\hat{\mathcal{H}} = \hat{\mathcal{H}}_0 + \hat{V}\hat{X} + \hat{\mathcal{H}}_{\text{bath}}(\hat{X}), \quad (2)$$

where $\hat{\mathcal{H}}_0 = \sum_n E_n |n\rangle \langle n|$ is the Hamiltonian of the system, $\hat{\mathcal{H}}_{\text{bath}}(\hat{X})$ the Hamiltonian of the environment, and $\hat{V}\hat{X}$ is the coupling between the system and the environment, where the operator $\hat{V} = \sum_{n,m} V_{nm} |n\rangle \langle m|$ acts on the system degrees of freedom, and \hat{X} is an environment variable which commutes with the system degrees of freedom.

To compute the OTOC (1), where the operators \hat{A} , \hat{B} , \hat{C} and \hat{D} act on the system variables, it is convenient to decompose it as $K = K_{m_1 m_2, n_1 n_2} A_{n_1 m_1} C_{n_2 m_2}$ (summation over repeated indices implied), where $A_{n_1 m_1}$ and $C_{n_2 m_2}$ are the matrix elements of the operators \hat{A} and \hat{C} , and

$$K_{m_1 m_2, n_1 n_2} = \left\langle |n_1\rangle \langle m_1| (t) \hat{B}(0) |n_2\rangle \langle m_2| (t) \hat{D}(0) \right\rangle, \quad (3)$$

where $\langle \dots \rangle$ is the averaging with respect to both the system and environment states.

In the limit of a vanishing system-environment coupling \hat{V} , the correlators (3) oscillate with time, $K_{m_1 m_2, n_1 n_2} \propto e^{i(E_{n_1} + E_{n_2} - E_{m_1} - E_{m_2})t}$. A finite coupling between the system and the environment leads to dissipation and relaxation processes and thus to the decay

of the elements $K_{m_1 m_2, n_1 n_2}$. For a weak coupling considered in this paper, the characteristic decay times of the OTOCs significantly exceed the correlation time of the environment degrees of freedom, i.e. of the function $S(t - t') = \langle \hat{X}(t) \hat{X}(t') \rangle_{\text{env}}$, and the evolution of the elements is described by a system of Markovian Bloch-Redfield¹⁶ equations of the form¹⁷

$$\begin{aligned} \partial_t K_{m_1 m_2, n_1 n_2} = & \\ i(E_{n_1} + E_{n_2} - E_{m_1} - E_{m_2}) K_{m_1 m_2, n_1 n_2} & \\ - \sum_{m'_1, m'_2, n'_1, n'_2} \Gamma_{m_1 m_2, n_1 n_2}^{m'_1 m'_2, n'_1 n'_2} K_{m'_1 m'_2, n'_1 n'_2}. & \end{aligned} \quad (4)$$

From the definition of the elements (3) it follows that

$$\sum_{m,n} K_{nm, nm} = \left\langle \hat{B}(0) \hat{D}(0) \right\rangle = \text{const.} \quad (5)$$

Eq. (5) may also be derived from the microscopic equations of evolution, as we show in Ref.¹⁷.

Due to the smallness of the decay rates $\Gamma_{m_1 m_2, n_1 n_2}^{m'_1 m'_2, n'_1 n'_2}$ in Eq. (4), the evolution of each element $K_{m_1 m_2, n_1 n_2}$ is affected only by the elements $K_{m'_1 m'_2, n'_1 n'_2}$ with the same oscillation frequency $E_{n_1} + E_{n_2} - E_{m_1} - E_{m_2}$ (secular approximation). In this paper we consider systems with sufficiently non-degenerate energy spectra; if two elements oscillate with the same frequency, they may be different only by permutations of indices n_1 and n_2 and/or m_1 and m_2 .

For a generic N -level system there are $2N^2 - N$ elements (3) with zero energy gaps $E_{n_1} + E_{n_2} - E_{m_1} - E_{m_2}$ (with $m_1 = n_1$, $m_2 = n_2$ and/or $m_1 = n_2$, $m_2 = n_1$). These elements are immune to dephasing, i.e. to the accumulation of random phases caused by slow fluctuations of the energies E_{n_i} . Such vanishing of dephasing is similar to that in decoherence-free subspaces^{18,19} of multiple-qubit systems. We emphasise, however, that even dephasing-immune correlators in general decay at long times due to the environment-induced inelastic transitions between the levels (relaxation processes).

A generic OTOC (1) includes components both sensitive and insensitive to dephasing, as well as a component independent of time, which exists due to the conservation law (5). For an environment with a smooth spectral function on the scale of the characteristic level splitting, the characteristic decay rate of the dephasing-immune components may be estimated as $1/\tau_{\text{rel}} \sim V_{\perp}^2 S(\Delta E)$, where V_{\perp} is the typical off-diagonal matrix element of the perturbation \hat{V} and ΔE is the characteristic level spacing. The other components decay with the characteristic rate $1/\tau_{\text{deph}} + 1/\tau_{\text{rel}}$, where $1/\tau_{\text{deph}} \sim V_{\parallel}^2 S(0)$ is the characteristic dephasing rate, where V_{\parallel} is the typical diagonal matrix element of the perturbation \hat{V} . As a result, the decay of the OTOC consist of three stages, corresponding to these characteristic times, as illustrated in Fig. 1.

Two-level system. In order to illustrate the meaning of these time scales and the related phenomena, we focus

below on the case of a two-level system, equivalent to a spin-1/2 in a random magnetic field (we provide microscopic analysis of OTOCs for a generic multi-level system in Ref.¹⁷), described by the Hamiltonian

$$\hat{\mathcal{H}} = \frac{1}{2}B\hat{\sigma}_z + \frac{1}{2}\hat{\boldsymbol{\sigma}}\mathbf{n}\hat{X} + \hat{\mathcal{H}}_{bath}(\hat{X}), \quad (6)$$

where $\hat{\boldsymbol{\sigma}}$ is a vector of Pauli matrices and \mathbf{n} is a constant unit vector, the direction of the fluctuations of the magnetic field.

The dissipative environment induces transitions $|\uparrow\rangle \rightarrow |\downarrow\rangle$ with the rate $\Gamma_\downarrow = \frac{1}{4}(n_x^2 + n_y^2)S(B)$, as well as the opposite transitions $|\downarrow\rangle \rightarrow |\uparrow\rangle$ with the rate $\Gamma_\uparrow = \frac{1}{4}(n_x^2 + n_y^2)S(-B)$, where $S(\omega)$ is the environment spectrum, the Fourier-transform of $S(t-t') = \langle \hat{X}(t)\hat{X}(t') \rangle_{\text{env}}$. Weak fluctuations of the magnetic field in the longitudinal direction lead to dephasing with the rate $\Gamma^\phi = \frac{1}{2}n_z^2 S(0)$. We focus below on the long-time dynamics of the system and assume for simplicity that the rate Γ^ϕ of pure dephasing significantly exceeds the rates Γ_\uparrow and Γ_\downarrow of inelastic transitions between the levels of the spin; in the opposite case, all OTOC decay rates are of the same order of magnitude.

The OTOCs $K_{\uparrow\uparrow,\downarrow\downarrow}$ and $K_{\downarrow\downarrow,\uparrow\uparrow}$ oscillate with frequencies $\pm 2(E_\downarrow - E_\uparrow) = \mp 2B$ and have dephasing rate $4\Gamma^\phi$, the same as ± 1 -projection states of a spin-1 in magnetic field B ,

$$K_{\uparrow\uparrow,\downarrow\downarrow}, K_{\downarrow\downarrow,\uparrow\uparrow} \propto e^{\mp 2iBt} e^{-4\Gamma^\phi t}, \quad (7)$$

where we have neglected the small relaxation rates $\Gamma_{\uparrow,\downarrow} \ll \Gamma^\phi$.

There are 8 elements (3) which correspond to 3 spin indices pointing in one direction and one spin index pointing in the opposite direction. These elements oscillate with frequencies $\pm B$ and have the same dephasing rate as a spin-1/2,

$$K_{\uparrow\downarrow,\downarrow\downarrow}, K_{\downarrow\downarrow,\uparrow\uparrow}, K_{\downarrow\downarrow,\uparrow\downarrow}, \dots \propto e^{-\Gamma^\phi t}. \quad (8)$$

The behaviour of OTOCs at long times $t \gg 1/\Gamma^\phi$ is determined by the components with a vanishing frequency $E_{n_1} + E_{n_2} - E_{m_1} - E_{m_2}$ of coherent oscillations, because such components are insensitive to dephasing. For a spin-1/2, their evolution is described by the system of equations (as follows from the generic master equations¹⁷)

$$\partial_t \begin{pmatrix} K_{\uparrow\uparrow,\uparrow\downarrow} \\ K_{\uparrow\downarrow,\downarrow\uparrow} \\ K_{\uparrow\downarrow,\uparrow\downarrow} \\ K_{\downarrow\uparrow,\downarrow\uparrow} \\ K_{\uparrow\uparrow,\uparrow\uparrow} \\ K_{\downarrow\downarrow,\downarrow\downarrow} \end{pmatrix} = \begin{pmatrix} -\Gamma_\downarrow - \Gamma_\uparrow & 0 & -\Gamma_\downarrow & -\Gamma_\downarrow & \Gamma_\downarrow & \Gamma_\downarrow \\ 0 & -\Gamma_\downarrow - \Gamma_\uparrow & -\Gamma_\uparrow & -\Gamma_\uparrow & \Gamma_\uparrow & \Gamma_\uparrow \\ -\Gamma_\uparrow & -\Gamma_\downarrow & -\Gamma_\downarrow - \Gamma_\uparrow & 0 & \Gamma_\downarrow & \Gamma_\uparrow \\ -\Gamma_\uparrow & -\Gamma_\downarrow & 0 & -\Gamma_\downarrow - \Gamma_\uparrow & \Gamma_\downarrow & \Gamma_\uparrow \\ \Gamma_\uparrow & \Gamma_\downarrow & \Gamma_\uparrow & \Gamma_\uparrow & -2\Gamma_\downarrow & 0 \\ \Gamma_\uparrow & \Gamma_\downarrow & \Gamma_\downarrow & \Gamma_\downarrow & 0 & -2\Gamma_\uparrow \end{pmatrix} \begin{pmatrix} K_{\downarrow\uparrow,\uparrow\downarrow} \\ K_{\uparrow\downarrow,\downarrow\uparrow} \\ K_{\uparrow\downarrow,\uparrow\downarrow} \\ K_{\downarrow\uparrow,\downarrow\uparrow} \\ K_{\uparrow\uparrow,\uparrow\uparrow} \\ K_{\downarrow\downarrow,\downarrow\downarrow} \end{pmatrix}. \quad (9)$$

The rates of the long-time decay of OTOCs are given by the eigenvalues of the matrix in Eq. (9) (with minus sign) and are shown (except for the zero eigenvalue) in Fig. 2. Such a matrix always has a zero eigenvalue, due to the conservation law (5). The system also has a triply degenerate decay rate $\Gamma_\uparrow + \Gamma_\downarrow$. The other two decay rates are given by $\frac{1}{2} \left[3\Gamma_\uparrow + 3\Gamma_\downarrow \pm \left(\Gamma_\uparrow^2 + 34\Gamma_\uparrow\Gamma_\downarrow + \Gamma_\downarrow^2 \right)^{\frac{1}{2}} \right]$.

At long times $t \rightarrow \infty$ the correlator (1) saturates to a constant value determined by the projection of the OTOC (1) on the zero-decay-rate mode,

$$K(t \rightarrow \infty) = \frac{1}{\sqrt{2\Gamma_\uparrow^2 + 2\Gamma_\downarrow^2}} (\Gamma_\downarrow A_{\uparrow\downarrow} C_{\downarrow\uparrow} + \Gamma_\uparrow A_{\downarrow\uparrow} C_{\uparrow\downarrow} + \Gamma_\uparrow A_{\uparrow\uparrow} C_{\uparrow\uparrow} + \Gamma_\downarrow A_{\downarrow\downarrow} C_{\downarrow\downarrow}). \quad (10)$$

While we assumed a small inelastic relaxation rate in comparison with the dephasing rate, we emphasise that the result (10) for the saturation value of the OTOC holds for an arbitrary ratio of dephasing and relaxation rates.




FIG. 2. Non-zero rates (in units Γ_\downarrow) of long-time decay of out-of-time-order correlators in a two-level system as a function of the ratio $\Gamma_\uparrow/\Gamma_\downarrow$ of the transition rates between the system levels.

Mapping to the evolution of two systems for a classical environment. The evolution of the OTOCs (3) is similar

to that of the density-matrix elements

$$\rho_{m_1 m_2, n_1 n_2} = \left\langle \left\langle |m_1\rangle \langle n_1| (t) \right\rangle_{\text{Sys}_1} \left\langle |m_2\rangle \langle n_2| (t) \right\rangle_{\text{Sys}_2} \right\rangle_X \quad (11)$$

of a compound system consisting of two identical subsystems (“Sys₁” and “Sys₂”) coupled to the same dissipative environment, where m_1 and n_1 and m_2 and n_2 in Eq. (11) are the states of the first and the second subsystems respectively, $|m_i\rangle \langle n_i| (t)$ is an operator in the interaction representation, and $\langle \dots \rangle_X$ is the averaging with respect to the environment degrees of freedom. The Hamiltonian of such a compound system is given by

$$\hat{H} = \hat{H}_0 \otimes \mathbb{1} + \mathbb{1} \otimes \hat{H}_0 + \left(\hat{V} \otimes \mathbb{1} + \mathbb{1} \otimes \hat{V} \right) \hat{X} + \hat{H}_{\text{bath}}(\hat{X}), \quad (12)$$

where $\dots \otimes \dots$ is the product of the subsystem subspaces; \hat{H}_0 and \hat{V} are the Hamiltonian of each subsystem and its coupling to the environment, and the environment variable \hat{X} commutes with all degrees of freedom of subsystems “Sys₁” and “Sys₂”.

The evolution of the elements (3) and (11) is described by similar Markovian master equations (see Ref.¹⁷ for a microscopic derivation). In particular, in the limit of a classical environment ($\langle \langle \hat{X}(t) \hat{X}(t') \rangle \rangle_{\text{env}} = \langle \langle \hat{X}(t') \hat{X}(t) \rangle \rangle_{\text{env}}$), the evolution of OTOCs (3) can be mapped exactly onto that of the density matrix (11) of two systems coupled to this environment, as follows from the definitions of these quantities. The conservation law (5) is mapped then onto the conservation of the trace of the density matrix of a compound system consisting of two subsystems.

In the limit of a classical environment, the spectral function is even, $S(\omega) = S(-\omega)$, the relaxation rate $i \rightarrow j$ for each pair of levels i and j in a system matches the reverse rate $j \rightarrow i$. In particular, in the case of a two-level system $\Gamma_{\uparrow} = \Gamma_{\downarrow} = \Gamma$, and the OTOC has three decay rates at long times $t \gg \Gamma^{-1}$: $\Gamma_1 = 6\Gamma$, $\Gamma_2 = 2\Gamma$ (triply degenerate) and $\Gamma_3 = 0$ (doubly degenerate), as shown in Fig. 2. Due to the mapping, these rates match the decay rates of pair-wise correlators of observables in, e.g., an ensemble of spins in a uniform random magnetic field and thus may be conveniently measured in such ensembles.

We emphasise that the mapping between an OTOC and the evolution of two subsystems coupled to the same classical environment holds for an arbitrary system-environment coupling but not only in the limit of a weak coupling considered in this paper. This mapping suggests a way for measuring OTOCs in generic systems in the presence of classical environments through observing correlators $\left\langle \left\langle \hat{A}(t) \right\rangle_{\text{Sys}_1} \left\langle \hat{C}(t) \right\rangle_{\text{Sys}_2} \right\rangle_X$ of observables \hat{A} and \hat{C} between two systems.

Discussion. We computed OTOCs in a finite system weakly coupled to a dissipative environment. The model

of such a finite open system may be used to understand OTOC behaviour in sufficiently strongly disordered interacting media (in the presence or in the absence of a phonon bath).

A material with weak short-range interactions and strong disorder, which leads to the localisation of all single-particle states, exhibits insulating behaviour at low temperatures¹⁵. Local physical observables in such a system are strongly correlated only on short length scales, and their properties may be understood by considering a single “localisation cell”, particle states in a region of space of the size of order of the (single-particle) localisation length ξ , which may be assumed weakly coupled to the rest of the system.

The energy spectrum of the localisation cell may be probed via response functions of local operators in the cell, e.g., the response function $\chi(\omega) = \sum_{\alpha, \beta} \frac{(f_{\alpha} - f_{\beta}) |Q_{\alpha\beta}|^2}{E_{\alpha} - E_{\beta} + \omega + i0}$ of the charge Q in a region inside the cell to the voltage in this region, where E_{α} and E_{β} are the energies of many-body states and f_{α} is their distribution function. For temperatures smaller than a critical value, quasiparticles in the system have zero decay rate²⁰ (“superinsulating” regime¹⁵), and the system thus responds only at a discrete set of frequencies $\omega = E_i - E_j$, determined by the energy gaps between many-body states, as shown in Fig. 3. The OTOC (1) in this regime oscillates $K(t) \propto \sum_n a_n e^{i\omega_n t}$ with a discrete set of frequencies $\omega_n = E_{i_n} + E_{j'_n} - E_{j_n} - E_{i'_n}$. In the limit of a very large number of levels in a localisation cell, the sum of such oscillating contributions approaches a smooth function with exponentially growing contributions describing the “single-particle” chaos of Ref.¹, so long as dynamics on time scales shorter than the Thouless scale $t_{\text{Th}} = \xi^2/D$ is neglected. However, the energy levels remain discrete in the “superinsulating” regime.

When the temperature (or the interaction strength at a given temperature) exceeds a critical value, the




FIG. 3. Response of a localisation cell in an interacting system for various temperatures T (or interaction strengths for $T > 0$). ΔE_i are the energy gaps between the many-body levels in the cell, and Γ is the level width.

levels and response functions get broadened (“metallic” phase¹⁵), as illustrated in Fig. 3, becoming smoother with increasing temperature and/or interactions. Near the superinsulator-metal transition the characteristic level width Γ is significantly smaller than the gaps between levels, and the localisation cell may be considered as an open system weakly coupled to a dissipative environment. The same model may be applied also to a strongly disordered material with an external bath, such as a system of phonons, which provide a finite level width Γ at all finite temperatures. The local operators \hat{A} , \hat{B} , \hat{C} and \hat{D} in Eq. (1) do not necessarily act on states in one localisation cell, but may involve states in several cells close to each other. These cells may still be considered as a single quantum dot in a noisy environment so long as the level spacing in the dot exceeds the level width. Such a model of an open quantum dot may be also realised directly, e.g., using superconducting qubits or trapped cold atoms.

The results of this paper suggest, in particular, that a strongly disordered system, with the level spacing δ_ξ exceeding the effective dissipation rates Γ , is non-chaotic. While our results applies to weakly-conducting and insulating materials, for which the system-environment cou-

pling may be considered small, we leave it for a future study whether non-chaotic behaviour persists in systems strongly coupled to the environment (corresponding to an effectively continuous energy spectrum of a localisation cell).

For a classical environment, the evolution of an OTOC in an open system matches the evolution of correlators of observables between two identical systems coupled to the same environment, which may be used for measuring OTOCs in open systems in classical environments. The possibility to develop a similar measurement method for the case of a quantum environment is another question which deserves further investigation.

ACKNOWLEDGMENTS

We have benefited from discussions with Yidan Wang. V.G. and S.V.S. were supported by US-ARO (contract No. W911NF1310172), NSF-DMR 1613029 and Simons Foundation; A.V.G. and S.V.S. acknowledge support by NSF QIS, AFOSR, NSF PFC at JQI, ARO MURI, ARO and ARL CDQI. S.V.S. also acknowledges the hospitality of School of Physics and Astronomy at Monash University, where a part of this work was completed.

-
- ¹ A. Larkin and Y. N. Ovchinnikov, “Quasiclassical method in the theory of superconductivity,” *Sov. Phys. JETP* **28**, 960 (1969).
 - ² G.P. Berman and G.M. Zaslavsky, “Condition of stochasticity in quantum nonlinear systems,” *Physica A* **91**, 450 – 460 (1978).
 - ³ A. Kitaev, “A simple model of quantum holography,” (2015), KITP Programme: Entanglement in Strongly-Correlated Quantum Matter.
 - ⁴ J. Maldacena, S. H. Shenker, and D. Stanford, “A bound on chaos,” *JHEP* **8**, 106 (2016).
 - ⁵ B. Swingle and D. Chowdhury, “Slow scrambling in disordered quantum systems,” *Phys. Rev. B* **95**, 060201 (2017).
 - ⁶ Igor L. Aleiner, Lara Faoro, and Lev B. Ioffe, “Microscopic model of quantum butterfly effect: out-of-time-order correlators and traveling combustion waves,” (2016), arXiv:1609.01251.
 - ⁷ Efim B. Rozenbaum, Sriram Ganeshan, and Victor Galitski, “Lyapunov exponent and out-of-time-ordered correlator’s growth rate in a chaotic system,” (2016), arXiv:1609.01707.
 - ⁸ A. A. Patel and S. Sachdev, “Quantum chaos on a critical Fermi surface,” *Proceedings of the National Academy of Science* **114**, 1844–1849 (2017).
 - ⁹ Because an OTOC involves evolution backwards in time, measuring it requires using a second copy of the system^{21,22}; effectively reverting the sign of the Hamiltonian^{23–29} (e.g., by means of spin-echo-type techniques) or using time evolution controlled by an ancilla qubit^{23–25,30}.
 - ¹⁰ S. V. Syzranov, A. V. Gorshkov, and V. M. Galitski, “Interaction-induced transition in the quantum chaotic dynamics of a disordered metal,” *ArXiv e-prints* (2017), arXiv:1709.09296.
 - ¹¹ R. Fan, P. Zhang, H. Shen, and H. Zhai, “Out-of-Time-Order Correlation for Many-Body Localization,” *ArXiv e-prints* (2016), arXiv:1608.01914.
 - ¹² Y. Chen, “Universal logarithmic scrambling in many body localization,” *ArXiv e-prints* (2016), arXiv:1608.02765 [cond-mat.dis-nn].
 - ¹³ Y. Huang, Y.-L. Zhang, and X. Chen, “Out-of-time-ordered correlators in many-body localized systems,” *Annalen der Physik* **529**, 1600318 (2017).
 - ¹⁴ A. A. Patel, D. Chowdhury, S. Sachdev, and B. Swingle, “Quantum Butterfly Effect in Weakly Interacting Diffusive Metals,” *Physical Review X* **7**, 031047 (2017).
 - ¹⁵ D. M. Basko, I. L. Aleiner, and B. L. Altshuler, “Metal insulator transition in a weakly interacting many-electron system with localized single-particle states,” *Ann. Phys.* **321**, 1126 (2006).
 - ¹⁶ C.P. Slichter, *Principles of Magnetic Resonance*, Springer Series in Solid-State Sciences (Springer Berlin Heidelberg, 1996).
 - ¹⁷ See Supplemental Material at [URL will be inserted by publisher] for a microscopic derivation of the Bloch-Redfield equations of OTOC evolution in a multi-level system.
 - ¹⁸ P. Zanardi and M. Rasetti, “Noiseless quantum codes,” *Phys. Rev. Lett.* **79**, 3306–3309 (1997).
 - ¹⁹ L.-A. Wu and D. A. Lidar, “Creating decoherence-free subspaces using strong and fast pulses,” *Phys. Rev. Lett.* **88**, 207902 (2002).
 - ²⁰ When studying response functions, a system in the insulating state should be assumed coupled to an external bath,

with the value of the coupling sent to zero at the end of the calculation¹⁵. The quasiparticle decay rate is then given by the bath strength.

- ²¹ N. Y. Yao, F. Grusdt, B. Swingle, M. D. Lukin, D. M. Stamper-Kurn, J. E. Moore, and E. A. Demler, “Interferometric Approach to Probing Fast Scrambling,” ArXiv e-prints (2016), arXiv:1607.01801 [quant-ph].
- ²² A. Bohrdt, C. B. Mendl, M. Endres, and M. Knap, “Scrambling and thermalization in a diffusive quantum many-body system,” *New Journal of Physics* **19**, 063001 (2017).
- ²³ B. Swingle, G. Bentsen, M. Schleier-Smith, and P. Hayden, “Measuring the scrambling of quantum information,” *Phys. Rev. A* **94**, 040302 (2016).
- ²⁴ Guanyu Zhu, Mohammad Hafezi, and Tarun Grover, “Measurement of many-body chaos using a quantum clock,” *Phys. Rev. A* **94**, 062329 (2016).
- ²⁵ I. Danshita, M. Hanada, and M. Tezuka, “Creating and probing the Sachdev-Ye-Kitaev model with ultracold gases: Towards experimental studies of quantum gravity,” *Progress of Theoretical and Experimental Physics* **2017**, 083I01 (2017).
- ²⁶ M. Gärttner, J. G. Bohnet, A. Safavi-Naini, M. L. Wall, J. J. Bollinger, and A. M. Rey, “Measuring out-of-time-order correlations and multiple quantum spectra in a trapped-ion quantum magnet,” *Nature Physics* **13**, 781–786 (2017).
- ²⁷ Naoto Tsuji, Philipp Werner, and Masahito Ueda, “Exact out-of-time-ordered correlation functions for an interacting lattice fermion model,” *Phys. Rev. A* **95**, 011601 (2017).
- ²⁸ J. Li, R. Fan, H. Wang, B. Ye, B. Zeng, H. Zhai, X. Peng, and J. Du, “Measuring Out-of-Time-Order Correlators on a Nuclear Magnetic Resonance Quantum Simulator,” *Phys. Rev. X* **7**, 031011 (2017).
- ²⁹ N. Yunger Halpern, B. Swingle, and J. Dressel, “The quasiprobability behind the out-of-time-ordered correlator,” ArXiv e-prints (2017), arXiv:1704.01971 [quant-ph].
- ³⁰ Nicole Yunger Halpern, “Jarzynski-like equality for the out-of-time-ordered correlator,” *Phys. Rev. A* **95**, 012120 (2017).