



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Lack of a thermodynamic finite-temperature spin-glass phase in the two-dimensional randomly coupled ferromagnet

Zheng Zhu, Andrew J. Ochoa, and Helmut G. Katzgraber

Phys. Rev. B **97**, 174425 — Published 24 May 2018

DOI: [10.1103/PhysRevB.97.174425](https://doi.org/10.1103/PhysRevB.97.174425)

Lack of a thermodynamic finite-temperature spin-glass phase in the two-dimensional randomly-coupled ferromagnet

Zheng Zhu,¹ Andrew J. Ochoa,¹ and Helmut G. Katzgraber^{1,2,3}

¹*Department of Physics and Astronomy, Texas A&M University, College Station, Texas 77843-4242, USA*

²*IQB Information Technologies (IQBit), Vancouver, British Columbia, Canada V6B 4W4*

³*Santa Fe Institute, 1399 Hyde Park Road, Santa Fe, New Mexico 87501, USA*

The search for problems where quantum adiabatic optimization might excel over classical optimization techniques has sparked a recent interest in inducing a finite-temperature spin-glass transition in quasi-planar topologies. We have performed large-scale finite-temperature Monte Carlo simulations of a two-dimensional square-lattice bimodal spin glass with next-nearest ferromagnetic interactions claimed to exhibit a finite-temperature spin-glass state for a particular relative strength of the next-nearest to nearest interactions [Phys. Rev. Lett. **76**, 4616 (1996)]. Our results show that the system is in a paramagnetic state in the thermodynamic limit, despite zero-temperature simulations [Phys. Rev. B **63**, 094423 (2001)] suggesting the existence of a finite-temperature spin-glass transition. Therefore, deducing the finite-temperature behavior from zero-temperature simulations can be dangerous when corrections to scaling are large.

PACS numbers: 75.50.Lk, 75.40.Mg, 05.50.+q, 64.60.-i

I. INTRODUCTION

The advent of analog quantum annealing machines^{1–14} and, in particular, the D-Wave Inc.¹⁵ D-Wave 2X quantum annealer has sparked a new interest in the study of (quasi-) planar Ising spin glasses^{16–19} with finite-temperature transitions. While there have been multiple attempts to discern if the D-Wave quantum annealers display an advantage over conventional technologies^{20–40}, to date there are only few “success stories”^{34,41} where the analog quantum optimizers show an advantage over current conventional silicon-based computers. Recent results^{26,34} suggest that problems with a more complex energy landscape are needed to discern if quantum annealers can outperform current digital computers. In particular, the search for salient features in the energy landscape³⁴, the careful construction of problems with particular features^{34,34,40,41}, as well as the attempt to induce a finite-temperature spin-glass transition for lattices restricted to the quasi-two-dimensional topologies of the quantum chips⁴² have gained considerable attention. The quest for a finite-temperature spin-glass transition in quasi-two-dimensional topologies stems from the interest in creating an energy landscape that becomes more complex and rugged already at finite temperatures, such that thermal (sequential) simulated annealing⁴³ has a harder time in determining the optimal solution to an Ising-spin-glass-like optimization problem. On the other hand, quantum annealing should, in principle, be able to tunnel through barriers if these are thin enough. We emphasize that the comparison between simulated annealing — a well-known poor optimizer — and quantum annealing is based on the fact that both methods are sequential in nature. Comparisons to state-of-the-art optimization techniques⁴⁴ have been performed and shed a more complete light on the current situation.

Here we want to study the thermodynamic properties of a model proposed by N. Lemke and I. A. Campbell⁴⁵ — later analyzed in much detail in Refs.^{46–48} — that might have the desired finite-temperature spin-glass transition and, most importantly, be of a mostly-planar topology that can easily be

constructed with current superconducting flux qubits. Our results show that, unfortunately, for large enough system sizes the model is in a paramagnetic phase at finite temperatures for a parameter range where it is predicted to be a spin glass. We do note that this would have been surprising, because there is solid evidence that the lower critical dimensions of spin glasses is believed to be between 2 and 3 space dimensions^{49–51} — a value below which any phase transition to a spin-glass state only occurs at zero temperature.

The paper is structured as follows: In Sec. II we describe the model and numerical details, as well as the current understanding of its properties, followed by results and concluding remarks in Sec. III.

II. MODEL AND NUMERICAL DETAILS

In their letter⁴⁵, Lemke and Campbell argue that a finite-temperature spin-glass transition can be induced in two-dimensional planar topologies with next-nearest interactions. To be precise, the model is a two-dimensional square-lattice Ising spin glass with uniform ferromagnetic next-nearest interactions of strength J , in addition to random bimodal nearest-neighbor interactions of strength $\pm\lambda J$. The Hamiltonian of the model is

$$\mathcal{H} = - \sum_{\langle i,j \rangle} J_{ij} S_i S_j - J \sum_{\langle\langle i,j \rangle\rangle} S_i S_j, \quad (1)$$

where in Eq. (1) $S_i \in \{\pm 1\}$ represent Ising spins on a square lattice with $N = L^2$ sites (L is the linear dimension of the lattice). $J = 1$ are ferromagnetic interactions between next-nearest neighbors (denoted by $\langle\langle i,j \rangle\rangle$) and $J_{ij} = \pm\lambda J$ are nearest-neighbor bimodally-distributed spin-glass interactions (denoted by $\langle i,j \rangle$). In our simulations we set $J = 1$. Depending on the relative strength of the interactions, i.e., the value of λ , Ref.⁴⁵ states that a finite-temperature spin-glass transition can be induced in two space dimensions. These results were further expanded in Ref.⁴⁷: A freezing temperature




FIG. 1: Binder cumulant g_q for the spin-glass order parameter as a function of the temperature T for the model described in Ref.⁴⁵ with $\lambda = 0.50$ (a) and $\lambda = 0.75$ (b) and system sizes $L > \ell$. In both cases the data show no crossing at any finite temperature studied, thus suggesting that there is no finite-temperature spin-glass phase. Square of the magnetization m^2 as a function of T for different system sizes for $\lambda = 0.50$ (c) and $\lambda = 0.75$ (d). The data decreases with increasing system size, i.e., the system is likely in a paramagnetic phase.

of $T_c \sim 2.1$ exists for $\lambda = 0.5$, a “slightly lower” freezing temperature for $\lambda = 0.7$, and a zero-temperature freezing for $\lambda = 1.5$. We do emphasize, that these results were produced by relatively small system sizes. Extensive numerical simulations by Parisi *et al.*⁴⁶ find a *crossover* in the critical behavior for large enough system sizes. First, from a seemingly ordered state to a spin-glass-like state, followed by a second crossover to a (possibly) paramagnetic state. This means that the true thermodynamic behavior can only be observed if the system sizes exceed a certain break up length ℓ .

However, a conclusive characterization of the critical behavior, as well as the λ -dependence of the break up length ℓ were not discussed in detail until the extensive zero-temperature study by Hartmann and Campbell⁴⁸. By computing ground-state configurations for intermediate system sizes and estimating the stiffness exponent that describes the scaling of energy excitations when a domain is introduced into the system, they argue—based on *zero-temperature* estimates of the spin stiffness—that there should be a finite-temperature spin-glass transition for certain values of λ and linear system

sizes L that fulfill $L > \ell$. In particular, they estimate that for $\lambda > \lambda_\infty = 0.27(8)$ no ferromagnetic order should be present. Because the break up length ℓ is large for $\lambda \sim 0.5$ ($\ell \gtrsim 45$), Ref.⁴⁸ suggests studying systems with $\lambda = 0.7$ where $\ell \approx 10$. On the other hand, for $\lambda = 0.90$, the stiffness exponent $\theta = 0.09(5)$ is very close to zero. Therefore, in this work we focus on the cases where (i) we can simulate system sizes $L \gg \ell$ and (ii) the stiffness exponent θ is clearly positive, thus implying a finite-temperature phase, i.e., $\lambda = 0.50$ and 0.75 . A summary of the properties of the model for these values of λ , as well as the simulation parameters are listed in Table I. The simulations were performed using parallel tempering Monte Carlo⁵² combined with isoenergetic cluster updates^{53,54}. Note that we determine the estimated value of θ for $\lambda = 0.75$ by performing a linear fit to the data of Ref.⁴⁸ (quality of fit ~ 0.58 ⁵⁵) and estimate $\theta(\lambda) \approx 1.083(3) - 1.12(4)\lambda$, valid in the window $\lambda \in [0.5, 1.1]$. Furthermore, by inspecting Fig. 7 in Ref.⁴⁸, we estimate that the break up length for $\lambda = 0.75$ is approximately $\ell \approx 9$.




FIG. 2: Finite-size scaling of the data shown in Fig. 1(b) for $\lambda = 0.75$ with $T_c = 0$.

TABLE I: Simulation parameters and estimates of the stiffness exponent θ and break up length ℓ for different values of λ . For both values of λ we studied different system sizes L using parallel tempering Monte Carlo. The lowest [highest] temperature simulated is $T_{\min} = 0.4$ [$T_{\max} = 2.8$] with $N_T = 50$ temperature steps. Thermalization is tested by a logarithmic binning; once the last three bins agree within error bars we deem the system to be thermalized. For all systems, this was the case after $N_{\text{sw}} = 2^{22}$ Monte Carlo sweeps. Furthermore, N_{sa} samples were computed for each parameter combination. Note that the estimate of θ for $\lambda = 0.50$ is taken from Ref.⁴⁸, whereas the value for $\lambda = 0.75$ is estimated from the published data (see text for details).

λ	θ	ℓ	L	N_{sw}	T_{\min}	T_{\max}	N_T	N_{sa}
0.50	0.59(8)	45	48	2^{22}	0.4	2.8	50	10^4
			64	2^{22}	0.4	2.8	50	10^4
			96	2^{22}	0.4	2.8	50	10^4
			128	2^{22}	0.4	2.8	50	10^4
0.75	0.23(1)	9	24	2^{22}	0.4	2.8	50	10^4
			32	2^{22}	0.4	2.8	50	10^4
			48	2^{22}	0.4	2.8	50	10^4
			64	2^{22}	0.4	2.8	50	10^4

To detect the existence of a spin-glass transition, we measure the Binder cumulant g^{56} of the spin-glass order parameter q via

$$g_q = \frac{1}{2} \left(3 - \frac{[\langle q^4 \rangle]_{\text{av}}}{[\langle q^2 \rangle]_{\text{av}}^2} \right). \quad (2)$$

In Eq. (2), $\langle \dots \rangle$ represents a thermal average over Monte Carlo steps and $[\dots]_{\text{av}}$ an average over N_{sa} realizations of the disorder (see Table I for details). The spin-glass order parameter q is given by

$$q = \frac{1}{N} \sum_{i=1}^N S_i^\alpha S_i^\beta, \quad (3)$$

where “ α ” and “ β ” represent two copies of the system with the same disorder. The Binder cumulant is dimensionless and

scales as $g_q = G[L^{1/\nu}(T - T_c)]$. Therefore, if $T = T_c$, data for different system sizes cross. If, however, there is no transition, data for different system sizes do not cross. To rule out a transition at a temperature not simulated, a finite-size scaling of the data can be used. Finally, we also measure the average of the square of the magnetization $m^2 \equiv [\langle m^2 \rangle]_{\text{av}}$ with

$$m = \frac{1}{N} \sum_{i=1}^N S_i^\alpha. \quad (4)$$

Note that we measure the square of the magnetization because, on average, $m \equiv [\langle m \rangle]_{\text{av}} = 0$. Furthermore, the magnetic susceptibility χ_m is related to m^2 via $\chi_m = Nm^2$.

III. RESULTS AND CONCLUSIONS

We have performed large-scale Monte Carlo simulations of the Hamiltonian in Eq. (1) for system sizes $L \gg \ell$ and $\lambda = 0.50$ and 0.75 . Our results for the Binder cumulant—which should display a crossing if there is a finite-temperature transition—are summarized in Fig. 1. The Binder cumulant for the spin-glass order parameter g_q does not show a crossing down to low temperatures for both values of λ studied. In addition, a finite-size scaling of the data for $\lambda = 0.75$ shown in Fig. 2 strongly suggests that $T_c = 0$. Furthermore, the magnetization m^2 as a function of the temperature T decreases with increasing system sizes for both values of λ studied (see Fig. 1.). Based on these results, we conclude that the system is in a *paramagnetic* state for both $\lambda = 0.50$ and 0.75 in the thermodynamic limit.

Our results show that the model introduced in Ref.⁴⁵ and studied in detail in subsequent publications^{46–48} does not exhibit a finite-temperature spin-glass transition in the thermodynamic limit for values of the parameter λ where it is expected to show such behavior. In agreement with the results of Ref.⁴⁶, however for larger system sizes and better statistics, we show that, indeed, the thermodynamic limit is a paramagnetic phase at finite temperature. This also means that deducing a finite-temperature behavior from zero-temperature simulations can be dangerous when the system sizes are not in the thermodynamic limit⁴⁸. Given recent interest in inducing finite-temperature spin-glass transitions in quasi-planar topologies²⁶, we conjecture that adding any set of interactions that do not grow with the system size to a nearest-neighbor lattice will likely not result in a finite-temperature spin-glass transition.

Acknowledgments

H.G.K. and Z.Z. acknowledge support from the National Science Foundation (Grant No. DMR-1151387). We would like to thank Texas A&M University for HPC resources. H.G.K. would also like to thank Warinomo (New Republic) for inspiration during the final stages of this manuscript. Part

of this work is supported in part by the Office of the Director of National Intelligence (ODNI), Intelligence Advanced Research Projects Activity (IARPA), via MIT Lincoln Laboratory Air Force Contract No. FA8721-05-C-0002. The views and conclusions contained herein are those of the authors and should not be interpreted as necessarily representing the of-

ficial policies or endorsements, either expressed or implied, of ODNI, IARPA, or the U.S. Government. The U.S. Government is authorized to reproduce and distribute reprints for Governmental purpose notwithstanding any copyright annotation thereon.

-
- ¹ M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2000).
- ² H. Nishimori, *Statistical Physics of Spin Glasses and Information Processing: An Introduction* (Oxford University Press, New York, 2001).
- ³ A. B. Finnila, M. A. Gomez, C. Sebenik, C. Stenson, and J. D. Doll, *Quantum annealing: A new method for minimizing multidimensional functions*, Chem. Phys. Lett. **219**, 343 (1994).
- ⁴ T. Kadowaki and H. Nishimori, *Quantum annealing in the transverse Ising model*, Phys. Rev. E **58**, 5355 (1998).
- ⁵ J. Brooke, D. Bitko, T. F. Rosenbaum, and G. Aeppli, *Quantum annealing of a disordered magnet*, Science **284**, 779 (1999).
- ⁶ E. Farhi, J. Goldstone, S. Gutmann, and M. Sipser, *Quantum Computation by Adiabatic Evolution* (2000), arXiv:quant-ph/0001106.
- ⁷ J. Roland and N. J. Cerf, *Quantum search by local adiabatic evolution*, Phys. Rev. A **65**, 042308 (2002).
- ⁸ G. Santoro, E. Martoňák, R. Tosatti, and R. Car, *Theory of quantum annealing of an Ising spin glass*, Science **295**, 2427 (2002).
- ⁹ A. Das and B. K. Chakrabarti, *Quantum Annealing and Related Optimization Methods* (Edited by A. Das and B.K. Chakrabarti, Lecture Notes in Physics 679, Berlin: Springer, 2005).
- ¹⁰ G. E. Santoro and E. Tosatti, *TOPICAL REVIEW: Optimization using quantum mechanics: quantum annealing through adiabatic evolution*, J. Phys. A **39**, R393 (2006).
- ¹¹ D. A. Lidar, *Towards Fault Tolerant Adiabatic Quantum Computation*, Phys. Rev. Lett. **100**, 160506 (2008).
- ¹² A. Das and B. K. Chakrabarti, *Quantum Annealing and Analog Quantum Computation*, Rev. Mod. Phys. **80**, 1061 (2008).
- ¹³ S. Morita and H. Nishimori, *Mathematical Foundation of Quantum Annealing*, J. Math. Phys. **49**, 125210 (2008).
- ¹⁴ S. Mukherjee and B. K. Chakrabarti, *Multivariable optimization: Quantum annealing and computation*, Eur. Phys. J. Special Topics **224**, 17 (2015).
- ¹⁵ URL <http://www.dwavesys.com>.
- ¹⁶ M. Mézard, G. Parisi, N. Sourlas, G. Toulouse, and M. Virasoro, *Nature of the Spin-Glass Phase*, Phys. Rev. Lett. **52**, 1156 (1984).
- ¹⁷ K. Binder and A. P. Young, *Spin Glasses: Experimental Facts, Theoretical Concepts and Open Questions*, Rev. Mod. Phys. **58**, 801 (1986).
- ¹⁸ A. P. Young, ed., *Spin Glasses and Random Fields* (World Scientific, Singapore, 1998).
- ¹⁹ D. L. Stein and C. M. Newman, *Spin Glasses and Complexity*, Primers in Complex Systems (Princeton University Press, Princeton NJ, 2013).
- ²⁰ N. G. Dickson, M. W. Johnson, M. H. Amin, R. Harris, F. Altomare, A. J. Berkley, P. Bunyk, J. Cai, E. M. Chapple, P. Chavez, et al., *Thermally assisted quantum annealing of a 16-qubit problem*, Nat. Commun. **4**, 1903 (2013).
- ²¹ K. L. Pudenz, T. Albash, and D. A. Lidar, *Error-corrected quantum annealing with hundreds of qubits*, Nat. Commun. **5**, 3243 (2014).
- ²² G. Smith and J. Smolin, *Putting “Quantumness” to the Test*, Physics **6**, 105 (2013).
- ²³ S. Boixo, T. Albash, F. M. Spedalieri, N. Chancellor, and D. A. Lidar, *Experimental signature of programmable quantum annealing*, Nat. Commun. **4**, 2067 (2013).
- ²⁴ T. Albash, T. F. Rønnow, M. Troyer, and D. A. Lidar, *Reexamining classical and quantum models for the D-Wave One processor* (2014), (arXiv:1409.3827).
- ²⁵ T. F. Rønnow, Z. Wang, J. Job, S. Boixo, S. V. Isakov, D. Wecker, J. M. Martinis, D. A. Lidar, and M. Troyer, *Defining and detecting quantum speedup* (2014), (arXiv:quant-ph/1401.2910).
- ²⁶ H. G. Katzgraber, F. Hamze, and R. S. Andrist, *Glassy Chimeras Could Be Blind to Quantum Speedup: Designing Better Benchmarks for Quantum Annealing Machines*, Phys. Rev. X **4**, 021008 (2014).
- ²⁷ T. Lanting, A. J. Przybysz, A. Y. Smirnov, F. M. Spedalieri, M. H. Amin, A. J. Berkley, R. Harris, F. Altomare, S. Boixo, P. Bunyk, et al., *Entanglement in a quantum annealing processor*, Phys. Rev. X **4**, 021041 (2014).
- ²⁸ S. Santra, G. Quiroz, G. Ver Steeg, and D. A. Lidar, *Max 2-SAT with up to 108 qubits*, New J. Phys. **16**, 045006 (2014).
- ²⁹ S. W. Shin, G. Smith, J. A. Smolin, and U. Vazirani, *How “Quantum” is the D-Wave Machine?* (2014), (arXiv:1401.7087).
- ³⁰ W. Vinci, T. Albash, A. Mishra, P. A. Warburton, and D. A. Lidar, *Distinguishing classical and quantum models for the D-Wave device* (2014), (arXiv:1403.4228).
- ³¹ S. Boixo, T. F. Rønnow, S. V. Isakov, Z. Wang, D. Wecker, D. A. Lidar, J. M. Martinis, and M. Troyer, *Evidence for quantum annealing with more than one hundred qubits*, Nat. Phys. **10**, 218 (2014).
- ³² T. Albash, W. Vinci, A. Mishra, P. A. Warburton, and D. A. Lidar, *Consistency Tests of Classical and Quantum Models for a Quantum Device*, Phys. Rev. A **91**, 042314 (2015).
- ³³ T. Albash, T. F. Rønnow, M. Troyer, and D. A. Lidar, *Reexamining classical and quantum models for the D-Wave One processor*, Eur. Phys. J. Spec. Top. **224**, 111 (2015).
- ³⁴ H. G. Katzgraber, F. Hamze, Z. Zhu, A. J. Ochoa, and H. Munoz-Bauza, *Seeking Quantum Speedup Through Spin Glasses: The Good, the Bad, and the Ugly*, Phys. Rev. X **5**, 031026 (2015).
- ³⁵ V. Martin-Mayor and I. Hen, *Unraveling Quantum Annealers using Classical Hardness* (2015), (arXiv:1502.02494).
- ³⁶ K. L. Pudenz, T. Albash, and D. A. Lidar, *Quantum Annealing Correction for Random Ising Problems*, Phys. Rev. A **91**, 042302 (2015).
- ³⁷ I. Hen, J. Job, T. Albash, T. F. Rønnow, M. Troyer, and D. A. Lidar, *Probing for quantum speedup in spin-glass problems with planted solutions*, Phys. Rev. A **92**, 042325 (2015).
- ³⁸ D. Venturelli, S. Mandrà, S. Knysh, B. O’Gorman, R. Biswas, and V. Smelyanskiy, *Quantum Optimization of Fully Connected Spin Glasses*, Phys. Rev. X **5**, 031040 (2015).
- ³⁹ W. Vinci, T. Albash, G. Paz-Silva, I. Hen, and D. A. Lidar, *Quantum annealing correction with minor embedding*, Phys. Rev. A **92**, 042310 (2015).

- ⁴⁰ Z. Zhu, A. J. Ochoa, F. Hamze, S. Schnabel, and H. G. Katzgraber, *Best-case performance of quantum annealers on native spin-glass benchmarks: How chaos can affect success probabilities*, Phys. Rev. A **93**, 012317 (2016).
- ⁴¹ V. S. Denchev, S. Boixo, S. V. Isakov, N. Ding, R. Babbush, V. Smelyanskiy, J. Martinis, and H. Neven, *What is the Computational Value of Finite Range Tunneling?* (2015), (arXiv:quant-ph/1512.02206).
- ⁴² P. Bunyk, E. Hoskinson, M. W. Johnson, E. Tolkacheva, F. Altomare, A. J. Berkley, R. Harris, J. P. Hilton, T. Lanting, and J. Whittaker, *Architectural Considerations in the Design of a Superconducting Quantum Annealing Processor*, IEEE Trans. Appl. Supercond. **24**, 1 (2014).
- ⁴³ S. Kirkpatrick, C. D. Gelatt, Jr., and M. P. Vecchi, *Optimization by simulated annealing*, Science **220**, 671 (1983).
- ⁴⁴ S. Mandrà, Z. Zhu, W. Wang, A. Perdomo-Ortiz, and H. G. Katzgraber, *Strengths and Weaknesses of Weak-Strong Cluster Problems: A Detailed Overview of State-of-the-art Classical Heuristics vs Quantum Approaches* (2016), (arXiv:1604.01746).
- ⁴⁵ N. Lemke and I. A. Campbell, *Two-Dimensional Ising Spin Glasses with Nonzero Ordering Temperatures*, Phys. Rev. Lett. **76**, 4616 (1996).
- ⁴⁶ G. Parisi, J. J. Ruiz-Lorenzo, and D. A. Stariolo, *Crossovers in the two-dimensional Ising spin glass with ferromagnetic next-nearest-neighbour interactions*, J. Phys. A **31**, 4657 (1998).
- ⁴⁷ N. Lemke and I. A. Campbell, *Finite-temperature phase transition in the two-dimensional randomly coupled ferromagnet*, J. Phys. A **32**, 7851 (1999).
- ⁴⁸ A. K. Hartmann and I. A. Campbell, *Ordered phase in the two-dimensional randomly coupled ferromagnet*, Phys. Rev. B **63**, 094423 (2001).
- ⁴⁹ A. J. Bray and M. A. Moore, *Lower critical dimension of Ising spin glasses: a numerical study*, J. Phys. C **17**, L463 (1984).
- ⁵⁰ A. K. Hartmann and H. Rieger, *Optimization Algorithms in Physics* (Wiley-VCH, Berlin, 2001).
- ⁵¹ S. Boettcher, *Stiffness of the Edwards-Anderson Model in all Dimensions*, Phys. Rev. Lett. **95**, 197205 (2005).
- ⁵² K. Hukushima and K. Nemoto, *Exchange Monte Carlo method and application to spin glass simulations*, J. Phys. Soc. Jpn. **65**, 1604 (1996).
- ⁵³ Z. Zhu, A. J. Ochoa, and H. G. Katzgraber, *Efficient Cluster Algorithm for Spin Glasses in Any Space Dimension*, Phys. Rev. Lett. **115**, 077201 (2015).
- ⁵⁴ J. Houdayer, *A cluster Monte Carlo algorithm for 2-dimensional spin glasses*, Eur. Phys. J. B. **22**, 479 (2001).
- ⁵⁵ W. H. Press, S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery, *Numerical Recipes in C* (Cambridge University Press, Cambridge, England, 1995).
- ⁵⁶ K. Binder, *Critical properties from Monte Carlo coarse graining and renormalization*, Phys. Rev. Lett. **47**, 693 (1981).