

This is the accepted manuscript made available via CHORUS. The article has been published as:

# Application of an ab initio S matrix to data analysis of transfer reactions to the continuum populating $^{11}\text{Be}$

A. Bonaccorso, F. Cappuzzello, D. Carbone, M. Cavallaro, G. Hupin, P. Navrátil, and S. Quaglioni

Phys. Rev. C **100**, 024617 — Published 19 August 2019

DOI: [10.1103/PhysRevC.100.024617](https://doi.org/10.1103/PhysRevC.100.024617)

# Application of an *ab-initio* S-matrix to data analysis of transfer to the continuum reactions populating $^{11}\text{Be}$

A. Bonaccorso<sup>1</sup>, F. Cappuzzello<sup>2,3</sup>, D. Carbone<sup>3</sup>, M. Cavallaro<sup>3</sup>, G. Hupin<sup>4</sup>, P. Navrátil<sup>5</sup> and S. Quaglioni<sup>6</sup>.

<sup>1</sup>INFN, Sez. di Pisa, Largo B. Pontecorvo 3, 56127 Pisa, Italy.

<sup>2</sup>Dipartimento di Fisica ed Astronomia "Ettore Majorana", Università degli Studi di Catania

<sup>3</sup>INFN, Lab. Naz. del Sud, Via S. Sofia, Catania, Italy.

<sup>4</sup>Institut de Physique Nucléaire, IN2P3, CNRS, Université Paris-Sud,  
Université Paris-Saclay, 91406 Orsay Cedex, France

<sup>5</sup>TRIUMF, 4004 Wesbrook Mall, Vancouver, British Columbia V6T 2A3, Canada.

<sup>6</sup>Lawrence Livermore National Laboratory, P.O. Box 808, L-414, Livermore, California 94551, USA.

Recently, the bound and continuum spectrum of  $^{11}\text{Be}$  has been calculated within the *ab-initio* no-core shell model with continuum (NCSMC) method successfully reproducing the parity inversion in the ground state. The continuum spectrum obtained is in agreement with known experimental levels. The S-matrix contained in the NCSMC continuum wave functions of the  $n+^{10}\text{Be}$  system is used in this work for the first time in a Transfer-to-the-Continuum (TC) reaction calculation. The TC approach is applied to study the excitation energy spectrum of  $^{11}\text{Be}$  measured in the  $^9\text{Be}(^{18}\text{O}, ^{16}\text{O})^{11}\text{Be}$  reaction at 84 MeV. Previously known levels are confirmed and theoretical and experimental evidence for a  $9/2^+$  state at  $E_x=5.8$  MeV is given, whose configuration is thought to be  $^{10}\text{Be}(2^+)+n(d_{5/2})$ .

## I. INTRODUCTION

This work presents the first application of an *ab-initio* structure model for continuum states [1] to the calculation of a transfer reaction to resonant states [2–6]. Thanks to the S-matrix obtained within the no-core shell model with continuum (NCSMC) method [1], the traditional reduction of the complicated many-body problem to a one-body problem via a phenomenological optical potential to treat the continuum neutron-target final state will not be necessary, as previously done for  $^{10}\text{Be}$  [6; 7]. We will show how various states of an exotic nucleus such as  $^{11}\text{Be}$  are built up adding one or two neutrons to a  $^9\text{Be}$  target.

The case of  $^9\text{Be}$  target is particularly interesting because one can access the  $^{10}\text{Be}$  and  $^{11}\text{Be}$  residual nuclei, which have been the subject of a large number of studies by the nuclear physics community in the past years. In particular  $^{11}\text{Be}$  is a weakly bound nucleus with a very rich and somehow surprising phenomenology [8; 9]. Some of the interesting characteristics are for example the role of correlations between core excited states and neutron orbitals [1; 10–12], as well as the neutron-neutron pairing, and even the molecular clustering [13]. They are all relevant for the discussions of the present work. Thus by comparing one- and two-neutron transfer results, as we have consistently done for some time now, one can understand how correlations arise in an exotic nucleus.

In our investigation we use one- and two-neutron transfer reactions induced by the  $^{18}\text{O}$  projectile. In this way states in neutron-rich nuclei are populated starting from stable targets [6], [14–21]. The observed selectivity of  $^{18}\text{O}$ -induced transfer reactions allows for a consistent exploration of both single particle features via the  $(^{18}\text{O}, ^{17}\text{O})$  reaction and two-neutron correlations via the  $(^{18}\text{O}, ^{16}\text{O})$ . In particular we argue that the  $(^{18}\text{O}, ^{16}\text{O})$  re-

action proceeds mainly via two different mechanisms: i) a two-step single neutron transfer, where the two neutrons are independently transferred to accessible orbitals in the target field, ii) a one-step transfer of a correlated pair of neutrons, populating mainly two-neutron configurations in the states of the residual nucleus. In the following we will use a theoretical model corresponding to mechanism i).

Here we would like to stress the fact that in the part of the  $^{11}\text{Be}$  spectrum that we aim at describing in this work, one neutron is transferred to a bound state while the other is transferred to the continuum thus the independence of the two mechanisms is better justified. At the moment there is no reaction model using a two-neutron correlated final state state with one neutron bound and the other unbound.

The state-of-the-art on this subject is different for (t,p) reactions populating bound states in the target. It can be resumed with the findings of Ref.[22] and [23] which agree with the whole previous literature. In the first paper it is stated that "the simultaneous and nonorthogonal contributions [of the two neutrons to transfer] are in antiphase, so that the contribution corresponding to the coherent superposition of these two amplitudes tend to cancel. The calculated total cross section thus essentially coincides with the successive process". On the other hand Ref.[23] shows that the two-step, successive, transfer also occurs in a correlated fashion. If we were to use the model of [23] for example, the different possible quantum mechanical paths associated with the population of different intermediate states of  $^{10}\text{Be}+^{17}\text{O}$  in our case, would be summed coherently. Also, the different terms (successive transfer, simultaneous transfer, and non-orthogonality contribution) would represent different amplitude

contributions to the second-order process, to be added coherently to get the second order cross section. Their relative importance in a coupled channel model such as that discussed in [23] depends on the arbitrary choice of the reaction representation (prior-prior, post-post, prior-post or post-prior). However in our case because we treat transfer between heavy ions we can use a semiclassical model in which the post and prior representations coincide. Furthermore the TC provides analytical probabilities for energy spectra in the continuum in heavy-ion reactions. Results for (p,t) or (t,p) to bound states can give only partial guidance to this work, due to the presence of a heavy core in the projectile and target and of a neutron final continuum state in our case.

Also or the ( $^{18}\text{O}$ ,  $^{16}\text{O}$ ) two-neutron transfer reactions the  $2^+$  excited state at 1.98 MeV in  $^{18}\text{O}$  plays a key role in the coupling scheme that would require a coupled channel theoretical treatment which at the moment is still missing [15]. On the other hand the successful results of a semiclassical, two-step treatment of the  $^{208}\text{Pb}(^{16}\text{O}, ^{18}\text{O})^{206}\text{Pb}$  reaction [24], have been used for long time as a justification [25] for neglecting interference effects and simultaneous pair transfer. In our case this is particularly true as we do not study angular distribution but energy spectra.

Using the one-neutron transfer reaction we have been able in the past to obtain both bound and continuum states of  $^{10}\text{Be}$  [6]. A very accurate description of the latter was achieved thanks to a previous work in which two different  $n+^9\text{Be}$  optical potentials, fitted to scattering data on a very large energy range [7], were compared. These potentials allowed some of us to calculate an S-matrix which is necessary for the correct definition of a continuum nucleon-nucleus state. This in turn is one of the ingredients of the Transfer to the Continuum (TC) model [2–4] which allows to reproduce the excitation energy spectrum of the target [5; 6]. The  $^9\text{Be}(^{18}\text{O}, ^{17}\text{O})^{10}\text{Be}$  experiment and relative theoretical description [6] constituted the first step in the study of of the  $^9\text{Be}(^{18}\text{O}, ^{16}\text{O})^{11}\text{Be}$  reaction that we discuss in this paper. Two-neutron transfer experiments on  $^9\text{Be}$  have been performed previously at higher incident energies and with different projectiles such as tritium [26]–[30],  $^6\text{He}$  [31],  $^{13}\text{C}$  [32; 33] and  $^{16}\text{O}$  [34]–[36]. From the experimental point of view the novelty in our case resides in the fact that for a heavy projectile the core spectator situation is realized and thanks to the large relative angular momentum of projectile and target, coupled with the initial state angular momentum, high spin states in the final nucleus can be reached [37]. From the theoretical point of view we are achieving here a unification of structure and reaction formalisms via the use of an *ab-initio* S-matrix.

## II. THE EXPERIMENT

The  $^{18}\text{O}^{6+}$  beam at 84 MeV incident energy was accelerated by the Tandem Van de Graaff facility of INFN-LNS. A self-supporting  $^9\text{Be}$   $200 \pm 10 \mu\text{g}/\text{cm}^2$  thick target was used. Supplementary runs with a  $50 \pm 3 \mu\text{g}/\text{cm}^2$  self-supporting  $^{12}\text{C}$  target and a  $260 \pm 10 \mu\text{g}/\text{cm}^2$   $\text{WO}_3$  target were recorded in order to estimate the background in the energy spectra from  $^{12}\text{C}$  and  $^{16}\text{O}$  impurities in the  $^9\text{Be}$  target. The MAGNEX magnetic spectrometer [38] was used to momentum analyse the  $^{16}\text{O}$  ejectiles, detected by its focal plane detector [39]. The angular range between  $3^\circ$  and  $10^\circ$  in the laboratory reference frame was explored. Details about the particle identification and the trajectory reconstruction techniques used for the reduction of the MAGNEX data can be found in Refs. [40–43]. An overall energy and angular resolution of about 500 keV (FWHM) and  $0.3^\circ$  was obtained. The absolute cross section was also estimated according to Ref. [41] with a total error of about 10% induced by the uncertainties in the target thickness and beam current integration. An example of an energy spectrum for the ( $^{18}\text{O}, ^{16}\text{O}$ ) reaction with  $^9\text{Be}$  target, in a limited angular range, is shown in Fig. 1 where the background contributions are also shown. The angle integrated absolute cross section spectrum obtained after background subtraction is shown in Fig. 2.

## III. MODEL INDEPENDENT ANALYSIS OF THE ENERGY SPECTRUM

A number of peaks signals the population of bound and resonant states of  $^{11}\text{Be}$  in this transfer reaction.

In our data the  $1/2^+$  ground and the  $1/2^-$  first excited state at 0.320 MeV are not resolved. Above the one-neutron separation energy  $S_n = 0.502$  MeV, a strong excitation of the  $5/2^+$  state at 1.783 MeV is observed, while the states at 2.654 MeV and 3.400 MeV are less populated. Interestingly these are the only states observed in one-neutron transfer reactions in the past [44], [45], [46], showing a similar pattern to that observed here. This could indicate that, for these states, the  $^{10}\text{Be}_{gs}$  is preferentially populated as an intermediate system in a first step of the reaction and a second neutron is then transferred to accessible single particle orbitals.

Beyond the  $^{10}\text{Be}$  core excitation threshold we observe a peak at 3.9 MeV, where we cannot separate the transitions to the known states at 3.889 MeV ( $5/2^-$ ) and 3.955 MeV ( $3/2^-$ ). Only in ref. [26] the two states are separated in a two-neutron transfer reaction, showing that they are both strongly excited by such a probe. Intense transitions are observed for the  $5/2^-$  state at 5.255 MeV and the state at 6.705 MeV, resembling the situation observed in all the reported two-neutron transfer reactions data [32], [33], [34]. These states are not populated in single neutron transfer and in  $^{11}\text{B}(^7\text{Li}, ^7\text{Be})^{11}\text{Be}$  charge exchange reaction [47], [50] suggesting a selective response

of  $^{11}\text{Be}$  to two-neutron transfer operators, likely populating  $^9\text{Be} \otimes (\text{sd})^2$  configurations.  $\beta$ -decay studies from  $3/2^-$   $^{11}\text{Li}$  parent nucleus show the transitions to the  $3/2^-$  state at 3.955 MeV and  $5/2^-$  at 5.255 MeV of  $^{11}\text{Be}$ . This does not contradict the above argument about the  $^9\text{Be} \otimes (\text{sd})^2$  configurations for these states, since  $\beta$ -decay can also occur within the  $^9\text{Li}$  core of  $^{11}\text{Li}$  parent nucleus. In addition,  $\beta$ -decay does not populate the state at 6.705 MeV, which would indicate either a different parity or a spin higher than that achievable by allowed Fermi and Gamow-Teller operators. In ref. [32], [33], [34] this state is described as belonging to a rotational band built on the  $3/2^-$  state at 3.955 MeV, together with the 5.255 MeV state, with  $7/2^-$  spin assignment.

Also debated are the states at 5.8 MeV and 8.813 MeV, which are both present in all reported two-neutron transfer experiments, despite the centroid energy and width are slightly different in the various studies. The former is not observed in  $\beta$ -decay, nor in single neutron transfer, but it is likely the same observed in  $^{11}\text{B}(^7\text{Li}, ^7\text{Be})^{11}\text{Be}$  charge exchange reaction at 6.05 MeV (FWHM 300keV) [47]. Due to the large momentum transfer, such reaction can easily excite states with high spin, not populated in  $\beta$ -decay. In ref. [34] the neutron decay of this state to the first excited  $2^+$  state of  $^{10}\text{Be}$  suggests a core-excited configuration. Recently a  $9/2^+$  spin-parity for a state with a  $[^{10}\text{Be}(2_1^+) \otimes (1d)^{5/2}]^{9/2^+}$  stretched configuration on the  $^{10}\text{Be}(2_1^+)$  excited state was predicted in the *ab-initio* NC-SMC approach [1], which is then in perfect agreement with the present finding and our interpretation of previous literature. For the state at 8.813 MeV,  $\beta$ -decay studies [48; 49] assign  $3/2^-$  spin parity, conflicting with the assumption of a high spin member of the rotational band built on the  $3/2^-$  state at 3.955 MeV proposed in ref. [34], [32], [33]. Our data, despite confirming the population of this state in two-neutron transfer reactions, do not add much to the previous debate.

#### IV. THEORETICAL APPROACH

From the theoretical point of view, the description of the  $n + ^{10}\text{Be}$  part of the spectrum above  $S_n$  is very challenging because no experimental data exist on the free  $n + ^{10}\text{Be}$  scattering.

However, the recent *ab-initio* calculation of the  $^{11}\text{Be}$  spectrum [1] over a wide energy range, allows for a description of both bound and continuum states, the continuum part providing a  $S$ -matrix. Thus we can apply the TC method overcoming the knowledge of the optical potential thanks to the NCSMC  $S$ -matrix. One particularly interesting aspect is that the  $n + ^{10}\text{Be}$  component of the spectrum has contributions from bound states of  $^{10}\text{Be}$ .

TABLE I: Structure parameters of the valence neutron, see text. The present work  $^{10}\text{Be}$  spectroscopic factors were obtained with the chiral  $\text{N}^2\text{LO}_{\text{SAT}}$   $\text{NN}+3\text{N}$  interaction [51], the same used in the  $^{11}\text{Be}$  calculations. The  $C^2S$  and  $C_i$  values for  $^{17}\text{O}$  have been used in the calculations of Eq.(1) while the others are given here for completeness.

State	$S_n$ (MeV)	$C_i$ ( $\text{fm}^{-\frac{1}{2}}$ )	$j$	$l$	$C^2S$
$^{18}\text{O}_{g.s.}$	8.04	1.73	5/2	2	1.7[15]
$^{17}\text{O}_{g.s.}$	4.145	0.69	5/2	2	0.945[15]
$^{10}\text{Be}_{g.s.}$	6.812	2.3	3/2	1	2.13[59], 2.6[60], 2.73 <sup>a</sup>
$^{10}\text{Be}(2_1^+)$	3.44		1/2	1	0.0226[59], 0.274[60] <sup>b</sup> , 0.036 <sup>a</sup>
$^{10}\text{Be}(2_1^+)$			3/2	1	0.268[59], 0.274[60] <sup>b</sup> , 0.24 <sup>a</sup>
$^{10}\text{Be}(2_2^+)$	0.854		1/2	1	0.616[59], 0.421[60] <sup>b</sup> , 0.406 <sup>a</sup>
$^{10}\text{Be}(2_2^+)$			3/2	1	0.13[59], 0.421[60] <sup>b</sup> , 0.076 <sup>a</sup>

<sup>a</sup>Present work,

<sup>b</sup>This reference does not distinguish the two  $j$ -components.

#### A. Reaction model:

We imagine the reaction going through two independent steps as anticipated in the Introduction and indicated as mechanism i). In the first step the reaction  $^9\text{Be}(^{18}\text{O}, ^{17}\text{O})^{10}\text{Be}$  takes place and neutron  $n1$  is transferred from  $^{18}\text{O}$  to populate the bound states of  $^{10}\text{Be}$ .

We then imagine that the second neutron  $n2$  is transferred from  $^{17}\text{O}$  to  $^{10}\text{Be}$  and since we just want to describe the continuum part of the spectrum (above  $E_x = S_n$ ) we use for the two-neutron transfer description the following formula:

$$\frac{d\sigma_{2n}}{d\varepsilon_{f2}} = C^2S P_{n1}^{phen}(R_s) \int_{b_{min}}^{b_{max}} db_c |S_{ct}(b_c)|^2 \frac{dP_{n2}(b_c)}{d\varepsilon_{f2}}, \quad (1)$$

$P_{n1}^{phen}$ , given by Eq.(3), is the bound state transfer probability for neutron  $n1$  from  $^{18}\text{O}$  to  $^9\text{Be}$ , at the strong absorption radius between core and target, defined as  $R_s = 1.4(A_c^{1/3} + A_t^{1/3})$ . Then in the next step,  $n2$  is the second neutron which is transferred to a continuum state from  $^{17}\text{O}$  and  $C^2S$  is its initial wave function spectroscopic factor.  $\varepsilon_{f2}$  is the continuum final state energy of  $n2$  and  $|S_{ct}(b_c)|^2$  has been calculated according to [52]. It is the elastic scattering probability between  $^{16}\text{O}$  and  $^{10}\text{Be}$ , the second-step core and target nuclei respectively. Following Ref.[6], the correspondence between the measured scattering angle  $3^\circ < \theta_{lab} < 10^\circ$  and the impact parameter has been obtained via a classical trajectory calculation according to Ref.[53] providing  $7 < b_c < 8$  fm to be used in Eq.(1).  $P_{n1}^{phen}$  is extracted as described below, from the integrated experimental one-neutron transfer cross section in the  $^9\text{Be}(^{18}\text{O}, ^{17}\text{O})^{10}\text{Be}$  reaction whose data were presented in [6]. Above the threshold for the

first excited state of  $^{10}\text{Be}$ ,  $P_{n1}^{phen}$  is obtained consistently using the parameters appropriate to take into account the excitation energy in  $^{10}\text{Be}$  given in Table I.

To take advantage of the fact that the experimental cross section for one-neutron transfer is already known and to minimize the dependence from parameter's uncertainty, we use the one-neutron transfer cross section between bound states formula from [54]:

$$\sigma_{1n} = \pi \frac{(R_s - a_c)}{\eta} P_{n1}(R_s) \quad (2)$$

which is obtained when the core-target S-matrix is calculated in the sharp cut off approximation:

$$|S_{ct}(b_c)|^2 = 1 \quad \text{if} \quad b_c > R_s$$

$$|S_{ct}(b_c)|^2 = 0 \quad \text{if} \quad b_c < R_s.$$

$\eta$  is a kinematical parameter depending on the initial and final neutron separation energies and the energy of relative motion [54],

$$\eta = \sqrt{\gamma_i^2 + k_1^2}$$

where  $\gamma_i^2 = 2mS_{i(1,2)}/\hbar^2$  and  $S_{i(1,2)}$  is the bound state initial separation energy of the first and second neutron respectively.  $k_1^2 = (Q + 1/2mv^2)/(\hbar v)^2$  where  $Q = \varepsilon_{i(1,2)} - \varepsilon_{f(1,2)}$  is the Q-value. The  $\varepsilon_{i(1,2)}$  are the negative initial binding energies of neutron n1 and n2 in their bound states, while  $\varepsilon_{f(1,2)}$  is the negative final energy of the first step neutron in  $^{10}\text{Be}$  and is the positive continuum energy of neutron n2 in  $^{11}\text{Be}$ . The  $\eta$ -value ranges from  $0.68\text{fm}^{-1}$  to  $0.88\text{fm}^{-1}$  for the three states bound  $^{10}\text{Be}$  states considered here.

$R_s$  is again the strong absorption radius and  $a_c$  is the Coulomb length parameter. Then we derive from Eq.(2) the following phenomenological transfer probability at the strong absorption radius to be used to describe the two-neutron transfer in Eq.(1):

$$P_{n1}^{phen}(R_s) = \frac{\sigma_{1n}^{exp}}{\pi \frac{(R_s - a_c)}{\eta}} \quad (3)$$

where  $\sigma_{1n}^{exp} = 2.1 \text{ mb}$  ( $^{10}\text{Be}_{g.s.}$ ),  $2.2 \text{ mb}$  ( $^{10}\text{Be}_{2_1^+}$ ),  $3.2 \text{ mb}$  ( $^{10}\text{Be}_{2_2^+}$ ) are the experimental values from Ref.[6]. By adopting this formula we assume that all structure information, such as spectroscopic factors, of both projectile ( $^{18}\text{O}$ ) and target ( $^9\text{Be}$ ) are included in the measured  $\sigma_{1n}^{exp}$ .

Our approach is equivalent to a treatment of a two step process via second order perturbation theory [53; 55] which is justified by the small first step average probability ( $P_{n1}^{phen}=0.01$ , cf. Fig.2). Furthermore the case of  $^{10}\text{Be}$ , as an intermediate system in the transition from  $^9\text{Be}$  to  $^{11}\text{Be}$ , is quite peculiar and makes our assumption of incoherent summation still approximately valid. The key point is that  $^{10}\text{Be}$  level density is quite low in the energy region of interest and in fact we only need to consider three states, i.e.  $^{10}\text{Be}_{g.s.}$  and the excited states at 3.4 MeV ( $^{10}\text{Be}_{2_1^+}$ ) and 6 MeV ( $^{10}\text{Be}_{2_2^+}$ ). As a consequence, the  $^{10}\text{Be}_{2_1^+}$  has an important role in  $^{11}\text{Be}$  spectra from the inelastic thresholds of ( $^{10}\text{Be}_{2_1^+}$ ) + n, located at about 3.9 MeV, while the  $^{10}\text{Be}_{2_2^+}$  from the inelastic thresholds of ( $^{10}\text{Be}_{2_2^+}$ ) + n, located at about 6.5 MeV. Thus for  $^{11}\text{Be}$  states up to 3.9 MeV the role of  $^{10}\text{Be}$  first and second excited states is expected to be weak. For states of  $^{11}\text{Be}$  between 3.9 and 6.5 MeV one can have the contribution from  $^{10}\text{Be}_{g.s.}$  and the first  $^{10}\text{Be}_{2_1^+}$ . However, the states of  $^{11}\text{Be}$  in that energy range with pronounced  $^{10}\text{Be}_{g.s.}$  + n configuration are quite broad since they are far from ( $^{10}\text{Be}_{g.s.}$ ) + n threshold, located at  $S_n = 0.5 \text{ MeV}$ . As an example a simulated pure single particle  $d_{3/2}$  state at 4 MeV excitation energy would have a width as large as 3 MeV and more. Therefore the role of such configurations, so much spread out in energy, is minor in the narrow region covered by the analyzed sharp resonances. In fact, the  $9/2^+$  state we claim at 5.9 MeV is built with the  $^{10}\text{Be}$  first excited state and not with the  $^{10}\text{Be}_{g.s.}$ , as the ab initio calculation of [1] correctly indicates. The same argument holds for  $^{11}\text{Be}$  states above 6.5 MeV where configurations with  $^{10}\text{Be}_{g.s.}$  and the first  $^{10}\text{Be}_{2_1^+}$  likely are not contributing significantly.

The second neutron n2 transfer probability to the continuum states of  $^{11}\text{Be}$  is given by

$$\frac{dP_{n2}(b_c)}{d\varepsilon_{f2}} = \frac{|C_i|^2}{2k_{f2}} \left[ \frac{\hbar}{mv} \right] \frac{e^{-2\eta b_c}}{2\eta b_c} \Sigma_{jf,\nu} (|1 - \bar{S}_{jf,\nu}|^2 + 1 - |\bar{S}_{jf,\nu}|^2) (2j_f + 1) (1 + R_{if}) M_{l_f l_i}. \quad (4)$$

Details of the TC method have been given in several previous publications, see for example Ref.[3] where the definitions of the parameters appearing in Eq.(4) can also be found.

In Eq.(4) for each continuum energy the sum is over

all possible n+ $^{10}\text{Be}$  total angular momenta. Above the thresholds for the first and second  $2^+$  excited states in  $^{10}\text{Be}$ , for each angular momentum there is also an incoherent sum over all channels  $\nu$  contributing to it. This is the same as when calculating total reaction cross sec-

tions and the observable is the final nucleus excitation energy spectrum. The situation is different (c.f. Eq.(6) of Ref.[55]) when two neutron transfer is discussed with the aim of calculating core angular distributions. In the latter case for each angle different channels can interfere and contribute coherently.

We have usually calculated  $\bar{S}_{j_f}$  from an optical potential [3]. The same procedure cannot be applied to the system  $n+^{10}\text{Be}$  because there are no data available. However the S-matrix calculated by some of us in Ref.[1] is perfectly suited to be used in Eq.(4) as it is given in terms of the  $n-^{10}\text{Be}$  continuum energy and angular momentum  $j_f$  appearing in Eq.(4) and it contains at each energy, all possible inelastic channel contributions. Notice that Eq.(4) contains two terms proportional to  $|1 - \bar{S}_{j_f}|^2$  and  $1 - |\bar{S}_{j_f}|^2$  giving the elastic and inelastic neutron breakup from the initial state in the projectile, respectively.

## B. Ab-initio S-matrix

Some of us investigated the structure of  $^{11}\text{Be}$  by studying the  $^{10}\text{Be}+n$  system within the NCSMC approach [1]. This approach [58] uses a basis expansion with two key components: one describing all nucleons close together, forming the  $^{11}\text{Be}$  nucleus, and a second one describing the separated neutron and  $^{10}\text{Be}$  clusters. The former part utilizes a square-integrable harmonic-oscillator basis expansion treating all 11 nucleons on the same footing. The latter part factorizes the wave function into products of  $^{10}\text{Be}$  and neutron components and their relative motion with proper bound-state or scattering boundary conditions. **We compute the S-matrix from the 11-body NCSMC calculations by employing the calculable R-matrix method [56; 57] and matching the many-body  $^{11}\text{Be}$  internal wave function with the asymptotic binary  $n-^{10}\text{Be}$  channels at about 18 fm, well beyond the range of the nuclear interaction.** The chiral  $\text{N}^2\text{LO}_{\text{SAT}}$  two-nucleon (NN) and three-nucleon (3N) interaction [51] served as input. For the  $^{10}\text{Be}$  cluster, in addition to the ground state, we also included the first and the second excited  $2^+$  states. The outcome of the NCSMC calculations are the energies and wave functions of the bound states, here  $1/2^+$  and  $1/2^-$  in the correct order compared to experiment, as well as of the continuum states. The latter include the S-matrix that we in turn apply in the present investigation of the  $^9\text{Be}(^{18}\text{O}, ^{16}\text{O})^{11}\text{Be}$  two-neutron transfer reaction. As seen in Figs. 2 and 3 and Table I of Ref. [1], the  $\text{N}^2\text{LO}_{\text{SAT}}$  interaction provides a quite reasonable description of the low-lying bound states and resonances of  $^{11}\text{Be}$ . However, to achieve a spectroscopic accuracy needed in the present study, we turn to the NCSMC-pheno approach that allows to reproduce the experimental thresholds and bound- and resonant state energies as presented in the right part of Table I in Ref. [1]. The NCSMC calculations with the  $\text{N}^2\text{LO}_{\text{SAT}}$  interaction predict a low-lying  $9/2^+$  resonance at 5.42 MeV and, with

the NCSMC-pheno approach, at 5.59 MeV after adjusting the  $^{10}\text{Be}$  thresholds to experiment. In this work we examine an experimental candidate for this resonance at 5.3 MeV. Notice that these energies are given with respect to the  $n+^{10}\text{Be}$  threshold. Consequently, we have performed a new NCSMC-pheno calculation to fit the calculated  $9/2^+$  resonance position to that energy. Resulting S-matrix is then used in the present study.




FIG. 1: (Color online) Inclusive excitation energy spectrum of the  $^9\text{Be}(^{18}\text{O}, ^{16}\text{O})^{11}\text{Be}$  reaction at 84 MeV incident energy and  $6^\circ < \theta_{\text{lab}} < 7^\circ$ . The background coming from C and O contaminations in the target is shown as red and blue hatched areas, respectively.

## V. DISCUSSION OF THE RESULTS

In Fig.2 we show the experimental inclusive energy spectrum of  $^{11}\text{Be}$  together with the theoretical calculations (red dashed curve). This calculation does not contain any fitting parameter. **On the other hand the green solid curve is the same calculation folded with the experimental resolution renormalized by a factor 3.2 to fit the data.** The first three continuum states are well reproduced as far as the position is concerned, indicating that the TC model contains the correct dynamics and the *ab-initio* S-matrix accurate structure information. Notice that also the order of magnitude is reproduced reasonably well within the uncertainty due to the description of the two-neutron transfer reaction mechanism. The calculated relative population of the three resonances at  $E_x = 1.783, 2.654, 3.400$  MeV with  $J^\pi = 5/2^-, 3/2^-, 3/2^+$ , respectively, compares well with the experimental results from  $^{10}\text{Be}(d,p)^{11}\text{Be}$  [44]. The state at 2.654 MeV is depleted in our model calculation because of the unfavourable one-neutron transfer reac-




FIG. 2: (Color online)  $^{11}\text{Be}$  inclusive excitation-energy spectrum from the  $^9\text{Be}(^{18}\text{O}, ^{16}\text{O})^{11}\text{Be}$  reaction at 84 MeV incident energy, integrated in the total measured angular range  $3^\circ < \theta_{lab} < 10^\circ$ . Black histogram: experimental data after background subtraction. Red dashed curve: total 2n transfer calculations resulting from the use of the S-matrix from Ref. [1]. Green full curve: the red curve calculation folded with the experimental resolution and renormalized by a factor 3.2. Blue dotted curve: theoretical free  $n+^{10}\text{Be}$  cross section in barn, see text for details.

tion matching, as it is in the data from [44]. However it is well seen in the present data as well as in all previous two-neutron transfer experiments [34] confirming for this state a relevant  $2n+^9\text{Be}$  configuration and a population via the simultaneous 2n-transfer reaction mechanism, type ii) discussed in the Introduction. On the other hand the calculated peak at  $E_x = 3.400$  MeV, has such a small width ( $\approx 0.02\text{MeV}$ ) from [1] that its presence has little physical significance and in fact it is hardly visible as a structure in the experiment. The transfer calculation probably misses the maximum value for the same reason. Notice that in the figures we indicate  $(3/2^+, 3/2^-)$  for this state because from the literature it is often quoted as  $3/2^-$  while from the present S-matrix calculations it appears to be  $3/2^+$ . As far as the absolute total cross sections are concerned we believe that due to the experimental and theoretical uncertainties the only comparable values are those for the 1.783 MeV  $d_{5/2}$  state for which we get  $\sigma_{exp} = 359 \pm 35(\mu\text{b})$  and  $\sigma_{th} = 175(\mu\text{b})$ . However one might argue that due to the indistinguishability of the two neutrons Eq.(1) should contain a factor 2 which would give an almost perfect agreement between theory and experiment. On the other hand as we have discussed before our transfer mechanism model is over-simplified and thus we conclude that the present absolute theoretical cross

section are reliable only at the order of magnitude level.

The cross-section of the resonance at 5.8 MeV is reasonably well reproduced by our model, thus strengthening the interpretation of a  $9/2^+$  stretched configuration, already discussed above and in Ref. [1]. It is the only state that can have a definite single particle nature in the high energy part of the spectrum, because it has an high spin which does not mix with underlying components of low angular momenta. It can be seen only in a reaction with heavy ions where the matching conditions allow to reach high resonance energy and high angular momentum. This state can be reproduced by the TC which contains explicitly the spin couplings between initial and final states besides the angular momentum couplings. Therefore the present work provides the first evidence for the existence of such a state. We believe our attribution is quite firm and well justified and unique so far in the literature. From the point of view of the structure part, we can see the  $9/2^+$  because it is built on the  $2_1^+$  state of  $^{10}\text{Be}$  that we include in our NCSMC calculation and thus it comes about as a  $9/2^+$  from the S-matrix calculation.

Finally, for comparison, the free  $n+^{10}\text{Be}$  cross section (elastic plus inelastic) calculated with the *ab-initio* S-matrix and rescaled by a factor 0.06, is shown by the blue dots in Fig. 2. For this cross section the y-axis scale should be read in units of barn as indicated in the legend. Obviously this is another observable with respect to the transfer cross section shown in the figure. Notice that the magnitude of this cross section is consistent with that of the  $n+^9\text{Be}$  data and calculation of Ref.[7]. This is a proof of the accuracy of the *ab-initio* model in providing the magnitude of the neutron-core couplings (interaction). The three resonance states coupled to the  $^{10}\text{Be}_{g.s.}$  appear indeed at the same positions as in the data and they scale as the simple  $1/k^2$  law. Above the first and second  $2^+$  excited state threshold respectively, the shown free cross section is the sum of the tails of the cross sections from lower neutron energies, in accordance with the hypothesis that  $^{10}\text{Be}$  excited states are populated by the first neutron.

## VI. CONCLUSIONS

In conclusion we have presented new experimental data on  $^9\text{Be}(^{18}\text{O}, ^{16}\text{O})^{11}\text{Be}$  two-neutron transfer reaction at 84 MeV, providing absolute cross section energy spectra. An *ab-initio* S-matrix coupled for the first time with the TC method is able to reproduce the position and widths of the  $n + ^{10}\text{Be}$  components in the experimental excitation energy spectrum of the exotic nucleus  $^{11}\text{Be}$  and to predict the order of magnitude of the absolute cross sections. We have given evidence, both theoretical as well as experimental, of the presence of a  $9/2^+$  state at  $E_x=5.8$  MeV with  $[^{10}\text{Be}(2_1^+) \otimes (1d)^{5/2}]^{9/2+}$  configuration.

It is very encouraging to see that very elaborated *ab-initio* structure calculations for continuum states can provide the correct ingredients to be easily incorporated in a reaction model which is simple and whose accuracy is within the present state-of-the-art of the literature. We have also shown that exotic nuclei can be successfully studied at stable beam facilities providing interesting complementary information to those obtained with RIBs (Radioactive Ion Beams).

### Acknowledgments

Two of us, D.C and M.C., have received funding from the European Research Council (ERC) under the European Unions Horizon 2020 research and innovation programme (grant agreement No 714625).

P.N. work was supported in part by the NSERC Grant No. SAPIN-2016-00033. TRIUMF receives federal funding via a contribution agreement with the National Research Council of Canada. Computing support for G.H., P.N. and S.Q. came from an INCITE Award on the Titan supercomputer of the Oak Ridge Leadership Computing Facility (OLCF) at ORNL, from Compute Canada, and from the LLNL institutional Computing Grand Challenge Program.

This article was prepared in part by LLNL (S.Q.) under Contract DE-AC52-07NA27344. This material is based in part upon work supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under Work Proposals No. SCW0498.

A.B. is grateful to J. Dohet-Eraly for enlightening discussions on the structure of the *ab-initio* S-matrix at the early stages of this work.

- 
- [1] A. Calci, P. Navrátil, R. Roth, J. Dohet-Eraly, S. Quaglioni, and G. Hupin, Phys. Rev. Lett. **117** (2016) 242501.
  - [2] A. Bonaccorso and D. M. Brink, Phys. Rev. C **38**, 1776 (1988).
  - [3] A. Bonaccorso and D. M. Brink, Phys. Rev. C **43** 299 (1991).
  - [4] A. Bonaccorso and D. M. Brink, Phys. Rev. C **44** (1991) 1559.
  - [5] G. Blanchon, A. Bonaccorso and N. Vinh Mau, Nucl. Phys. A **739** 259 (2004).
  - [6] D. Carbone et al., Phys. Rev. C **90** 064621 (2014).
  - [7] A. Bonaccorso and R. J. Charity, Phys. Rev. C **89** 024619 (2014).
  - [8] H. T. Fortune, Eur. Phys. J. A **54** (2018).
  - [9] J.H.Kelley, J.E.Purcell, C.G.Sheu, Nucl. Phys. A **968** 71 (2017).
  - [10] N. Vinh Mau, Nucl. Phys. A **592** 33 (1995).
  - [11] H. Lenske et al., Rep. Prog. Nucl. Part. Phys. **46** (2001) 187, nucl-th/0012082.
  - [12] F. Barranco, G. Potel, R. A. Broglia, and E. Vigezzi, Phys. Rev. Lett. **119** 082501 (2017).
  - [13] W. von Oertzen, Z. Phys. A **354**, 37 (1996).
  - [14] F. Cappuzzello et al., Phys. Lett. B **711**, 347 (2012).
  - [15] Cavallaro,M., Cappuzzello,F., Bondi,M., Carbone, D., Garcia,V.N., Gargano, A., Lenzi,S.M., Lubian,J.,Agodi,C., Azaiez,F., DeNapoli,M., Foti,A., Franchoo,S., Linares,R., Nicolosi,D., Niikura,M., Scarpaci,J.A., Tropea,S., , Phys. Rev. C **88**, 054601 (2013) and references therein.
  - [16] M. Cavallaro et al., Phys Rev. C **93** 064323 (2016).
  - [17] D. Carbone et al., Phys. Rev. C **95** 034603 (2017).
  - [18] F. Cappuzzello et al., Nature Comm. **6**, 6743 (2015).
  - [19] Ermamatov,M.J., Cappuzzello,F., Lubian, J., Cubero,M., Agodi,C., Carbone,D., Cavallaro,M., Ferreira,J.L., Foti,A., Garcia,V.N., Gargano,A., Lay,J.A., Lenzi,S.M., Linares,R., Santagati,G., Vitturi,A., Phys. Rev. C **94**, 024610 (2016).
  - [20] B. Paes et al., Phys. Rev. C **96**, 044612 (2017).
  - [21] Agodi,C., Giuliani,G., Cappuzzello,F., Bonasera,A., Carbone,D., Cavallaro,M., Foti,A., Linares,R., Santagati,G., , Phys. Rev. C **97**, 034616 (2018).
  - [22] G. Potel et al., EPJ Web of Conferences **17**, 01004 (2011);Potel,G., Barranco,F., Vigezzi,E., Broglia,R.A., , Phys. Rev. Lett. **105** 172502 (2010) and references therein.
  - [23] Thompson, I.J. Reaction mechanism of pair transfer, in 50 Years of Nuclear BCS p. 455, World Scientific (2013).
  - [24] M. A. Franey, B. F. Bayman, J. S. Lilley and W. R. Phillips, Phys. Rev. Lett. **41** 837 (1978).
  - [25] W. von Oertzen and A. Vitturi, Rep. Prog. Phys. **64** 1247 (2001).
  - [26] G.-B. Liu and H. T. Fortune, Phys. Rev. C **42** 167 (1990).
  - [27] S. Hinds and R. Middleton, Phys. Lett. **9** 149 (1964).
  - [28] D.J. Pullen et al., Nucl. Phys. **36** 1 (1962).
  - [29] F. Ajzenberg-Selove et al., Phys. Lett. B **40** 205 (1972).
  - [30] Ajzenberg-Selove,F., Flynn,E.R., Hansen,O., Phys. Rev. C **17** 1283 (1978).
  - [31] M. Majer, R. Raabe, M. Milin, C. Angulo, J. Cabrera, E. Casarejos et al., Eur. Phys. J. A **43** 153 (2010).
  - [32] H. G. Bohlen et al., Phys. At. Nucl. **66** 1495 (2003).
  - [33] H. G. Bohlen, R. Kalpakchieva, W. von Oertzen, TN. Massey, C. B. Gebauer et al, Nuclear Physics A **722** 3c (2003).
  - [34] P. J. Haigh et al., Phys. Rev. C **79**, 014302 (2009).
  - [35] H.G. Bohlen et al., Nucl. Phys. A **734** 345 (2004).
  - [36] M. Freer et al., Nucl. Phys. A **834** 621c (2010).
  - [37] D. M. Brink, Phys. Lett. B **40** 37 (1972).
  - [38] F. Cappuzzello, C. Agodi, D. Carbone and M. Cavallaro, Eur. Phys. J. A **52**, 167 (2016).
  - [39] M. Cavallaro et al., Eur. Phys. Jour. A **48** 59 (2012).
  - [40] F. Cappuzzello et al., Nucl. Instr. Methods A **621** 419 (2010).
  - [41] D. Carbone, Eur. Phys. J. Plus **130**143 (2015).
  - [42] M. Cavallaro et al., Nucl. Instr. and Meth. A **637** 77 (2011).
  - [43] F. Cappuzzello, D. Carbone, and M. Cavallaro, Nucl. Instr. Methods A **638** 74 (2011).
  - [44] Schmitt,K.T., Jones,K.L., Ahn,S., Bardayan,D.W., Bey,A., Blackmon,J.C., Brown,S.M., Chae,K.Y., Chipps,K.A./Cizewski,J.A., Hahn,K.I., Kolata,J.J., Kozub,R.L., Liang,J.F., Matei,C., Matos,M.,



- Matyas,D. , Moazen,B. , Nesaraja,C.D. , Nunes,F.M. , OMalley,P.D. , Pain,S.D. , Peters,W.A. , Pittman,S.T. , Roberts,A. , Shapira,D. , Shriner,J.F. , Smith,M.S. , Spassova,I. , Stracener,D.W. , Upadhyay,N.J. , Villano,A.N. , Wilson,G.L. , Phys Rev. C **88** 064612 (2013) and references therein.
- [45] D.L. Auton, Nucl. Phys. A 157 (1970) 305.
- [46] B. Zwieglinski, W. Benenson, R. G. H. Robertson, and W. R. Coker, Nucl. Phys. A 315, 124 (1979).
- [47] F. Cappuzzello et al., Phys. Lett. B 516, 21 (2001).
- [48] Y. Hirayama et al., Phys. Lett. B **611** 239 (2005).
- [49] H.O.U. Fynbo et al., Nucl. Phys. A **736** 39 (2004).
- [50] F. Cappuzzello et al., Nucl. Phys. A 739 30 (2004).
- [51] A. Ekström et al., Phys. Rev. C **91**, 051301(R) (2015).
- [52] A. Bonaccorso, F. Carstoiu, R. J. Charity, Phys. Rev. C **94** 034604 (2016).
- [53] R. A. Broglia and A. Winther, *Heavy-Ion Reactions* (Addison- Wesley Publishing Company, Redwood City, California, 1991).
- [54] A. Bonaccorso, D. M. Brink and L. Lo Monaco, J. Phys. **G 13** 1407 (1987).
- [55] G. Potel, F. Barranco, E. Vigezzi, R. A. Broglia, Phys. Rev. Lett. **105** 172502 (2010).
- [56] Hesse M, Sparenberg J M, Van Raemdonck F and Baye D, Nucl. Phys. A 640 37 (1998).
- [57] Hesse M, Roland J and Baye D, Nucl. Phys. A 709 184 (2002).
- [58] P. Navrátil, S. Quaglioni, G. Hupin, C. Romero-Redondo, and A. Calci, Phys. Scr. 91, 053002 (2016).
- [59] R. B. Wiringa, private communication.
- [60] S. Cohen and D. Kurath, Nucl. Phys. A101 1 (1967).