

This is the accepted manuscript made available via CHORUS. The article has been published as:

β and γ bands in $N=88, 90$, and 92 isotones investigated with a five-dimensional collective Hamiltonian based on covariant density functional theory: Vibrations, shape coexistence, and superdeformation

S. N. T. Majola *et al.*

Phys. Rev. C **100**, 044324 — Published 30 October 2019

DOI: [10.1103/PhysRevC.100.044324](https://doi.org/10.1103/PhysRevC.100.044324)

β and γ bands in $N = 88, 90, 92$ isotones investigated with a five-dimensional collective Hamiltonian based on covariant density functional theory: vibrations, shape coexistence and superdeformation

S. N. T. Majola,^{1,2,3,4} Z. Shi,⁵ B. Y. Song,⁶ Z. P. Li,⁶ S. Q. Zhang,⁷ R. A. Bark,² J. F. Sharpey-Schafer,⁸ D.G. Aschman,⁴ S. P. Bvumbi,³ T. D. Bucher,^{2,9} D. M. Cullen,^{10,11} T. S. Dinoko,^{2,12} J. E. Easton,^{2,8} N. Erasmus,^{2,8} P. T. Greenlees,¹⁰ D. J. Hartley,¹³ J. Hirvonen,¹⁰ A. Korichi,¹⁴ U. Jakobsson,¹⁰ P. Jones,² S. Jongile,^{1,2,9} R. Julin,¹⁰ S. Juutinen,¹⁰ S. Ketelhut,¹⁰ B. V. Kheswa,^{2,3} N. A. Khumalo,^{2,8} E. A. Lawrie,^{2,8} J. J. Lawrie,⁸ R. Lindsay,⁸ T. E. Madiba,^{2,8} L. Makhathini,^{2,9} S. M. Maliage,^{2,8} B. Maqabuka,^{2,8} K. L. Malatji,^{2,9} P. L. Masiteng,^{2,3,8} P. I. Mashita,^{2,8} L. Mdletshe,^{1,2} A. Minkova,¹⁵ L. Msebi,^{2,8} S. M. Mullins,² J. Ndayishimye,² D. Negi,^{2,16} A. Netshiya,^{2,8} R. Newman,⁹ S. S. Ntshangase,¹ R. Ntshodu,² B. M. Nyakó,¹⁷ P. Papka,^{2,9} P. Peura,¹⁰ P. Rahkila,¹⁰ L. L. Riedinger,¹⁸ M. A. Riley,¹⁹ D. G. Roux,²⁰ P. Ruotsalainen,¹⁰ J. J. Saren,¹⁰ C. Scholey,¹⁰ O. Shirinda,^{2,9} M. A. Sithole,^{2,8} J. Sorri,^{10,21} M. Stankiewicz,^{2,4} S. Stolze,^{10,22} J. Timár,¹⁷ J. Uusitalo,¹⁰ P. A. Vymers,^{2,9} M. Wiedeking,² and G. L. Zimba^{2,3,10}

¹Department of Physics, University of Zululand, Private Bag X1001, KwaDlangezwa, 3886, South Africa

²iThemba LABS, National Research Foundation, P.O. Box 722, Somerset-West 7129, South Africa

³Department of Physics, University of Johannesburg,
P.O. Box 524, Auckland Park 2006, South Africa

⁴Department of Physics, University of Cape Town, P/B X3, Rondebosch 7701, South Africa

⁵School of Physics and Nuclear Energy Engineering, Beihang University, Beijing 100191, China

⁶School of Physical Science and Technology, Southwest University, Chongqing, 400715 China

⁷State Key Laboratory of Nuclear Physics and Technology,
School of Physics, Peking University, Beijing 100871, China

⁸Department of Physics, University of the Western Cape, P/B X17, Bellville 7535 South Africa

⁹Department of Physics, Stellenbosch University, P/B X1, Matieland, 7602, South Africa

¹⁰Department of Physics, P.O. Box 35, FI-40014 University of Jyväskylä, Finland

¹¹Schuster Laboratory, University of Manchester, Manchester M13 9PL, United Kingdom

¹²National Metrology Institute of South Africa, P/B x34, Lynnwood Ridge, Pretoria, 0040, South Africa

¹³Department of Physics, U.S. Naval Academy, Annapolis, Maryland 21402, USA

¹⁴CSNSM-IN2P3-CNRS, F-91405 Orsay Campus, France

¹⁵University of Sofia, Faculty of Physics, Sofia 1164, Bulgaria

¹⁶UM-DAE Centre for Excellence in Basic Sciences, Kalina, Mumbai 400098, India

¹⁷MTA Atomki, P.O. Box 51, H-4001 Debrecen, Hungary

¹⁸University of Tennessee, Department of Physics and Astronomy, Knoxville, Tennessee 37996, USA

¹⁹Department of Physics, Florida State University, Tallahassee, Florida 32306, USA

²⁰Department of Physics, Rhodes University, P.O. Box 94, Grahamstown 6140, South Africa

²¹Sodankylä Geophysical Observatory, University of Oulu, Tähteläntie 62, FI-99600 Sodankylä

²²Physics Division, Argonne National Laboratory, Argonne, Illinois 60439, USA

A comprehensive systematic study is made for the collective β and γ bands in even-even isotopes with neutron numbers $N = 88$ to 92 and proton numbers $Z = 62$ (Sm) to 70 (Yb). Data, including excitation energies, $B(E0)$, $B(E2)$ values and branching ratios from previously published experiments are collated with new data presented for the first time in this study. The experimental data are compared to calculations using a five-dimensional collective Hamiltonian (5DCH) based on the PC-PK1 covariant density functional theory (CDFT). A realistic potential in the quadrupole shape parameters $V(\beta, \gamma)$ is determined from potential energy surfaces (PES) calculated using the CDFT. The parameters of the 5DCH are fixed and contained within the RMF. Overall, a satisfactory agreement is found between the data and the calculations. In line with the energy staggering $S(I)$ of the levels in the 2^+_γ bands, the potential energy surfaces of the CDFT calculations indicate γ -soft shapes in the $N = 88$ nuclides, which become γ -rigid for $N = 90$ and $N = 92$. The nature of the 0^+_β bands changes with atomic number. In the isotopes of Sm to Dy, they can be understood as β -vibrations, but in the Er and Yb isotopes the 0^+_β bands have wavefunctions with a large components in a triaxial superdeformed minimum. In the vicinity of ^{152}Sm , the present calculations predict a soft potential in the β -direction but do not find two coexisting minima. This is reminiscent of ^{152}Sm exhibiting an $X(5)$ behaviour. The model also predicts that the 0^+_β bands be of two-phonon nature, having an energy of twice the 0^+_β band. This is in contradiction with the data and implies that other excitation modes must be invoked to explain their origin.

PACS numbers: 21.10.Re, 21.60.Ev, 27.70.+q, 21.60.Jz

I. INTRODUCTION

clear shape which have been associated with the first ex-

The Bohr Hamiltonian [1, 2] predicts the existence of time-dependent β and γ quadrupole vibrations of the nu-

cited $K^\pi = 0_2^+$ and $K^\pi = 2_\gamma^+$ intrinsic states respectively. The nature of these states has been studied extensively over the years [3–24]. However, despite many decades of research, a full understanding of these levels in even-even deformed nuclei remains elusive. In particular, low-lying rotational bands based on the first excited 0_2^+ state, which are traditionally understood as β vibrational bands show properties at odds with this interpretation [15]. This may be due to the interplay of other modes of excitations contributing to their formation.

The most common competing low-lying 0_2^+ configuration occurs when the nucleus exhibits shape coexistence [21]. Another mode of excitation that may further contribute to the formation of the first excited 0_2^+ states is quadrupole pairing. Pairing is the well-known residual interaction that gives rise to the 0^+ ground states in all even-even nuclei. In the simplest approximation, the strength of the interaction is independent of the orbitals near the Fermi surface, but in a more refined approximation, it is configuration dependent and may lead to the formation of low lying first excited 0_2^+ states that can compete with β vibrations [25–32]. Experimentally, the challenge is to determine which of the three aforementioned excitations best describes the nature of the first-excited 0_2^+ states in the $A \approx 160$ mass region. While there is a long history of doubt being cast on the axial β vibration interpretation of the first excited $K^\pi = 0_2^+$ rotational bands, the $K^\pi = 2_\gamma^+$ bands arise naturally due to axial symmetry breaking [33, 34], be it static or dynamic.

In a search for a more accurate description of the so-called quadrupole vibrational bands, an extensive systematic is carried out for the nuclides in the $A \approx 160$ mass region, between $N = 88$ and 92 and Sm to Yb . To this end we have performed in-beam γ -ray spectroscopy measurements of $K^\pi = 0_2^+$ and $K^\pi = 2_\gamma^+$ bands in the even-even transitional deformed nuclei with neutron numbers $N = 88, 90$ and 92 with proton numbers $Z = 62$ to 70 . In many instances the 0_2^+ band and γ bands have been extended or observed for the first time. The determination of a comprehensive set of level energies and branching ratios between bands allows their electromagnetic properties to be compared to nuclear models.

The theoretical approach adopted here, to come to an understanding of the properties of these bands, is to use a modern form of the Bohr Hamiltonian, a five-dimensional collective Hamiltonian (5DCH) based on the covariant density functional theory (CDFT) [35, 36]. Rather than use a harmonic oscillator potential in β and γ , a realistic potential in $V(\beta, \gamma)$ is determined from the potential energy surfaces calculated using relativistic mean field theory. Thereafter, the inertial parameters of the model are determined and a five-dimensional Bohr Hamiltonian is solved to give the resulting level scheme. The advantages of this approach are that the potential energy surfaces can automatically incorporate any shape-coexisting minima, allowing vibrational and shape-coexisting bands to be calculated on the same footing. An important point is that the parameters of pairing and the relativistic mean-

field are fixed.

In the next section, we present the experimental details. In section III we present our data on $K^\pi = 0_2^+$ and $K^\pi = 2_\gamma^+$ bands comprising new level schemes, γ -ray angular and polarization data, together with ratios of out-of-band to in-band $B(E2)$ ratios. These, together with literature values that fill in gaps in our data, including absolute $B(E2)$ values and $E0$ transitions rates, are then compared to the 5DCH-CDFT calculations in section V.

II. EXPERIMENTAL DETAILS

In total, data from thirteen $\gamma - \gamma$ coincidence measurements have been analyzed to study the low spin spectroscopy of twelve different nuclides. Measurements for two species were carried out using the JUROGAM II arrays [38], while the rest were conducted using the AFRODITE array of iThemba LABS [39]. The experimental details including reactions, beam energies, statistics and arrays are shown in Table I.

Of the thirteen nuclides studied, we present here substantial revisions or additions only to the level schemes of $^{158,160}\text{Yb}$ and ^{158}Dy , with an emphasis on bands relevant to this study, namely the first-excited 0^+ (denoted as 0_2^+ in this manuscript), ground and γ bands. Other level schemes deduced during the course of our investigations have been presented elsewhere [40–45], however, in general, spectroscopic information such as DCO ratios, polarization anisotropies and ratios of out-of-band to in-band $B(E2)$ s have not previously been reported. Table II contains this spectroscopic information. In this study, the technique of directional correlation from oriented states (DCO or R_{DCO}) has been used in order to assist in determining the multipolarities of new transitions and to confirm those of transitions deduced from previous studies. DCO matrices were prepared in such a way that transitions detected at a forward and/or backward angle θ_1 are placed on one axis and the coincident transitions detected at an angle θ_2 , close to 90° , are placed on the other axis. The R_{DCO} ratio is then defined by

$$R_{DCO} = \frac{I(\gamma_1(\theta_2)\gamma_2(\theta_1))}{I(\gamma_1(\theta_1)\gamma_2(\theta_2))} \quad (1)$$

where one of the transitions in the ratio is chosen to be of known stretched $E2$ character. In this work, R_{DCO} ratios for all three data sets give values close 0.6 and 1 when the second transition is of stretched pure dipole or quadrupole character, respectively. The DCO ratios for the JUROGAM II array were deduced using detectors in rings at 158° and $86^\circ + 94^\circ$. Similarly, detectors at angles 135° and 90° were used to determine DCO ratios for data collected using the AFRODITE array. In order to determine the magnetic or electric nature of the transitions, linear polarization measurements have been performed.

TABLE I. Experimental details showing apparatus and target-beam combinations that were used for the experiments analyzed in this work. The acquired statistics for each experiment is also shown in the table.

Nucleus	Reaction(s)	Beam energy (MeV)	Target thickness mg/cm ²	events ×10 ⁹	Spectrometer(s)
<i>N</i> = 88					
¹⁵⁸ Yb	¹⁴⁴ Sm(¹⁸ O,4n)	78	3	2.0 $\gamma\gamma$	AFRODITE
¹⁵⁶ Er	¹⁴⁷ Sm(¹² C,3n)	65	6	1.4 $\gamma\gamma$	AFRODITE
¹⁵⁴ Dy	¹⁵⁵ Gd(³ He,4n)	37.5	3.2	0.4 $\gamma\gamma$	AFRODITE
¹⁵² Gd	¹⁵² Sm(α ,4n)	45	5	0.5 $\gamma\gamma$	AFRODITE
¹⁵⁰ Sm	¹³⁶ Xe(¹⁸ O,4n)	75	5	0.5 $\gamma\gamma$	AFRODITE
	¹⁴⁸ Nd(α ,2n)	25	5	2.0 $\gamma\gamma\gamma$	JUROGAM II
<i>N</i> = 90					
¹⁶⁰ Yb	¹⁴⁷ Sm(¹⁶ O,3n)	73	4	2.0 $\gamma\gamma$	AFRODITE
¹⁵⁸ Er	¹⁵⁰ Sm(¹² C,4n)	65	1	0.4 $\gamma\gamma$	AFRODITE
¹⁵⁶ Dy	¹⁵⁵ Gd(α ,3n)	25	0.98	14 $\gamma\gamma$	JUROGAM II
¹⁵⁴ Gd	¹⁵² Sm(α ,2n)	25	4	0.5 $\gamma\gamma$	AFRODITE
<i>N</i> = 92					
¹⁶² Yb	¹⁵⁰ Sm(¹⁶ O,4n)	83	3	7.4 $\gamma\gamma$	AFRODITE
¹⁶⁰ Er	¹⁵² Sm(¹² C,4n)	64	5	2.7 $\gamma\gamma$	AFRODITE
¹⁵⁸ Dy	¹⁵⁶ Gd(α ,2n)	27	11	1.1 $\gamma\gamma$	AFRODITE

In effect, the polarization sensitivity possessed by both the AFRODITE and JUROGAM II arrays has allowed us to determine the electromagnetic nature of transitions reported in this work. In both cases, this was achieved by using clover detectors close to 90°. Here the clovers are treated as Compton polarimeter apparatus. The polarization anisotropy A_p can then be obtained using

$$A_p = \frac{aN_v - N_h}{aN_v + N_h}, \quad (2)$$

where N_h and N_v represent the number of γ rays, which respectively scatter perpendicular or parallel to the beam direction between the crystals of a clover detector. The relative efficiency parameter a is a normalization constant used to account for the asymmetry of a configuration. Pure stretched electric transitions such as $E1$ s and $E2$ s, preferably scatter in the perpendicular direction with respect to the beam axis. As a result, a polarization anisotropy measurement A_p yields a value with a positive sign for a stretched pure electric transition. Conversely, a value with a negative sign is obtained for a stretched pure magnetic dipole.

III. LEVEL SCHEMES

A. *N* = 88 isotones

The spectroscopy of low spin structures in ¹⁵⁰Sm [40], ¹⁵²Gd [40] and ¹⁵⁴Dy [41, 91] has been reported in our previous in-beam works. Here a couple of levels have been added to both the odd and even spin γ -bands of ¹⁵⁰Sm and ¹⁵²Gd. In this work only ratios of transition rates extracted for the above mentioned isotones, relating

to the decays out of the 0_2^+ and 2_7^+ bands, are reported for the first time.

For ¹⁵⁸Yb, a completely new sequence of rotational levels built on the (2_7^+) state, as illustrated in Fig. 1, is observed. A spectrum supporting the placements of the transitions associated with this band is shown in Fig. 2a. The spectrum gated on the 486 keV doublet, depopulating the 4^+ and 8^+ members, clearly shows the in-band transitions, namely the 486, 488, 555 and 634 keV γ rays. It also shows numerous interband transitions connecting this structure to the ground band. The 579 and 937 keV transitions decaying out of the 937 keV level of this band (to the 0^+ and 2^+ members of the ground band) confine the possible spin and parity assignment of the 937 keV level to either $I^\pi = 0^+, 1^-, 1^+$ or 2^+ . The DCO value for the 937 keV transition is consistent with it being a stretched $E2$ transition, and this leaves the 2^+ as the most probable assignment for the 937 keV level. The DCO measurement that has been carried out for the 579 keV transition is indicative of this transition being a dipole, thus validating our spin and parity assignment for the 937 keV level. Similar decay patterns are observed for the transitions (i.e. 589, 1065, 508, 1077, 349, 994 and 903 keV transitions) decaying out of the 1423, 1911 and 2951 keV levels. The DCO measurements were successfully performed for most of these transitions. By applying analogous arguments (used to infer spin and parity assignment for the 937 keV level, the 1423, 1911 and 2951 keV levels have been respectively assigned to $I^\pi = 4^+, 6^+, 8^+$ and 10^+ . The 3585 and 4300 keV levels are assumed to be additional members of this sequence connected by stretched $E2$ transitions.

The spin and parity assignment and excitation energies of levels in this band relative to the ground band as well as its decay pattern identify it as the even spin sequence

of the 2_{γ}^{+} band.

TABLE II: Experimentally determined properties for the nuclei investigated in this study with the exception of ^{152}Sm , ^{154}Dy , ^{158}Er and ^{162}Yb . Data includes excitation levels E_x (in keV), γ -ray energies E (in keV), spins for the initial and final states, polarization anisotropy A_p , DCO ratios and assigned multipolarities. All DCO ratios were deduced by gating on stretched $E2$ transitions with the exception of those marked with *, which were measured by gating on $E1$ transitions. The symbol $^{\zeta}$ is used to denote DCO ratios that were deduced using the alpha induced reaction data in ^{156}Dy . The branching ratios for out-of-band to in-band transitions, $(BE2)_{out}/(BE2)_{in}$, for the 0_2^{+} and 2^{+} bands are also listed. Empty entries refer to information that could not be obtained.

Nucleus/band name	E_i	I_i	I_f	$E_{\gamma}(keV)$	DCO	A_p	Assign	$(BE2)_{out}/(BE2)_{in}$
^{150}Sm								
ground								
	334	2	0	333.9(1)	0.98(10)	0.09(10)	E2	
	773	4	2	439.3(1)	0.94(10)	0.09(10)	E2	
	1279	6	4	505.6(1)	0.97(10)	0.07(10)	E2	
	1837	8	6	558.1(1)	0.97(10)	0.08(20)	E2	
	2433	10	8	596.1(1)	1.09(20)	0.08(20)	E2	
	3048	12	10	615.2(1)	0.94(50)	0.06(60)	E2	
	3676	14	12	627.5(1)	0.97(11)		E2	
	4305	16	14	629.6(2)			E2	
0_2^{+} band								
	740	0	2	406.5(2)	0.67(36)	0.07(54)	E2	
	1046	2	0	305.6(2)	0.78(17)	0.06(24)	E2	
			4	272.8(2)			E2	
			2	712.1(2)	1.41(16)	-0.002(10)	M1/E2	
			0	1046.0(2)			E2	0.009(2)
	1449	4	2	403.0(2)	1.02(20)	0.10(20)	E2	
			4	676.0(3)	1.27(12)	-0.04(11)	M1/E2	
			2	1115.3(2)			E2	0.001(10)
	1822	6	4	372.7(2)	0.82(90)	0.12(17)	E2	
			6	543.1(3)	0.66(30)	-0.07(24)	M1/E2	
			4	1049.1(3)			E2	0.002(1)
	2247	8	6	424.9(3)	0.86(11)	0.11(31)	E2	
			6	967.9(2)	1.31(22)		E2	0.003(1)
	2746	10	8	499.0(2)	0.64(33)		E2	
	3306	(12)	10	560.3(3)			E2	
(even)								
	1194	2	0	453.3(2)			E2	
			2	859.8(2)			M1/E2	
			0	1193.7(2)			E2	
	1642	4	2	448.9(1)	1.24(36)		E2	
			4	869.4(3)	1.36(31)	-0.04(30)	M1/E2	
			2	1308.7(2)		0.15(76)	E2	0.033(4)
	2107	6	4	464.8(2)	1.31(21)	0.39(25)	E2	
			6	285.6(2)			M1/E2	
			6	828.5(3)			M1/E2	
			4	1334.0(3)			E2	0.027(4)
	2664	(8)	6	557.2(3)			E2	
			6	1385.7(3)			E2	
	3200	10	(8)	535.3(1)			E2	
			(8)	1362.9(1)			E2	
(odd)								
	1505	3	4	731.4(2)			M1/E2	
			2	1170.7(2)	1.08(33)	-0.00(2)	M1/E2	
	2020	5	3	515.8(1)	0.91(48)		E2	
			4	377.8(1)			M1/E2	

Continued on next page

TABLE II –continued from previous table

Nucleus/band name	E_i	I_i	I_f	$E_\gamma(keV)$	DCO	A_p	Assign	$(BE2)_{out}/(BE2)_{in}$
^{152}Gd ground			6	741.8(2)			M1/E2	
			4	1247.1(1)	1.03(36)	-0.03(21)	M1/E2	0.036(2)
	2570	(7)	5	550.0(1)	0.74(13)	0.05(20)	E2	
			(6)	463.0(1)			M1/E2	
			8	748.6(2)			M1/E2	
	3155	(9)	(7)	1291.5(2)	1.02(62)	-0.09(27)	M1/E2	0.019(1)
			(7)	585.0(2)	0.91(21)		E2	
	345	2	0	344.73(10)	1.08(1)	0.05(1)	E2	
	756	4	2	411.58(10)	1.05(1)	0.06(1)	E2	
	1229	6	4	472.40(10)	1.01(1)	0.06(1)	E2	
	1749	8	6	520.02(10)	1.03(1)	0.06(1)	E2	
	2303	10	8	554.19(10)	1.07(1)	0.06(1)	E2	
	2888	12	10	585.2(1)	0.98(2)	0.05(1)	E2	
	3504	14	12	616.1(1)	0.87(3)	0.06(2)	E2	
	4147	16	14	642.4(2)	0.93(7)	0.11(4)	E2	
0_2^+ band (even) (odd)	616	0	2	271.50(11)			E2	
	931	2	0	315.54(12)			E2	
			2	586.77(13)			M1/E2	
			0	931.35(17)		0.16(6)	E2	0.011(1)
	1283	4	2	352.12(13)	1.13(4)	0.10(2)	E2	
			4	527.36(13)	1.00(3)	-0.03(3)	M1/E2	
			2	938.8(1)			E2	
	1669	6	4	386.37(13)	1.02(3)	0.10(3)	E2	
			6	441.08(16)	1.12(5)	-0.04(5)	M1/E2	
	2141	8	8	392.2(2)			M1/E2	
			6	472.01(6)	1.28(3)	0.06(1)	E2	
	2695	10	10	392.1(4)			M1/E2	
			8	553.94(13)	1.04(2)	0.08(2)	E2	
	1110	2	2	765.93(16)			M1/E2	
			0	1110.3(2)			E2	
	1552	4	2	441.44(17)	1.23(13)		E2	
			4	795.52(10)	0.62(13)	-0.09(10)	M1/E2	
			2	1207.02(15)			E2	0.014(10)
	1999	6	4	448.20(11)			E2	
			6	771.33(12)			M1/E2	
^{154}Dy ground			4	1243.44(23)			E2	0.006(2)
	2464	8	6	464.60(12)	1.23(11)	0.08(7)	E2	
			6	1236.20(19)			E2	0.004(1)
	2969	10	8	504.56(19)			E2	
	1435	3	4	679.08(21)	0.79(10)	-0.02(11)	M1/E2	
			2	1090.73(10)			M1/E2	
	1863	5	3	427.94(17)	1.11(16)	0.05(9)	E2	
			6	634.30(19)	0.55(11)	-0.09(5)	M1/E2	
			4	1107.21(16)			M1/E2	0.019(11)
	2304	7	5	440.64(15)	1.29(14)	0.11(9)	E2	
			6	1075.38(16)		-0.02(5)	M1/E2	0.014(5)
	2780	9	7	476.73(15)		0.12(9)	E2	
			8	1030.93(19)			M1/E2	0.007(5)
	3299	11	9	519.11(19)			E2	
			10	995.96(18)			M1/E2	
	3857	13	11	557.9(3)			E2	
	335	2	0	334.7(1)			E2	
	747	4	2	412.5(1)			E2	

Continued on next page

TABLE II –continued from previous table

Nucleus/band name	E_i	I_i	I_f	$E_\gamma(keV)$	DCO	A_p	Assign	$(BE2)_{out}/(BE2)_{in}$
0_2^+ band	1224	6	4	477.2(1)			E2	
	1748	8	6	523.7(1)			E2	
	2305	10	8	557.1(1)			E2	
	2894	12	10	589.1(1)			E2	
	3511	14	12	616.33(12)			E2	
	4175	16	14	664.49(12)			E2	
	4871	18	16	696.00(11)			E2	
	5567	20	18	695.81(14)			E2	
	661	0	2	326.22(12)			E2	
	905	2	0	245.26(13)			E2	
			2	570.71(13)			M1/E2	
			0	905.29(14)			E2	0.008(3)
	1252	4	2	346.71(13)			E2	
			4	504.86(13)			M1/E2	
	1659	6	4	407.07(13)			E2	
			6	435.13(14)			M1/E2	
	2164	8	8	416.30(14)			M1/E2	
			6	504.59(13)			E2	
	2760	10	8	595.73(13)			E2	
	3291	12	10	531.54(14)			E2	
$\gamma(\text{even})$			10	985.01(15)			E2	
	1028	2	0	367.1(13)			E2	
	1028		2	692.82(15)			M1/E2	
	1443	4	2	415.26(16)			E2	
	1443		4	695.82(13)			M1/E2	
	1443		2	1108.05(15)			E2	0.007(1)
	1886	6	4	443.35(13)			E2	
	1886		6	661.60(14)			M1/E2	
	1886		4	1138.69(16)			E2	0.003(1)
	2371	8	6	485.43(13)			E2	
$\gamma(\text{odd})$	2371		8	622.81(16)			M1/E2	
	2371		6	1147.12(19)			E2	0.001(1)
	2913	10	8	541.66(14)			E2	
	3515	12	10	602.22(25)			E2	
	1334	3	4	587.75(14)			M1/E2	
			2	999.82(14)			M1/E2	
	1740	5	3	405.68(14)			E2	
			4	993.14(13)			M1/E2	0.025(2)
	2183	7	5	443.47(13)			E2	
			6	959.36(14)			M1/E2	0.016(1)
^{156}Er ground	2678	9	7	495.01(13)			E2	
			8	930.47(18)			M1/E2	0.013(1)
	3223	11	9	545.11(14)			E2	
			10	918.12(15)			M1/E2	
	3810	13	11	586.63(17)			E2	
	345	2	0	344.6(1)	1.11(5)		E2	
	798	4	2	453.1(1)	1.30(6)		E2	
	1341	6	4	543.5(1)	1.16(6)		E2	
	1959	8	6	618.2(1)	1.17(9)		E2	
	2634	10	8	674.31(13)	0.98(19)		E2	
$\gamma(\text{even})$	3315	12	10	682.02(13)			E2	
	3838	14	12	522.47(13)			E2	
	4383	16	14	544.88(13)			E2	
	1221	2	2	876.49(11)			M1	
			0	1220.93(25)			E2	

Continued on next page

TABLE II –continued from previous table

Nucleus/band name	E_i	I_i	I_f	$E_\gamma(keV)$	DCO	A_p	Assign	$(BE2)_{out}/(BE2)_{in}$
0_2^+ band	1547	4	2	325.71(19)			E2	
			4	749.10(16)			M1	
			2	1202.24(15)			E2	0.004(1)
	1970	6	4	423.39(14)			E2	
			6	629.09(14)			M1	
			4	1172.62(15)			E2	0.004(1)
	2482	8	6	511.34(13)			E2	
			8	521.80(14)			M1	
			6	595.91(16)			E2	
			6	1140.62(17)			E2	0.002(1)
	3044	10	8	562.55(13)			M1/E2	
			8	1084.74(17)			E2	0.001(1)
	3653	12	10	609.29(14)			E2	
			10	708.69(17)			E2	
	930	2	2	585.81(14)			M1/E2	
			0	930.51(13)			E2	
	1406	4	2	475.72(14)			E2	
			4	608.30(14)			M1/E2	
			2	1061.52(17)			E2	0.008(1)
	1886	6	4	339.27(19)			E2	
			4	480.41(13)			E2	
			6	544.80(13)			M1/E2	
			4	1089.17(14)			E2	0.009(1)
	2378	8	8	418.28(17)			M1/E2	
			6	490.99(13)			E2	
			6	1036.88(14)			E2	0.009(1)
	2945	10	8	566.44(13)			E2	
			8	985.0(10)			E2	0.004(1)
	3592	12	10	645.80(14)			E2	
	4284	14	12	692.32(17)			E2	
$\gamma(\text{odd})$	1352	3	2	420.72(16)			M1/E2	
			4	554.1(10)			M1/E2	
			2	1006.89(15)			M1/E2	
	1836	5	3	484.12(14)			E2	
			4	289.3(10)			M1/E2	
			4	430.00(16)			M1/E2	
			4	1038.30(17)			M1/E2	0.021(2)
	2369	7	5	533.38(14)			E2	
			6	1028.07(18)			M1/E2	0.007(1)
	2963	9	7	593.66(15)			E2	
			8	1003.4(2)			M1/E2	0.036(3)
^{158}Yb ground	358	2	0	358.02(10)			E2	
	834	4	2	476.41(10)			E2	
	1404	6	4	569.36(10)			E2	
	2047	8	6	643.44(10)			E2	
	2744	10	8	697.29(10)			E2	
	3427	12	10	682.87(10)			E2	
	3936	14	12	508.67(12)			E2	
	4504	16	14	567.81(13)			E2	
$\gamma(\text{even})$ band	937	2	2	579.2(1)	0.67(8)		M1/E2	
			0	937.2(1)	0.97(6)		E2	
	1423	4	2	486.1(1)			E2	
			4	589.3(1)	0.59(7)		M1/E2	
			2	1065.3(2)	1.15(6)		E2	0.020(4)
	1911	6	4	488.0(1)			E2	

Continued on next page

TABLE II –continued from previous table

Nucleus/band name	E_i	I_i	I_f	$E_\gamma(keV)$	DCO	A_p	Assign	$(BE2)_{out}/(BE2)_{in}$
^{154}Gd ground			6	507.7(2)			M1/E2	
			4	1077.3(2)	1.14(10)		E2	0.019(2)
	2397	8	8	350.3(1)	0.61(11)		M1/E2	
			6	485.9(2)	0.91(8)		E2	
			6	993.6(2)	1.12(12)		E2	0.018(2)
	2951	10	8	554.6(2)	1.11(19)		E2	
			8	903.4(3)			E2	0.021(5)
	3585	12	10	634.3(2)			E2	
			10	840.2(2)			E2	0.09(2)
	4300	14	12	714.1(3)			E2	
	123	2	0	122.86(10)	1.01(10)		E2	
	370	4	2	247.44(10)	1.02(11)	0.15(10)	E2	
	716	6	4	345.9(1)	1.01(10)	0.15(5)	E2	
	1142	8	6	426.0(1)	1.00(10)	0.14(15)	E2	
0_2^+ band	1634	10	8	491.8(1)	1.013(20)	0.13(10)	E2	
	2180	12	10	546.64(13)	1.01(15)	0.14(10)	E2	
	2772	14	12	591.98(14)	0.97(3)	0.12(30)	E2	
	3398	16	14	625.87(18)	1.05(15)		E2	
	814	2	0	134.80(20)			E2	
			4	443.61(14)			E2	
			2	690.97(14)			M1/E2	
			0	813.88(19)			E2	
	1045	4	2	231.66(14)			E2	
			6	329.20(16)			E2	
			4	675.17(13)	0.85(12)	-0.04(10)	M1/E2	
			2	922.66(14)	0.88(30)	0.11(20)	E2	0.002(1)
	1363	6	4	317.63(13)	0.99(10)	0.12(10)	E2	
			6	646.86(13)		-0.05(10)	M1/E2	
$\gamma(\text{even})$			4	993.05(14)	0.95(20)	0.03(10)	E2	0.001(1)
	1753	8	6	389.96(13)	1.00(10)	0.16(10)	E2	
			8	610.85(14)			M1/E2	
			6	1037.23(17)			E2	0.001(1)
	2190	10	8	436.89(13)	1.01(10)		E2	
			10	555.90(15)	0.86(5)	-0.01(5)	M1/E2	
			8	1047.98(18)			E2	0.001(1)
	2617	12	10	427.11(13)	1.00(1)		E2	
			12	435.80(13)	1.01(1)		M1/E2	
			10	983.5(3)			E2	0.001(1)
	3022	14	12	404.76(14)	0.98(4)	0.09(30)	E2	
	3484	16	14	462.73(17)	1.11(12)		E2	
	995	2	2	871.42(14)			M1/E2	
			0	994.32(14)	0.95(4)		E2	
$\gamma(\text{odd})$	1261	4	2	266.91(17)	0.96(5)	0.07(5)	E2	
			4	890.97(13)	0.91(3)	-0.13(5)	M1/E2	
			2	1138.56(14)	0.89(8)		E2	0.008(1)
	1603	6	4	341.96(15)			E2	
			6	886.86(13)	0.77(20)	-0.02(1)	M1/E2	
			4	1233.10(15)			E2	0.004(1)
	2014	8	6	411.15(14)	0.99(5)	0.16(5)	E2	
			7	207.8(5)			M1/E2	
			6	651.3(10)			E2	
			8	872.18(14)			M1/E2	
			6	1298.36(16)	0.93(7)		E2	0.004(1)
	2485	10	8	470.97(14)	0.97(13)	0.19(6)	E2	
			10	851.21(15)	0.6(4)	-0.13(3)	M1/E2	
			8	1343.55(19)	1.04(11)	0.19(24)	E2	0.004(1)

Continued on next page

TABLE II –continued from previous table

Nucleus/band name	E_i	I_i	I_f	$E_\gamma(keV)$	DCO	A_p	Assign	$(BE2)_{out}/(BE2)_{in}$
$\gamma(\text{odd})$	3005	12	10	519.65(12)			E2	
			12	824.7(14)			M1/E2	
			10	1372.14(18)			E2	0.006(1)
	1125	3	4	755.20(14)	0.88(3)	-0.05(2)	M1/E2	
			2	1002.78(13)	0.93(2)	-0.05(4)/	M1/E2	
	1429	5	3	303.86(17)	1.02(7)	0.18(5)	E2	
			6	713.18(14)	0.69(5)	-0.01(5)	M1/E2	
			4	1059.31(13)	0.72(1)	0.01(2)	M1/E2	0.040(3)
	1807	7	5	377.23(14)		0.24(4)	E2	
			8	664.45(14)	0.92(4)	-0.11(3)/	M1/E2	
			6	1090.65(13)	0.53(2)	0.06(3)	M1/E2	0.023(1)
	2247	9	7	440.22(14)		0.08(4)	E2	
			10	613.31(18)	0.81(5)	-0.01(4)	M1/E2	
			8	1104.88(14)	0.57(1)		M1/E2	0.017(1)
	2741	11	9	494.13(15)		0.16(5)	E2	
^{156}Dy ground			10	1107.38(18)	0.57(1)	0.16(8)	M1/E2	0.011(1)
	3278	13	11	537.31(19)	1.23(15)	0.15(9)	E2	
	3278	13	12	1098.1(12)	0.57(1)		M1/E2	
	138	2	0	137.63(10)	0.92(10) $^\zeta$		E2	
	404	4	2	266.23(10)	1.00(10) $^\zeta$	0.14(10)	E2	
	770	6	4	366.07(10)	1.02(10) $^\zeta$	0.13(10)	E2	
	1215	8	6	445.14(10)	1.04(15) $^\zeta$	0.096(30)	E2	
	1724	10	8	508.89(10)	1.04(10) $^\zeta$	0.098(12)	E2	
	2285	12	10	560.67(10)	1.00(13) $^\zeta$	0.089(12)	E2	
	2886	14	12	601.84(11)	0.90(10) $^\zeta$	0.06(2)	E2	
	3522	16	14	635.41(11)	0.99(3) $^\zeta$	0.099(2)	E2	
0_2^+ band	828	2	4	424.04(22)			E2	
			2	690.69(11)			M1/E2	
	1088	4	2	259.47(12)			E2	
			6	317.70(11)			E2	
			4	683.84(12)			M1/E2	
			2	950.44(12)			E2	0.0010(10)
	1437	6	4	348.81(10)	0.98(3)		E2	
			6	666.70(12)	0.68(10)	-0.05(41)	M1/E2	
			4	1032.91(12)			E2	0.0010(10)
	1858	8	6	421.31(10)			E2	
			8	642.91(11)	0.71(3)	-0.10(30)	M1/E2	
			6	1088.14(15)			E2	0.001(1)
	2315	10	8	456.64(13)			E2	
			10	590.72(15)	0.71(10)		M1/E2	
			8	1099.71(14)			E2	0.001(1)
$\gamma(\text{even})$	891	2	2	752.41(14)			M1/E2	
	1168	4	2	277.18(21)			E2	
			4	763.92(13)	0.708(41) $^\zeta$	-0.060(40)	M1/E2	
			2	1030.40(12)			E2	0.004(2)
	1524	6	4	356.53(15)			E2	
			6	754.45(16)			M1/E2	
			4	1120.60(17)			E2	0.004(1)
	1956	8	6	431.86(14)		0.101(70)	E2	
			8	740.95(15)	0.443(44) $^\zeta$	-0.033(60)	M1/E2	
			8	1186.34(15)			E2	0.003(1)
	2447	10	8	490.51(14)			E2	
			10	1231.54(16)		0.028(150)	E2	0.009(1)
	2969	12	10	522.00(15)			E2	
			12	683.90(10)			M1/E2	

Continued on next page

TABLE II –continued from previous table

Nucleus/band name	E_i	I_i	I_f	$E_\gamma(keV)$	DCO	A_p	Assign	$(BE2)_{out}/(BE2)_{in}$
$\gamma(\text{odd})$	3523	14	10	1244.96(17)			E2	0.014(1)
			12	556.00(16)			E2	
			14	637.99(13)			M1/E2	
			12	1238.16(18)			E2	
	1022	3	4	617.73(8)			M1/E2	
			2	884.25(11)			M1/E2	
	1335	5	3	312.86(18)			E2	
			6	565.38(19)			M1/E2	
	1728	7	4	931.14(13)	0.59(5) $^\zeta$	0.04(6)	M1/E2	0.031(2)
			5	393.04(13)			E2	
			8	512.37(14)			M1/E2	
			6	958.11(13)	0.451(30) $^\zeta$	0.02(10)	M1/E2	
	2190	9	7	462.46(13)		0.093(20)	E2	0.020(1)
			10	466.52(15)			M1/E2	
	2711	11	8	975.42(14)	0.562(4) $^\zeta$	0.042(5)	M1/E2	0.014(1)
			9	520.07(13)		0.069(4)	E2	
			10	264.58(18)			M1/E2	
			10	986.61(17)			M1/E2	
	3274	13	11	563.01(14)			E2	0.011(10)
			12	989.18(19)			M1/E2	
	3861	15	13	587.16(14)			E2	
^{160}Yb ground	243	2	0	243.20(10)			E2	
	639	4	2	395.39(10)			E2	
	1148	6	4	508.73(10)			E2	
	1737	8	6	589.27(10)			E2	
	2374	10	8	637.03(10)			E2	
	2960	12	10	586.50(10)			E2	
	3364	14	12	404.02(10)			E2	
	3848	16	14	484.08(10)			E2	
0_2^+ band	1293	2	2	1048.65(15)			M1/E2	
			0	1292.01(21)			E2	
	1592	4	2	299.33(21)			E2	
			4	953.34(15)	0.95(12)		M1/E2	
	1958	6	2	1348.65(15)			E2	0.0037(10)
			4	365.60(11)	1.10(18)		E2	
			6	809.89(12)	0.54(5)		M1/E2	
			4	1318.74(11)	0.92(9)		E2	
	2365	8	6	406.81(10)	1.20(11)		E2	0.0047(10)
			6	1216.91(11)	0.97(10)		E2	
$\gamma(\text{even})$	2841	10	8	476.22(11)			E2	0.0036(10)
			8	566.18(10)			E2	
			8	1104.52(33)			E2	
	821	2	2	577.16(10)	0.77(6)	-0.05(16)	M1/E2	
			0	820.44(10)	0.98(10)	0.84(25)	E2	
	1256	4	2	435.15(10)	1.02(9)		E2	
			4	616.71(10)	0.78(7)	-0.15(12)	M1/E2	
	1744	6	2	1012.67(11)		-0.20(27)	E2	0.007(1)
			4	488.04(10)	0.99(9)		E2	
			6	596.37(10)	0.72(6)		M1/E2	
			4	1104.52(33)		0.17(40)	E2	
$\gamma(\text{even})$	2275	8	6	530.90(10)	1.15(10)	0.22(14)	E2	0.0019(2)
			8	537.45(15)			M1/E2	
			6	1127.35(16)			E2	
	2790	10	8	425.55(10)	0.80(9)		E2	
			8	515.63(10)	0.92(8)		E2	

Continued on next page

TABLE II –continued from previous table

Nucleus/band name	E_i	I_i	I_f	$E_\gamma(keV)$	DCO	A_p	Assign	$(BE2)_{out}/(BE2)_{in}$
$\gamma(\text{odd})$	3319	12	8	1053.14(11)			E2	0.048(4)
			10	478.37(10)	0.68(69)		E2	
			10	528.84(10)	0.98(9)		E2	
	3870	14	12	550.79(10)			E2	
	1113	3	2	292.35(11)	0.68(8)	-0.07(32)	M1/E2	
			4	474.15(11)	0.72(7)		M1/E2	
			2	869.61(10)	0.69(6)	0.07(12)	M1/E2	
	1574	5	3	461.33(10)	1.02(12)	0.58(14)	E2	
			4	318.05(11)			M1/E2	
			6	427.08(11)	0.98(10)		M1/E2	
	2109	7	4	935.43(10)	0.58(5)	0.24(8)	M1/E2	0.00366(26)
			5	534.62(10)	1.09(10)		E2	
			6	365.55(12)			M1/E2	
	2701	9	6	961.51(11)		0.44(15)	M1/E2	0.013(1)
			7	592.47(10)			E2	
			8	963.71(15)		0.44(15)	M1/E2	
^{158}Dy ground	3331	11	9	629.71(10)	1.03(9)		E2	0.0086(11)
	4017	13	11	686.00(12)			E2	
	99	2	0	98.58(10)			E2	
	317	4	2	217.98(10)			E2	
	637	6	4	320.34(10)			E2	
	1043	8	6	405.97(10)			E2	
	1519	10	8	475.75(10)			E2	
	2048	12	10	529.07(10)			E2	
	2611	14	12	563.28(10)			E2	
	3189	16	14	577.81(13)			E2	
0_2^+ band	1086	2	4	769.2(4)			E2	
			2	987.1(5)			M1/E2	
			0	1086.1(2)			E2	
	1279	4	2	193.7(2)			E2	
			6	642.50(18)			E2	
			4	962.5(3)			M1/E2	
	1554	6	2	1180.9(3)			E2	0.020(4)
			4	274.59(17)			E2	
			8	510.86(13)			E2	
	1901	8	6	917.13(12)			M1/E2	0.008(3)
			4	1237.75(11)			E2	
			6	346.63(8)			E2	
	2267	10	6	424.79(17)			E2	0.004(1)
			8	857.76(15)			M1/E2	
			6	1263.79(11)			E2	
$\gamma(\text{even})$	2698	12	8	367.5(4)			E2	0.015(3)
			8	404.58(17)			E2	
			10	748.59(15)			M1/E2	
	3259	14	8	1223.99(19)			E2	
			10	430.55(22)			E2	
			12	650.17(7)			M1/E2	
	946	2	10	1179.10(13)			E2	
			12	561.01(13)			E2	
			12	1211.17(25)			E2	
	1163	4	(2)	216.69(11)			E2	
			3	118.8(10)			M1/E2	
			6	526.08(22)			E2	

Continued on next page

TABLE II –continued from previous table

Nucleus/band name	E_i	I_i	I_f	$E_\gamma(keV)$	DCO	A_p	Assign	$(BE2)_{out}/(BE2)_{in}$
$\gamma(\text{odd})$	1476	6	4	846.39(5)			M1/E2	
			2	1064.43(15)			E2	0.005(3)
			4	312.92(6)			E2	
			5	161.78(15)			M1/E2	
			6	839.00(4)			M1/E2	
			4	1159.31(20)			E2	0.003(1)
	1863	8	6	386.69(6)			E2	
			7	188.1(1)			M1/E2	
			6	308.5(1)			E2	
			8	819.66(15)			M1/E2	
			6	1225.69(20)			E2	0.010(1)
			8	486.33(7)			E2	
	2349	10	8	448.2(1)			E2	
			10	830.30(18)			M1/E2	
			8	1306.01(10)			E2	0.001(1)
		12	10	516.97(15)			E2	
			12	818.34(18)			M1/E2	
			(2)	97.90(10)			M1/E2	
	1044	3	4	727.63(9)			M1/E2	
			2	945.67(5)			M1/E2	
			4	151.1(4)			M1/E2	
	1314	5	3	269.90(7)			E2	
			6	677.22(5)			M1/E2	
			4	997.53(3)			M1/E2	0.034(2)
	1675	7	5	360.39(4)			E2	
			6	198.61(13)			M1/E2	
			8	631.59(6)			M1/E2	
			6	1037.61(4)			M1/E2	0.020(1)
	2111	9	7	436.85(4)			E2	
			10	592.75(15)			M1/E2	
			8	1068.44(5)			M1/E2	0.015(1)
	2610	11	9	498.38(4)			E2	
			12	562.21(12)			M1/E2	
			10	1091.1(10)			M1/E2	0.013(1)
	3147	13	11	537.59(15)			E2	
			12	1099.8(10)			M1/E2	0.003(1)
	3685	15	13	537.5(2)			E2	
			14	1074.1(2)			M1/E2	
^{160}Er ground	125	2	0	125.43(6)			E2	
	389	4	2	263.87(6)			E2	
	765	6	4	375.71(6)			E2	
	1229	8	6	464.08(6)			E2	
	1761	10	8	531.86(6)			E2	
	2340	12	10	579.22(6)			E2	
	2932	14	12	592.21(6)			E2	
	3466	16	14	534.04(6)			E2	
0_2^+ band	1008	2	0	1008.0(1)			E2	
	1230	4	4	840.31(17)			M1/E2	
			2	1104.30(24)			E2	
	1542	6	4	312.48(20)			E2	
			6	777.2(3)			M1/E2	
			4	1152.64(12)			E2	0.003(2)
	1921	8	6	379.20(11)			E2	
			6	1156.47(13)			E2	0.006(1)
	2360	10	8	438.69(14)			E2	
			8	409.6(10)			E2	

Continued on next page

TABLE II –continued from previous table

Nucleus/band name	E_i	I_i	I_f	$E_\gamma(keV)$	DCO	A_p	Assign	$(BE2)_{out}/(BE2)_{in}$
$\gamma(\text{even})$	2846	12	10	599.20(10)			M1/E2	
			8	1131.01(10)			E2	0.005(1)
			10	485.79(14)			E2	
			10	1084.99(10)			E2	
			14	526.23(14)			E2	
			16	594.07(12)			E2	
	3372	14	14	1033.84(22)			E2	
			2	728.9(10)			M1/E2	
			0	854.21(15)			E2	
			2	274.1(10)			E2	
			4	739.13(5)			M1/E2	
			2	1003.09(9)			E2	0.005(3)
	1499	6	4	370.66(8)			E2	
			6	734.26(5)			M1/E2	
			6	451.18(10)			E2	
	1950	8	7	209.6(20)			M1/E2	
			6	408.3(12)			E2	
			8	721.36(10)			M1/E2	
			6	1185.44(14)			E2	0.003(1)
			8	486.27(7)			E2	
			9	194.5(19)			M1/E2	
	2437	10	8	515.3(25)			E2	
			10	675.82(11)			M1/E2	
			8	1207.63(14)			E2	0.003(1)
			10	561.52(8)			E2	
			10	1237.45(9)			E2	0.001(1)
			12	568.21(9)			E2	
$\gamma(\text{odd})$	987	3	4	597.77(5)			M1/E2	
			2	861.73(11)			M1/E2	
			3	329.21(9)			E2	
	1316	5	4	187.41(39)			M1/E2	
			6	511.50(11)			M1/E2	
			4	926.99(5)			M1/E2	0.034(3)
			5	424.36(4)			E2	
	1741	7	6	241.6(10)			M1/E2	
			8	511.50(11)			M1/E2	
			6	975.66(5)			M1/E2	0.019(1)
			7	501.35(5)			E2	
	2242	9	8	291.72(25)			M1/E2	
			8	1013.09(6)			M1/E2	0.013(1)
			12	459.96(20)			M1/E2	
			9	557.91(6)			E2	
	2800	11	10	1039.19(10)			M1/E2	0.010(1)
			11	562.92(9)			E2	
			12	1022.9(4)			M1/E2	

B. $N = 90$ isotones

The spectroscopic information (such as level energies and branching ratios) of the first excited 0^+ and 2^+ bands in ^{152}Sm are taken from [25, 46].

The level schemes of ^{154}Gd and ^{156}Dy have been reported in our recent work, published in Ref. [42, 47, 48] and [43], respectively. Here, we report for the first time the DCO and polarization observables from some of these measurements, as well as $B(E2)$ ratios, see Table II. A detailed paper on ^{158}Er has been completed and results will be published elsewhere [49].

A partial level scheme of low-lying positive-parity bands obtained in ^{160}Yb is shown in Fig. 3. It is worth noting that a study of the negative-parity levels in ^{160}Yb from the same experiment has been published in Ref. [50]. Spectra demonstrating the positive-parity bands in ^{160}Yb are shown in Figs. 2 b-d. The 0^+ state shown dashed on the level scheme at 1086 keV, is the 0_2^+ level identified in β -decay [51, 52]. The levels at 821 and 1113 keV, assigned as $I^\pi = 2^+$ and 3^+ in the β -decay work, are placed in the 2_γ^+ band which is now extended to spin $I = 11\hbar$. Support for a positive-parity assignment arises from our limits on mixing ratios which have

^{158}Yb (N=88)




FIG. 1. (Color online) Partial level scheme showing the ground band and the even spin member of the 2^+ band in ^{158}Yb . Levels and transitions marked with asterisk (*) symbols and colored in red have been established in this work.

been extracted for some of the transitions depopulating the levels in the 2^+ band to the ground state band. In particular, those for the 577 and 870 keV transitions are consistent with the pure $E2$ assignment favoured in the angular correlation analysis of Garrett *et al.* [51]. However, the level at 1256 keV in the 2^+ band was assigned spin and parity of $I^\pi = 3^-$ by Auer *et al.* [52], based on measured conversion coefficients for the 617 and 435 keV transitions. This assignment is rejected here because our DCO ratio for the 435 keV transition, 1.02(9) is consistent with a stretched $E2$ transition, and if a $I^\pi = 3^-$

assignment were adopted, M2 multipolarity would be required for the 318 and 366 keV transitions linking the signature partners of the 2^+ band. Also visible in Fig. 2 b are the 566 and 478 keV transitions, which link the 0^+ band to 2^+ band. Apparently, these transitions and the 426 keV transition are the result of mixing due to the near degeneracy of the levels at 2790 and 2841 keV. On this basis the 2790 keV level of the 0^+ band is assigned as a $I^\pi = 10^+$ state, which in turn fixes the spins and parities of all the members of the 0^+ band, which is observed down to the 2^+ state. Members of this band are




FIG. 2. (Color online) Spectra showing transitions confirming the placements of the newly built structures in (a) ^{158}Yb and (b-d) ^{160}Yb . Photo peak energies marked with asterisk (*) symbols and colored in red represent new transitions established in this work, previously known γ -rays are highlighted in blue and unmarked. Plus symbols (+) are used for identifying contaminants from other reaction channels and/or other bands, not associated with the cascade of interest.

also visible in Fig. 2d, which shows the spectrum produced by gating on the 1319 keV line. Due to the loss of intensity through $I \rightarrow I-2$ and $I \rightarrow I$ transitions to the ground band, the 0^+ bandhead could not be observed. The assignment of the 1086 keV level to the bandhead is justified by the smooth continuation of states when it is

plotted as a member of the band in Fig. 6b, which shows the energies of the levels of the bands of ^{160}Yb , minus a rigid rotor reference, as a function of spin.


^{160}Yb (N=90)

FIG. 3. (Color online) Partial level scheme of ^{160}Yb showing the ground band, 0_2^+ and 2_γ^+ bands. Levels and transitions marked with asterisk (*) symbols and colored in red have been established in this work.

C. $N = 92$ isotones

The experimental levels and transition rates for ^{154}Sm and ^{156}Gd are extracted from Ref. [22, 53] and [54], respectively.

Low-lying positive-parity states in ^{158}Dy , deduced in this work, are presented in Fig. 4. According to the literature [55], the states built on the 0_2^+ and even spin sequence of the 2_γ^+ bands were known up to be spins $I = 6$ and 8, with the 6^+ member of the 0_2^+ band placed at 1547 keV. However, this study positions the 6^+ level at 1554 keV. In addition, five new in-band transitions (including the γ -ray depopulating the newly revised level) have been added to this structure. The 6^+ and 8^+ members of the even spin 2_γ^+ band are listed in the data sheets at 1486 and 1893 keV [55]. This study has revised the positioning of both levels in the level scheme; they are now placed at 1476 and 1863 keV, respectively. The re-arrangement of these levels is supported by the newly established inter-band transitions connecting both the odd and even spin members of the 2_γ^+ band. Therefore, when taking into account the re-arrangements made in this study, the latter has been extended by five rotational levels from $I = 4$ to $I = 14$. In addition, the odd spin sequence of the 2_γ^+ band has also been extended by four in-band transitions from $I = 7$ to $I = 15$. Fig. 5, shows a spectrum with inter-band transitions decaying out of the newly established rotational levels of 0_2^+ and 2_γ^+ bands, to the ground band of ^{158}Dy . The DCO and polarization mea-

surements could not be determined for the level scheme of ^{158}Dy . The ordering of the transitions observed is solely based on the spin and parity selection rule, the interacting behavior between levels (from different bands) that lie energetically close together as well as the systematics of the neighboring nuclei.

The low lying positive-parity states in ^{160}Er , deduced from this work, confirm all the placements that were reported in the in-beam works of [56, 57], which studied the spectroscopy of this nucleus previously. In this study we report for the first time, the ratios of transition rates extracted for the decays out of the 0_2^+ and 2_γ^+ bands, which are listed in Table II. The $^{150}\text{Sm}(^{16}\text{O}, 4n)^{162}\text{Yb}$ reaction, was recently performed at iThemba LABS, using the AFRODITE array. Though partial results of ^{162}Yb are shown in this work, the complete spectroscopy pertaining to the low lying positive-parity bands of this nucleus will be published elsewhere [58].

IV. ENERGY SYSTEMATICS

The energy systematics of the lowest-lying positive-parity bands are summarized, in Fig. 6, by plotting the energies of the levels of the ground, 0_2^+ and 2_γ^+ bands, less a rigid-rotor reference. At higher spins, ($I > 8\hbar$), the moments-of-inertia are altered by interactions with higher-lying bands. To ascertain the level of mixing due to crossings with high-lying quasiparticle bands such as

^{158}Dy (N=92)




FIG. 4. (Color online) Partial level scheme of ^{158}Dy showing the low-lying positive-parity bands, namely the ground, 0^+_2 and 2^+_1 bands. Levels and transitions marked with asterisk (*) symbols and colored in red have been established in this work.

the S-band (attributed to the alignment of $i_{13/2}$ neutrons) [59], the ground bands (blue squares) have been plotted together with those of the S-band (black open diamonds). The maximum strength of the interaction V at the first crossing of the ground band with the S-band is, in a two-band mixing approximation, given by half of the closest separation of the states of the observed bands. This is evidently limited to $E_x < 250$ keV in all cases, with the possible exception of the lighter $N = 92$ isotones, where the yrare states have not been confirmed and hence the interactions may be stronger. Using the band mixing equations listed in [60] and solving unperturbed states, the 250 keV limit implies a negligible perturbation of less than 10 keV at spin $8\hbar$ due to the crossing with the S-band.

In general, the slopes of the ground bands decrease with increasing N , indicating an increasing moment-of-inertia, and by implication, an increasing deformation.

The opposite is true with increasing proton number, Z , the slope increases, implying a decreasing moment-of-inertia and a decreasing deformation.

In Fig. 7, the moment-of-inertia is examined quantitatively in the region below spin 10, which is relatively free of mixing with the S-band. The quantity $2\mathfrak{I}/\hbar^2$ has been calculated as a function of spin, using $2\mathfrak{I}/\hbar^2 = (4I + 2)/E_\gamma$, where E_γ is the transition energy between

states of spin $I+1$ and $I-1$. In agreement with the above discussion, $2\mathfrak{I}/\hbar^2$ increases with N and decreases with Z . In Fig. 7, it can also be seen that the moments-of-inertia are not constant, but increase with spin, which has been discussed in terms of Coriolis Anti-Pairing [59]. The next striking feature of the data is observed in the comparison of the 2^+_1 bands with the ground-state bands. As a general rule, the odd-spin member of the 2^+_1 bands track the ground bands as a function of spin, indicating simi-




FIG. 5. (Color online) Summed spectra showing peaks associated with some of the low-lying positive parity bands in ^{158}Dy , namely the (a) even and (b) odd γ bands. Photo peak energies colored in red and marked with asterisk (*) symbols represent new transitions established in this work. The in-band members of the ground band are highlighted in blue and unmarked. Transition belonging to 0_2^+ and 2_7^+ bands, known from previous studies are highlighted in black and marked with a hash symbol (#). Transitions marked with a dollar sign (\$) are amongst the transitions that were included in the gating that produced the coincident spectra. Plus symbols (+) are used for identifying contaminants from other reaction channels and/or other bands, not associated with the cascades of interest.

lar moments-of-inertia. The 2_7^+ even-spin band members also has a similar moment-of-inertia to that of the ground bands, with the main exception being at $N = 88$, where the best agreement is in ^{154}Dy and ^{152}Gd . By contrast, the 0_2^+ bands often have moments-of-inertia greater than their ground bands and the 2_7^+ bands (see Fig. 7). As a result, crossings are observed between the 2_7^+ and 0_2^+ bands, particularly in Yb and Er isotopes, where the 0_2^+ bands have the right bandhead energy to cause crossings with the even members of the 2_7^+ bands (Fig. 6).

Our assignment of the new rotational structure in ^{158}Yb , shown in Fig. 1, to the even spin 2_7^+ band can now be understood in the context of the systematics presented in Fig. 6. Firstly, the heads of all 2_7^+ bands have energies just below 1 MeV, consistent with the newly established structure, whereas the band-head energy of the 0_2^+ band appears to increase gradually in excitation energy, (with either N or Z), such that its band-head energy can be expected to be above 1 MeV in ^{158}Yb . This is not compatible with an extrapolated 0_2^+ energy close to 500 keV and therefore leaves the band in question as the ideal candidate for the 2_7^+ band in ^{158}Yb . Finally, the moment-of-inertia for this band is similar to those of well established 2_7^+ bands observed in the neighboring Yb isotopes.

V. ANALYSIS OF THE LOW-LYING SPECTROSCOPY USING A FIVE-DIMENSIONAL COLLECTIVE HAMILTONIAN BASED ON COVARIANT DENSITY FUNCTIONAL THEORY

The observed energy and moment-of-inertia systematics are the starting point for our theoretical analysis of the bands. The covariant density functional theory has been used as input to a five-dimensional collective Hamiltonian. The model is compared to the energies and the in-band to out-of-band branching-ratios determined in our measurements. In addition, comparisons are also made for absolute and relative transition $E0$ and $E2$ strengths using experimental quantities obtained from the literature.

Here we present a brief introduction to the five-dimensional collective Hamiltonian based on the covariant density functional theory (5DCH-CDFT), which could simultaneously treat the quadrupole vibrational and rotational excitations with the collective parameters self-consistently determined by the microscopic CDFT calculations [35]. The collective Hamiltonian is expressed in terms of the two deformation parameters β and γ , and three Euler angles $(\phi, \theta, \psi) \equiv \Omega$ that define the orientation of the intrinsic principal axes in the laboratory frame,

$$\hat{H}_{\text{coll}}(\beta, \gamma) = \hat{T}_{\text{vib}}(\beta, \gamma) + \hat{T}_{\text{rot}}(\beta, \gamma, \Omega) + V_{\text{coll}}(\beta, \gamma). \quad (3)$$




FIG. 6. (Color online) Energies, less a rigid-rotor reference, of ground, 0_2^+ , γ , and S-bands for $N = 88$ to 92 .




FIG. 7. (Color online) Moment-of-inertia parameter, $1/A = 2\mathfrak{I}/\hbar^2$ as a function of spin, deduced from the experimental data.

The three terms in $\hat{H}_{\text{coll}}(\beta, \gamma)$ are the vibrational kinetic energy

$$\begin{aligned}
 \hat{T}_{\text{vib}} = & -\frac{\hbar^2}{2\sqrt{wr}} \left\{ \frac{1}{\beta^4} \left[\frac{\partial}{\partial \beta} \sqrt{\frac{r}{w}} \beta^4 B_{\gamma\gamma} \frac{\partial}{\partial \beta} - \frac{\partial}{\partial \beta} \sqrt{\frac{r}{w}} \beta^3 B_{\beta\gamma} \frac{\partial}{\partial \gamma} \right] \right. \\
 & + \frac{1}{\beta \sin 3\gamma} \left[-\frac{\partial}{\partial \gamma} \sqrt{\frac{r}{w}} \sin 3\gamma B_{\beta\gamma} \frac{\partial}{\partial \beta} \right. \\
 & \left. \left. + \frac{1}{\beta} \frac{\partial}{\partial \gamma} \sqrt{\frac{r}{w}} \sin 3\gamma B_{\beta\beta} \frac{\partial}{\partial \gamma} \right] \right\}, \quad (4)
 \end{aligned}$$

the rotational kinetic energy

$$\hat{T}_{\text{rot}} = \frac{1}{2} \sum_{k=1}^3 \frac{\hat{J}_k^2}{\mathcal{I}_k}, \quad (5)$$

and the collective potential V_{coll} , respectively. Here, $\hat{J}_k$ denote the components of the total angular momentum in the body-fixed frame and all the collective parameters, including the mass parameters $B_{\beta\beta}$, $B_{\beta\gamma}$, $B_{\gamma\gamma}$, the moments-of-inertia $\mathcal{I}_k$ and the collective potential V_{coll} , depend on the quadrupole deformation variables β and γ . Two additional quantities that appear in the $\hat{T}_{\text{vib}}$ term, $r = B_1 B_2 B_3$ and $w = B_{\beta\beta} B_{\gamma\gamma} - B_{\beta\gamma}^2$, determine the volume element in the collective space.

In the 5DCH-CDFT [35], which has provided successful descriptions for low-lying nuclear structure along with isotopic and isotonic chains in a variety of mass regions [61–63], the collective parameters of 5DCH are all determined from the triaxial CDFT calculations. In the present investigation, the moments-of-inertia are calculated with the Inglis-Belyaev formula [64, 65] and the mass parameters with the cranking approximation [66]. The collective potential V_{coll} is obtained by subtracting the zero-point energy corrections [66] from the total energy of the constrained triaxial CDFT.

The eigenvalue problem of the Hamiltonian (3) is solved using an expansion of eigenfunctions in terms of a complete set of basis functions that depend on the five collective coordinates β, γ and Ω [35]

$$\Psi_{\alpha}^{IM}(\beta, \gamma, \Omega) = \sum_{K \in \Delta I} \psi_{\alpha K}^I(\beta, \gamma) \Phi_{MK}^I(\Omega). \quad (6)$$

Then the various observables can be calculated with the obtained collective wave functions, for example the $E2$ transition probabilities

$$B(E2; \alpha I \rightarrow \alpha' I') = \frac{1}{2I+1} |\langle \alpha' I' | \hat{M}(E2) | \alpha I \rangle|^2, \quad (7)$$

where $\hat{M}(E2)$ is the electric quadrupole operator.

The analysis of low-lying states in this mass region starts by performing constrained self-consistent relativistic mean-field plus BCS (RMF+BCS) calculations for triaxial shapes (i.e., including both β and γ deformations). The energy density functional PC-PK1 [67] determines the effective interaction in the particle-hole channel, and a finite-range force that is separable in momentum space is used in the particle-particle channel [68]. The resulting self-consistent solutions (i.e. single-particle wave functions, occupation probabilities, and quasiparticle energies that correspond to each point on the binding energy surface) are used to calculate the parameters that determine the five-dimensional collective Hamiltonian: three mass parameters, three moments-of-inertia, and collective potential, as functions of the deformations β and γ . The diagonalization of the Hamiltonian yields the excitation energies and collective wave functions that are used to calculate observables. Detailed formalism of the 5DCH

based on covariant density functional theory can be found in Refs. [35, 36]. The model has been extensively applied to describe the nuclear collective properties, such as the phase transitions, shape coexistence, as well as the low-lying spectra along the isotopic and isotonic chains in different mass regions [37, 62, 69–71]. It should be emphasized that the model includes the full quadrupole deformed configurations, namely the whole $\beta - \gamma$ plane is adopted and for each deformed configuration, the whole single-particle space is included. Therefore, *the model is parameter-free* and accommodates transitional and shape-coexisting nuclei.

In the following, the theoretical calculations for the $N = 88$, $N = 90$ and $N = 92$ isotones are presented, including the potential energy surfaces (PESs) and the probability density distributions in the $\beta - \gamma$ plane, to obtain a better understanding of the calculated energy spectra and transition probabilities, and the comparison with the experimental data.

A. Potential energy surfaces

Fig. 8 displays the potential energy surfaces (PESs) of $N = 88$, $N = 90$, and $N = 92$ isotones in $\beta - \gamma$ plane calculated by constrained triaxial RMF+BCS with PC-PK1 density functional [67]. It can be seen that all the nuclei exhibit prolate-like shape in their ground states but show more or less different softness/rigidity in the β and γ directions. One can summarize the behavior as follows:

1. For the $N = 88$ isotones, the global minima are all located at $\beta \approx 0.2$ and $\gamma \approx 0^\circ$. As the proton number increases, the PES around the global minimum becomes more rigid in the β direction and a little bit softer in the γ direction.
2. For the $N = 90$ isotones, with increase in proton number, the global minima move to smaller prolate deformation, with $\beta \approx 0.3$ for ^{152}Sm and $\beta \approx 0.25$ for ^{160}Yb . The PES around the global minimum is rather soft in the β -direction and stiff in γ direction in the $N = 90$ isotones, ^{152}Sm , ^{154}Gd , and ^{156}Dy . This is consistent with the findings of Ref. [36], where both the microscopic calculations and the data show that there is an abrupt change of structure at $N = 90$, close to the proposed $X(5)$ critical point [72, 73].
3. For the $N = 92$ isotones, well-deformed prolate global minima are found in ^{154}Sm , ^{156}Gd , and ^{158}Dy . As the proton number increases, the PES around the global minimum becomes softer in both β and γ directions.
4. A secondary local minimum develops in the Er and Yb isotopes at $\beta \approx 0.45$ and $\gamma \approx 10^\circ$. These minima have a shape that corresponds to that of the




FIG. 8. (Color online) Potential energy surfaces (PESs) of $N = 88$, $N = 90$, and $N = 92$ isotones in the β - γ plane. Minima are marked with red symbols, circles and triangles represent the global and secondary minima, respectively. The energy spacing in the contour lines is 0.25 MeV.

well-known triaxial bands observed in nearby Lu isotopes, such as $^{163-165}\text{Lu}$ [74, 75]. It becomes very close in energy to the global minimum with the increase of neutron number, and therefore the effect of shape coexistence could be very important in ^{162}Yb . The counterpart of the Yb minimum is present in the PES of the Er isotopes, although it is less well developed.

In total, for each isotopic chain, from $N = 88$ to 92, the quadrupole deformation β of the global minimum increases and the PES around it becomes a little bit more rigid in the γ direction. For each isotonic chain, from Sm to Yb, the PES around the global minimum becomes softer in the γ direction, while in the β direction it is largely influenced by the development of a secondary local minimum.

B. Energy spectra

Based on the PESs and inertial parameters, one can construct a 5DCH Hamiltonian. Diagonalization of the Hamiltonian yields the excitation energies and collective wave functions that are used to calculate the observables such as $\langle r_c^2 \rangle^{1/2}$, $B(E0)$, $B(E2)$ and so on. For each nucleus, the collective band structure is constructed according to the calculated $E2$ transition probabilities between different states. Figs. 9–11 display the comparison of the calculated energy spectra to the available experimental data for the $N = 88$, $N = 90$, and $N = 92$ isotones, respectively. Here, the comparisons have been performed for the band structures built on the ground state 0_1^+ , the second 0_2^+ , third 0_3^+ states and the γ bands. Normally in the classical picture of the collective model, the band built on the 0_2^+ state is considered to be the β vibrational band, and the 0_3^+ state, to be the second β band. The bandhead of the γ band is taken to be the second excited 2_γ^+ state or the third excited 2_3^+ state, depending on which has the larger $K = 2$ component in the 5DCH-CDFT calculations.

In Figs. 9–11 the theoretical energy levels, minus a rigid rotor reference $7.7I(I+1)$ keV are plotted for the $N = 88$, 90 and 92 isotones and compared to the corresponding experimental quantities. The most apparent observation is that the theoretical bands rise faster in energy than the experimental ones. This discrepancy is consistent with what is observed between the theoretical moments-of-inertia, plotted in Fig. 12 and their experimental counterpart in Fig. 7. These deviations may come from the assumption of adiabatic approximation of the collective Hamiltonian, where the collective parameters are calculated in the vicinity of zero collective angular momenta, such that the moments-of-inertia are calculated in the vicinity of zero rotational frequency [35, 76].

It is worth noting that the transitional nuclei like the $N = 88$, 90 and 92 isotones discussed here, whose soft potentials are susceptible to the increasing angular momentum, this approximation becomes less satisfying [63, 77].

As shown in Ref. [63], one possibility to improve the agreement between theoretical results and experimental data is by introducing an empirical *ab* formula for the moments-of-inertia [78] which is responsible for the fourth order effect of collective momentum in the collective Hamiltonian. Nevertheless, despite the minor discrepancies mentioned above, the overall agreement in the trends between the theoretical energy spectra and the available data, is satisfactory.

The main features of the theoretical energy spectra are summarized as follows:

1. Ground state band: The trends of moments-of-inertia in both N and Z are consistent with the variation of the quadrupole deformation of the global minima in the PESs in Fig. 8. The larger deformations generally correspond to larger moments-of-inertia and result in a more compact ground-state band, rising slower in energy with spin. The general trends are in agreement with experiment.
2. γ -band: The theoretical moments-of-inertia of the γ -band mimic the behaviour of the experimental bands - their moments-of-inertia are very close to those of the ground bands. In Fig. 13 the theoretical and experimental bandhead energies of the γ -band are plotted and compared as a function of the atomic number. In both theory and experiment, the bandhead of the γ band shifts towards lower energy as Z increases, consistent with the observation that the PES around the global minimum becomes softer in the γ direction from Sm to Yb for each isotonic chain.
3. 0_2^+ band: Like the data, the calculated moments-of-inertia of the 0_2^+ band are often higher than those of the calculated ground-state band. The agreement for this feature between theory and experiment is moderate along $N = 88$ (see Figs. 7 and 12), best along $N = 90$, and moderate along the $N = 92$ isotonic chain. The experimental and calculated energies of the 0_2^+ bandheads are plotted as a function of the proton number in Fig. 14. For each isotope a minimum in experimental bandhead energy is found between $Z = 64$ and $Z = 68$. The calculated minima agree for $N = 88$ and $N = 90$, but for $N = 92$, the minimum is too soft, leading to the energy of the head of the 0_2^+ band being underestimated by over 400 keV in ^{162}Yb . Nevertheless, the calculated bandhead energies and moments-of-inertia are accurate enough to reproduce most of the band crossings observed in the data. At $N = 88$ (Fig. 9) the β -band in ^{158}Yb is predicted to lie above the γ -band and to cross it only at high spins. Experiment is consistent with this picture, the 0_2^+ band is not being observed, probably because it is too high in energy. In ^{156}Er , in both theory and experiment, the β -band comes lower in energy and crosses the γ -band. In ^{154}Dy and lighter iso-




FIG. 9. (Color online) The experimental energy spectra of the bands built on the ground state 0_1^+ , the second and third 0^+ states, and the γ bands for the $N = 88$ isotones (left panels), in comparison with the theoretical results (right panels). For the γ bands, the odd and even-spin sequences are shown with different symbols.




FIG. 10. (Color online) The experimental energy spectra of the bands built on the ground state 0_1^+ , the second and third 0^+ states, and the γ bands for the $N = 90$ isotones (left panels), in comparison with the theoretical results (right panels). For the γ bands, the odd and even-spin sequences are shown with different symbols.




FIG. 11. (Color online) The experimental energy spectra of the bands built on the ground state 0_1^+ , the second and third 0^+ states, and the γ bands for the $N = 92$ isotones (left panels), in comparison with the theoretical results (right panels). For the γ bands, the odd and even-spin sequences are shown with different symbols.

tones the calculations correctly predict that the β -band lies lower than the γ -band; consequently, no crossings are observed. At $N = 90$ (Fig. 10) in agreement with the data, crossings are predicted in ^{160}Yb and ^{158}Er , and none in the lighter isotones, due to the β -band lying lower than the γ -band. At $N = 92$ (Fig. 11) experimental crossings are observed in all but ^{154}Sm . The calculations predict that the γ - and β -bands should be close in energy as observed experimentally, but in the calculations, the moments-of-inertia of the two bands are too similar to reliably predict nuclei in which crossings occur. It is in Er and Yb isotopes that the theoretical moments-of-inertia of the 0_2^+ bands are sufficiently different to cause crossings, see Fig. 12, while in Sm to Dy isotopes, the moments-of-inertia of the 0_2^+ bands are very similar to those of the ground bands. As we shall see in the following sections, this points to a structural change in the predicted nature of the 0_2^+ bands in Er and Yb isotopes. In these isotopes, the 0_2^+ bands correspond to the superdeformed triaxial minima in Fig. 8, while in Sm to Dy, the 0_2^+ bands align more closely to a vibrational excitation.

4. 0_3^+ band: In the Sm, Gd and Yb isotopes, the calculated bandhead of the 0_3^+ band is nearly twice that of the 0_2^+ band, as would be expected for a two-phonon β -vibration. The 5DCH-CDFT calculations cannot reproduce the very low experimental excitation energy of the 0_3^+ state in some nuclei, especially in ^{154}Sm and ^{156}Gd . This may suggest that the 0_3^+ states observed in these nuclei are not collective quadrupole excitation states, but based on other excitation modes, such as pairing isomers [20, 28, 43], which have not been taken into account in the present 5DCH-CDFT calculations. For Er and Yb isotopes the calculated excitation energy of the 0_3^+ band is much less than twice of that of the 0_2^+ band. According to the PESs in Fig. 8, the triaxial local minimum plays an important role in the 0_2^+ and 0_3^+ states in these nuclei. In subsection E of section V, the 0_2^+ band is identified with superdeformed minimum while the 0_3^+ band has mixed, vibrational character.

C. Energy staggering in the γ band

It is interesting to further investigate the odd-even energy staggering in the γ bands, to probe the γ deformation. The energy staggering is defined as:

$$S(I) = \frac{[E(I) - E(I-1)] - [E(I-1) - E(I-2)]}{E(2_1^+)} \quad (8)$$

Within the γ band, $S(I)$ has been suggested as an important measure to distinguish soft or rigid triaxiality. For

a γ -rigid rotor, the γ band should exhibit pairs of levels close in energy, $(2^+, 3^+)$, $(4^+, 5^+)$, $(6^+, 7^+)$, $\dots$ (even spins lower in energy) [79], while for a γ -soft collective structure a grouping of levels 2^+ , $(3^+, 4^+)$, $(5^+, 6^+)$, $\dots$ (odd spins lower) should be observed [80]. Thus in both cases $S(I)$ shows an odd-even staggering with the increase of spin, and $S(4) > 0$ in the former case while $S(4) < 0$ in the latter case. In addition, for an ideal axially symmetric rotor, this staggering parameter is a constant $S(I) = 0.33$; while for a harmonic vibrator, it exhibits a staggering behavior with an absolute value equal to 1.0 and $S(4) = -1$.

Fig. 15 displays the experimental and theoretical energy staggering parameter $S(I)$ as function of spin for the γ bands in the $N = 88$, $N = 90$, and $N = 92$ isotones. The calculations can reproduce the experimental data not only for the staggering behavior but also for the variations in the absolute amplitude in the $N = 88$ and $N = 90$ isotones. For the $N = 92$ isotones, although the agreement between the calculated and experimental $S(I)$ is not as impressive as the $N = 88$ and $N = 90$ isotones, the amplitudes of $S(I)$ are close to each other.

For the $N = 88$ isotones, the $S(I)$ exhibits an obvious odd-even staggering with $S(4) \approx -1$, indicating a prominent vibrational character of the γ bands. This is consistent with their potential energy surfaces with slightly quadrupole deformed minimum and γ -soft character. The amplitude of $S(I)$ decreases with spin I for ^{152}Gd and ^{154}Dy but increases for ^{156}Er and ^{158}Yb which may reflect the delicate interplay with the second local minimum.

For the $N = 90$ and 92 isotones, with two or four more neutrons being added to $N = 88$ isotones, the odd-even staggering of $S(I)$ becomes very weak with a negative or positive $S(4)$ tending to 0, indicating large rotational compositions inside the γ bands. This is consistent with their potential energy surfaces having relatively large deformed minimum and γ -stiff character. Further experimental and theoretical investigations may be needed for the small difference in the comparisons of $S(I)$ for the $N = 92$ isotones ^{154}Sm , ^{158}Dy and ^{160}Er .

D. Transition ratios and probabilities

1. In-band $B(E2)$

The experimental in-band $B(E2; I \rightarrow I-2)$ values for the yrast and 0_2^+ bands in the $N = 88$, $N = 90$, and $N = 92$ isotones, taken from the literature [46, 53–55, 81–86] are summarized in Fig. 16. The calculations reproduce the increasing trend of the in-band $B(E2; I \rightarrow I-2)$ rates with spin for all the nuclei in the low spin region, and also the evolution of $B(E2; 2 \rightarrow 0)$ with neutron number for all the isotopic chains. A decrease in the experimental $B(E2; I \rightarrow I-2)$ values in the high spin region can be seen in several nuclei, which is due to




FIG. 12. (Color online) Calculated values of $1/A = 2\Im/\hbar^2$ for the ground band (blue), 0_2^+ band (green) and 2_γ^+ band (brown). Solid lines have even spin.




FIG. 13. (Color online) Bandhead energies of the 2_2^+ bands as a function of Z .




FIG. 14. (Color online) Bandhead energies of the 0_2^+ bands as a function of Z .

the influence of the band crossings and goes beyond the scope of present 5DCH-CDFT calculations. It should be emphasized that the calculation is completely parameter free, an important advantage of using collective models based on self-consistent mean-field single-particle solu-

tions. Here physical observables, such as transition probabilities and spectroscopic quadrupole moments, are calculated in the full configuration space and there is no need for effective charges. Considering this the systematic agreement between the theoretical $B(E2; I \rightarrow I - 2)$




FIG. 15. (Color online) The experimental staggering parameter $S(I)$ for the γ bands in the $N = 88$, $N = 90$, and $N = 92$ isotones (solid circles), in comparison with the calculated results by the 5DCH-CDFT (dash-dotted lines).

values and the available data is reasonably good.

There are relatively few experimental data points for the in-band $B(E2; I \rightarrow I - 2)$ values of the 0_2^+ bands. These values are reasonably reproduced by the microscopic calculations, with the exception of the measured values for the Gd isotopes. The 5DCH-CDFT calculations predict much larger in-band $B(E2; I \rightarrow I - 2)$ values for the 0_2^+ bands compared to the ground bands in $^{156,158,160}\text{Er}$, and $^{160,162}\text{Yb}$. This supports the interpretation that the deformations of the two bands are rather different, pointing to possible shape coexistence in these nuclei. To this end, in the future more lifetime measurements for these nuclei would be welcome.

2. $B(E2)$ branching ratios

A systematic comparison between theory and experiment is also made for the branching ratios of both the 0_2^+ and γ bands with respect to the ground state bands, see Figs. 17-20. As can be seen in Figs. 17 and 18 the systematic comparison for the 0_2^+ bands in the $N = 88, 90$ and 92 isotones, shows a very good agreement between the experimental and theoretical branching ratios for the out-of-band to in-band transitions.

A striking feature of the branching ratio shown in Fig. 17 is the rapid increase of the predicted $B(E2; I_{0_2^+} \rightarrow (I - 2)_g)/B(E2; I_{0_2^+} \rightarrow (I - 2)_{0_2^+})$ ratios, around spin $I \approx 10\hbar$, in ^{154}Dy , $^{156,158}\text{Er}$, and ^{160}Yb . It indicates configuration mixing between the ground and 0_2^+ bands, which can also be clearly seen as the band interaction in the calculated energy spectra of these nuclei in Figs. 9 and 10. Such a rapid increase of the $B(E2; I_{0_2^+} \rightarrow (I - 2)_g)/B(E2; I_{0_2^+} \rightarrow (I - 2)_{0_2^+})$ ratio is seen experimentally in ^{160}Er , but in this spin region, ($I > 10\hbar$), mixing with the S-band is likely to also play a role. Evidence of such an increase can also be found experimentally in ^{158}Er and ^{160}Yb , but more experimental data are needed to confirm the observations.

Fig. 19 displays the out-of-band to in-band branching ratios $B(E2; I_\gamma \rightarrow (I - 2)_g)/B(E2; I_\gamma \rightarrow (I - 2)_\gamma)$ for even I and $B(E2; I_\gamma \rightarrow (I - 1)_g)/B(E2; I_\gamma \rightarrow (I - 2)_\gamma)$ for odd I , for the γ bands in the $N = 88, 90$ and 92 isotones. Fig. 20 further displays the branching ratios $B(E2; I_\gamma \rightarrow I_g)/B(E2; I_\gamma \rightarrow (I - 2)_\gamma)$ for these γ bands. The vertical lines in Fig. 19 correspond to a range of possible $B(E2)$ ratios depending on the actual value of the mixing ratio δ associated with the $I_\gamma \rightarrow (I - 1)_g$ transitions. The lowest value of the vertical lines correspond to $\delta = 1$ while the highest correspond to pure $E2$. In general, the agreement between the experimental and theoretical results is impressive. In all cases the ratios appear to be within the same order of magnitude and this suggests *structural similarities amongst these nuclei*. In addition, in both the experimental and theoretical results, there is also a fair amount of staggering between $B(E2; I_\gamma \rightarrow (I - 1)_g)/B(E2; I_\gamma \rightarrow (I - 2)_\gamma)$ for odd I and $B(E2; I_\gamma \rightarrow$

$(I - 2)_g)/B(E2; I_\gamma \rightarrow (I - 2)_\gamma)$ for even I .

One of the most decisive observables that can be used to characterize the first excited $K^\pi = 0_2^+$ bands is $B(E2; I_2 \rightarrow I_g + 2)$. In Table III, the experimental $\frac{B(E2; I_2 \rightarrow I_g + 2)}{B(E2; I_2 \rightarrow I_2 - 2)}$ ratios deduced from this work are compared with those from previous studies. The experimental data are well reproduced by the calculations.

TABLE III. Results of the five-dimensional collective Hamiltonian (5DDCH) for the $\frac{B(E2; I_2 \rightarrow I_g + 2)}{B(E2; I_2 \rightarrow I_2 - 2)}$ ratios are shown below and they are compared with the experimental ratios deduced from the current and previous works, as per reference. The ratios deduced from Nuclear Data Sheets (NDS) for both ^{152}Sm and ^{154}Gd are from Refs. [46] and [53], respectively.

Nucleus	I_i	Current work	NDS[46, 53]	5DCH
^{152}Sm	2 ₂		0.106(10)	0.0948
	4 ₂		0.063(15)	0.0570
^{154}Gd	2 ₂	0.071(21)	0.20(3)	0.1277
	4 ₂	0.098(6)	0.076(7)	0.0845
^{156}Dy	4 ₂	0.025(9)		0.1206

3. Electric monopole transitions

Strong electric monopole transitions are often considered to be indicative of shape coexistence and configuration mixing between different deformations [21]. The absolute transition strength of the $E0$ transitions between $0_2^+ \rightarrow 0_1^+$ states can be defined as

$$\rho(E0; 0_2^+ \rightarrow 0_1^+) = \left| \frac{\langle 0_2^+ | \hat{T}(E0) | 0_1^+ \rangle}{eR^2} \right|, \quad (9)$$

where $R = 1.2A^{1/3}$ fm, $\hat{T}(E0) = \sum_k e_k r_k^2$ is the electric monopole transition operator, with e_k the electric charge of the k_{th} nucleon, r_k is its relative position in the center-of-mass frame. The absolute transition strengths of other $E0$ transitions can be calculated in a similar way.

In Figs. 21 and 22, the calculated transition strengths $\rho^2(E0; I_{0_2^+} \rightarrow I_{gs})$ and the relative transition strengths between $E0$ and $E2$ transitions $X(E0/E2)$ [87] as a functions of the spin I are plotted for the $N = 88, N = 90$, and $N = 92$ isotones, respectively, in comparison with the available data [21, 22, 46, 53, 54, 84, 85, 87, 88]. The value from the 5DCH-CDFT calculations are overall larger than the available data for $\rho^2(E0; I_{0_2^+} \rightarrow I_{gs})$ and $X(E0/E2)$. This suggests that the experimental 0_2^+ levels may involve other configurations, such as two-quasiparticle excitations or pairing isomers [20, 28, 43], which are not included in the present 5DCH-CDFT calculations.

The detailed trend of the calculated transition strength $\rho^2(E0; I_{0_2^+} \rightarrow I_{gs})$ with increase of spin can be qualitatively understood by the collective wave functions in the




FIG. 16. (Color online) In-band $B(E2; I \rightarrow I-2)$ values for the yrast (solid lines) and 0_2^+ bands (dashed lines) in the $N = 88$, $N = 90$, and $N = 92$ isotones, calculated with PC-PK1 (lines), in comparison with the available data, shown as filled circles for the yrast bands and as empty circles for the 0_2^+ bands [46, 53–55, 81–86].

(β, γ) plane of the initial $I_{0_2^+}$ and final I_{gs} states. For example, it is found that due to a relatively soft potential in ^{152}Gd , the centers of the collective wave functions of the $I_{0_2^+}$ and I_{gs} states move to large quadrupole deformation with the increase of spin from 0^+ to 8^+ . On the contrary, due to a relatively rigid potential in ^{154}Gd , the centers

of collective wave functions of the $I_{0_2^+}$ and I_{gs} states are nearly stable in the (β, γ) plane with the increase of spin. As a result, the $E0$ strength increases with the increase of spin from 0^+ to 8^+ in ^{152}Gd while it remains constant with spin in ^{154}Gd .

Another trend visible, at $N = 90$ and $N = 92$, is the increase in $E0$ strength with Z . The $E0$ strength be-




FIG. 17. (Color online) The calculated branching ratios for out-of-band to in-band transitions, $B(E2; I_{0_2^+} \rightarrow (I-2)_g) / B(E2; I_{0_2^+} \rightarrow (I-2)_{0_2^+})$, for the 0_2^+ bands in the $N = 88$, $N = 90$, and $N = 92$ isotones, in comparison with the available data.




FIG. 18. (Color online) The calculated branching ratios for out-of-band to in-band transitions, $B(E2; I_{0_2^+} \rightarrow I_g) / B(E2; I_{0_2^+} \rightarrow (I-2)_{0_2^+})$, for the 0_2^+ bands in the $N = 88$, $N = 90$, and $N = 92$ isotones, in comparison with the available data.




FIG. 19. (Color online) The calculated branching ratios for out-of-band to in-band transitions, $B(E2; I_\gamma \rightarrow (I-2)_g)/B(E2; I_\gamma \rightarrow (I-2)_\gamma)$ for even I and $B(E2; I_\gamma \rightarrow (I-1)_g)/B(E2; I_\gamma \rightarrow (I-2)_\gamma)$ for odd I , for the γ bands in the $N = 88$, $N = 90$, and $N = 92$ isotones, in comparison with the available data. The lowest values of the vertical lines, for $B(E2)$ values of odd spin members of the γ bands correspond to $\delta = 1$ while the highest correspond to a pure $E2$.

comes especially strong in $^{158,160}\text{Er}$ and $^{160,162}\text{Yb}$, where the secondary minimum lies lowest in energy, supporting the picture in which the $I_{0_2^+}$ bands are actually shape-coexisting triaxial bands. To study this possibility further, we study the calculated wavefunctions in the next section.

E. Probability density distribution

To analyze the configuration mixing in the collective excitation states, one can evaluate the probability density distributions in the β - γ plane, which are defined by

$$\rho_{I\alpha}(\beta, \gamma) = \sum_{K \in \Delta I} |\psi_{\alpha K}^I(\beta, \gamma)|^2 \beta^3, \quad (10)$$




FIG. 20. (Color online) The calculated branching ratios for out-of-band to in-band transitions, $B(E2; I_\gamma \rightarrow (I)_g)/B(E2; I_\gamma \rightarrow (I-2)_\gamma)$, for the γ bands in the $N = 88$, $N = 90$, and $N = 92$ isotones, in comparison with the available data.

with the normalization

$$\int_0^\infty \beta d\beta \int_0^{2\pi} \rho_{I\alpha} |\sin 3\gamma| d\gamma = 1. \quad (11)$$

The probability density distributions for the ground states, 0_2^+ states, 0_3^+ states, and bandheads of γ bands for the $N = 88$, $N = 90$ and $N = 92$ isotones are plotted in Figs. 23–26, respectively.

It is clearly seen in Fig. 23 that the probability density distributions of the ground states for the $N = 88$,

$N = 90$, and $N = 92$ isotones all concentrate on the prolate sides with one pronounced peak. The location of the peak for each nucleus is consistent with the global energy minimum of the nucleus shown in Fig. 8. The feature that the density distributions are very concentrated means that the ground states are almost pure with little configuration mixing with other states. Furthermore, the probability density distributions extend along the β and γ directions in accordance with the softness of PESs around the ground states, i.e., the PES around the global




FIG. 21. (Color online) Absolute transition strengths of $E0$ transitions $\rho^2(E0; I_{0_2^+} \rightarrow I_{gs})$ in the $N = 88$, $N = 90$, and $N = 92$ isotones obtained by the 5DCH-CDFT calculations, in comparison with the available data in [21, 22, 46, 53, 54, 84, 85, 87, 88, 90].




FIG. 22. (Color online) The relative transition strength between $E0$ and $E2$ transitions in the $N = 88$, $N = 90$, and $N = 92$ isotones obtained by the 5DCH-CDFT calculations, in comparison with the data in [21, 22, 46, 53, 54, 84, 85, 87–90].




FIG. 23. (Color online) Probability density distribution in the β - γ plane for the ground states of $N = 88$, $N = 90$, $N = 92$ isotones, predicted by PC-PK1 density functional.




FIG. 24. (Color online) Probability density distribution in the β - γ plane for the 0_2^+ states of $N = 88$, $N = 90$, $N = 92$ isotones, predicted by PC-PK1 density functional.




FIG. 25. (Color online) Probability density distribution in the β - γ plane for the 0_3^+ states of $N = 88$, $N = 90$, $N = 92$ isotones, predicted by PC-PK1 density functional.




FIG. 26. (Color online) Probability density distribution in the β - γ plane for the γ -band heads states of $N = 88$, $N = 90$, $N = 92$ isotones, predicted by PC-PK1 density functional.




FIG. 27. (Color online) Probability density distributions in the β - γ plane for the ground state band (0_1^+ , 2_1^+ , 4_1^+ , 6_1^+ , and 8_1^+), γ band (2_2^+ , 4_3^+ , 6_3^+ , and 8_3^+), 0_2^+ (0_2^+ , 2_3^+ , 4_2^+ , 6_2^+ , and 8_2^+) and 0_3^+ (0_3^+ , 2_5^+ , 4_6^+ , 6_6^+ , and 8_5^+) bands in ^{162}Yb , obtained by the 5DCH-CDFT calculations with PC-PK1 density functional.

minimum becomes γ -softer with the increase of the proton number and become more rigid with the increase of the neutron number. Note that the probability density distributions of the ground states for ^{162}Yb and neighboring nuclei are influenced by the local minimum developing in the PES.

In Figs. 24 and 25, the probability density distributions are shown for the 0_2^+ and 0_3^+ states, respectively. For most nuclei that are investigated, such as Sm, Gd and Dy isotopes, the density distributions of the 0_2^+ and 0_3^+ states possess two and three maxima respectively along

the β direction in the β - γ plane. In Bohr's RVM model, the 0_2^+ and 0_3^+ states are considered as the bandheads of the β and the second β bands, which correspond to one-phonon and two-phonon vibrational excitations in the β collective degree of freedom. Therefore, the behavior of the density distributions suggests that for most nuclei investigated, the 0_2^+ and 0_3^+ states originate mainly from the one-phonon and two-phonon β vibrations and could be regarded as the bandheads of the β and the second β bands.

Exceptional cases are found in the Er and Yb iso-

topes. The density distributions of 0_2^+ states in ^{156}Er and $^{158,160}\text{Yb}$ are different from those discussed above. For ^{156}Er and ^{158}Yb , there also exist two maxima; but for ^{156}Er the distribution with larger deformation is much more extended in the β and γ direction, while for ^{158}Yb the maximum with larger deformation no longer concentrates on the prolate side but has considerable triaxial deformation. For ^{160}Yb , there appear three maxima in the density distributions of 0_2^+ state. The density distributions of 0_3^+ states in all the Er and Yb isotopes investigated are somewhat different from the general feature of two-phonon β vibration, where a maximum with considerable triaxial deformation is shown. As discussed above, the PESs of the Er and Yb isotopes are softer than those of the Sm, Gd and Dy isotopes, meanwhile a local triaxial minimum starts to emerge with the increase of the proton number and manifests itself obviously in $^{158,160,162}\text{Yb}$. The exceptional probability density distributions of the Er and Yb isotopes can be therefore understood, as arising from the softer potential and the influence of the local triaxial minimum. In this sense, the 0_2^+ states in ^{156}Er , $^{158,160}\text{Yb}$ and the 0_3^+ states in all the Er and Yb isotopes investigated cannot be simply interpreted as the one-phonon and two-phonon β vibrations, which is also consistent with the fact that the calculated energy differences between 0_3^+ and 0_2^+ states are much smaller than those between 0_3^+ and 0_1^+ states in the isotopes like $^{156,158}\text{Er}$ and $^{158,160}\text{Yb}$.

In Fig. 26, the probability density distributions for the bandhead of γ bands are presented. It is noted that in some nuclei the head of the γ band is the second 2^+ state while in the other nuclei it corresponds to the third 2^+ state. It can be seen in Fig. 26 that generally the probability density distributions for the bandhead of γ bands has a peak with a considerable triaxiality. In the $N = 88$ isotones, the distributions all concentrate around $\beta \approx 0.25$ and $\gamma \approx 25^\circ$. In ^{150}Sm , due to the configuration mixing with the 2_2^+ state, there also exists a little peak around $\beta \approx 0.4, \gamma \approx 0^\circ$. With the increase of proton number, such distribution is not obvious in ^{152}Gd and ^{154}Dy , and disappears in ^{156}Er and ^{158}Yb . In the $N = 90$ isotones, the probability density distributions are quite similar to those in the $N = 88$ isotones. The distributions are all concentrated around $\beta \approx 0.3, \gamma \approx 20^\circ$. With the increase of proton number, the density distributions become more concentrated. In the $N = 92$ isotones, apart from ^{162}Yb , the probability density distributions are all concentrated around $\beta \approx 0.35, \gamma \approx 15^\circ$. This is to say, with the increase of neutron number from $N = 88$ to 92, the probability density distributions for the bandhead of γ bands become closer to prolate side, which reflects the shape transition from the near spherical to well deformed case. In ^{162}Yb , besides the peak around $\beta \approx 0.3, \gamma \approx 20^\circ$, there is another peak near $\beta \approx 0.4, \gamma \approx 0^\circ$. This can be understood from the well-developed shape coexistence in this nucleus where two minima in the PES are very close in energy as shown in Fig. 8.

In Fig. 27, for the shape coexistence nucleus ^{162}Yb , we

further plot the probability density distributions in the β - γ plane for the ground state, γ , 0_2^+ and 0_3^+ bands. These bands are organized according to the calculated electromagnetic transitional probabilities. It can be seen that due to the existence of the second minimum in the PES, the levels in ground state bands have important contributions from the second minimum with a large deformation ($\beta_2 \approx 0.45$), which becomes more important with the increase of spin I (there is a crossing near spin 8). The levels in γ bands also have two important contributions from both the global and the second minima, and the latter contribution becomes more important with the increase of spin I . The probability distributions for the levels in the 0_2^+ band exhibit two maxima along the prolate side, the characteristic of a β vibration, meanwhile the maximum with large deformation is predominant, reflecting the mixing with the second minimum. The probability distributions for the levels in the 0_3^+ band exhibit three maxima, however, two maxima along the prolate side and the other with obvious triaxial degree of freedom. Such distributions are different from the typical distribution of a β vibration with two phonons for a well-deformed nucleus.

VI. SUMMARY AND CONCLUSION

An extensive set of data on the low-lying positive-parity bands in even-even nuclei for $N = 88$, $N = 90$ and $N = 92$ from Sm to Yb has been collected. The following observations can be made:

1. The 0_2^+ bands all have the same or larger moment-of-inertia than the ground-state bands. By contrast, the γ bands, particularly the odd spin members, track the ground band as a function of spin. This appears to be a general feature throughout the $A = 150$ to 160 mass region.
2. The splitting $S(I)$ of the γ bands varies considerably over the nuclides studied.
3. The even-spin 0_2^+ bands cross the γ bands in Er and Yb isotopes.

Overall, the comparison of the data to the results obtained using the 5DCH-CDFT calculations with PC-PK1 density functional yields a satisfactory agreement. The trends in energy and moment-of-inertia of all bands are well-reproduced, including the staggering $S(I)$ of the 2_γ^+ bands. The model does well with electromagnetic transition rates and branching ratios, with the exception of the $E0$ rates which are generally overestimated. Within the model, the following conclusions can be drawn:

1. The potential energy surfaces of the RMF calculations indicate γ -soft nuclei at $N = 88$ becoming γ -rigid along $N = 90$ and $N = 92$. This is in agreement with the staggering $S(I)$ of the levels in the γ bands.

2. The nature of the 0_2^+ band changes with atomic number. In the isotopes of Sm to Dy, they can be understood as β -vibrations, but in the Er and Yb isotopes, the 0_2^+ bands can be understood as having large components in a triaxial superdeformed minimum. They are better understood as shape-coexisting states.
3. Shape coexistence has been suggested to describe the nuclei in the vicinity of ^{152}Sm . The present calculations predict a soft potential in the β -direction but do not find two coexisting minima. This is reminiscent of ^{152}Sm exhibiting an $X(5)$ behaviour.
4. In the Sm, Gd and Dy isotopes, the model predicts that the 0_3^+ bands are of two-phonon nature, having an energy of twice the 0_2^+ band. This is in contradiction with the experimental data and implies that other excitation modes must be invoked

to explain their origin.

ACKNOWLEDGMENTS

We would like to thank the iThemba LABS technical staff for their support in these experiments. This work is supported by the National Research Foundation of South Africa grant under grants (No. 109711, 92791, 106012 and 92792), the Natural Science Foundation of China under Grants (No. 11461141002, No. 11475140 and No. 11375015), the U.S. National Science Foundation under Grant Nos. PHY-1401574 (USNA) and PHY-0754674 (FSU), the National Research, Development and Innovation Fund of Hungary, financed under the K18 funding scheme with Grant no. K128947, as well as by the European Regional Development Fund (Grant No. GINOP-2.3.3-15-2016-00034). JYFL research is supported by the Academy of Finland under the Finnish Centre of Excellence Programme 20062011, Contract No. 213503.

-
- [1] A. Bohr, Dan. Mat. Fys. Medd. **27**, 16 (1952).
 - [2] A. Bohr and B.R. Mottelson, Mat. Fys. Medd., Dan. Vid. Selsk. **27**, 16 (1953).
 - [3] D.R. Bes and R.A. Broglia, Nucl. Phys. **80**, 289 (1966).
 - [4] O. Mikoshiba, Nucl. Phys. A **101**, 202 (1967).
 - [5] A. Kuliev and N.I. Pyatov, Nucl. Phys. A **106**, 689 (1968).
 - [6] S.T. Belyaev and B.A. Rumiantsev, Phys. Lett. B **30**, 444 (1969).
 - [7] J.V. Maher, J.R. Erskine, A.M. Friedman, J.P. Schiffer, and R.H. Siemssen, Phys. Rev. C **5**, 1380 (1972).
 - [8] P. Kleinheinz, R. K. Sheline, M. R. Maier, R. M. Diamond, and F. S. Stephens, Phys. Rev. Lett. **32**, 68 (1974).
 - [9] J. J. Kolata and M. Oothoudt, Phys. Rev. C **15**, 1947 (1977).
 - [10] M. H. Mortensen, R. R. Betts and C. K. Bockelman, Phys. Rev. C **21**, 2275 (1980).
 - [11] S. W. Yates, I. Y. Lee, N. R. Johnson, E. Eichler, L. L. Riedinger, M. W. Guidry, A. C. Kahler, D. Cline, R. S. Simon, P. A. Butler, P. Colombani, F. S. Stephens, R. M. Diamond, R. M. Ronningen, R. D. Hichwa, J. H. Hamilton and E. L. Robinson, Phys. Rev. C **21**, 2366 (1980).
 - [12] D. R. Zolnowski, M. B. Hughes, J. Hunt and T. T. Sugihara, Phys. Rev. C **21**, 2556 (1980).
 - [13] A. Bohr and B. R. Mottelson, Phys. Scripta **25**, 28 (1982).
 - [14] P. E. Garrett, M. Kadi, Min Li, C. A. McGrath, V. Sorokin, Minfang Yeh and S. W. Yates, Phys. Lett. B **400**, 250 (1997).
 - [15] P.E. Garrett, J Phys. G: Nucl. Part. Phys. **27**, R1 (2001).
 - [16] H. F. Wirth, G. Graw and S. Christen, D. Cutoiu, Y. Eisermann, C. Günther, R. Hertenberger, J. Jolie, A. I. Levon, O. Möller, G. Thiamova, P. Thierolf, D. Tonev and N. V. Zamfir, Phys. Rev. C **69**, 044310 (2004).
 - [17] N. Pietralla and O. M. Gorbachenko, Phys. Rev. C **70**, 011304 (2004).
 - [18] D. Bucurescu, G. Graw, R. Hertenberger, H.-F. Wirth, N. Lo Iudice, A. V. Sushkov, N. Yu. Shirikova, Y. Sun, T. Faestermann, R. Krücken, M. Mahgoub, J. Jolie, P. von Brentano, N. Braun, S. Heinze, O. Möller, D. Muecher, C. Scholl, R. F. Casten and D. A. Meyer, Phys. Rev. C **73**, 064309 (2006).
 - [19] E. A. McCutchan, Dennis Bonatsos, N.V. Zamfir, and R. F. Casten, Phys. Rev. C **76**, 024306 (2007).
 - [20] W. D. Kulp, J. L. Wood, P. E. Garrett, C. Y. Wu, D. Cline, J. M. Allmond, D. Bandyopadhyay, D. Dashdorj, S. N. Choudry, A. B. Hayes, H. Hua, M. G. Mynk, M. T. McEllistrem, C. J. McKay, J. N. Orce, R. Teng, and S. W. Yates, Phys. Rev. C, **77** 061301 (2008).
 - [21] K. Heyde and J. L. Wood, Reviews of Modern Physics **83**, 1467 (2011).
 - [22] J. Smallcombe, P.J. Davies, C.J. Bartona, D.G. Jenkins, L.L. Andersson, P.A. Butler, D.M. Cox, R.-D. Herzberg, A. Mistry, E. Parr, T. Grahn, P.T. Greenlees, K. Hauschild, A. Herzan, U. Jakobsson, P. Jones, R. Julin, S. Juutinen, S. Ketelhut, M. Leino, A. Lopez-Martens, P. Nieminen, J. Pakarinen, P. Papadakis, P. Peura, P. Rahkila, S. Rinta-Antila, P. Ruotsalainen, M. Sandzelius, J. Sarén, C. Scholey, J. Sorri, and J. Uusitalo, Phys. Lett. B **732**, 161 (2014).
 - [23] F. -Q. Chen and J. Luis Egido, Phys. Rev. **93**, 064313 (2016).
 - [24] F. -Q. Chen and J. Luis Egido, Phys. Rev. **95**, 024307 (2017).
 - [25] W. D. Kulp, J. L. Wood, P. E. Garrett, J. M. Allmond, D. Cline, A. B. Hayes, H. Hua, K. S. Krane, R.-M. Larimer, J. Loats, E. B. Norman, P. Schmelzenbach, C. J. Stapels, R. Teng, and C. Y. Wu, Phys. Rev. C **71**, 041303 (R) (2005).
 - [26] W. D. Kulp, J. L. Wood, J. M. Allmond, J. Eimer, D. Furse, K. S. Krane, J. Loats, P. Schmelzenbach, C. J. Stapels, R.-M. Larimer, E. B. Norman, and A. Piechaczek, Phys. Rev. C **76**, 034319 (2007).

- [27] I. Ragnarsson and R. A. Broglia, Nucl. Phys. A **263**, 315 (1976).
- [28] J.F. Sharpey-Schafer, TE Madiba, SP Bvumbi, EA Lawrie, JJ Lawrie, A Minkova, SM Mullins, P Papka, DG Roux, and J Timár, Eur. Phys. J. A **47**, 6 (2011).
- [29] S.K. Abdulvagabova, S.P. Ivanova, and N.I. Pyatov, Phys. Lett. **38**, 215 (1972).
- [30] R.E. Griffin, A.D. Jackson, and A.B. Volkov, Phys. Lett. B **36** 281 (1971).
- [31] W.I. Van Rij and S.H. Kahana, Phys. Rev. Lett. **28**, 50 (1972).
- [32] J.F. Sharpey-Schafer, R.A. Bark, S.P. Bvumbi, T.R.S. Dinoko and S.N.T. Majola, Eur. Phys. J. A **55** 15 (2019).
- [33] J. L. Wood, A-M. Oros-Peusquens, R. Zaballa, J. M. Allmond, and W. D. Kulp, Phys. Rev. C **70**, 024308 (2004).
- [34] J. M. Allmond, R. Zaballa, A. M. Oros-Peusquens, W. D. Kulp and J. L. Wood, Phys. Rev. C **78**, 014302 (2008).
- [35] T. Nikšić, Z P Li, Dario Vretenar, L Próchniak, Jie Meng, and Peter Ring, Phys. Rev. C **79**, 034303 (2009).
- [36] Z. P. Li, T. Nikšić, D. Vretenar, J. Meng, G. A. Lalazissis, and P. Ring, Phys. Rev. C **79**, 054301 (2009).
- [37] Z P Li, Tamara Nikšić, Dario Vretenar, and Jie Meng, Phys. Rev. C **80**, 061301 (2009).
- [38] G. Duchene, Nucl. Inst. Meth. **432**, 90 (1999).
- [39] J. F. Sharpey-Schafer, Nucl. Phys. News **14**, 5 (2004).
- [40] S. P. Bvumbi, J. F. Sharpey-Schafer, P. M. Jones, S. M. Mullins, B. M. Nyakó, K. Juhász, R. A. Bark, L. Bianco, D. M. Cullen, D. Curien, P. E. Garrett, P. T. Greenlees, J. Hirvonen, U. Jakobsson, J. Kau, F. Komati, R. Julin, S. Juutinen, S. Ketelhut, A. Korichi, E. A. Lawrie, J. J. Lawrie, M. Leino, T. E. Madiba, S. N. T. Majola, P. Maine, A. Minkova, N. J. Ncapayi, P. Nieminen, P. Peura, P. Rakhila, L. L. Riedinger, P. Ruotsalainen, J. Saren, C. Scholey, J. Sorri, S. Stolze, J. Timár, J. Uusitalo, and P. A. Vymers. Phys. Rev. C **87**, 044333 (2013).
- [41] G. L. Zimba, J. F. Sharpey-Schafer, P. Jones, S. P. Bvumbi, L. P. Masiteng, S. N. T. Majola, T. S. Dinoko, E. A. Lawrie, J. J. Lawrie, D. Negi, P. Papka, D. Roux, O. Shirinda, J. E. Easton, and N. A. Khumalo, Phys. Rev. C **94**, 054303 (2016).
- [42] S.N.T. Majola, DJ Hartley, LL Riedinger, JF Sharpey-Schafer, JM Allmond, C Beausang, MP Carpenter, CJ Chiara, N Cooper, D Curien, *et al.* Phys. Rev. C **91**, 034330 (2015).
- [43] J.F. Sharpey-Schafer, T. E. Madiba, S. P. Bvumbi, E. A. Lawrie, J. J. Lawrie, A. Minkova, S. M. Mullins, P. Papka, D. G. Roux, and J. Timár, Eur. Phys. J. A **47**, 5 (2011).
- [44] R. A. Bark, Z. P. Li, S. N. T. Majola, J. F. Sharpey-Schafer, Z. Shi, S. Q. Zhang, EPJ Web of Conferences **178**, 02012 (2018)
- [45] J.F. Sharpey-Schafer et al., Nuclear Physics A **834**, 45c (2010).
- [46] M.J. Martin, Nuclear Data Sheets **114**, 1497 (2013).
- [47] D. J. Hartley, L. L. Riedinger, R. V. F. Janssens, S. N. T. Majola, M. A. Riley, J. M. Allmond, C. W. Beausang, M. P. Carpenter, C. J. Chiara, N. Cooper, D. Curien, B. J. P. Gall, P. E. Garrett, F. G. Kondev, W. D. Kulp, T. Lauritsen, E. A. McCutchan, D. Miller, S. Miller, J. Piot, N. Redon, J. F. Sharpey-Schafer, J. Simpson, I. Stefanescu, X. Wang, V. Werner, J. L. Wood, C.-H. Yu, S. Zhu, and J. Dudek, Phys. Rev. C **95**, 014321 (2017).
- [48] S. Jehangir, G. H. Bhat, J. A. Sheikh, S. Frauendorf, S. N. T. Majola, P. A. Ganai, and J. F. Sharpey-Schafer, Phys. Rev. C **97**, 014310 (2018).
- [49] T. S. Dinoko *et al.*, Phys. Rev. C, submitted (2019).
- [50] R. A. Bark, J. F. Sharpey-Schafer, S. M. Maliage, T. E. Madiba, F. S. Komati, E. A. Lawrie, J. J. Lawrie, R. Lindsay, P. Maine, S. M. Mullins, S. H. T. Murray, N. J. Ncapayi, T. M. Ramashidza, F. D. Smit, and P. Vymers, Phys. Rev. Lett. **104**, 022501 (2010).
- [51] C. Garrett, S. Rastikerdar, W. Gelletly, and D.D. Warner. Physics Lett. **118B**, 292 (1982).
- [52] H. Auer, J. Fernandez-Niello, H. Puchta, and F. Riess, Z. Phys. A **318**, 323 (1984).
- [53] C. W. Reich, Nuclear Data Sheets **110**, 2257 (2009).
- [54] C.W. Reich, Nuclear Data Sheets **113**, 2537 (2012).
- [55] R.G. Helmer, Nuclear Data Sheets **101**, 325 (2004).
- [56] K. Dusling, N. Pietralla, G. Rainovski, T. Ahn, B. Bochev, A. Costin, T. Koike, T. C. Li, A. Linnemann, S. Pontillo, and C. Vaman, Phys. Rev. C **73**, 014317 (2006).
- [57] J. Ollier, J. Simpson, M.A. Riley, E.S. Paul, X. Wang, A. Aguilar, M.P. Carpenter, I.G. Darby, D.J. Hartley, R.V. F. Janssens, F.G. Kondev, T. Lauritsen, P.J. Nolan, M. Petri, J.M. Rees, S.V. Rigby, C. Teal, J. Thomson, C. Unsworth, S. Zhu, A. Kardan, and I. Ragnarsson, Phys. Rev. C **83**, 044309 (2011).
- [58] L. Mdletshe *et al.*, Eur. Phys. J. A **54** 176 (2018).
- [59] F. S. Stephens and R. S. Simon, Nucl. Phys. A **179**, 753 (1972).
- [60] P.J. Brussaard and P.W. Glaudemans, *Shell Model Applications in Nuclear Spectroscopy* (Elsevier, 1977).
- [61] Z P Li, C Y Li, J Xiang, J M Yao, and J Meng, Phys. Lett. B **717**, 470 (2012).
- [62] Z P Li, Tamara Nikšić, Dario Vretenar, Peter Ring, and Jie Meng, Phys. Rev. C **81**, 064321 (2010).
- [63] Y. Y. Wang, Z. Shi, Q. B. Chen, S. Q. Zhang, and C. Y. Song, Phys. Rev. C **93**, 044309 (2016).
- [64] D. R. Inglis, Phys. Rev. **103**, 1786 (1956).
- [65] S. T. Beliaev, Nucl. Phys. **24**, 322 (1961).
- [66] M. Girod and B. Grammaticos, Nucl. Phys. A **330**, 40 (1979).
- [67] Peng-Wei Zhao, Zhi Pan Li, Jiang-Ming Yao, J Meng, *et al.*, Phys. Rev. C **82**, 054319 (2010).
- [68] J. Xiang, Z. P. Li, Z. X. Li, J. M. Yao, and J. Meng, Nucl. Phys. A **873**, 1 (2012).
- [69] Z P Li, Tamara Nikšić, Dario Vretenar, and Jie Meng, Phys. Rev. C **81** 034316 (2010).
- [70] Z P Li, J M Yao, Dario Vretenar, Tamara Nikšić, H Chen, and Jie Meng, Phys. Rev. C, **84**, 054304 (2011).
- [71] S. Quan, Q. Chen, Z. P. Li, T. Nikšić, and D. Vretenar. Phys. Rev. C, **95** 054321 (2017).
- [72] F. Iachello. Phys. Rev. Lett. **87**, 052502 (2001).
- [73] F. Iachello, Phys. Rev. Lett. **85**, 3580 (2000).
- [74] Ødegård, G.B. Hagemann, M. Bergström, B. Herskind, D.R. Jensen, G. Sletten, J. Wilson, K. Spohr, H. Hübel, A. Bracco, S. Leoni, A. Maj, C. Petrache, P. Bednarczyk, D. Curien, and I. Hamamoto, Nucl. Phys. A **682**, 427 (2001).
- [75] G. Schönwaßer, H. Hübel, G.B. Hagemann, H. Amro, R.M. Clark, M. Cromaz, R.M. Diamond, P. Fallon, B. Herskind, G. Lane, W.C. Ma, A.O. Macchiavelli, S.W. Ødegård, G. Sletten, D. Ward, and J.N. Wilson, Eur. Phys. J A **15**, 435 (2002).
- [76] M. Baranger, Annals of Physics **114**, 123 (1978).
- [77] J J. Sun, Z. Shi, X Q. Li, H. Hua, C. Xu, Q B. Chen, S Q. Zhang, C Y. Song, J. Meng, X G. Wu, S.P. Hu, H.Q.

- Zhang, W.Y. Liang, F.R. Xu, Z.H. Li, G.S. Li, C.Y. He, Y. Zheng, Y.L. Ye, D.X. Jiang, Y.Y. Cheng, C. He, R. Han, Z.H. Li, C.B. Li, H.W. Li, J.L. Wang, J.J. Liu, Y.H. Wu, P.W. Luo, S.H. Yao, B.B. Yu, X.P. Cao, and H.B. Sun, *Phys. Lett. B* **734**, 313 (2014).
- [78] C. S. Wu and J. Y. Zeng, *Commun. Theor. Phys* **8**, 51 (1987).
- [79] A. S. Davydov and G.F. Filippov, *Nucl. Phys.* **8**, 237 (1958).
- [80] L. Wilets and M. Jean, *Phys. Rev.* **102**, 788 (1956).
- [81] O. Möller, A. Dewald, P. Petkov, B. Saha, A. Fitzler, K. Jessen, D. Tonev, T. Klug, S. Heinze, J. Jolie, P. von Brentano, D. Bazzacco, C. A. Ur, E. Farnea, M. Axiotis, S. Lunardi, G. de Angelis, D. R. Napoli, N. Marginean, T. Martinez, M. A. Caprio, and R. F. Casten, *Phys. Rev. C* **74**, 024313 (2006).
- [82] N.R. Johnson, I.Y. Lee, F.K. McGowan, T.T. Sugihara, S.W. Yates, and M.W. Guidry, *Phys. Rev. C* **26**, 1004 (1982).
- [83] M.P. Fewell, N.R. Johnson, F.K. McGowan, J.S. Hattula, I.Y. Lee, C. Baktash, Y. Schutz, J.C. Wells, L.L. Riedinger, M.W. Guidry, and S.C. Pancholi, *Phys. Rev. C* **37**, 101 (1988).
- [84] C.W. Reich, *Nuclear Data Sheets* **105**, 557 (2005).
- [85] C.W. Reich, *Nuclear Data Sheets* **108**, 1807 (2007).
- [86] D. Tonev, A. Dewald, T. Klug, P. Petkov, J. Jolie, A. Fitzler, O. Möller, S. Heinze, P. von Brentano, and R. F. Casten, *Phys. Rev. C* **69**, 034334 (2004).
- [87] T. Kibedi and R.H. Spear, *Atomic Data and Nuclear Data Tables* **89**, 77 (2005).
- [88] J.L. Wood, E.E. Zganjar, C. De Coster, and K. Heyde, *Nucl. Phys. A* **651**, 323 (1999).
- [89] N. Blasi, L. Guerro, A. Saltarelli, O. Wieland, and L. Fortunato, *Phys. Rev. C* **88**, 014318 (2013).
- [90] S.K. Basu and A.A. Sonzogn, *Nuclear Data Sheets* **114**, 435 (2013).
- [91] G. L. Zimba, S. P. Bvumbi, L. P. Masiteng, P. Jones, J. F. Sharpey-Schafer, S. N. T. Majola, T. S. Dinoko, O. Shirinda, J. J. Lawrie, J. E. Easton, N. A. Khumalo, L. Msebi, P.I. Mashita, P. Papka, D. Roux, D. Negi, *Eur. Phys. J. A* **54**, 59 (2018).