



# CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

## $^9\text{Li}(d,p)$ reaction as a specific probe of $^{10}\text{Li}$ , the paradigm of parity-inverted nuclei around the $N=6$ closed shell

F. Barranco, G. Potel, E. Vigezzi, and R. A. Broglia

Phys. Rev. C **101**, 031305 — Published 24 March 2020

DOI: [10.1103/PhysRevC.101.031305](https://doi.org/10.1103/PhysRevC.101.031305)

# ${}^9\text{Li}(d,p)$ reaction as a specific probe of ${}^{10}\text{Li}$ , paradigm of parity–inverted nuclei around $N = 6$ closed shell

F. Barranco,<sup>1</sup> G. Potel,<sup>2</sup> E. Vigezzi,<sup>3</sup> and R. A. Broglia<sup>4,5</sup>

<sup>1</sup>*Departamento de Física Aplicada III, Escuela Superior de Ingenieros, Universidad de Sevilla, Camino de los Descubrimientos, Sevilla, Spain*

<sup>2</sup>*National Superconducting Cyclotron Laboratory, Michigan State University, East Lansing, Michigan 48824, USA*

<sup>3</sup>*INFN Sezione di Milano, Via Celoria 16, I-20133 Milano, Italy*

<sup>4</sup>*The Niels Bohr Institute, University of Copenhagen, DK-2100 Copenhagen, Denmark*

<sup>5</sup>*Dipartimento di Fisica, Università degli Studi Milano, Via Celoria 16, I-20133 Milano, Italy*

(Dated: January 14, 2020)

We show, within the framework of renormalized nuclear field theory and of the induced reaction surrogate formalism, that the highly debated  ${}^{10}\text{Li}$  structure, studied in a recent high statistics  ${}^9\text{Li}(d,p){}^{10}\text{Li}$  one–neutron transfer experiment, is consistent with, or better, requires, the presence of a virtual  $1/2^+$  state of similar single–particle strength than that of the  $1/2^-$  resonance at  $0.45\pm 0.03$  MeV. Based on continuum spectroscopy self-energy techniques, we find that the physical mechanism responsible for parity inversion in  ${}^{10}_3\text{Li}$  is the same as that at the basis of the similar phenomenon observed in  ${}^{11}_4\text{Be}$  and as that needed in  ${}^{11}\text{Li}$  to have an important  $s$ -wave ground state component. In particular the strong dynamical coupling between the  $s_{1/2}$  and the  $d_{5/2}$  states, mediated by the quadrupole vibration of the core  ${}^9\text{Li}$ . A phenomenon which also affects the strength distribution of the  $d_{5/2}$  state, in particular in the energy range 3–4.5 MeV. Furthermore, this mechanism is also consistent with the (normal) sequence of the  $1p_{1/2}$  and  $2s_{1/2}$  levels in the  $N = 7$  isotones  ${}^{12}_5\text{B}$  and  ${}^{13}_6\text{C}$ . The main aim of the present paper is that of treating structure and reactions on equal footing and in a common language. In other words, the calculation of  ${}^9\text{Li}(d,p){}^{10}\text{Li}$  reaction as a single conceptual step, from individual single-particle motion and collective vibrations to absolute double differential cross sections of renormalised virtual and resonant final states, which can be directly compared with experiment.

## Introduction.

Seven decades have elapsed since the seminal papers in which Mayer [1–3] (see also [4]) and Jensen [5] introduced the shell model of the atomic nucleus. Much work on the subject of the magic numbers has been dedicated ever since [6–11]. In spite of this, the quest for these pillars of nuclear structure, is far from completed, being very much an open question reserving surprises and challenges [12–16].

*Novel magic numbers: parity inversion.* The first two Meyer–Jensen magic numbers are 2 and 8 for both protons and neutrons. Increasing the number of neutrons of a normal nucleus, Pauli principle forces them into states of higher momentum. When the core becomes neutron saturated, the nucleus expels most of the wave function of the last neutrons outside to form a halo which, because of its large size, can have a lower momentum, that is less kinetic energy of confinement. The system  ${}^{11}_4\text{Be}_7$  ( $(N - Z)/A \approx 0.36$ ) constitutes a much studied example of one neutron halo nucleus ([17–19] and refs. therein). In principle one could have expected that because the  $1s_{1/2}$  and  $1p_{3/2}$  levels are filled, the last neutron occupies a substate of the  $1p_{1/2}$  orbital. However, the ground state of  ${}^{11}\text{Be}$  has spin and parity  $1/2^+$ , implying inversion in the sequence of the  $1p_{1/2}$  and  $2s_{1/2}$  orbitals. Because the  $1/2^+$  (–0.50 MeV) and  $1/2^-$  (–0.180 MeV) levels are very close to each other, and separated from the  $3/2^-$  level by about 3 MeV the  $N = 8$  role of magic number has been taken over by  $N = 6$ . In other words,  ${}^{11}\text{Be}$  can be viewed as a one–neutron system outside closed

shell, the reaction  ${}^{10}\text{Be}(d,p){}^{11}\text{Be}$  being thus the specific probe of such a system. It is furthermore of notice that closely associated with the parity inversion phenomenon, the dipole transition between the  $1/2^+$  and  $1/2^-$  states carries about one Weisskopf unit, being the strongest  $E1$ -transition between bound states of the whole mass table [20]. A piece of information which can be used at profit in connection with the position of the  $1/2^+$  and  $1/2^-$  states in  ${}^{11}\text{Li}$ .

A substantial set of experimental data [21–42] and theoretical insight [43–60] exists on the unbound isotope of  ${}^{11}_4\text{Be}$ , namely  ${}^{10}_3\text{Li}$  which indicates parity inversion also in this case. This scenario is furthermore consistent with –required by – the bound, two neutron halo system  ${}^{11}_3\text{Li}_8$  ([61] and refs. therein). The presence of a low-lying dipole resonance with  $\approx 6 - 8\%$  of the dipole Energy Weighted Sum Rule (EWSR),  $\approx 0.5$  MeV width and centroid energy  $\lesssim 1$  MeV ([39] and refs. therein), and thus carrying about one Weisskopf unit, implies the presence of a particle–hole dipole excitation with energy not much larger than 0.3–0.5 MeV. Furthermore, the value of the absolute differential two–neutron pickup cross section associated with the reaction  ${}^1\text{H}({}^{11}\text{Li}, {}^9\text{Li}(\text{gs})){}^3\text{H}$  [42] implies that the  $|s_{1/2}^2(0)\rangle$  and  $|p_{1/2}^2(0)\rangle$  configurations enter the  ${}^{11}\text{Li}$  ground state with about the same amplitude (0.45  $|s_{1/2}^2(0)\rangle + 0.55|p_{1/2}^2(0)\rangle$  [55], the  $1p_{3/2}(\pi)$  odd proton being considered as a frozen spectator, is not explicitly written). The above requirement implies that the energies of the two configurations are not too different, likely within 0.5–0.6 MeV.




FIG. 1. (color online) The experimental energies of the low-lying states in the  $N = 7$  isotones  $^{11}\text{Be}$ ,  $^{12}\text{B}$  and  $^{13}\text{C}$  are shown by solid lines. The corresponding theoretical energies are displayed by dashed lines. Also reported are the predictions for  $^{10}\text{Li}$ . States based on  $1/2^-$ ,  $1/2^+$  and  $5/2^+$  neutron configurations are shown by blue, red and green lines respectively.

The generally accepted picture was recently set in doubt as a result of a one-nucleon transfer experiment [62] which led to the conclusion that "... the level sequence in the  $^{10}\text{Li}$  system may not show the shell inversion features observed in other  $N=7$  isotopes such as  $^{11}\text{Be}$ ". The specificity of the  $^9\text{Li}(d, p)$  reaction to provide insight into the level sequence of a nuclear system combined with the high statistics of the experiment, implied that the above conclusions constituted a serious question mark on the validity of the entire picture of shell evolution leading to the  $N=6$  magic number and associated parity inversion in exotic halo nuclei at the neutron drip line. And thus of crucial importance to find the missing link. This is the challenge we take up in the present paper. And to do so we use theoretical tools in which structure and reactions and thus bound states and continuum dynamics become unified.

*Scope and outcome.* In this letter we will show that the data of Cavallaro *et al.* [62] are consistent with the presence of  $1/2^+$  strength at threshold - and thus with the findings of [21]-[42] - of similar magnitude as that of the  $1/2^-$  resonance observed at  $0.45 \pm 0.03$  MeV. But to observe it one has to look into a rather different angular region than that used in the experiment ( $5.5^\circ$ - $16.5^\circ$ ) in which the  $s$ -strength ( $2^-, 1^-$ ) has a minimum, while the  $p$  one ( $1^+, 2^+$ ) displays a maximum. Namely conduct the search for angles  $\theta_{cm} > 40^\circ$  and likely centred at backward angles, implying a different region of momentum transfer. Detailed predictions of the full angular distribution and of the absolute energy differential cross sections associated with the  $s_{1/2}$ ,  $p_{1/2}$ ,  $p_{3/2}$  and  $d_{5/2}$  virtual and resonant states are given. The theoretical framework

used provides, at the same time, a unified account of the experimental findings regarding the  $N = 7$  isotones  $^{13}\text{C}$ ,  $^{12}\text{B}$ ,  $^{11}\text{Be}$  and  $^{10}\text{Li}$ , and closes the issue concerning the missing  $s$ -strength at threshold in  $^{10}\text{Li}$ .

Below, the methods used and the results obtained in the description of the different virtual and resonant states are discussed, following the study reported in the ArXiv manuscript [63], extended to higher excitation energies. and to include  $5/2^+$  scattering states. Within this context we note also [64, 65].

*Methods and results.* In the calculation of  $^{11}\text{Be}$  [19], the four parameters characterizing the bare mean field  $U(r)$  (to be used in connection with an effective mass  $m_k(r)$  ( $m_k = 0.7m$  ( $0.91m$ ) for  $r = 0$  ( $r = \infty$ )), were determined imposing the self-consistent condition that the dressed single-particle levels resulting from the coupling to the quadrupole vibration of the core  $^{10}\text{Be}$  reproduce the experimental energies, in particular those of the parity inverted  $1/2^+$  and  $1/2^-$  states, but also equally well that of the  $5/2^+$  resonance. We have extended this approach to the normal sequence of  $N = 7$  isotones  $^{12}\text{B}$  and  $^{13}\text{C}$  (see Supplemental Material at ... for more details), obtaining again an accurate account of the corresponding experimental spectra (see Fig. 1). The effect of the neutron-proton interaction, leading to the observed doublet splitting in  $^{12}\text{B}$ , is discussed below. With this global potential and  $k$ -mass, together with the quadrupole vibration of  $^9\text{Li}$  ( $\hbar\omega_2 = 3.37$  MeV,  $\beta_2 = 0.72$ ), we have calculated the self-energy matrix  $\Sigma_{ik}^a(E)$  for the single particle levels  $i, k$  ( $50 \text{ MeV} > \epsilon_i, \epsilon_k > \epsilon_F$ ) of  $^{10}\text{Li}$  in a box of radius 60 fm, carrying the quantum numbers  $a = \{lj\}$ . Renormalization processes associated with the coupling to the quadrupole vibrations of the core were considered<sup>1</sup>, in a similar manner to our previous work on  $^{11}\text{Be}$ . The dressed  $\tilde{j}^\pi$  neutron states were determined by diagonalizing  $\widetilde{\Sigma}_{ik}^a(E)$ .

$\widetilde{1/2^+}$  and  $\widetilde{1/2^-}$  waves

In the following we will discuss the results associated with the  $1/2^+$  and the  $1/2^-$  waves, which essentially de-

<sup>1</sup> Within this context, it has been stated (see e.g. [66]) that exotic (halo) nuclei being much less bound than nuclei lying along the stability valley, offer a unique framework to study mean field properties without the complications of polarization from valence particles (separability issue). In particular one could argue that valence neutrons of  $^{210}\text{Pb}$ , can exert a much larger polarisation of the core  $^{208}\text{Pb}$ , than those of  $^{11}\text{Li}$  regarding the  $^9\text{Li}$  core, let alone in the case of  $^{209}\text{Pb}$  as compared with the extreme case of  $^{10}\text{Li}$ . Inverting the issue one could e.g., posit that the dressing of the valence neutron in  $^{10}\text{Li}$  due to the quadrupole vibration of the core is poorer than that of the valence neutron of  $^{209}\text{Pb}$  due to a similar mechanism. Now, the total zero point amplitude associated with the lowest mode of  $^{208}\text{Pb}$  is  $\beta_2 \approx 0.06$  (see e.g. [67] Eqs. (6-52) and (6-390)) while that of  $^9\text{Li}$  is  $\beta_2 \approx 0.66$ , namely one order of magnitude larger. Because the dressing (self-energy) of single-particle states is proportional to  $\beta_2^2$ , the assumption of separability, essentially not applicable in the case of  $^{209}\text{Pb}$ , can hardly be used in treating  $^{10}\text{Li}$  or, for that sake, in treating any of the  $N = 7$  isotopes considered in the present paper (see also [68])

termine the spectrum up to about 2 MeV. The  $1/2^+$  scattering length is  $\alpha = -\lim_{k \rightarrow 0} tg(\delta_{1/2^+})/k = -8$  fm, corresponding to the energy [69, 70]  $\epsilon_{1/2^+} = \frac{\hbar^2 \kappa^2}{2m} = 0.32$  MeV, where  $\kappa = 1/\alpha$ . The eigenfunction of a state lying close to  $\epsilon_{1/2^+}$  and thus representative of this virtual state is,

$$|\widetilde{1/2^+}\rangle = \sqrt{0.98}|s_{1/2}\rangle + \sqrt{0.02}|(d_{5/2} \otimes 2^+)_{1/2^+}\rangle. \quad (1)$$

Similarly, the resonant  $1/2^-$  state can be written as

$$|\widetilde{1/2^-}\rangle = \sqrt{0.94}|p_{1/2}\rangle + \sqrt{0.07}|((p_{1/2}, p_{3/2}^{-1})_{2^+} \otimes 2^+)_{0^+}, p_{1/2}; 1/2^-\rangle. \quad (2)$$

The peak and the width of the resonance are  $\epsilon_{1/2^-} = 0.50$  MeV, and  $\Gamma_{1/2^-} = 2 \left( \frac{d\delta_{1/2^-}}{dE} \Big|_{\epsilon_{1/2^-}} \right)^{-1} = 0.35$  MeV respectively. The parallel between these results and those shown in Eqs (1)–(3) of ref. [19] for  $^{11}\text{Be}$ , let alone with those displayed in Fig. 1 of ref. [61] and in Eqs (1)–(4) of ref. [55] is apparent. The dressed neutron couples to the  $1p_{3/2}^{-1}(\pi)$  proton hole leading to the doublets  $(1^-, 2^-)$  ( $\widetilde{1/2^+} \otimes p_{3/2}^{-1}(\pi)$ ) and  $(1^+, 2^+)$  ( $\widetilde{1/2^-} \otimes p_{3/2}^{-1}(\pi)$ ), both in  $^{10}\text{Li}$  and  $^{12}\text{B}$ . In keeping with a well established approach [71–73], we assume that the proton interacts with the odd neutron through a spin-dependent contact interaction,  $V_{np} = F_0 \delta(\mathbf{r}_1 - \mathbf{r}_2) (1 - \alpha(1 - \boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2))$ . The two parameters  $F_0$  and  $\alpha$  were determined by fitting the experimental splittings in  $^{12}\text{Be}$  (see Fig. 1) and optimising the agreement with the transfer cross sections measured in  $^{10}\text{Li}$  as discussed below see Supplemental Material at ... for more details). The scattering lengths of the resulting  $2^-, 1^-$  states are equal to -19 fm and -5 fm respectively ( $\epsilon_{2^-} \approx 0.05$  MeV,  $\epsilon_{1^-} \approx 0.8$  MeV). For the positive parity doublet one finds instead ( $\epsilon_{1^+} \approx 0.3$  MeV,  $\epsilon_{2^+} \approx 0.6$  MeV) (Fig. 1).

The phase shifts associated with these states in comparison with those corresponding to the bare neutron states are shown in Figs. 2(a) and 2(b). From these phase shifts, one can derive the spectral functions,  $\bar{\rho}_{j\pi}(\omega) = -\frac{(2j+1)}{\pi} \text{Im}[G_{j\pi}(\omega + i0^+) - G_{0,j\pi}(\omega + i0^+)] = \frac{(2j+1)}{\pi} \frac{d\delta_{j\pi}}{d\omega}$  where  $G$  and  $G_0$  denote the Green's functions of the renormalized and of the bare particles respectively [74–76] (see also [77], p.226). The differences between the corresponding spectral functions associated with the  $2^-, 1^-$  and  $2^+, 1^+$  states are shown in Figs. 2(c) and 2(d). The slopes of phase shifts associated with the  $2^-, 1^-$  states change sign in going from bare to renormalized states, resulting in a conspicuous modification of the strength functions (Fig. 2(c)). The positive parity states, which are bound in the bare potential, are moved into the continuum (notice the corresponding change of the value of the phase shift, from  $\pi$  to 0 for  $E_x \approx 0$ ) and acquire a resonant character (Fig. 2(b) and 2(d); the values of  $\delta_{1^+}$  and  $\delta_{2^+}$  being equal to  $\pi/2$  for  $E_x \approx 0.3$  and

0.7 MeV). Summing up, the importance of many-body effects is apparent.




FIG. 2. Upper panels: Phase shifts associated with the renormalized and the unperturbed single-particle states of negative (a) and positive (b) parity in  $^{10}\text{Li}$ . Lower panels: differences between the spectral functions of the renormalized low-lying negative (c) and positive (d) parity states and the spectral functions of the corresponding unperturbed single-particle states.

Based on the non-local self-energy matrices  $\Sigma^a(r, r'; E)$ , whose configuration space representation corresponds to  $\Sigma_{ik}^a(E)$ , in conjunction with the optical parameters of ref. [78], we have calculated the absolute double differential cross section  $d^2\sigma/dE d\Omega$  within the framework of induced-surrogate reaction formalism ([79, 80] and refs. therein). The associated absolute single differential cross sections  $(d\sigma/dE)_{5.5^\circ-16.5^\circ}$  and  $(d\sigma/d\Omega)_{0.2-1 \text{ MeV}}$  were obtained by integrating  $d^2\sigma/dE d\Omega$  in the angular and energy ranges within which data was recorded [62]. In keeping with the experimental energy resolution, the theoretical results were folded with Lorentzian functions of FWHM of 170 keV. As seen from Figs. 3 (a) and (b), theory provides a quantitative account of the experimental findings, confirming the (apparent) lack of any relevant contribution associated with  $s_{1/2}$  strength.

The picture changes radically when looking at a quite different angular region, this time centered around more backward angles ( $\theta_{CM} \geq 40^\circ$ ), and thus corresponding to larger momentum transfer, as testified by the absolute cross sections  $(d\sigma/dE)$  integrated in the angular range  $50^\circ - 180^\circ$  (Fig. 3 (c)) as well as by the absolute differential cross section at angles  $\theta_{CM} > 40^\circ$  (Fig. 3 (d)). These results unarguably demonstrate the presence of a virtual  $s_{1/2}$  state, which appears absent (non observable) from  $(d\sigma/dE)_{5.5^\circ-16.5^\circ}$ . To make such a definite statement it is required that one is able to predict absolute one-particle transfer cross sections within experimental errors. To fulfil such requirements one has to be able to calculate continuum self-energy processes. That is, the




FIG. 3. (a) Theoretical prediction (continuous thick solid curve) of the  $^{10}\text{Li}$  strength function for the  $d(^9\text{Li},p)^{10}\text{Li}$  reaction at 100 MeV incident energy and  $\theta_{CM} = [5.5^\circ, 16.5^\circ]$  in comparison with the experimental data (solid dots with errors) [62]. The contributions associated with states of different angular momenta are also shown; (b) Corresponding angular distributions associated with the states in the energy interval 0.2–1 MeV in comparison with the experimental data; (c) Predicted strength function integrated in the angular range  $\theta_{CM} = [50^\circ, 180^\circ]$ , compared to the result obtained neglecting the contributions from the s-wave ( $1^-$  and  $2^-$  states); (d) Predicted angular distributions integrated in the energy interval 0–0.2 MeV.




FIG. 4. (Color online) (a) Theoretical prediction (continuous solid curve) of the  $^{10}\text{Li}$  strength function for the  $d(^9\text{Li},p)^{10}\text{Li}$  reaction at 21.4 MeV incident energy and  $\theta_{CM} = [98^\circ, 134^\circ]$  in comparison with the experimental data. (solid dots with errors) [26]. The data expressed as counts/MeV in [26] have been converted into mb/MeV using the appropriate acceptance function (A.M. Moro, private communication). (b) Corresponding angular distributions associated with the states in the energy interval 0–1 MeV in comparison with the experimental data.

dressing of a particle state through the coupling of the quadrupole vibration of the core  $^9\text{Li}$ , renormalising in the process energies, single-particle spectroscopic amplitudes and wavefunctions (form factors) of virtual and resonant states.

Similar calculations to the ones discussed above were carried out but this time for the reaction  $^9\text{Li}(d,p)$  investigated at 2.36 MeV/A at the REX-ISOLDE facility (see




FIG. 5. Theoretical prediction (continuous solid curve) of the  $^{10}\text{Li}$  strength function for the  $d(^9\text{Li},p)^{10}\text{Li}$  reaction at 100 MeV incident energy and  $\theta_{CM} = [5.5^\circ, 16.5^\circ]$  in comparison with the experimental data (solid dots with errors) [62] (see also Fig. 3(a)).

[26] where the presence of the  $1s_{1/2}$  is apparent), making use of the optical potentials of this reference (see also [81]). The outcome of such calculations provide an overall quantitative description of the experimental findings (Fig. 4). Summing up, the results shown in Figs. 3 and 4 dissipate the possible doubts concerning the presence of a virtual  $s_{1/2}$  state in the low-energy continuum spectrum of  $^{10}\text{Li}$ , and confirm the soundness of the picture at the basis of the description of  $^{11}\text{Li}$  provided in [61] and [55] (see also [42]). Within this scenario, Fig. 3(c) and (d) constitute absolute strength function and differential cross sections predictions with an estimated error of 10%.

#### $\widetilde{5/2^+}$ and $\widetilde{3/2^-}$ waves

Theory predicts the existence of a resonant, many-body  $\widetilde{5/2^+}$  state with centroid at  $\approx 3.5$  MeV which splits into four states  $(\widetilde{5/2^+} \otimes 1p_{3/2}(\pi))_{1^-, 2^-, 3^-, 4^-}$  spanning the energy interval 2–6 MeV. In the measured energy interval, the main contribution originates from the  $4^-$  state. The calculation gives an overall account of the strength function, as shown in Fig. 5. The  $\widetilde{5/2^+}$  resonance state has a pronounced many-body character, similarly the virtual  $1/2^+$  and resonant  $1/2^-$  states discussed above. In particular, configurations including two quadrupole phonons and associated anharmonic effects play an important role in the renormalization of all three states but especially in the  $\widetilde{5/2^+}$  case, and have been calculated as outlined in (see [19], see also Supplemental Material at ... for more details; see also [82]). These anharmonicities have been found to be more important in the present case in connection with the  $\widetilde{5/2^+}$ , and in turn also for the  $\widetilde{1/2^+}$  and  $\widetilde{1/2^-}$  states, than in the case of  $^{11}\text{Be}$ . This is because the energy of the  $\widetilde{5/2^+}$  resonance in  $^{10}\text{Li}$  lies closer to

the  $\widetilde{1/2^+} \otimes 2^+$  configuration than in  $^{11}\text{Be}$ . Theory also predicts the presence of a  $3/2^-$  component, which splits into four states  $(\widetilde{3/2^-} \otimes 1p_{3/2}(\pi))_{0^+,1^+,2^+,3^+}$  with energies within the range 3–6 MeV. This component however only produces a small and smooth background, included in Figs. 3 and 5. Another  $3/2^-$  contribution, not included in our calculation, is expected at  $\approx 5.4$  MeV, based on the state  $|(p_{3/2}^{-1} \otimes 0_a^+)_{3/2^-}\rangle$  where  $|0_a^+\rangle = |gs(^{11}\text{Li})\rangle$  is the pair addition mode of the core  $^9\text{Li}$ , that is the ground state of  $^{11}\text{Li}$ . We estimate the coupling between the  $3/2^-$  resonances to occur mainly through the  $(p_{1/2} \otimes 2^+)_{3/2^-}$  configuration, and to be weak. Within this context we recall a similar situation, this time for bound states, concerning the two  $3/2^+$  states found in connection with the study of the septuplet of states  $|h_{9/2} \otimes 3^-(^{208}\text{Pb}); I\rangle$  ( $I = 3/2^+, 5/2^+, \dots, 15/2^+$ ) of  $^{209}\text{Bi}$ , the second  $3/2^+$  being connected with the  $2p-1h$  state  $|d_{3/2}^{-1} \otimes gs(^{210}\text{Po}); 3/2^+\rangle$  (see [83] and references therein). In this case the mixing between the two states is much larger due mainly to the fact that the unperturbed energies of the two  $3/2^+$  states are almost degenerate.

*Conclusions.* Structure and reactions, in particular when referred to a specific elementary mode of excitation (e.g. single-particle motion) and its specific probe (one-nucleon transfer), are two aspects of the same physics. Renormalized energies and wavefunctions (effective  $Q$ -values, spectroscopic amplitudes and formfactors) are the “observables”, the meeting point between theory and experiment being absolute differential cross sections.

For normal nuclei, structure essentially refers to bound states, reactions to continuum asymptotic waves. A distinction which becomes blurred in the case of exotic light bound halo nuclei like  $^{11}\text{Be}$  and  $^{11}\text{Li}$ . Just think in the  $5/2^+$  resonance in the first case (centroid  $E_x = 1.28$  MeV, width  $\Gamma = 100$  keV) and in the soft-dipole mode in the second ( $E_x \lesssim 1$  MeV,  $\Gamma = 0.5$  MeV), let alone on the virtual and resonant states in the case of  $^{10}\text{Li}$  and of its specific probe,  $^9\text{Li}(d,p)^{10}\text{Li}$ . Making use of renormalized nuclear field theory of structure and reactions we find it similarly possible (trying) to provide a “complete” description of the structure and reaction process associated with  $^{11}\text{Be}$  and  $^{11}\text{Li}$  than with  $^{10}\text{Li}$ , in which case one is referring exclusively to continuum spectroscopy (structure) and reactions. The parameters used to calculate the bare single-particle levels of  $^{10}\text{Li}$  were obtained by extrapolating those determined following the protocol presented in [19] in connection with the calculation of the  $^{11}\text{Be}$  spec-

trum applied also to the isotones  $^{13}_6\text{C}_7$  and  $^{12}_5\text{B}_7$  which display Mayer–Jensen sequence. The apparent puzzle–(hieroglyphic–) like position (see [62] and refs [7,8,10,11] therein, i.e. [30, 32–34] of the present paper) of the continuous structure and reaction aspects of  $^{10}\text{Li}$  within the  $N = 6$ , parity inverted scenario becomes readily understandable as the consequence of the choice of a restricted angular window associated with low linear momentum transfer.

Because of the strong mixing of the  $s_{1/2}$  and  $d_{5/2}$  virtual and resonant states through the quadrupole vibrations of the core  $^9\text{Li}$ , either one reproduces both the  $d\sigma/dE(1/2^+)$  and  $d\sigma/dE(5/2^+)$  or likely none of them. To be remembered furthermore is the fact that these couplings renormalize single-particle content, energies and widths (spectral functions), and equally importantly for what concerns  $d\sigma/dE$ , the radial dependence of the wave functions (form factors), all elements which have to be calculated selfconsistently. The role played by such requirements is likely emphasised in the case of continuous spectroscopy, in which structure and reactions are just two aspects of the same physics. From here the requirement to treat them both on a common basis and in a unified fashion.

Within the unified view adopted in the present paper,  $^{10}\text{Li}$ ,  $^{11}\text{Be}$  and  $^{11}\text{Li}$  can be viewed as the top, middle and bottom texts of a rosetta-like stone dealing with parity inverted halo nuclei poised to acquire a permanent dipole moment. The uniqueness of the apparently different phenomena (“texts”) is due to the fact that they all emerge from the same underlying physics, namely: a) a quantal phase transition [84] close to the crossing point (parity inversion); b) spontaneous symmetry breaking phenomenon (dipole instability) [85], and of their interplay. Another example of the relation existing between physical correct collective variables, emergent properties, transferability and effective lower dimensionality of many-body systems ([86–88] and refs. therein).

Discussions with H. Lenske, A. Moro and H. T. Fortune are gratefully acknowledged. F.B and E.V. acknowledge funding from the European Union Horizon 2020 research and innovation program under Grant Agreement No. 654002. F.B. thanks the Spanish Ministerio de Economía y Competitividad and FEDER funds under project FIS2017-88410-P. This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics, under the FRIB Theory Alliance award DE-SC0013617.

---

[1] Maria G. Mayer. On closed shells in nuclei. *Phys. Rev.*, 74:235, 1948.  
 [2] Maria Goeppert Mayer. On closed shells in nuclei. ii. *Phys. Rev.*, 75:1969, 1949.  
 [3] Maria G. Mayer and Edward Teller. On the origin of elements. *Phys. Rev.*, 76:1226, 1949.  
 [4] W. Elsasser. Sur le principe de Pauli dans les noyaux. *J.*

*Phys. Radium*, 4:549, 1933.  
 [5] Otto Haxel, J. Hans D. Jensen, and Hans E. Suess. On the “Magic Numbers” in nuclear structure. *Phys. Rev.*, 75:1766, 1949.  
 [6] M. Mayer and J. Jensen. *Elementary Theory of Nuclear Shell Structure*. Wiley, New York, NY, 1955.  
 [7] A. De Shalit and I. Talmi. *Nuclear Shell Theory*. Aca-

- dem Press, New York, 1963.
- [8] A. Bohr and B. R. Mottelson. *Nuclear Structure, Vol.I*. Benjamin, New York, 1969.
- [9] G. F. Bertsch. *The practitioner's Shell Model*. North Holland, Amsterdam, 1972.
- [10] P. Navrátil, S. Quaglioni, I. Stetcu, and B. Barrett. Recent developments in no-core shell-model calculations. *Journal Phys. G*, 36:083101, 2009.
- [11] J. P. Vary, P. Maris, H. Potter, and M. Caprio. Ab Initio No Core Shell Model—Recent Results and Further Prospects. *arxiv:0904.0463v1[nucl-th]*, 2015.
- [12] O. Sorlin and M.-G. Porquet. Nuclear magic numbers: new features far from stability. *Prog. Part. Nucl. Phys.*, 61:602, 2008.
- [13] K. Heyde and J.L. Wood. Shape coexistence in atomic nuclei. *Rev. Mod. Phys.*, 83:1467, 2011.
- [14] R. Kruecken. Introduction to shell structure in exotic nuclei. *Cont. Phys.*, 52:101, 2012.
- [15] Hagen G., M. Hjorth-Jensen, G.R. Jansen, R. Machleidt, and T. Papenbrock. Evolution of shell structure in neutron-rich calcium isotopes. *Phys. Rev. Lett.*, 109:032502, 2012.
- [16] T. Otsuka and Y. Tsunoda. The role of shell evolution in shape coexistence. *J. Phys. G*, 43:024009, 2016.
- [17] J.S. Winfield, S. Fortier, W.N. Catford, S. Pita, N.A. Orr, J. Van de Wiele, Y. Blumenfeld, R. Chapman, S.P.G. Chappell, N.M. Clarke, N. Curtis, M. Freer, S. Gals, H. Langevin-Joliot, H. Laurent, I. Lhenry, J.M. Maison, P. Roussel-Chomaz, M. Shawcross, K. Spohr, T. Suomijrvi, and A. de Vismes. Single-neutron transfer from  $^{11}\text{Be}_{gs}$  via the  $(p, d)$  reaction with a radioactive beam. *Nucl. Phys. A*, 683:48, 2001.
- [18] Angelo Calci, Petr Navrátil, Robert Roth, Jérémy Dohet-Eraly, Sofia Quaglioni, and Guillaume Hupin. Can ab initio theory explain the phenomenon of parity inversion in  $^{11}\text{Be}$ ? *Phys. Rev. Lett.*, 117:242501, 2016.
- [19] F. Barranco, G. Potel, R. A. Broglia, and E. Vigezzi. Structure and reactions of  $^{11}\text{Be}$ : many-body basis for single-neutron halo. *Phys. Rev. Lett.*, 119:082501, 2017.
- [20] E. Kwan, C.Y. Wu, N.C. Summers, G. Hackman, T.E. Drake, C. Andreoiu, R. Ashley, G.C. Ball, P.C. Bender, A.J. Boston, H.C. Boston, A. Chester, A. Close, D. Cline, D.S. Cross, R. Dunlop, A. Finlay, A.B. Garnsworthy, A.B. Hayes, A.T. Laffoley, T. Nano, P. Navrátil, C.J. Pearson, J. Pore, S. Quaglioni, C.E. Svensson, K. Starosta, I.J. Thompson, P. Voss, S.J. Williams, and Z.M. Wang. Precision measurement of the electromagnetic dipole strengths in  $^{11}\text{Be}$ . *Phys. Lett. B*, 732:210, 2014.
- [21] K.H. Wilcox, R.B. Weisenmiller, G.J. Wozniak, N.A. Jelley, D. Ashery, and J. Cerny. First measurement of  $^{10}\text{Li}$  from  $^9\text{Be}(^9\text{Be}, ^8\text{B})^{10}\text{Li}$ . *Phys. Lett. B*, 59:142, 1975.
- [22] B.M. Young, W. Benenson, J.H. Kelley, N.A. Orr, R. Pfaff, B.M. Sherrill, M. Steiner, M. Thoennessen, J.S. Winfield, J.A. Winger, S.J. Yennello, and A. Zeller. Low-lying structure of  $^{10}\text{Li}$  in the reaction  $^{11}\text{B}(^7\text{Li}, ^8\text{B})^{10}\text{Li}$ . *Phys. Rev. C*, 49:279, 1994.
- [23] H.G. Bohlen, W. von Oertzen, T. Stolla, R. Kalpakchieva, B. Gebauer, M. Wilpert, T. Wilpert, A.N. Ostrowski, S.M. Grimes, and T.N. Massey. Study of Weakly Bound and Unbound States of Exotic Nuclei with Binary Reactions. *Nucl. Phys. A*, 616:254c, 1997.
- [24] J.A. Caggiano, D. Bazin, W. Benenson, B. Davids, B.M. Sherrill, M. Steiner, J. Yurkon, A.F. Zeller, and B. Blank. Spectroscopy of the  $^{10}\text{Li}$  nucleus. *Phys. Rev. C*, 60:064322, 1999.
- [25] P. Santi, J.J. Kolata, V. Guimaraes, D. Peterson, R. White-Stevens, E. Rischette, D. Bazin, B.M. Sherrill, A. Navin, DeYoung P.A., Jolivet P.L., G.F. Peaslee, and R.T. Guray. Structure of the  $^{10}\text{Li}$  nucleus investigated via the  $^9\text{Li}(d, p)^{10}\text{Li}$  reaction. *Phys. Rev. C*, 67:024606, 2003.
- [26] H.B. Jeppesen, A.M. Moro, U.C. Bergmann, M.J.G. Borge, J. Cederkäll, L.M. Fraile, H.O.U. Fynbo, J. Gómez-Camacho, H.T. Johansson, B. Jonson, M. Meister, T. Nilsson, G. Nyman, M. Pantea, K. Risager, A. Richter, G. Schrieder, T. Sieber, O. Tengblad, E. Tengborn, M. Turrión, and F. Wenander. Study of  $^{10}\text{Li}$  via the  $^9\text{Li}(^2\text{H}, p)$  reaction at REX-ISOLDE. *Phys. Lett. B*, 642:449, 2006.
- [27] J.K. Smith, T. Baumann, J. Brown, P.A. DeYoung, N. Frank, J. Hinnefeld, Z. Kohley, B. Luther, B. Marks, A. Spyrou, S.L. Stephenson, M. Thoennessen, and S.J. Williams. Spectroscopy of the  $^{10}\text{Li}$  nucleus. *Nucl. Phys. A*, 940:235, 2015.
- [28] A. Sanetullaev, R. Kanungo, J. Tanaka, M. Alcorta, C. Andreoiu, P. Bender, A.A. Chen, G. Christian, B. Davids, J. Fallis, J.P. Fortin, N. Galinski, A.T. Galant, P.E. Garrett, G. Hackman, B. Hadinia, S. Ishimoto, M. Keefe, R. Krcken, J. Lighthall, E. McNeice, D. Miller, J. Purcell, J.S. Randhawa, T. Roger, A. Rojas, H. Savajols, A. Shotter, I. Tanihata, I.J. Thompson, C. Unsworth, P. Voss, and Z. Wang. Investigation of the role of  $^{10}\text{Li}$  resonances in the halo structure of  $^{11}\text{Li}$  through the  $^{11}\text{Li}(p, d)^{10}\text{Li}$  transfer reaction. *Physics Letters B*, 755:481, 2016.
- [29] R.A. Kryger, A. Azhari, A. Galonsky, J.H. Kelley, R. Pfaff, E. Ramakrishnan, D. Sackett, B.M. Sherrill, M. Thoennessen, J.A. Winger, and S. Yokoyama.  $^9\text{Li}$  and n detected in coincidence from a stripping of  $^{18}\text{O}$  beam at 80MeV/n on a C target, obtaining relative velocity spectra. *Phys. Rev. C*, 47:R2439, 1993.
- [30] M. Zinser, F. Humbert, T. Nilsson, W. Schwab, Th. Blaich, M. J. G. Borge, L. V. Chulkov, H. Eickhoff, Th. W. Elze, H. Emling, B. Franzke, H. Freiesleben, H. Geissel, K. Grimm, D. Guillemaud-Mueller, P. G. Hansen, R. Holzmann, H. Irnich, B. Jonson, J. G. Keller, O. Klepper, H. Klingler, J. V. Kratz, R. Kulesa, D. Lambrecht, Y. Leifels, A. Magel, M. Mohar, A. C. Mueller, G. Münzenberg, F. Nickel, G. Nyman, A. Richter, K. Risager, C. Scheidenberger, G. Schrieder, B. M. Sherrill, H. Simon, K. Stelzer, J. Stroth, O. Tengblad, W. Trautmann, E. Wajda, and E. Zude. Study of the Unstable Nucleus  $^{10}\text{Li}$  in Stripping Reactions of the Radioactive Projectiles  $^{11}\text{Be}$  and  $^{11}\text{Li}$ . *Phys. Rev. Lett.*, 75(9):1719, 1995.
- [31] T. Kobayashi, K. Yoshida, A. Ozawa, I. Tanihata, A.A. Korshennikov, E. Nikolski, and T. Nakamura. Quasifree nucleon-knockout reactions from neutron-rich nuclei by a proton target:  $p(^6\text{He}, pn)^5\text{He}$ ,  $p(^{11}\text{Li}, pn)^{10}\text{Li}$ ,  $p(^6\text{He}, 2p)^5\text{H}$ , and  $p(^{11}\text{Li}, 2p)^{10}\text{He}$ . *Nucl. Phys. A*, 616:223, 1997.
- [32] M. Thoennessen, S. Yokoyama, A. Azhari, T. Baumann, J. A. Brown, A. Galonsky, P. G. Hansen, J. H. Kelley, R. A. Kryger, E. Ramakrishnan, and P. Thiof. Population of  $^{10}\text{Li}$  by fragmentation. *Phys. Rev. C*, 59:111, 1999.
- [33] M. Chartier, J.R. Beene, B. Blank, L. Chen, A. Galonsky,

- N Gan, K Govaert, P.G Hansen, J Kruse, V Maddalena, M Thoennesen, and R.L Varner. Identification of the  $^{10}\text{Li}$  ground state. *Phys. Lett. B*, 510:24, 2001.
- [34] H. Simon, M. Meister, T. Aumann, M.J.G. Borge, L.V. Chulkov, U. Datta Pramanik, Th.W. Elze, H. Emling, C. Forssn, H. Geissel, M. Hellstrm, B. Jonson, J.V. Kratz, R. Kulesa, Y. Leifels, K. Markenroth, G. Mnzenberg, F. Nickel, T. Nilsson, G. Nyman, A. Richter, K. Riisager, C. Scheidenberger, G. Schrieder, O. Tengblad, and M.V. Zhukov. Systematic investigation of the drip-line nuclei  $^{11}\text{Li}$  and  $^{14}\text{Be}$  and their unbound subsystems  $^{10}\text{Li}$  and  $^{13}\text{Be}$ . *Nucl. Phys. A*, 791:267, 2007.
- [35] Yu. Aksyutina, T. Aumann, K. Boretzky, M.J.G. Borge, C. Caesar, A. Chatillon, L.V. Chulkov, D. Cortina-Gil, U. Datta Pramanik, H. Emling, H.O.U. Fynbo, H. Geissel, G. Ickert, H.T. Johansson, B. Jonson, R. Kulesa, C. Langer, T. LeBleis, K. Mahata, G. Münzenberg, T. Nilsson, G. Nyman, R. Palit, S. Paschalis, W. Prokopowicz, R. Reifarh, D. Rossi, A. Richter, K. Riisager, G. Schrieder, H. Simon, K. Sümmerer, O. Tengblad, H. Weick, and M.V. Zhukov. Momentum profile analysis in one-neutron knockout from Borromean nuclei. *Phys. Lett. B*, 718:1309, 2013.
- [36] A.I. Amelin, M.G. Gornov, Yu. B. Gurov, A.L. Ilin, P.V. Morokhov, V.A. Pechkurov, V.I. Savelev, F.M. Sergeev, S.A. Smirnov, B.A. Chernyshev, R.R. Shafigullin, and A.V. Shishkov. Production of  $^{10}\text{Li}$  in absorption of stopped  $\pi$  mesons by  $^{11}\text{B}$  nuclei. *Sov.J. Nucl. Phys.*, 52:782, 1990.
- [37] M.G. Gornov, Y.B. Gurov, S.V. Lapushkin, P.V. Morokhov, V.A. Pechkurov, K. Seth, T. Pedlar, J. Wise, and D. Zhao. Spectroscopy of  $^7\text{Li}$ ,  $^8\text{He}$ ,  $^{10}\text{Li}$ , and  $^{13}\text{Be}$  nuclei in stopped  $\pi^-$  meson absorption reactions. *Bull. Rus. Acad. Sci.*, 62:1781, 1998.
- [38] B.A. Chernysev, Yu. B. Gurov, L. Yu. Korotkova, S.V. Lapushkin, R.V. Pritula, and V.G. Sandukovsky. Study of the level structure of the lithium isotope  $^{10}\text{Li}$  in stopped pion absorption. *Int. J. Mod. Phys. E*, 24:1550004, 2015.
- [39] R. Kanungo, A. Sanetullaev, J. Tanaka, S. Ishimoto, G. Hagen, T. Myo, T. Suzuki, C. Andreoiu, P. Bender, A. A. Chen, B. Davids, J. Fallis, J. P. Fortin, N. Galinski, A. T. Gallant, P. E. Garrett, G. Hackman, B. Hadinia, G. Jansen, M. Keefe, R. Krücken, J. Lighthall, E. McNeice, D. Miller, T. Otsuka, J. Purcell, J. S. Randhawa, T. Roger, A. Rojas, H. Savajols, A. Shotter, I. Tanihata, I. J. Thompson, C. Unsworth, P. Voss, and Z. Wang. Evidence of soft dipole resonance in  $^{11}\text{Li}$  with isoscalar character. *Phys. Rev. Lett.*, 114:192502, 2015.
- [40] I. Tanihata, H. Savajols, and R. Kanungo. Recent experimental progress in nuclear halo structure studies. *Progress in Particle and Nuclear Physics*, 68:215, 2013.
- [41] H.T. Fortune. Structure of exotic light nuclei:  $Z=2,3,4$ . *Eur. Phys. J. A*, 54:51, 2018.
- [42] Tanihata, I., M. Alcorta, D. Bandyopadhyay, R. Bieri, L. Buchmann, B. Davids, N. Galinski, D. Howell, W. Mills, S. Mythili, R. Openshaw, E. Padilla-Rodal, G. Ruprecht, G. Sheffer, A. C. Shotter, M. Trinczek, P. Walden, H. Savajols, T. Roger, M. Caamano, W. Mitig, P. Roussel-Chomaz, R. Kanungo, A. Gallant, M. Notani, G. Savard, and I. J. Thompson. Measurement of the two-halo neutron transfer reaction  $^1\text{H}(^{11}\text{Li}, ^9\text{Li})^3\text{H}$  at 3A MeV. *Phys. Rev. Lett.*, 100:192502, 2008.
- [43] I. Thompson and M.V. Zhukov. Effect of a virtual state on the momentum distributions of  $^{11}\text{Li}$ . *Phys. Rev. C*, 49:1904, 1994.
- [44] G.F. Bertsch, K. Hencken, and H. Esbensen. Analysis of nuclear-induced breakup reactions of halo nuclei with the eikonal model. *Phys. Rev. C*, 57:1366, 1998.
- [45] G. Blanchon, A. Bonaccorso, D.M. Brink, and N.V. Mau.  $^{10}\text{Li}$  spectrum from  $^{11}\text{Li}$  fragmentation. *Nucl. Phys. A*, 791:303, 2007.
- [46] M. Gomez-Ramos, J. Casal, and A.M. Moro. Linking structure and dynamics in (p, pn) reactions with Borromean nuclei: The  $^{11}\text{Li}(p, pn)^{10}\text{Li}$  case. *Phys. Lett. B*, 772:115, 2017.
- [47] F.C. Barker and G.T. Hickey. Ground-state configurations of  $^{10}\text{Li}$  and  $^{11}\text{Li}$ . *J. Phys. G*, 3:L23, 1977.
- [48] N.A.F.M. Poppelier, A.A. Wolters, and P.W.M. Glaudemans. Properties of exotic light nuclei. *Z. Phys. A*, 346:11, 1993.
- [49] H. Kitagawa and H. Sagawa. Isospin dependence of kinetic energies in neutron-rich nuclei. *Nucl. Phys. A*, 551:16, 1993.
- [50] P. Descouvemont. Simultaneous study of the  $^{11}\text{Li}$  and  $^{10}\text{Li}$  nuclei in a microscopic cluster model. *Nucl. Phys. A*, 626:647, 1997.
- [51] J. Wurzer and H.M. Hofmann. Microscopic multi-channel calculations for the  $^{10}\text{Li}$  system. *Zeit. Phys. A*, 354:135, 1996.
- [52] K. Kato and K. Ikeda. Analysis of  $^9\text{Li}+n$  Resonances in  $^{10}\text{Li}$  by Complex Scaling Method. *Prog. Theor. Phys.*, 89:623, 1993.
- [53] E. Garrido, D.V. Fedorov, and A.S. Jensen. Structure of exotic light nuclei:  $Z=2,3,4$ . *Nucl. Phys. A*, 770:117, 2002.
- [54] E. Garrido, D.V. Fedorov, and A.S. Jensen. Spin-dependent effective interactions for halo nuclei. *Phys. Rev. C*, 68:014002, 2003.
- [55] G. Potel, F. Barranco, E. Vigezzi, and R. A. Broglia. Evidence for phonon mediated pairing interaction in the halo of the nucleus  $^{11}\text{Li}$ . *Phys. Rev. Lett.*, 105:172502, 2010.
- [56] J. Casal, M. Gomez-Ramos, and A.M. Moro. Description of the  $^{11}\text{Li}(p, d)^{10}\text{Li}$  transfer reaction using structure overlaps from a full three-body model. *Phys. Lett. B*, 767:307, 2017.
- [57] N Vinh Mau. Particle-vibration coupling in one neutron halo nuclei. *Nuclear Physics A*, 592:33, 1995.
- [58] F.M. Nunes, I.J. Thompson, and R.C. Johnson. Core excitation in one neutron halo systems. *Nuclear Physics A*, 596:171, 1996.
- [59] H. Sagawa, B. A. Brown, and H. Esbensen. Parity inversion in the  $N=7$  isotones and the pairing blocking effect. *Phys. Lett. B*, 309:1, 1993.
- [60] T. Myo et al. Systematic study of  $^9,^{10},^{11}\text{Li}$  with the tensor and pairing correlations. *Prog. Theor. Phys.*, 110:561, 2008.
- [61] Barranco, F., P. F. Bortignon, R. A. Broglia, G. Colò, and E. Vigezzi. The halo of the exotic nucleus  $^{11}\text{Li}$ : a single Cooper pair. *Eur. Phys. J. A*, 11:385, 2001.
- [62] M. Cavallaro, M. De Napoli, F. Cappuzzello, S. E. A. Orrigo, C. Agodi, M. Bondí, D. Carbone, A. Cunsolo, B. Davids, T. Davinson, A. Foti, N. Galinski, R. Kanungo, H. Lenske, C. Ruiz, and A. Sanetullaev. Investigation of the  $^{10}\text{Li}$  shell inversion by neutron continuum transfer reaction. *Phys. Rev. Lett.*, 118:012701, 2017.
- [63] F. Barranco, G. Potel, R.A. Broglia, and E. Vigezzi.



- The  ${}^9\text{Li}(d,p)$  reaction, specific probe of  ${}^{10}\text{Li}$ , paradigm of parity-inverted nuclei around  $N = 6$  closed shell. *arXiv:1812.01761 [nucl-th]*, 2018.
- [64] A.M. Moro, J. Casal, and M. Gòmez-Ramos. Investigating the  ${}^{10}\text{Li}$  continuum through  ${}^9\text{Li}(d, p){}^{10}\text{Li}$  reactions. *arXiv:1904.0422v1 [nucl-th]*, 2019.
- [65] A.M. Moro, J. Casal, and M. Gòmez-Ramos. Investigating the  ${}^{10}\text{Li}$  continuum through  ${}^9\text{Li}(d, p){}^{10}\text{Li}$  reactions. *Phys. Lett. B*, 793:13, 2019.
- [66] T. T. S. Kuo, F. Krmpotic, and Y. Tzeng. Suppression of core polarization in halo nuclei. *Phys. Rev. Lett.*, 78:2708, 1997.
- [67] Bohr, A. and B. R. Mottelson. *Nuclear Structure, Vol.II*. Benjamin, New York, 1975.
- [68] R. A. Broglia, G. Potel, F. Barranco, and E. Vigezzi. Difference between stable and exotic nuclei: medium polarization effects. *J. Phys. G*, 37:064022, 2010.
- [69] L.D Landau and L.M. Lifshitz. *Quantum Mechanics, 3rd ed.* Pergamon Press, London, 1965.
- [70] H. Friedrich. *Scattering theory, 2nd ed.* Springer, Berlin, 2016.
- [71] A. De-Shalit. The energy levels of odd-odd nuclei. *Phys. Rev.*, 91:1479, 1953.
- [72] I. Talmi and I. Unna. Order of levels in the shell model and the spin of  ${}^{11}\text{Be}$ . *Phys. Rev. Lett.*, 4:469, 1960.
- [73] K. Heyde. *The nuclear shell model*. Springer, Berlin, 1990.
- [74] E. Beth and G.E. Uhlenbeck. The quantum theory of the non-ideal gas. II. Behaviour at low temperatures. *Physica*, 4:915, 1937.
- [75] S. Shlomo, V.M. Kolomietz, and H. Deibaksh. Single particle level density in a finite depth potential well. *Phys. Rev. C*, 55:1972, 1997.
- [76] K. Mizuyama, Colò G., and E. Vigezzi. Continuum particle-vibration coupling method in coordinate-space representation for finite nuclei. *Phys. Rev. C*, 86:034318, 2012.
- [77] K Huang. *Statistical mechanics, 2nd ed.* Wiley, New York, 1987.
- [78] K. T. Schmitt, K. L. Jones, S. Ahn, D. W. Bardayan, A. Bey, J. C. Blackmon, S. M. Brown, K. Y. Chae, K. A. Chipps, J. A. Cizewski, K. I. Hahn, J. J. Kolata, R. L. Kozub, J. F. Liang, C. Matei, M. Matos, D. Matyas, B. Moazen, C. D. Nesaraja, F. M. Nunes, P. D. O’Malley, S. D. Pain, W. A. Peters, S. T. Pittman, A. Roberts, D. Shapira, J. F. Shriner, M. S. Smith, I. Spassova, D. W. Stracener, N. J. Upadhyay, A. N. Villano, and G. L. Wilson. Reactions of a  ${}^{10}\text{Be}$  beam on proton and deuteron targets. *Phys. Rev. C*, 88:064612, 2013.
- [79] G. Potel, F. M. Nunes, and I. J. Thompson. Establishing a theory for neutron-induced surrogate reactions. *Phys. Rev. C*, 92:034611, 2015.
- [80] G. Potel, G. Perdikakis, B. V. Carlson, M. C. Atkinson, W. H. Dickhoff, J. E. Escher, M. S. Hussein, J. Lei, W. Li, A. O. Macchiavelli, A. M. Moro, F. M. Nunes, S. D. Pain, and J. Rotureau. Toward a complete theory for predicting inclusive deuteron breakup away from stability. *Eur. Phys. J. A*, 53:178, 2017.
- [81] Orrigo, S. E. A. and H. Lenseke. Pairing resonances and the continuum spectroscopy of  ${}^{10}\text{Li}$ . *Phys. Lett. B*, 677:214, 2009.
- [82] F. Barranco, G. Potel, R.A. Broglia, and E. Vigezzi. Structure and reactions of  $N=7$  isotones: parity inversion and transfer processes. *Proc. of the NSD2019 Conference, to be published*, 2019.
- [83] Bortignon, P. F., R. A. Broglia, D. R. Bès, and R. Liotta. Nuclear field theory. *Phys. Rep.*, 30:305, 1977.
- [84] S. Sachdev. *Quantum Phase Transitions*. Cambridge University Press, Cambridge, 2001.
- [85] P. W. Anderson. *Basic Notions of Condensed Matter*. Addison-Wesley, Reading, Mass, 1984.
- [86] M. Buchanan. Wheat from the chaf. *Nature Physics*, 11:296, 2015.
- [87] Mark K. Transtrum, Benjamin B. Machta, Kevin S. Brown, Bryan C. Daniels, Christopher R. Myers, and James P. Sethna. Perspective: Sloppiness and emergent theories in physics, biology, and beyond. *The Journal of Chemical Physics*, 143:010901, 2015.
- [88] Tamara Nikšić and Dario Vretenar. “Sloppy” nuclear energy density functionals: Effective model reduction. *Phys. Rev. C*, 94:024333, 2016.